

Math 341: Midterm 2

Daniel Ko

Spring 2020

§1

Let

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ and } \mathbf{b} = \begin{pmatrix} e \\ f \end{pmatrix} \quad (1)$$

- a. Suppose that $a \neq 0$, compute the solution of $\mathbf{Ax} = \mathbf{b}$ using row reduction and provide the conditions on a, b, c, d such that your computations are valid. Express the result as a simplified expression. (**Hint:** recall that you can not divide by zero)

Proof. We perform reduced row echelon form (rref) on the augmented matrix

$$\begin{aligned} (A|b) &= \left[\begin{array}{cc|c} a & b & e \\ c & d & f \end{array} \right] \\ R_2 &\leftarrow R_2 - \frac{c}{a}R_1 \left[\begin{array}{cc|c} a & b & e \\ 0 & d - \frac{cb}{a} & f - \frac{ce}{a} \end{array} \right] \\ &\left[\begin{array}{cc|c} a & b & e \\ 0 & \frac{ad-cb}{a} & \frac{af-ce}{a} \end{array} \right] \\ R_2 &\leftarrow \frac{a}{ad-cb}R_2 \quad \text{Assuming that } ad-cb \neq 0 \quad \left[\begin{array}{cc|c} a & b & e \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow R_1 - bR_2 \left[\begin{array}{cc|c} a & 0 & e - b\frac{af-ce}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow \frac{R_1}{a} \left[\begin{array}{cc|c} 1 & 0 & \frac{1}{a}(e - b\frac{af-ce}{ad-cb}) \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ &\left[\begin{array}{cc|c} 1 & 0 & \frac{de-bf}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \end{aligned}$$

$$\mathbf{x} = \begin{bmatrix} \frac{de-bf}{ad-cb} \\ \frac{af-ce}{ad-cb} \end{bmatrix} \quad \text{where } ad-cb \neq 0$$

□

- b. If $a = 0$, and $c \neq 0$, is your above computation still valid? How would you modify it? (explain briefly) (**Hint:** recall that you can swap the equations and the result is the same)

Proof. If $a = 0$, and $c \neq 0$, then the above computation will not be valid as we divided by a multiple times when we computed the rref. I would swap the first and second rows so that it would look like

$$\left[\begin{array}{cc|c} c & d & f \\ 0 & b & e \end{array} \right]$$

and compute the rref, using c as □

- c. If $a = 0$, $c = 0$, but $b \neq 0$, $d \neq 0$, what are the conditions on e and f such that the system $\mathbf{Ax} = \mathbf{b}$ has a solution? Is the solution unique? (**Hint:** recall that $\mathbf{Ax} = \mathbf{b}$ has a solution if and only if \mathbf{b} can be written as a linear combination of the columns of \mathbf{A})

- d. Solve the system

$$\begin{bmatrix} \sqrt{2} & 3\sqrt{2} \\ 2\sqrt{2} & \sqrt{2} \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 5\sqrt{2} \\ 5\sqrt{2} \end{pmatrix}. \quad (2)$$

(**Hint:** You may want to use the formula you just deduced)

Proof.

$$\begin{aligned} x_1 &= \frac{de - bf}{ad - cb} \\ &= \frac{\sqrt{2}(5\sqrt{2}) - 3\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\ &= \frac{10 - 30}{2 - 12} \\ &= \frac{-20}{-10} \\ &= 2 \end{aligned}$$

$$\begin{aligned} x_2 &= \frac{\sqrt{2}(5\sqrt{2}) - 2\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\ &= \frac{10 - 20}{-10} \\ &= 1 \end{aligned}$$

□

§2

§3

§4