

Math 341: Midterm 2

Daniel Ko

Spring 2020

§1

Let

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ and } \mathbf{b} = \begin{pmatrix} e \\ f \end{pmatrix} \quad (1)$$

- a. Suppose that $a \neq 0$, compute the solution of $\mathbf{Ax} = \mathbf{b}$ using row reduction and provide the conditions on a, b, c, d such that your computations are valid. Express the result as a simplified expression. (**Hint:** recall that you can not divide by zero)

Proof. We perform reduced row echelon form (rref) on the augmented matrix

$$\begin{aligned} (A|b) &= \left[\begin{array}{cc|c} a & b & e \\ c & d & f \end{array} \right] \\ R_2 &\leftarrow R_2 - \frac{c}{a}R_1 \left[\begin{array}{cc|c} a & b & e \\ 0 & d - \frac{cb}{a} & f - \frac{ce}{a} \end{array} \right] \\ &\left[\begin{array}{cc|c} a & b & e \\ 0 & \frac{ad-cb}{a} & \frac{af-ce}{a} \end{array} \right] \\ R_2 &\leftarrow \frac{a}{ad-cb}R_2 \quad \text{Assuming that } ad-cb \neq 0 \left[\begin{array}{cc|c} a & b & e \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow R_1 - bR_2 \left[\begin{array}{cc|c} a & 0 & e - b\frac{af-ce}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ R_1 &\leftarrow \frac{R_1}{a} \left[\begin{array}{cc|c} 1 & 0 & \frac{1}{a}(e - b\frac{af-ce}{ad-cb}) \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ &\left[\begin{array}{cc|c} 1 & 0 & \frac{de-bf}{ad-cb} \\ 0 & 1 & \frac{af-ce}{ad-cb} \end{array} \right] \\ x &= \begin{bmatrix} \frac{de-bf}{ad-cb} \\ \frac{af-ce}{ad-cb} \end{bmatrix} \quad \text{where } ad-cb \neq 0 \end{aligned}$$

□

- b. If $a = 0$, and $c \neq 0$, is your above computation still valid? How would you modify it? (explain briefly) (**Hint:** recall that you can swap the equations and the result is the same)

Proof. If $a = 0$, and $c \neq 0$, then the above computation will not be valid as we divided by a multiple times when we computed the rref. I would swap the first and second rows so that it would look like

$$\left[\begin{array}{cc|c} c & d & f \\ 0 & b & e \end{array} \right]$$

and compute the rref, assuming that $b \neq 0$. We obtain the rref,

$$\left[\begin{array}{cc|c} 1 & 0 & \frac{bf-de}{bc} \\ 0 & 1 & \frac{e}{b} \end{array} \right]$$

□

- c. If $a = 0$, $c = 0$, but $b \neq 0$, $d \neq 0$, what are the conditions on e and f such that the system $\mathbf{Ax} = \mathbf{b}$ has a solution? Is the solution unique? (**Hint:** recall that $\mathbf{Ax} = \mathbf{b}$ has a solution if and only if \mathbf{b} can be written as a linear combination of the columns of \mathbf{A})

Proof. If $a = 0$, $c = 0$, $b \neq 0$, $d \neq 0$, we get the augmented matrix

$$\left[\begin{array}{cc|c} 0 & b & e \\ 0 & d & f \end{array} \right]$$

Performing row reduction,

$$\left[\begin{array}{cc|c} 0 & 1 & \frac{e}{d} \\ 0 & 1 & \frac{f}{d} \end{array} \right]$$

So the condition of the solution is,

$$x_2 = \frac{e}{b} = \frac{f}{d}$$

Thus, there exists a infinite amount of solution.

□

- d. Solve the system

$$\begin{bmatrix} \sqrt{2} & 3\sqrt{2} \\ 2\sqrt{2} & \sqrt{2} \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 5\sqrt{2} \\ 5\sqrt{2} \end{pmatrix}. \quad (2)$$

(**Hint:** You may want to use the formula you just deduced)

Proof.

$$\begin{aligned} x_1 &= \frac{de - bf}{ad - cb} \\ &= \frac{\sqrt{2}(5\sqrt{2}) - 3\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\ &= \frac{10 - 30}{2 - 12} \\ &= \frac{-20}{-10} \\ &= 2 \end{aligned}$$

$$\begin{aligned} x_2 &= \frac{\sqrt{2}(5\sqrt{2}) - 2\sqrt{2}(5\sqrt{2})}{\sqrt{2}\sqrt{2} - 3\sqrt{2}(2\sqrt{2})} \\ &= \frac{10 - 20}{-10} \\ &= 1 \end{aligned}$$

□

§2

Let

$$\mathbf{A} = \begin{bmatrix} 0 & -\alpha & 2 & 0 \\ 1 & 1 & 0 & 1 \\ 2 & 2 & 2 & 3 \\ -2 & -2 & 4 & 2\alpha \\ 0 & \alpha & -1 & 2\alpha + 1/2 \end{bmatrix}, \quad \text{and } \mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ 4 \\ 2 + \alpha \\ 2\beta + \alpha - 2 \end{bmatrix} \quad (3)$$

What are the conditions on α and β such that the system $\mathbf{Ax} = \mathbf{b}$:

a. Has no solution?

Proof. We begin by putting the augmented matrix $(\mathbf{A}|\mathbf{b})$ in its reduced form.

$$\begin{aligned}
 (\mathbf{A}|\mathbf{b}) &= \left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 2 & 2 & 2 & 3 & 4 \\ -2 & -2 & 4 & 2\alpha & 2 + \alpha \\ 0 & \alpha & -1 & 2\alpha + 1/2 & 2\beta + \alpha - 2 \end{array} \right] \\
 R_5 \leftarrow R_5 + R_1 &\left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 2 & 2 & 2 & 3 & 4 \\ -2 & -2 & 4 & 2\alpha & 2 + \alpha \\ 0 & 0 & 1 & 2\alpha + 1/2 & 2\beta + \alpha \end{array} \right] \\
 R_3 \leftarrow R_3 - 2R_2 &\left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ -2 & -2 & 4 & 2\alpha & 2 + \alpha \\ 0 & 0 & 1 & 2\alpha + 1/2 & 2\beta + \alpha \end{array} \right] \\
 R_4 \leftarrow R_4 + 2R_2 &\left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 4 & 2\alpha + 2 & 4 + \alpha \\ 0 & 0 & 1 & 2\alpha + 1/2 & 2\beta + \alpha \end{array} \right] \\
 R_4 \leftarrow R_4 - 2R_3, R_5 \leftarrow R_5 - \frac{1}{2}R_3 &\left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 2\alpha & 2\beta + \alpha + 1 \end{array} \right] \\
 R_5 \leftarrow R_5 - R_4 &\left[\begin{array}{cccc|c} 0 & -\alpha & 2 & 0 & 2 \\ 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 2\beta - 1 \end{array} \right] \\
 R_1 \leftrightarrow R_2 &\left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & -\alpha & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 2\beta - 1 \end{array} \right]
 \end{aligned}$$

Thus this system will have no solution if $2\beta - 1 \neq 0$, which is when $\beta \neq \frac{1}{2}$. explain more... \square

- b. Has an unique solution? Find the solution. (**Hint:** you will need to row reduce the augmented system to echelon form, and then use the theorems seen in class to impose the conditions on α and β).

Proof. Combining Theorem 3.10 and the corollary to Theorem 4.7, we get that a system has a unique solution if and only if $\det(\mathbf{A}) \neq 0$. The determinant of an upper triangular matrix is the product of its diagonal entries by property 4 of determinants in section 4.4. So, this is when

$$\begin{aligned} 1 * -\alpha * 2 * 2\alpha &\neq 0 \\ -4\alpha^2 &\neq 0 \\ \alpha &\neq 0 \end{aligned}$$

and from (a), $\beta = \frac{1}{2}$. Combining these two conditions we get the following system,

$$\left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & -\alpha & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 2\alpha & \alpha \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

So the generic solution will be

$$\mathbf{x} = \begin{bmatrix} \frac{1}{2} + \frac{1}{2\alpha} \\ -\frac{1}{2\alpha} \\ \frac{3}{4} \\ \frac{1}{2} \\ 0 \end{bmatrix}$$

□

- c. Has infinite amount of solutions? Find the solution set in parametric form. (**Hint:** You may have one equations for α and one for β that have to be satisfied simultaneously).

Proof. Having an infinite amount of solutions is by definition another way of saying that a system that is consistent and that the solutions are not unique. A system is consistent if $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}|\mathbf{b})$ by Theorem 3.11. A solution is not unique if $\det(\mathbf{A}) = 0$ by Theorem 3.10 and the corollary to Theorem 4.7.

If $\det(\mathbf{A}) = 0$,

$$\begin{aligned} 1 * -\alpha * 2 * 2\alpha &= 0 \\ -4\alpha^2 &= 0 \\ \alpha &= 0 \end{aligned}$$

Given that $\alpha = 0$, and that $\beta = \frac{1}{2}$ from part (a) we get the following system,

$$\left[\begin{array}{cccc|c} 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 2 & 0 & 2 \\ 0 & 0 & 2 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

It is clear that $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}|\mathbf{b})$ because \mathbf{b} is a linear combination of the second and third column from \mathbf{A} . □

§3

§4