

**OPD ASSESSMENT
PHYSIOTHERAPY DEPARTMENT**

Patient Name
Date of Birth (DD/MM/YYYY):
Age
HN : VN /AN :
Visit/ Admit Date : Gender:
Allergies:.....

Date..... Time..... Doctor..... VN.....
Transportation ☐ Walk ☐ Wheelchair ☐ Stretcher ☐ Other
PR.....bpm RR.....breaths/min BP...../.....mmHg SpO2.....%

Chief complaint

Diagnosis

Physiotherapy Precaution

Underlying Disease ☐ None ☐ Heart Disease ☐ Cancer
☐ Diabetes Mellitus ☐ Hypertension ☐ Other.....

Pain Score Location of Pain

Pain assessment Tool ☐ newborn - 1 year (NIPS) ☐ >1-3 years (FLACC)

☐ 3-8 years (FRS) ☐ >8years (NRS) ☐ CPOT (Critical care Pain Observation Tool)

Characteristic ☐ Prick ☐ Sharp ☐ Dull ☐ Burning ☐ Tight

☐ Throbbing ☐ Radiating ☐ Pin & Needles ☐ Other.....

Duration of PainFrequency: ☐ Constant ☐ Intermittent

Fall risk : ☒ Standard fall (NO ABCS) ☐ Strict fall (A,B,C) ☐ Highly Strict fall(S)

☐ Age: ผู้ป่วยเด็กที่อายุ ≤ 2 ปี , ผู้ป่วยสูงอายุที่อายุ ≥ 85 ปี ☐ Bone (Fracture risk or history)

☐ Balance: มีปัญหาเรื่องการทรงตัว วิ่งเวียนบ้านหมุน ใช้อุปกรณ์ช่วยเดิน ใช้รถเข็นนั่ง รถเข็นนอน ☐ ตั้งครวกร ≥20 สัปดาห์

☐ Coagulation ได้รับยากกลุ่มต้านการแข็งตัวของเลือด (On Anticonvulsants or Coagulopathy or Bleeding disorder)

☐ Surgery ผู้ป่วยได้รับการทำหัตถการ (ใช้ยาสมบ) ทำให้สถิติสัมประญะไม่สมบูรณ์ สับสน ซึม ☐ ได้รับยาขยายม่านตา

Precaution: ☐ Standard Precaution ☐ Airborne ☐ Droplet
☐ drug-Resistance ☐ Contact ☐ Protective

Barthel Index ☐ Yes ☐ No

Physical Examination: Visit.....

☐ Orthopedics (if condition applied to) ☐ NA

Observation

Palpation.....

Muscle Power.....

Range of Motion.....

Sensation.....

Functional Movement.....

☐ Cardiopulmonary (if condition applied to) ☐ NA

Observation.....

Palpation.....

Percussion.....

Auscultation.....

☐ Neurology (if condition applied to) ☐ NA

Observation.....

Muscle Tone.....

Balance.....

Bed mobility and Transferring.....

Muscle Power.....

Sensation.....

Treatment Plan:

.....
.....
.....
.....
.....

Participatory Goal Setting:

Short term goal

.....

Long term goal.....

.....

Physiotherapy Treatment

☐ Ultrasound: ☐ Pulse ☐ Continuousminutes area

☐ Manual Therapy ☐ Paraffin ☒ Tilt Table.....minutes Degree

☐ Electrical Stimulationminutes area

☐ Hydrocollator:minutes area

☐ Cryotherapy:minutes area

☐ Traction: ☐ Cervical ☐ Lumbarkilogramsminutes

☒ Laser: ☐ Pulse ☐ Continuousminutes areaJul/cm²

☒ Shockwave Diathermy :area

☒ Peripheral Diathermy: Modeminutes area

☐ Shortwave Diathermy/Microwave Diathermy
☐ Pulse ☐ Continuousminutes area

☐ Range of Motion Exercise ☐ Hand Exercise ☐ Strengthening Exercise

☐ Balance Training ☐ Balance Test ☒ Bicycle Training : ...Intensity...minutes

☒ Treadmill Training: Speed.....km/hminutesDistance

☒ Continuous Passive Motion:.....minutes Flexion/ExtensionDegree

☐ Percussion ☒ Vibration ☐ Chest Mobilization ☐ Breathing Exercise

☐ Suction

☐ Ambulation: with

☐ Partial Weight Bearing ☐ Non Weight Bearing ☐ Full Weight Bearing

☒ Other.....

Treatment Detail:

Re-assessment: Time..... Pain Score:.... Frequency: ☐ Constant ☐ Intermittent

PR....bpm RR....breaths/min BP...../.....mmHg SpO2.....%

Duration of Pain: Location of Pain :.....

Pain assessment Tool ☐ newborn - 1 year (NIPS) ☐ >1-3 years (FLACC)

☐ 3-8 years (FRS) ☐ >8years (NRS) ☐ CPOT (Critical care Pain Observation)

Characteristic : ☐ Prick ☐ Sharp ☐ Dull ☐ Burning ☐ Tight

☐ Throbbing ☐ Radiating ☐ Pin & Needles ☐ Other.....

☐ Orthopedics (if condition applied to) ☐ NA

Observation

Palpation.....

Muscle Power.....

Range of Motion.....

Sensation.....

Functional Movement.....

☐ Cardiopulmonary (if condition applied to) ☐ NA

Observation.....

Palpation.....

Percussion.....

Auscultation.....

☐ Neurology (if condition applied to) ☐ NA

Observation.....

Muscle Tone.....

Balance.....

Bed mobility and Transferring.....

Muscle Power.....

Sensation.....

Fall risk: ☐ Standard fall (NO ABCS) ☐ Strict fall (A,B,C) ☐ Highly Strict fall(S)

คำแนะนำที่ได้ รับ ☐ การบริหารร่างกาย (P/I-BSI-024.1) ☐ การลดปวด (P/I-BSI-025.1) ☐ Fall

☐ การฝึกเดิน (P/I-BSI-023.1) ☐ การเกาะปอด (P/I-BSI-022.1) ☐ การออกกำลังกายในผู้ ป่วยอัมพาต (P/I-

BSI-017)
Discharge status: ☐ รู้สึกดี ☐ ซึม ☐ สับสน ☐ ไม่รู้สึกตัว ☐ อื่นๆ.....

PhysioTherapistDate.....Time.....