

# E theory

- ▶ Following Jack's work
- ▶ An analogue to nematic Q tensor for smectics
- ▶ Replace the real order parameters  $S$  with complex  $\psi$
- ▶ Mainly thinking about uniaxial case

$$\underline{\underline{Q}} = S_1(\underline{\underline{N}}\underline{\underline{N}} - \frac{\delta}{d}) + \boxed{S_2(\underline{\underline{M}}\underline{\underline{M}} - \frac{\delta}{d})}$$

$S_2 = 0$  makes it uniaxial

$$\underline{\underline{E}} \sim \psi_1(\underline{\underline{N}}\underline{\underline{N}} - \frac{\delta}{d})$$

# Thinking about constraints

- ▶  $Q$  is real, so symmetry makes it diagonalizable – this leads to the biaxial form
- ▶ If a complex matrix is normal ( $\underline{\underline{E}}\underline{\underline{E}}^\dagger = \underline{\underline{E}}^\dagger \underline{\underline{E}}$ ) it can be diagonalized by a unitary matrix

$$\underline{\underline{E}} = \underline{\underline{U}}^\dagger \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & -\lambda_1 - \lambda_2 \end{pmatrix} \underline{\underline{U}} = \dots = \psi_1(\underline{\underline{N}}\underline{\underline{N}} - \frac{\delta}{d}) + \psi_2(\underline{\underline{M}}\underline{\underline{M}} - \frac{\delta}{d})$$

also using symmetry

- ▶ With  $\underline{\underline{N}}$ ,  $\underline{\underline{M}}$  being real, orthogonal, unit vectors, and

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} = \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} - \lambda_3$$

- ▶ Overall  $2*2 + 2 + 1 = 7$  dof, 4 if uniaxial

# The free energy

- ▶ Using the simplest terms
- ▶ We need to match  $\underline{\underline{E}}$  and  $\underline{\underline{E}}^*$  to make it real

$$f_{\text{bulk}}(E_{ij}E_{ij}^* = \text{Tr}(\underline{\underline{E}}\underline{\underline{E}}^*)) = \frac{A}{2}E_{ij}E_{ij}^* + \frac{C}{4}(E_{ij}E_{ij}^*)^2$$

- ▶ Elastic terms need gradients – below are "single elastic constant" terms

$$|\underline{\nabla}\underline{\underline{E}}|^2 = E_{ij,k}E_{ij,k}^* \quad \text{the main term}$$

$$|\underline{\nabla} \cdot \underline{\underline{E}}|^2 = E_{ij,j}E_{ik,k}^* \quad \text{only a surface contribution (as per Jack's work)}$$

$$|\underline{\nabla}^2 \underline{\underline{E}}|^2 = E_{ij,kk}E_{ij,ll}^* \quad \text{the only double gradient term considered}$$

# Projection operators

- ▶ Gradients in different directions have different energy costs
- ▶ Working with uniaxial  $\underline{\underline{E}}$  (or nearly so) – special direction is  $\underline{N}$
- ▶ Projection operator  $\underline{\underline{\Pi}} = \underline{N}\underline{N}$
- ▶ Perpendicular projections are then  $\underline{\underline{T}} = \underline{\underline{\delta}} - \underline{\underline{\Pi}}$
- ▶ Adapt the free energies by  $\underline{\nabla} \rightarrow \underline{\underline{\Pi}} \cdot \underline{\nabla} + \underline{\underline{T}} \cdot \underline{\nabla}$

$$E_{ij,k} E_{ij,k}^* \rightarrow f_{\text{comp}} = b_1^{\parallel} \Pi_{kl} E_{ij,k} E_{ij,l}^* + b_1^{\perp} T_{kl} E_{ij,k} E_{ij,l}^*$$

$$E_{ij,kk} E_{ij,ll}^* \rightarrow f_{\text{curv}} = b_2^{\parallel} \Pi_{kl} E_{ij,lk} \Pi_{mn} E_{ij,nm}^* + b_2^{\perp} T_{kl} E_{ij,lk} T_{mn} E_{ij,nm}^* \\ + b_2^{\parallel\perp} (\Pi_{kl} E_{ij,lk} T_{mn} E_{ij,nm}^* + T_{kl} E_{ij,lk} \Pi_{mn} E_{ij,nm}^*)$$

# Projection operators

- ▶ Need a form for  $\underline{\underline{\Pi}}$  in terms of  $\underline{\underline{E}}$
- ▶ Have 2 forms which work for uniaxial  $\underline{\underline{E}}$

$$\underline{\underline{\Pi}} = \sqrt{\frac{d-1}{d\underline{\underline{E}}:\underline{\underline{E}}}} \underline{\underline{E}} + \frac{\delta}{d}$$
$$\underline{\underline{\Pi}} = \frac{d-1}{d-2} \left( \frac{\underline{\underline{E}} \cdot \underline{\underline{E}}^*}{\underline{\underline{E}}:\underline{\underline{E}}^*} - \frac{\delta}{d(d-1)} \right)$$

- ▶ Lead to seemingly different functional derivatives – why?
- ▶ First form only has  $\underline{\underline{E}}$ , how about  $\underline{\underline{E}} \rightarrow \underline{\underline{E}}^*$ ?
- ▶ How well do they work for biaxial  $\underline{\underline{E}}$ ?

# Dynamics and functional derivatives

- ▶ Starting from  $\mu \frac{\partial E_{ij}}{\partial t} = -\frac{\delta F}{\delta E_{ij}^*}$ , but need to preserve constraints
- ▶ If  $\frac{\delta F}{\delta E_{ij}^*}$  is symmetric and traceless, then so will  $\underline{\underline{E}}$
- ▶ Either treat  $\underline{\underline{E}}$  as symmetric, or symmetrize after
- ▶ Normality is more complicated, Lagrange multiplier from Djourde's work
- ▶ It might be nice to constrain it to be uniaxial too

# Functional derivatives

- Results using the square root version of  $\underline{\Pi}$

$$\frac{\delta F_{\text{bulk}}}{\delta E_{ij}^*} = \frac{1}{2}(A + CE_{ab}E_{ab}^*)E_{ij}$$

$$\frac{\delta F_{\text{comp}}}{\delta E_{ij}^*} = -(b_1^{\parallel} - b_1^{\perp})(\Pi_{kl,l}E_{ij,k} + \Pi_{kl}E_{ij,kl}) - b_1^{\perp}E_{ij,kk}$$

$$\begin{aligned}\frac{\delta F_{\text{curv}}}{\delta E_{ij}^*} = & (b_2^{\parallel} + b_2^{\perp} - 2b_2^{\parallel\perp})\left((\Pi_{kl}\Pi_{po,po} + 2\Pi_{kl,o}\Pi_{po,p} + \Pi_{kl,po}\Pi_{po})E_{ij,lk}\right. \\ & \left.+ 2(\Pi_{kl,o}\Pi_{po} + \Pi_{kl}\Pi_{po,o})E_{ij,lkp} + \Pi_{kl}\Pi_{po}E_{ij,lkpo}\right) \\ & + (b_2^{\parallel\perp} - b_2^{\perp})\left(\Pi_{po,po}E_{ij,kk} + 2\Pi_{po,o}E_{ij,kkp} + \Pi_{po}E_{ij,kkpo}\right. \\ & \left.+ \Pi_{kl,oo}E_{ij,lk} + 2\Pi_{kl,o}E_{ij,lko} + \Pi_{kl}E_{ij,lkoo}\right) \\ & + b_2^{\perp}E_{ij,kkoo}\end{aligned}$$

# Functional derivatives

- Results using the square root version of  $\Pi$

$$\Pi_{kl} = \frac{sE_{kl}}{\sqrt{E_{ab}E_{ab}}} + \frac{\delta_{kl}}{d}$$

$$\Pi_{kl,m} = \frac{s}{\sqrt{E_{ab}E_{ab}}} \left( E_{kl,m} - \frac{E_{kl}E_{cd}E_{cd,m}}{E_{ab}E_{ab}} \right)$$

$$\begin{aligned} \Pi_{kl,mn} = \frac{s}{\sqrt{E_{ab}E_{ab}}} & \left( E_{kl,mn} \right. \\ & - \frac{E_{kl,n}E_{cd}E_{cd,m} + E_{kl,m}E_{cd}E_{cd,n} + E_{kl}(E_{cd,n}E_{cd,m} + E_{cd}E_{cd,mn})}{E_{ab}E_{ab}} \\ & \left. + 3 \frac{E_{kl}E_{cd}E_{cd,m}E_{ef}E_{ef,n}}{(E_{ab}E_{ab})^2} \right) \end{aligned}$$