

# Ones

What are all the different (as in they lead to different behaviours) 2 digit cycles we can implement in the model? In the diagrams as shown in fig. 1 we first note that the diagrams lead to the same behaviour if they are the same under reflection along the top-left to bottom-right axis which corresponds to reflecting the sequence (physically the same system if working with a loop geometry). We first focus on such cycles that visit all possible 2 digit sequences. We either get diagrams with or without a crossing, and there is only one diagram without a crossing which is fig. 1a. For diagrams with a crossing that visit all four 2 digit sequences we always get the motif as in figs. 1b and 1c which are the only such diagrams up to the mentioned reflection. Also note that the two cycles with crossings are reverses of each other!




Figure 1: All the different 2 digit simple cycles that visit all four such sequences, these are unique up to right-left reflection.

Following the same logic, we only get four cycles which visit 3 of the sequences, shown in fig. 2. With the last two also being reverses of each other. We don't examine cycles which visit only 2 of the sequences as it's difficult to imagine anything interesting taking place.




Figure 2: All the different 2 digit simple cycles that do not visit all such sequences, these are unique up to right-left reflection.

As reverses of cycles will give exactly the same behaviour just reversed in time, we can focus on only one of each such cycles. In addition, it is reasonable to suspect that replacing all 0s by 1s and vice versa should also be some symmetry of the system and so we only look at one cycle for each of such classes. This gives only the cycles shown in fig. 3 to look at.




Figure 3: The 2 digit simple cycles that are worth investigating.