Doutorado em Matemática - IMPA

LECTURE NOTES ON DYNAMICS I

Alejandro Kocsard
IMPA
March 24, 2026

Contents

1 Introduction **3**

2 Some examples of Dynamical Systems results **7**

3 The Fundamental Theorem of Dynamical Systems **8**

3.1 Different forms of recurrence 9

1 Introduction

In this course we shall deal with *autonomous deterministic dynamical systems.* They can be used to represent or modelize any phenomenon that changes along the time in such a way that the current state of the system completely determines the future evolution and the law that drives this evolution does not change along the time. Mathematically, an **autonomous deterministic dynamical system** (a **dynamical system**, for short) is given by a set M, usually called the **phase space,** which represents the set of possible states of the system, a subsemigroup $\mathbb{T} \subset \mathbb{R}$ that represents the **time set,** and *"a law"* $\Phi\colon \mathbb{T} \times M \to M$ that determines the evolution of the system in such a way that given the current state of our system $x_0 \in M$, at time $t \in \mathbb{T}$ the corresponding state will be $x_t = \Phi(x_0, t)$. The autonomy of the system is reflected in the so called **flow** (or **semi-flow**) condition function Φ satisfies:

$$\Phi(t+s, x) = \Phi\big(t, \Phi(s, x)\big) = \Phi\big(t, \Phi(s, x)\big), \quad \forall t, s \in \mathbb{T},\ \forall x \in M. \tag{1}$$

We will consider two different kinds of dynamical systems: **discrete time** and **continuous time** ones, i.e. where $\mathbb{T} = \mathbb{N}_0$ and $\mathbb{T} = \mathbb{R}_{\geqslant 0}$, respectively. Moreover, we will also consider **invertible systems** where the current state does not just determine the future but the past as well. When dealing with invertible discrete dynamical systems, the time set will be $\mathbb{T} = \mathbb{Z}$, and in the case of invertible continuous time ones, we will have $\mathbb{T} = \mathbb{R}$.

The main goal of Dynamical Systems consists in describing the **orbits of the systems,** i.e. the sets

$$\mathcal{O}_\Phi(x) := \{\Phi(t, x) : t \in \mathbb{T}\}, \quad \forall x \in M.$$

Exercise 1.1 (Discrete time generator)**.** Given a discrete time dynamical system $\Phi\colon \mathbb{T} \times M \to M$, show that Φ satisfies the flow condition (1) if and only if there is $f\colon M \circlearrowleft$ such that

$$\Phi(n, x) = f^n(x), \quad \forall x \in M,\ \forall n \in \mathbb{N}_0,$$

where the iteration of f is inductively defined by $f^0 := id_M$ and $f^{n+1} = f \circ f^n$, for every $n \geqslant 0$. Show that Φ is invertible if and only if f is bijective and, in such a case, one can define $f^{-n} := \left(f^{-1}\right)^n$, for all $n \geqslant 0$.

On the other hand, one can find *infinitesimal generators* for continuous time smooth dynamical systems:

Exercise 1.2 (Continuous time generator)**.** Let M be a differentiable closed manifold (i.e. compact and boundaryless) and $\Phi\colon \mathbb{R} \times M \to M$ a continuous time dynamical system (that satisfies the flow condition 1). If we assume Φ is a C^2 map and we define $X(p) := \partial_t \Phi(t,p)\big|_{t=0} \in T_pM$, for every $p \in M$, then the function $\Phi(\cdot, x_0)\colon \mathbb{R} \to M$ is the unique solution of the *initial value problem*

$$\dot{x}(t) = X\big(x(t)\big), \quad x(0) = x_0,$$

for every $x_0 \in M$. In other words, the vector field X is the infinitesimal generator of the flow Φ.

As previously mentioned, the primary objective of Dynamical Systems is to understand the orbits of various systems, particularly their asymptotic behavior as time goes to infinity. From a historical perspective, there are three pivotal moments in the development of this mathematical theory. The first can be attributed to Isaac Newton, who laid the groundwork for Calculus and employed its principles to solve differential equations. During this era, the approach to studying a specific continuous-time dynamical system involved solving the corresponding differential equation, followed by an analysis of the obtained solutions.

The so-called *"restricted two-body problem"* is a remarkable question that was answered at that time:

Problem 1.3. The *"restricted two-body problem"* refers to a simplified model used to analyze the motion of two celestial bodies under the influence of their mutual gravitational attraction, where one body is significantly more massive than the other. In this framework, the more massive body is assumed to be stationary, while the second body is treated as a point mass that moves in the gravitational field created by the first one. This approach allows for the derivation of equations of motion for the second body without needing to account for its gravitational effects on the primary, simplifying the analysis. This model is particularly useful in celestial mechanics, where it can be applied to scenarios such as a satellite orbiting a planet or a comet passing by a star.

More precisely, the *restricted two-body problem* consider in describing the orbit of massive point p, whose location in the space at time $t \in \mathbb{R}$ is given by $x(t) \in \mathbb{R}^3$, moving in a central field force in $\mathbb{R}^3$ that is generated by a massive point of mass $M > 0$ which is fixed at the origin. According to Newton's laws, the trajectory $x(t) \in \mathbb{R}^3$ is the solution of the following initial value problem:

$$\begin{cases} \ddot{x}(t) = -\dfrac{GM}{\|x\|^3} x(t), \\ x(0) = x_0, \\ \dot{x}(0) = v_0, \end{cases} \tag{2}$$

where $G > 0$ is the universal constant of gravitation, $x_0 \in \mathbb{R}^3$ and $v_0 \in \mathbb{R}^3$ are the initial position and velocity, respectively.

Using classical *quadrature techniques* one can show that the solutions $x(t)$ of (2) are either straight lines through the origin (when x_0 and v_0 are collinear vectors in $\mathbb{R}^3$), or quadratic curves (i.e. ellipses, parabolas or hyperbolas).

The second pivotal moment in the development of Dynamical Systems can be attributed to Henri Poincaré. At the end of XIXth century it was already pretty clear that there were many ordinary differential equations that could not be solved with quadrature methods. A very simple and well-known example is the following one:

Example 1.1. *There is no* elementary (smooth) function $f\colon \mathbb{R} \to \mathbb{R}$ *such that*

$$f'(x) = e^{-x^2}, \quad \forall x \in \mathbb{R}.$$

Another example, which was fundamental in development of so called *qualitative theory of differential equations* is the 3-*body problem:*

Problem 1.4 (n-body problem)**.** The *n-body problem* refers to the challenge of predicting the individual motions of a group of n celestial objects that interact with each other through gravitational forces. The complexity arises from the fact that each body exerts a gravitational force on every other body, leading to a system of coupled differential equations. If $x_i(t) \in \mathbb{R}^3$ represents the position of the i-th object at time $t \in \mathbb{R}$ and $m_i > 0$ denotes its mass, then Newton's second law determines that these quantities are related by the following system of ordinary differential equations

$$\begin{cases} \ddot{x}_i(t) = \displaystyle\sum_{j \neq i} G \frac{m_j}{\|x_i(t) - x_j(t)\|^3} \big(x_i(t) - x_j(t)\big), \\ x_i(0) = x_{i,0}, \\ \dot{x}_i(0) = v_{i,0}, \end{cases} \tag{3}$$

for $1 \leqslant i \leqslant n$, and where $x_{i,0}, v_{i,0} \in \mathbb{R}^3$ denotes the position and the velocity of the i-th object at time $t = 0$, respectively.
It is pretty remarkable that, in general, the system of ODEs given by (3) cannot be solved by "quadrature methods" when $n \geqslant 3$. In fact, their solutions are not elementary functions in general.

The n-body problem, as outlined in Problem 1.4, served as a primary motivation for Henri Poincaré in developing the Geometric Theory of Differential Equations. Poincaré posited that if explicit solutions to a given differential equation cannot be readily obtained, one should at least strive to describe some interesting geometric properties of those solutions.

Indeed, many questions regarding the solutions of differential equations are more qualitative than quantitative, or more geometric than analytic. For instance, a fundamental qualitative inquiry concerning the system of ordinary differential equations in (3) is whether every solution is periodic. In some cases, even when solutions can be expressed using elementary functions, addressing certain qualitative questions based solely on the explicit formulas can prove to be quite challenging.

The third pivotal moment in the evolution of Dynamical Systems can be attributed to the schools of Steve Smale in the US and Aleksandr Andronov in the USSR. At this juncture, it became evident that the geometric properties of orbits, trajectories, or solutions to ODEs could change dramatically with variations in system parameters. For example, in Problem 1.4, there are $3n$ parameters—comprising the n masses, n initial positions, and n initial velocities. In certain instances, even minor changes to these parameters can lead to completely different geometric properties for the respective solutions. Thus, it becomes crucial to identify geometric properties of orbits that are robust or persistent under small perturbations of system parameters.

These three historical stages in the development of dynamical systems are cleverly (and provocatively) summarized by Étienne Ghys in his laudation for Artur Avila's Fields Medal: In Newton's time, you were given an ODE and tasked with finding its solutions. In Poincaré's era, you were given an ODE and asked to elucidate interesting geometric properties of its solutions. In Smale's time, you were given no ODE at all and challenged to say something interesting!

2 Some examples of Dynamical Systems results

A very first example of a result in Dynamical Systems might be the following classical

Theorem 2.1 (Banach fixed point theorem)**.** Let (M, d) be a complete metric space and $f\colon M \circlearrowleft$ be **uniform contraction,** i.e. f is a Lipschitz function and

$$\lambda = \mathrm{Lip}(f) := \sup_{x \neq y} \frac{d\big(f(x), f(y)\big)}{d(x, y)} < 1.$$

Then there exists a unique point $p \in M$ such that $f(p) = p$ and it holds

$$\lim_{n \to +\infty} f^n(x) = p, \quad \forall x \in M.$$

Proof. Given any $x \in M$, one can show by induction on n that

$$d\big(f^n(x), f^{n+1}(x)\big) \leqslant \lambda^n d(x, f(x)), \quad \forall n \in \mathbb{N}_0.$$

So, for every $m, n \in \mathbb{N}$ it holds

$$\begin{aligned} d\big(f^m(x), f^{m+n}(x)\big) &\leqslant \sum_{i=0}^{n-1} d\big(f^{m+i}(x), f^{m+i+1}(x)\big) \\ &\leqslant \sum_{i=0}^{n-1} \lambda^{m+i} d\big(x, f(x)\big) \leqslant \frac{\lambda^m}{1-\lambda} d\big(x, f(x)\big). \end{aligned}$$

Hence, the sequence $\big(f^n(x)\big)_{n \geqslant 0}$ is Cauchy, and since M is complete, it converges to a point $p \in M$. By continuity of f, p is a fixed point of f. On the other hand, since f is a contraction, it has at most one fixed point. $\square$

Exercise 2.1 (Perturbations of contractions)**.** Let $f\colon \mathbb{R}^d \circlearrowleft$ be a uniform contraction and $r > 0$ be arbitrary. We define the following balls

$$B_r^0(f) := \left\{ g \in C^0(\mathbb{R}^d, \mathbb{R}^d) : \|f - g\|_{C^0} := \sup_{x \in \mathbb{R}^d} \|f(x) - g(x)\| < r \right\},$$

and

$$B_r^L(f) := \left\{ g \in C^0(\mathbb{R}^d, \mathbb{R}^d) : \|f - g\|_L := \|f - g\|_{C^0} + \mathrm{Lip}(f - g) < r \right\}.$$

Show that given any $r > 0$, there is a $g \in B_r^0(f)$ which is not a contraction and such that $\mathrm{Fix}(g) := \{x \in \mathbb{R}^d : g(x) = x\}$ is an infinite set.
On the other hand, show that given the uniform contraction f, there exists $r_0 > 0$ such that every $g \in B_{r_0}^L(f)$ is a uniform contraction, and hence it exhibits a unique fixed point $p_g \in \mathbb{R}^d$. Finally, prove that given any $\varepsilon > 0$, there exists $\delta \in (0, r_0)$ such that

$$\|p_f - p_g\| < \varepsilon, \quad \forall g \in B_\delta^L(f).$$

Another important example is the following one:

Example 2.1 (Gradient flows)**.** *Let M be the unit sphere $\mathbb{S}^2 := \{v \in \mathbb{R}^3 : \|v\| = 1\}$ and consider the function $L\colon \mathbb{S}^2 \to \mathbb{R}$ given by*

$$L(x, y, z) := -z, \quad \forall (x, y, z) \in \mathbb{S}^2,$$

and the flow $\Phi\colon \mathbb{R} \times \mathbb{S}^2 \to \mathbb{S}^2$ generated by the vector field $X(v) := -\nabla L(v)$, for every $v \in \mathbb{S}^2$ and where ∇ denotes the gradient operator on $\mathbb{S}^2$.
One can easily show that $(0, 0, -1)$ and $(0, 0, 1)$ are the only singularities of X (i.e. the critical points of L), or equivalently, the fixed points of Φ, and it holds

$$\Phi^t(v) \to (0, 0, -1), \quad \text{and} \quad \Phi^{-t}(v) \to (0, 0, 1),$$

as $t \to +\infty$, for every $v \in \mathbb{S}^2 \backslash \{(0, 0, \pm 1)\}$.
Generalizing this example, one can consider an arbitrary compact differentiable manifold M endowed with a Riemannian structure $\langle \cdot, \cdot \rangle$ and a smooth function $L\colon M \to \mathbb{R}$. Then we can define the gradient of L as the only vector field ∇L such that

$$YL = dL(Y) = \langle \nabla L, Y \rangle,$$

for every vector field Y on M.
So, if we consider the flow $\Phi\colon \mathbb{R} \times M \to M$ induced by the vector field $X := -\nabla L$, one can easily check that

$$\partial_t (L \circ \Phi^t)(p)\big|_{t=0} = -\|\nabla L(p)\|^2 = -\langle \nabla L(p), \nabla L(p) \rangle, \quad \forall p \in M.$$

That means, given any point $p \in M$ the map $\mathbb{R} \ni t \mapsto L \circ \Phi^t(p) \in \mathbb{R}$ is not increasing, and it is strictly decreasing if and only if p is a regular point for L is a regular point of L, i.e. $dL_p\colon T_pM \to \mathbb{R}$ is not identically zero.
Moreover, if L has a finite number of critical points (i.e. points where the derivative dL vanishes), then one can show that for every $p \in M$, there exist critical points of L $\alpha_p, \omega_p \in M$ such that

$$\Phi^{-t}(p) \to \alpha_p, \qquad \Phi^t(p) \to \omega_p,$$

as $t \to +\infty$.

3 The Fundamental Theorem of Dynamical Systems

Motivated by Example 2.1, we would like to find functions decreasing along the orbits of arbitrary topological dynamical systems.

Definition 3.1 (Lyapunov function)**.** Given a continuous dynamical systems $f\colon M \circlearrowleft$ on a metric space M, a *Lyapunov function* for f is a continuous function $L\colon M \to \mathbb{R}$ such that

$$L\big(f(x)\big) \leqslant L(x), \quad \forall x \in M.$$

Of course, according to this definition every constant function is a Lyapunov one for every topological dynamical system, which does not seem to be very interesting. So, we are going to look for the "best possible" Lyapunov function.

3.1 Different forms of recurrence

It would be desirable to find a Lyapunov function such that $L\big(f(x)\big) < L(x)$ for every $x \in M$. However this is clearly impossible when f has, for instance, a **fixed point,** i.e. a point $p \in M$ such that $f(p) = p$. The set of fixed points of f will be denoted by $\mathrm{Fix}(f)$.

More generally, we write $p \in \mathrm{Per}(f)$ and say that p is a **periodic point** when there is $n \geqslant 1$ such that $f^n(p) = p$. We define $\mathrm{Per}(f) := \bigcup_{n\geqslant 1} \mathrm{Fix}(f^n)$.

Another form of recurrence that represents an obstruction for the existence of a Lyapunov functions satisfying the strict inequality for every $p \in M$ is the following: we say that a point $p \in M$ is **positively recurrent** for f when there exists strictly increasing sequence of natural numbers $n_j \uparrow +\infty$ such that $f^{n_j}(p) \to p$, as $n_j \to +\infty$. The set of positively recurrent points of f shall be denoted by $\mathrm{Rec}^+(f)$. When $f\colon M \circlearrowleft$ is bijective, one says that $p \in M$ is **negatively recurrent** for f when it is positively recurrent for f^{-1}. The set of negatively recurrent points will be denoted by $\mathrm{Rec}^-(f)$.

Given a point $p \in M$, we define its **ω-limit set** by

$$\omega_f(p) := \{x \in M : \exists n_j \uparrow \infty,\ f^{n_j}(p) \to x,\ \text{as } n_j \to +\infty\},$$

and the **positive limit set** of f by

$$L^+(f) := \bigcup_{p\in M} \omega_f(p).$$

When f is invertible, we also define the **α-limit set** of $p \in M$ by $\alpha_f(p) := \omega_{f^{-1}}(p)$, and the so called **limit set** of f by

$$L(f) := \bigcup_{p\in M} \alpha_f(p) \cup \omega_f(p).$$

We say that a point $p \in M$ is **non-wandering** when for every neighborhood U of p, there exists $n \geqslant 1$ such that $U \cap f^{-n}(U) \neq \varnothing$. The set of non-wandering points for f is usually denoted by $\Omega(f)$.

We say that a subset $K \subset M$ is **f-invariant** when $f(K) = K$, and it is said to be **totally f-invariant** when $f(K) = K = f^{-1}(K)$. Observe that if $f\colon M \circlearrowleft$ is bijective, then $K \subset M$ is totally f-invariant if and only if it is f-invariant.

Exercise 3.1. Let M denote a compact metric space and $f\colon M \circlearrowleft$ be a homeomorphism. Show that

$$\mathrm{Fix}(f) \subset \mathrm{Per}(f) \subset \mathrm{Rec}^+(f) \subset L(f) \subset \Omega(f),$$

and all these sets are f-invariant.

1. Show that $\mathrm{Fix}(f)$ and $\Omega(f)$ are compact.
2. Show that $\mathrm{Rec}^+(f) \cap \mathrm{Rec}^-(f) \neq \varnothing$. (Hint: show that there exists a non-empty compact subset $K \subset M$ which is **minimal** for f, i.e. K is f-invariant and it does not contain any proper compact f-invariant subset).
3. Show that if $L\colon M \to \mathbb{R}$ is a Lyapunov function for f and $p \in \Omega(f)$, then it holds $L\big(f(p)\big) = L(p)$.

Finally we introduce the notion of *chain recurrence.* Given a homeomorphism $f\colon M \circlearrowleft$ on the compact metric space (M, d) and an arbitrary $\varepsilon > 0$, a sequence $(x_n)_{n\in\mathbb{Z}}$ of points of M is said to be an **ε-chain** or **ε-pseudo-orbit** when

$$d\big(f(x_i), x_{i+1}\big) < \varepsilon, \quad \forall n \in \mathbb{Z}.$$

A point $p \in M$ is said to be **chain recurrent** when for every $\varepsilon > 0$ there exists a periodic ε-chain $(x_n)_{n\in\mathbb{Z}}$ with $p = x_0$. The set of chain recurrent points is denoted by $\mathscr{R}(f)$.

Exercise 3.2. Show that $\mathscr{R}(f)$ is a compact f-invariant set and $\Omega(f) = \Omega(f^{-1}) \subset \mathscr{R}(f) = \mathscr{R}(f^{-1})$.

Given any $p \in M$ and any $\varepsilon > 0$, we define the **ε-chainable set from** p as the set

$$\mathscr{R}_f(p, \varepsilon) := \{x \in M : \exists (x_n)_{n\in\mathbb{Z}}\ \varepsilon\text{-chain}, \exists k \in \mathbb{N}, x_0 = p,\ x_k = x\},$$

and the **chainable set from** p by

$$\mathscr{R}_f(p) := \bigcap_{\varepsilon>0} \mathscr{R}_f(p, \varepsilon).$$

So, we have $p \in \mathscr{R}(f)$ if and only if $p \in \mathscr{R}_f(p)$.

There is a fundamental and elementary notion that will play a crucial role in the understanding of the orbit structure of a given dynamical system, which is the concept of *trapping region.* We say that a subset $U \subset M$ is a **trapping region** for f when it is open and $f\big(\overline{U}\big) \subset U$.

Observe that if $U \subset M$ is a trapping region for f if and only if the set $M \backslash \overline{U}$ is a trapping region for f^{-1}.

Given a trapping region U for f, we can define the **dual pair of attractor/repellor** asso-

ciated to U as the pair of compact f-invariant sets

$$A_U^+ := \bigcap_{n\geqslant 0} f^n\big(\overline{U}\big), \quad \text{and} \quad A_U^- := \bigcap_{n\geqslant 0} f^{-n}\big(M\backslash U\big).$$

Exercise 3.3. Show that A_U^+ and A_U^- are compact, f-invariant and contained in U and $M\backslash\overline{U}$, respectively. Moreover, they are non-empty if and only if U and $M\backslash\overline{U}$ are non-empty.

We have the following fundamental result relating chain recurrence and trapping regions:

Proposition 3.1. Let $f\colon M \circlearrowleft$ be a homeomorphism on the compact metric space (M,d). Then, given any point $p\in M$ and any $\varepsilon>0$, the ε-chainable set from p is a trapping region for f.

Proof. Let us start showing that $\mathscr{R}_f(p,\varepsilon)$ is open. To do that, let us inductively defined

$$U_1 := B_\varepsilon(p) = \{y\in M : d(x,y)<\varepsilon\},$$

and

$$U_n := \bigcup_{x\in U_{n-1}} B_\varepsilon\big(f(x)\big), \quad \forall n\geqslant 2.$$

One can easily check that U_n is open for every $n\geqslant 1$ and $\mathscr{R}_f(p,\varepsilon) = \bigcup_{n\geqslant 1} U_n$. So, $\mathscr{R}_f(p,\varepsilon)$ is open.

By the very same definition, it is clear that $f\big(\mathscr{R}_f(p,\varepsilon)\big)\subset \mathscr{R}_f(p,\varepsilon)$. Now, let y be an arbitrary point of $\overline{\mathscr{R}_f(p,\varepsilon)}$. Since M is compact and f is continuous, there is $\delta>0$ such that $d\big(f(x_1),f(x_2)\big)<\varepsilon$, whenever $d(x_1,x_2)<\delta$. Then, there is $x\in\mathscr{R}_f(p,\varepsilon)\cap B_\delta(y)$. So there is a finite segment of ε-pseudo orbit $x_0=p,x_1,\dots,x_n=x$. Since $x\in B_\delta(y)$, we conclude $x_0=p,x_1,\dots,x_n=x,x_{n+1}:=f(y)$ is a finite segment of ε-pseudo orbit, too. Hence, $f(y)\in\mathscr{R}_f(p,\varepsilon)$, and we conclude that $f\big(\overline{\mathscr{R}_f(p,\varepsilon)}\big)\subset\mathscr{R}_f(p,\varepsilon)$. $\square$

Exercise 3.4. Let $f\colon M\circlearrowleft$ be a homeomorphism on the compact metric space (M,d) and $U\subset M$ be a trapping region for f. Show that

$$\mathscr{R}(f)\subset A_U^+\sqcup A_U^-,$$

where $\sqcup$ denotes the disjoint union and A_U^+, A_U^- is the dual pair of attractor/repellor associated to U.

Now we can define the notion of **chain recurrence classes.** If $f\colon M\circlearrowleft$ is a homeomorphism on the compact metric space (M,d) and $x,y\in\mathscr{R}(f)$, we say that they are **chain equivalent** when $y\in\mathscr{R}_f(x)$ and $x\in\mathscr{R}_f(y)$, i.e. $x,y\in\mathscr{R}_f(x,\varepsilon)\cap\mathscr{R}_f(y,\varepsilon)$, for every $\varepsilon>0$.

Then we have the following

Exercise 3.5 (Chain-recurrence classes)**.** If f and M are as above, let us show that *chain equivalence* determines an equivalence relation on $\mathscr{R}(f)$. Their equivalence classes are called **chain-recurrence classes.** Show that every chain-recurrent class is compact and f-invariant.
Moreover, two points $x, y \in \mathscr{R}(f)$ are chain equivalent if and only if either $x, y \in A_U^+$, or $x, y \in A_U^-$, for every trapping region U for f.

Exercise 3.6. If f and M are as above, show that the following set is at mots countable:

$$\left\{(A_U^+, A_U^-) : U \subset M \text{ is a trapping region for } f\right\}.$$

Using all previous exercises and results, one can easily prove the so-called "Fundamental Theorem of Dynamical Systems" due to Conley [Con78]:

Theorem 3.2 (Existence of complete Lyapunov functions)**.** Given a homeomorphism $f\colon M \circlearrowleft$ on the compact metric space (M, d), there exists a *complete Lyapunov function* for f, i.e. a continuous function $L\colon M \to \mathbb{R}$ such that the following conditions hold:

1. $L\big(f(x)\big) \leqslant L(x)$, for every $x \in M$,
2. $L\big(f(x)\big) = L(x)$ if and only if $x \in \mathscr{R}(f)$,
3. $L(x) = L(y)$, with $x, y \in \mathscr{R}(f)$ if and only if x and y are chain equivalent, and,
4. $L\big(\mathscr{R}(f)\big)$ is a compact totally disconnected set, i.e. it does not contain any open interval.

References

[Con78] C. Conley, *Isolated invariant sets and the Morse index*, CBMS Regional Conference Series in Mathematics, vol. 38, American Mathematical Society, Providence, RI, 1978.