

1. **ANA 222 Carpal Tunnel**
By Dr. Mbaka G.O.
The carpal bones are deeply concave on their flexor surface. This bony gutter is converted into the carpal tunnel by flexor retinaculum. The bony flexors of the thumb and fingers run through the tunnel. The four tendons of the superficial flexor are separate from each other and pass through in two rows, middle and ring finger tendons lying in front of index and little finger tendons.
2. The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.
3. The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.
4. The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.
5. The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.
6. The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.

ANA 222 Carpal Tunnel

By Dr. Mbaka G.O.

The carpal bones are deeply concave on their flexor surface. This bony gutter is converted into the carpal tunnel by flexor retinaculum. The bony flexors of the thumb and fingers run through the tunnel. The four tendons of the superficial flexor are separate from each other and pass through in two rows, middle and ring finger tendons lying in front of index and little finger tendons.

The tendons of flexor digitorum profundus on the other hand lie all on the same plane deep on the carpal bones with the tendon of the index finger separated while the other three are adherent to each other in the palm as they pass through the tunnel to palm.

FILE

HOME

INSERT

DESIGN


TRANSITIONS

ANIMATIONS


SLIDE SHOW

REVIEW


VIEW



Clipboard



Slides



Font



Paragraph



Drawing



Editing

1 **And... Carpal Tunnel**
By Dr. Niles G.O.

The carpal bones are deeply concave on two faces which they form a tunnel into the carpal tunnel. The tendons of the flexor digitorum profundus and flexor digitorum superficialis pass through the tunnel. The tendons of the flexor pollicis longus and the tendons of the flexor pollicis longus pass through the tunnel. The tendons of the flexor pollicis longus and the tendons of the flexor pollicis longus pass through the tunnel.

2 **all tendons of superficial and deep flexors share a common synovial sheath which does not invest them completely but reflected from their radial sides where artery of supply gain entrance to the tendon. It is as though from the radial side.**

3 **The tendon of flexor pollicis longus lies on its radial side as it passes through the carpal tunnel. The carpal bones are deeply concave on two faces which they form a tunnel into the carpal tunnel. The tendons of the flexor digitorum profundus and flexor digitorum superficialis pass through the tunnel.**

4 **Carpal tunnel syndrome is the result of increased pressure within the carpal tunnel. The pressure is increased by the swelling of the tendons, the swelling of the synovial sheath, and the swelling of the carpal bones. The pressure is increased by the swelling of the tendons, the swelling of the synovial sheath, and the swelling of the carpal bones.**

5 **the symptoms of the syndrome are numbness, tingling, burning, and pain—primarily in the thumb and index, middle, and ring fingers.**

Occasional shock-like sensations that radiate to the thumb and index, middle, and ring fingers.

Awakening at night due to numbness, tingling, and pain.


Weakness and clumsiness in the hand—may make it difficult to perform fine motor tasks such as buttoning your clothes.

Swelling of the hand—due to inflammation, edema, or a loss of proprioceptive awareness of where your hand is going.


6 **No symptoms just feeling weakness for the arm and a slight tingling sensation in the hand.**

All 8 tendons of superficial and deep flexors share a common synovial sheath which does not invest them completely but reflected from their radial sides where artery of supply gain entrance to the tendon. It is as though from the radial side.

Because these boundaries are very rigid, the carpal tunnel has little capacity to "stretch" or increase in size.



Simplified diagram of the Carpal Tunnel.
© MedPrepOnline.Com



Clipboard



Slides



Font



Paragraph



Drawing



Editing

1 **And 222 Carpal Tunnel**
By Dr. M. S. G. O.

The carpal bones are deeply concave on their base surface. This deep surface is converted into the carpal tunnel by the transverse carpal ligament. The base of the thumb and fingers pass through the tunnel. The base of the middle finger and the base of the ring finger pass through the tunnel. The base of the little finger passes through the tunnel.

The tendons of these digits are on the other hand on all on the other side of the carpal tunnel. The tendons of the middle finger are on the other side of the carpal tunnel. The tendons of the ring finger are on the other side of the carpal tunnel. The tendons of the little finger are on the other side of the carpal tunnel.

2

3

4

5

What is the cause of the carpal tunnel?

Numbness, tingling, burning, and pain—primarily in the thumb and index, middle, and ring fingers.

Chronic, disabling symptoms that relate to the thumb and index, middle, and ring fingers.


Swelling and tenderness in the hand may make it difficult to perform fine movements, such as buttoning your clothes.




Trapping fingers—due to swelling, tenderness, or a loss of proprioception (awareness of where your hand is in space).

6

The symptoms of carpal tunnel syndrome are further aggravated by the use of a computer mouse.

The tendon of flexor pollicis longus lies on its own synovial sheath as it passes through the fibro-osseous tunnel beside the trapezium. The median nerve passes beneath the flexor retinaculum on the lateral side of flexor digitorum superficialis.





1 **Just 1.0 Carpal Tunnel**
By Dr. N. S. G. D.

The carpal bones are deep, concave on their flexor surface. The deep groove is surrounded by the carpal tunnel. The median nerve and the tendons of the flexor digitorum profundus and flexor digitorum superficialis pass through the tunnel. The tendons of the flexor digitorum profundus are on the outer side of the tunnel and the tendons of the flexor digitorum superficialis are on the inner side of the tunnel.

The tendons of the flexor digitorum profundus are on the outer side of the tunnel and the tendons of the flexor digitorum superficialis are on the inner side of the tunnel.

2

The carpal tunnel is a narrow passage in the wrist. It is formed by the carpal bones and the transverse carpal ligament. The median nerve and the tendons of the flexor digitorum profundus and flexor digitorum superficialis pass through the tunnel.

3

The tendons of the flexor digitorum profundus and flexor digitorum superficialis are on the outer side of the tunnel and the tendons of the flexor digitorum profundus are on the inner side of the tunnel.

4 **Carpal tunnel syndrome** is the tightly crowded flexor tunnel, the median nerve can be compressed especially by long continued swelling in the synovial sheaths. The symptoms include wasting and weakness of thenar muscles with loss of power of opposition of thumb. Anaesthesia over three and half digits on thumb side

5

Thenar (flexor) muscle - Numbness, tingling, burning, and pain - especially in the thumb, index, middle, and ring fingers.


Discomfort, stiffness, and pain in the thumb and index, middle, and ring fingers.

Weakness and numbness in the thumb and index, middle, and ring fingers.

Weakness and numbness in the thumb and index, middle, and ring fingers.

Weakness and numbness in the thumb and index, middle, and ring fingers.

Carpal tunnel syndrome: In the tightly crowded flexor tunnel, the median nerve can be compressed especially by long continued swelling in the synovial sheaths. The symptoms include wasting and weakness of thenar muscles with loss of power of opposition of thumb. Anaesthesia over three and half digits on thumb side



FILE

HOME

INSERT

DESIGN


TRANSITIONS

ANIMATIONS


SLIDE SHOW

REVIEW

VIEW



Clipboard



Slides



Font






Paragraph



Drawing



Editing



No anaesthesia over thenar eminence for the area of skin is supplied by palmar branch of median nerve.


- Carpal tunnel syndrome must be distinguished from median nerve damage at higher level.
- In the latter case the palmar cutaneous branch will be affected and in addition weakening the relevant flexor muscles in forearm. In the carpal tunnel syndrome the terminal phalanx of thumb can be flexed with normal power but at higher power can be lost.

- 3
- 4
- 5
- 6
- 7
- 8

Other risk factors for carpal tunnel syndrome include:

Heredity. This is likely an important factor. The carpal tunnel may be smaller in some people or there may be anatomic differences that change the amount of space for the nerve—and these traits can run in families.

Repetitive hand use. Repeating the same hand and wrist motions or activities over a prolonged period of time may aggravate the tendons in the wrist, causing swelling that puts pressure on the nerve.



Hand and wrist position. Doing activities that involve extreme flexion or extension of the hand and wrist for a prolonged period of time can increase pressure on the nerve.

Pregnancy. Hormonal changes during pregnancy can cause swelling.

Health conditions. Diabetes, rheumatoid arthritis, and thyroid gland imbalance are conditions that are associated with carpal tunnel syndrome.