

Started on	Friday, 10 January 2025, 1:29 AM
State	Finished
Completed on	Friday, 10 January 2025, 1:29 AM
Time taken	20 secs
Marks	0.00/15.00
Grade	0.00 out of 108.00 (0%)

Question 1

Incorrect

Mark 0.00 out of 1.00

EN: Find (if any) three prime numbers x , y , and z such that $xyz = 6215$. If such prime numbers do not exist write **"None"** without quotation marks.

ID: Carilah (jika ada) tiga bilangan prima x , y , dan z yang memenuhi $xyz = 6215$. Jika bilangan-bilangan yang dimaksud tidak ada tuliskan **"None"** tanpa tanda kutip.

Answer: (penalty regime: 20,40, ... %)

[Reset answer](#)

```
x = ...
y = ...
z = ...

# erase ... and put a number or type "None" without quotation marks
```

Your code failed one or more hidden tests.

Your code must pass all tests to earn any marks. Try again.

▼ Show/hide question author's solution (Python3)

```
1 x = 5
2 y = 11
3 z = 113
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Incorrect

Marks for this submission: 0.00/1.00.

Question 2

Incorrect

Mark 0.00 out of 1.00

EN: Find (if any) three prime numbers $x < y < z$ such that $xyz = 3531$. If such prime numbers do not exist write **"None"** without quotation marks.

ID: Carilah (jika ada) tiga bilangan prima $x < y < z$ yang memenuhi $xyz = 3531$. Jika bilangan-bilangan yang dimaksud tidak ada tuliskan **"None"** tanpa tanda kutip.

Answer: (penalty regime: 20,40, ... %)

[Reset answer](#)

```
1 x = ...
2 y = ...
3 z = ...
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Your code failed one or more hidden tests.

Your code must pass all tests to earn any marks. Try again.

▼ Show/hide question author's solution (Python3)

```
1 x = 3
2 y = 11
3 z = 107
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Incorrect

Marks for this submission: 0.00/1.00.

Question 3

Incorrect

Mark 0.00 out of 1.00

EN: Find (if any) three integers x , y , and z such that $x^2 + y^2 + z^2 = 3$ but $x \neq y \neq z$ (all x , y , and z are different). If such integers do not exist write **"None"** without quotation marks.

ID: Carilah (jika ada) tiga bilangan bulat x , y , dan z yang memenuhi $x^2 + y^2 + z^2 = 3$ tetapi $x \neq y \neq z$ (semua x , y , dan z berbeda). Jika bilangan-bilangan yang dimaksud tidak ada tuliskan **"None"** tanpa tanda kutip.

Answer: (penalty regime: 20,40, ... %)

[Reset answer](#)

```
1 x = ...
2 y = ...
3 z = ...
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Your code failed one or more hidden tests.

Your code must pass all tests to earn any marks. Try again.

▼ Show/hide question author's solution (Python3)

```
1 x = None
2 y = None
3 z = None
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Incorrect

Marks for this submission: 0.00/1.00.

Question 4

Incorrect

Mark 0.00 out of 1.00

EN: Find (if any) three integers x , y , and z such that $x^2 + y^2 + z^2 = 2xy$ but $x \neq y \neq z$ (all x , y , and z are different). If such integers do not exist write "**None**" without quotation marks.

ID: Carilah (jika ada) tiga bilangan bulat x , y , dan z yang memenuhi $x^2 + y^2 + z^2 = 2xy$ tetapi $x \neq y \neq z$ (semua x , y , dan z berbeda). Jika bilangan-bilangan yang dimaksud tidak ada tuliskan "**None**" tanpa tanda kutip.

Answer: (penalty regime: 20,40, ... %)

[Reset answer](#)

```
1 x = ...
2 y = ...
3 z = ...
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Your code failed one or more hidden tests.

Your code must pass all tests to earn any marks. Try again.

▼ Show/hide question author's solution (Python3)

```
1 x = None
2 y = None
3 z = None
4
5 # erase ... and put a number or type "None" without quotation marks
6 # hapus ... dan berikan bilangan atau ketik "None" tanpa tanda kutip
```

Incorrect

Marks for this submission: 0.00/1.00.

Question 5

Not answered

Mark 0.00 out of 1.00

EN: An integer m is divisible by an integer d if there is an integer k such that $m = k \cdot d$. Suppose a and b are integers such that a is divisible by 4 and b is divisible by 7.

Suppose we consider the following statements:

1. $S1: a + b$ is always divisible by 15.
2. $S2: ab$ is always divisible by 14.
3. $S3: 2a + b$ is always even.
4. $S4: a(b + 1)$ is always divisible by 2.
5. $S5: 3a - b$ is always even.

Choose all statements that are true based on the previous assumption. Write your answer in the following Python format:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

For example, if you think that $S1$, $S2$, and $S3$ are the correct choices while $S4$ and $S5$ are incorrect, then you need to write

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

ID: Sebuah bilangan bulat m habis dibagi d jika terdapat bilangan bulat k sehingga $m = k \cdot d$. Misalkan a dan b adalah bilangan bulat dengan sifat a habis dibagi 9 dan b habis dibagi 6.

Misalkan kita meninjau pernyataan-pernyataan berikut:

1. $S1: a + b$ selalu habis dibagi 15.
2. $S2: ab$ selalu habis dibagi 14.
3. $S3: 2a + b$ selalu genap.
4. $S4: a(b + 1)$ selalu habis dibagi 2.
5. $S5: 3a - b$ selalu genap.

Pilihlah semua pernyataan yang benar berdasarkan asumsi sebelumnya. Tuliskan jawaban Anda dalam format Python berikut:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

Sebagai contoh, jika Anda berpikir bahwa $S1$, $S2$, dan $S3$ adalah jawaban yang benar sedangkan $S4$ dan $S5$ salah, maka Anda perlu menulis

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

Answer: (penalty regime: 20,40, ... %)

Reset answer

```
1 S1 = True/False
2 S2 = True/False
3 S3 = True/False
4 S4 = True/False
5 S5 = True/False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

▼ Show/hide question author's solution (Python3)

```
1 S1 = False
2 S2 = True
3 S3 = False
4 S4 = True
5 S5 = False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

Question 6

Not answered

Mark 0.00 out of 1.00

EN: An integer m is divisible by an integer d if there is an integer k such that $m = k \cdot d$. Suppose a and b are integers such that a is divisible by 10 and b is divisible by 8.

Suppose we consider the following statements:

1. $S1: a + b$ is always divisible by 18.
2. $S2: ab$ is always divisible by 20.
3. $S3: 2a + b$ is always odd.
4. $S4: a(b + 1)$ is always divisible by 10.
5. $S5: a - 5b$ is always divisible by 30.

Choose all statements that are true based on the previous assumption. Write your answer in the following Python format:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

For example, if you think that $S1$, $S2$, and $S3$ are the correct choices while $S4$ and $S5$ are incorrect, then you need to write

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

ID: Sebuah bilangan bulat m habis dibagi d jika terdapat bilangan bulat k sehingga $m = k \cdot d$. Misalkan a dan b adalah bilangan bulat dengan sifat a habis dibagi 10 dan b habis dibagi 8.

Misalkan kita meninjau pernyataan-pernyataan berikut:

1. $S1: a + b$ selalu habis dibagi 18.
2. $S2: ab$ selalu habis dibagi 20.
3. $S3: 2a + b$ selalu ganjil.
4. $S4: a(b + 1)$ selalu habis dibagi 10.
5. $S5: a - 5b$ selalu habis dibagi 30.

Pilihlah semua pernyataan yang benar berdasarkan asumsi sebelumnya. Tuliskan jawaban Anda dalam format Python berikut:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

Sebagai contoh, jika Anda berpikir bahwa $S1$, $S2$, dan $S3$ adalah jawaban yang benar sedangkan $S4$ dan $S5$ salah, maka Anda perlu menulis

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

Answer: (penalty regime: 20,40, ... %)

Reset answer

```
1 S1 = True/False
2 S2 = True/False
3 S3 = True/False
4 S4 = True/False
5 S5 = True/False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

▼ Show/hide question author's solution (Python3)

```
1 S1 = False
2 S2 = True
3 S3 = False
4 S4 = True
5 S5 = False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```


Question 7

Not answered

Mark 0.00 out of 1.00

EN: An integer m is divisible by an integer d if there is an integer k such that $m = k \cdot d$. Suppose $n = a \cdot b \cdot c$ and n is divisible by 7.

Suppose we consider the following statements:

1. $S1$: a , b , and c are always divisible by 7.
2. $S2$: at least one of a , b , and c is not divisible by 7.
3. $S3$: at least two of a , b , and c are not divisible by 7.
4. $S4$: at least one of a , b , and c is divisible by 7.
5. $S5$: at least two of a , b , and c are divisible by 7.

Choose all statements that are true based on the previous assumption. Write your answer in the following Python format:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

For example, if you think that $S1$, $S2$, and $S3$ are the correct choices while $S4$ and $S5$ are incorrect, then you need to write

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

ID: Sebuah bilangan bulat m habis dibagi d jika terdapat bilangan bulat k sehingga $m = k \cdot d$. Misalkan $n = a \cdot b \cdot c$ dan n habis dibagi 7.

Misalkan kita meninjau pernyataan-pernyataan berikut:

1. $S1$: a , b , dan c semuanya selalu habis dibagi 7.
2. $S2$: setidaknya satu dari a , b , dan c tidak habis dibagi 7.
3. $S3$: setidaknya dua dari a , b , dan c tidak habis dibagi 7.
4. $S4$: setidaknya satu dari a , b , dan c habis dibagi 7.
5. $S5$: setidaknya dua dari a , b , dan c habis dibagi 7.

Pilihlah semua pernyataan yang benar berdasarkan asumsi sebelumnya. Tuliskan jawaban Anda dalam format Python berikut:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

Sebagai contoh, jika Anda berpikir bahwa $S1$, $S2$, dan $S3$ adalah jawaban yang benar sedangkan $S4$ dan $S5$ salah, maka Anda perlu menulis

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

Answer: (penalty regime: 20,40, ... %)

Reset answer

```
1 S1 = True/False
2 S2 = True/False
3 S3 = True/False
4 S4 = True/False
5 S5 = True/False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

▼ Show/hide question author's solution (Python3)

```
1 S1 = False
2 S2 = False
3 S3 = False
4 S4 = True
5 S5 = False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

Question 8

Not answered

Mark 0.00 out of 1.00

EN: An integer m is divisible by an integer d if there is an integer k such that $m = k \cdot d$. An integer is even if it is divisible by 2. An integer is odd if it is not even. Suppose $n = a + b + c$ and n is odd.

Suppose we consider the following statements:

1. $S1$: a , b , and c are always odd.
2. $S2$: at least one of a , b , and c is odd.
3. $S3$: a is odd, b , and c are even.
4. $S4$: a and b are even, and c is odd.
5. $S5$: at least two of a , b , and c are even.

Choose all statements that are true based on the previous assumption. Write your answer in the following Python format:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

For example, if you think that $S1$, $S2$, and $S3$ are the correct choices while $S4$ and $S5$ are incorrect, then you need to write

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

ID: Sebuah bilangan bulat m habis dibagi d jika terdapat bilangan bulat k sehingga $m = k \cdot d$. Sebuah bilangan dikatakan genap bila habis dibagi 2. Sebuah bilangan dikatakan ganjil jika tidak genap. Misalkan $n = a + b + c$ dan n ganjil.

Misalkan kita meninjau pernyataan-pernyataan berikut:

1. $S1$: a , b , dan c selalu ganjil.
2. $S2$: setidaknya satu dari a , b , dan c ganjil.
3. $S3$: a ganjil, b , dan c ganjil.
4. $S4$: a dan b genap, dan c ganjil.
5. $S5$: setidaknya dua dari a , b , dan c genap.

Piihlah semua pernyataan yang benar berdasarkan asumsi sebelumnya. Tuliskan jawaban Anda dalam format Python berikut:

S1 = <True/False>

S2 = <True/False>

S3 = <True/False>

S4 = <True/False>

S5 = <True/False>

Sebagai contoh, jika Anda berpikir bahwa $S1$, $S2$, dan $S3$ adalah jawaban yang benar sedangkan $S4$ dan $S5$ salah, maka Anda perlu menulis

S1 = True

S2 = True

S3 = True

S4 = False

S5 = False

Answer: (penalty regime: 20,40, ... %)

Reset answer

```
1 S1 = True/False
2 S2 = True/False
3 S3 = True/False
4 S4 = True/False
5 S5 = True/False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

▼ Show/hide question author's solution (Python3)

```
1 S1 = False
2 S2 = False
3 S3 = True
4 S4 = True
5 S5 = False
6
7 # assign either True or False for each statement, True and False start with uppercase letter
8 # berikan nilai True atau False untuk masing-masing statement True dan False dimulai dengan huruf kapital
```

Question 9

Not answered

Marked out of 1.00

EN: In a jar, there are 91 candies as follows:

- 21 chocolate candies,
- 3 strawberry candies,
- 7 vanilla candies,
- 11 coffee candies.
- 14 mint candies,
- 13 milk candies,
- 5 orange candies,
- 17 banana candies,

All candies have **the same and identical wrappers**. You are asked to **take a number of candies so that at least you get four different flavors** (for example, you get one chocolate candy, one strawberry candy, one vanilla candy, and one coffee candy; or one chocolate candy, one vanilla candy, one mint candy, and one milk candy; or one strawberry candy, one coffee candy, one orange candy, and one mint candy). What is the minimum number of candies should you take if the candies are taken **randomly**? (You must ensure that you get four different flavors.)

ID: Di sebuah toples terdapat 91 permen dengan rincian:

- 21 permen cokelat,
- 3 permen stroberi,
- 7 permen vanila,
- 11 permen kopi,
- 14 permen mint,
- 13 permen susu,
- 5 permen jeruk,
- 17 permen pisang.

Semua permen memiliki **bungkus yang sama dan identik**. Anda diminta untuk **mengambil sejumlah permen dengan syarat Anda memperoleh setidaknya empat rasa berbeda** (misalnya Anda memperoleh satu permen cokelat, satu permen stroberi, satu permen vanila, dan satu permen kopi; atau satu permen cokelat, satu permen vanila, satu permen mint, dan satu permen susu; atau satu permen stroberi, satu permen kopi, satu permen jeruk, dan satu permen mint). Paling sedikit, berapa banyak permen yang harus Anda ambil jika pengambilan dilakukan secara **acak**? (Anda harus memastikan bahwa Anda memperoleh empat rasa permen yang berbeda.)

Answer:



The correct answer is: 53

Question 10

Not answered

Marked out of 1.00

EN: In a jar, there are 60 candies as follows:

- 7 chocolate candies,
- 13 strawberry candies,
- 9 vanilla candies,
- 10 coffee candies.
- 7 mint candies,
- 6 milk candies,
- 8 orange candies,

All candies have **the same and identical wrappers**. You are asked to **take a number of candies so that you get at least four flavors and each flavor is represented by at least two candies**. For example, you get two chocolate candies, two strawberry candies, two vanilla candies, and two coffee candies; or two mint candies, two milk candies, two orange candies, and two banana candies. What is the minimum number of candies should you take if the candies are taken **randomly**? (You must ensure that you get at least four different flavors and each flavor is represented by at least two candies.)

ID: Di sebuah toples terdapat 60 permen dengan rincian:

- 7 permen coklat,
- 13 permen stroberi,
- 9 permen vanila,
- 10 permen kopi,
- 7 permen mint,
- 6 permen susu,
- 8 permen jeruk,

Semua permen memiliki **bungkus yang sama dan identik**. Anda diminta untuk **mengambil sejumlah permen dengan syarat Anda memperoleh setidaknya empat rasa dan setiap rasa diwakili oleh setidaknya dua permen**. Sebagai contoh, Anda memperoleh dua permen coklat, dua permen stroberi, dua permen vanila, dan dua permen kopi; atau dua permen mint, dua permen susu, dua permen jeruk, dan dua permen pisang. Paling sedikit, berapa banyak permen yang harus Anda ambil jika pengambilan dilakukan secara **acak**? (Anda harus memastikan bahwa Anda memperoleh setidaknya empat rasa permen yang berbeda dan setiap rasa diwakili oleh dua permen.)

Answer:



The correct answer is: 37

Question 11

Not answered

Marked out of 1.00

EN: Find the result of $3 + 9 + 15 + 21 + \dots + 999\,999\,993 + 999\,999\,999 + 1\,000\,000\,005$. You may use a Python interpreter or the formula of arithmetic or geometric series you learn in high school. (Hint: use Python interpreter or MS Excel to ease your calculation. The answer is an integer consisting of 17 digits.)

Note: do not use periods, commas, or spaces as digit separators. For example, if your answer is 123 456 789 987 654 321 then write 123456789987654321.

ID: Carilah nilai dari $3 + 9 + 15 + 21 + \dots + 999\,999\,993 + 999\,999\,999 + 1\,000\,000\,005$. Anda dapat memakai interpreter Python atau formula untuk deret aritmetika atau geometri yang Anda pelajari di sekolah menengah. (Petunjuk: gunakan interpreter Python atau MS Excel untuk mempermudah kalkulasi Anda. Jawaban adalah bilangan bulat yang terdiri dari 17 digit.)

Catatan: jangan menggunakan titik, koma, atau spasi untuk pemisah digit. Sebagai contoh, jika jawaban Anda adalah 123 456 789 987 654 321 maka tulis 123456789987654321.

Answer:



The correct answer is: 83333334666666672

Question 12

Not answered

Marked out of 1.00

EN: Find the result of $3 + 8 + 13 + 18 + \dots + 999\,999\,993 + 999\,999\,998 + 1\,000\,000\,003$. You may use a Python interpreter or the formula of arithmetic or geometric series you learn in high school. (Hint: use Python interpreter or MS Excel to ease your calculation. The answer is an integer consisting of 18 digits.)

Note: do not use periods, commas, or spaces as digit separators. For example, if your answer is 123 456 789 987 654 321 then write 123456789987654321.

ID: Carilah nilai dari $3 + 8 + 13 + 18 + \dots + 999\,999\,993 + 999\,999\,998 + 1\,000\,000\,003$. Anda dapat memakai interpreter Python atau formula untuk deret aritmetika atau geometri yang Anda pelajari di sekolah menengah. (Petunjuk: gunakan interpreter Python atau MS Excel untuk mempermudah kalkulasi Anda. Jawaban adalah bilangan bulat yang terdiri dari 18 digit.)

Catatan: jangan menggunakan titik, koma, atau spasi untuk pemisah digit. Sebagai contoh, jika jawaban Anda adalah 123 456 789 987 654 321 maka tulis 123456789987654321.

Answer:



The correct answer is: 100000001100000000

Question 13

Not answered

Marked out of 1.00

EN: A sequence s_n is defined recursively as follows:

$$s_1 = 5, s_2 = 9, \text{ and } s_n = 2 \cdot s_{n-1} - s_{n-2} \quad \text{for } n \geq 3.$$

What is the value of s_5 ?

ID: Sebuah barisan s_n didefinisikan secara rekursif sebagai berikut:

$$s_1 = 5, s_2 = 9, \text{ dan } s_n = 2 \cdot s_{n-1} - s_{n-2} \quad \text{untuk } n \geq 3.$$

Berapakah nilai dari s_5 ?

Answer:



The correct answer is: 21

Question 14

Not answered

Marked out of 1.00

EN: A sequence r_n is defined recursively as follows:

$$r_1 = 4, r_2 = 7, \text{ and } r_n = r_{n-1} + 3 \cdot r_{n-2} \quad \text{untuk } n \geq 3.$$

What is the value of r_5 ?

ID: Sebuah deret r_n didefinisikan secara rekursif sebagai berikut:

$$r_1 = 4, r_2 = 7, \text{ dan } r_n = r_{n-1} + 3 \cdot r_{n-2} \quad \text{untuk } n \geq 3.$$

Berapakah nilai dari r_5 ?

Answer:



The correct answer is: 97

Question 15

Incorrect

Mark 0.00 out of 1.00

EN: Construct a Python 3 function **sum(n)** that takes a positive integer n as an input and performs the following computation:

$sum(n) = x(n) - y(n)$, where

$$x(n) = 9 + 16 + 23 + \dots + (7n - 5) + (7n + 2)$$

$$y(n) = 8 + 13 + 18 + \dots + (5n - 2) + (5n + 3),$$

for example:

1. $sum(1) = x(1) - y(1) = 9 - 8 = 1$,
2. $sum(2) = x(2) - y(2) = (9 + 16) - (8 + 13) = 4$,
3. $sum(3) = x(3) - y(3) = (9 + 16 + 23) - (8 + 13 + 18) = 9$.

The value of n is between 1 and 10^{12} . The time limit for the computation is 1 second per test case. The memory limit for the computation is 16 MB. To make your code efficient, derive an explicit formula (closed form) of $sum(n)$ using your knowledge learned in high school. You may further justify the correctness of your formula using induction.

Python hint: if a , b , and c are integers and c divides ab , then the integer expression $(ab)/2$ is represented as $(a * b) // 2$. The integer expression a^b is written as $a * b$.

ID: Buatlah sebuah fungsi Python 3 **sum(n)** yang mengambil bilangan bulat positif n sebagai masukan dan melakukan komputasi berikut:

$sum(n) = x(n) - y(n)$, where

$$x(n) = 9 + 16 + 23 + \dots + (7n - 5) + (7n + 2)$$

$$y(n) = 8 + 13 + 18 + \dots + (5n - 2) + (5n + 3),$$

sebagai contoh:

1. $sum(1) = x(1) - y(1) = 9 - 8 = 1$,
2. $sum(2) = x(2) - y(2) = (9 + 16) - (8 + 13) = 4$,
3. $sum(3) = x(3) - y(3) = (9 + 16 + 23) - (8 + 13 + 18) = 9$.

1.

Nilai dari n antara 1 dan 10^{12} . Batas waktu komputasi adalah 1 detik per kasus uji. Batas memori adalah 16 MB. Untuk membuat kode program Anda efisien, buatlah formula (bentuk tertutup) dari $sum(n)$ menggunakan pengetahuan Anda yang dipelajari di SMA. Anda dapat membuktikan kebenaran dari formula yang diperoleh menggunakan induksi.

Petunjuk Python: jika a , b , dan c adalah bilangan bulat dan c membagi ab , maka ekspresi $(ab)/2$ direpresentasikan sebagai $(a * b) // 2$. Ekspresi bilangan bulat a^b ditulis sebagai $a * b$.

For example:

Test	Input	Result
<code>print(sum(1))</code>	<code>print(sum(1))</code>	1
<code>print(sum(2))</code>	<code>print(sum(2))</code>	4
<code>print(sum(3))</code>	<code>print(sum(3))</code>	9
<code>print(sum(4))</code>	<code>print(sum(4))</code>	16
<code>print(sum(100))</code>	<code>print(sum(100))</code>	10000

Answer: (penalty regime: 20,40, ... %)

Reset answer

```

1 def sum(n):
2     # fix the following code
3     if n == 1: return (5 - 5)
4     elif n == 2: return (5 + 9) - (5 + 8)
5     elif n == 3: return (5 + 9 + 13) - (5 + 8 + 11)
6     else: return 0 # you may fix this line

```

	Test	Input	Expected	Got	
✖	print(sum(1))	print(sum(1))	1	0	✖
✖	print(sum(2))	print(sum(2))	4	1	✖
✖	print(sum(3))	print(sum(3))	9	3	✖
✖	print(sum(4))	print(sum(4))	16	0	✖
✖	print(sum(100))	print(sum(100))	10000	0	✖

[Show differences](#)

▼ Show/hide question author's solution (Python3)

```
1 ▼ def sum(n):  
2     # fix the following code  
3     if n == 1: return (9 - 8)  
4     elif n == 2: return (9 + 16) - (8 + 13)  
5     elif n == 3: return (9 + 16 + 23) - (8 + 13 + 18)  
6     else: return (n*(2*n))//2 # you may fix this line
```

Incorrect

Marks for this submission: 0.00/1.00.