



رويال حياة دنتل
ROYALE HAYAT DENTAL

POST-PROCEDURE CARE GUIDE



مستشفى رويال حياة
ROYALE HAYAT HOSPITAL



royalehayat



royale_hayat



/royalehayathospital

2536 0000
www.royalehayat.com

1. SURGICAL EXTRACTION CARE:

- Keep the gauze in place for 30 - 45 minutes to stop bleeding.
- Avoid touching the extraction site with your tongue or fingers.
- Do not rinse or spit for the first 24 hours.
- Avoid smoking and drinking with straw for a week.
- Avoid physical activity for at least 3 - 5 days.



2. IMPLANT CARE:

- Avoid touching the implant area with your tongue or fingers.
- Follow post-op instructions provided by your dentist.
- Rest and avoid physical activity for 48 hours.
- When lying down, use pillows to keep your head elevated.



3. ROOT CANAL TREATMENT CARE:

- Avoid chewing on the treated tooth until a permanent restoration is placed.
- If you have a temporary filling, take extra care to avoid chewing on the side of the treated tooth until the permanent crown is placed.
- It's common to experience mild discomfort or sensitivity for a few days after the procedure
- Delaying your permanent restoration can lead to reinfection or breakage of the tooth.

4. POST-BRIDGE CARE:

- Clean under and around your bridge with floss (use water floss) or a special interdental brush.
- Schedule regular checkups to ensure your bridge is in good condition.

5. POST CROWN CARE:

- Numbness: After the procedure, your mouth may still be numb. Avoid eating or drinking until the numbness subsides to prevent injury.
- Sensitivity: Some sensitivity to hot or cold is normal for the first few days. This should diminish as your tooth adjusts to the new crown.
- Avoid chewing on the side where the crown was placed.
- Use a soft-bristled toothbrush to prevent irritation.
- Floss Carefully: When flossing, slide the floss out instead of lifting it up to avoid loosening the crown

- Mild discomfort or soreness around the crown is normal and should subside within a few days.
- If you grind your teeth at night, consider getting a night guard to protect your crown and other teeth from excessive wear



COMMON INSTRUCTIONS:

- DIET:

- Eat soft foods (yogurt, mashed potatoes, soup) after the effect of local anesthetic agents.
- Avoid hot and spicy foods and drinks.
- Stick to soft foods during the adjustment period.
- Avoid sticky, hard and crunchy foods such as chewing gum, caramel, and nuts, as they could dislodge or damage the crown / bridge.
- Drink plenty of water to stay hydrated, but avoid hot drinks that can cause discomfort.

- PAIN MANAGEMENT:

- Take prescribed pain medications as directed.
- Apply an ice pack or a cold compress to the affected area (outside of the face) in 10 - 15 minutes intervals during the first 24 hours to reduce swelling and numb the area.

- ORAL HYGIENE:

- Brush gently, avoiding the surgical site.
- After 24 hours, rinse with warm salt water (1 tsp salt in a glass of water) 2 - 3 times a day.