Chapter 2

Creating graphs from data

Please read Chapter 6 from *Learning Statistics with R* before starting these assignments.

Content	Pag
Assignment 2.1. Creating a histogram by hand	
3 3 3	gram in R
Assignment 2.3: Creating and modifying a scatt	er plot in R
Assignment 2.4: Creating and modifying a line p	olot in R
Assignment 2.5. Creating a hox plot in R	26

Chapter 2: Creating graphs from data

Learning objectives of this chapter:

- Creating a histogram by hand and in R
- Creating and modifying a scatter plot in R
- Creating and modifying a line plot in R
- Creating a box plot in R

Assignment 2.1: Creating a histogram by hand




Suppose that you have the following set of fifteen numbers:

 $1.5 \ 5.5 \ 1.7 \ 7.2 \ 1.2 \ 7.9 \ 1.4 \ 3.6 \ 3.1 \ 3.8 \ 5.9 \ 3.6 \ 5.1 \ 3.2 \ 7.1$

2.1 a) For each of the ranges given below, find out their **frequency** in the data above.

Answer 2.1a:			
0 to 2	2 to 4	4 to 6	6 to 8
Frequency:	Frequency:	Frequency:	Frequency:

2.1 b) Draw the histogram for these data by hand. The name of the *x-axis* should represent the range of values, and the name of the *y-axis* should represent the **frequency** of the values that fall in the ranges of assignment 2.1a.



Assignment 2.2: Creating and modifying a histogram in R



2.2 a) Take the numbers from assignment 2.1a and use the c() function to enter these numbers in a vector called dataset4. Next, run the following code in R and explain what you see.

nist(dataset4)		
Answer 2.2a:		
		_

2.2 b) The graph resulting from the command in 2.2a is not the same as the histogram that you drew in assignment 2.1b. What is the difference between the graph from assignment 2.2a and your graph from assignment 2.1b?

Answer 2.2b:		
_		


2.2 c) Try to replicate your <u>exact</u> histogram from assignment 2.1b in R. This requires that you specify the **breaks** argument after a comma in the **hist()** function.



Hint 2.1: You can find more information on the <code>hist()</code> function by running <code>?hist</code> .

R code 2.2c:

2.2 d) Give your histogram a different color by adding and changing the col argument after the comma. Next, improve your histogram further by changing the x-axis name, the main title, and the rotation of the y-axis labels by adding more arguments.



R code 2.2d:		

2.2 e) Calculate the **mean** and **median** of the values in **dataset4**. Is the distribution of **dataset4 negatively skewed** or **positively skewed**? Explain your answer using the relation between the **mean** and the **median**.

Answer 2.2e:
The distribution is positively / negatively / not skewed.
Explanation:

Assignment 2.3: Creating and modifying a scatter plot in R



In this assignment you are going to work with a data set that is built into R.

The data() function gives you access to all the data sets that are built into R, or that are included with downloaded R packages. For example, you can load a data set called swiss into the environment by running the R code below:

data(swiss)

The data are imported as an object called **swiss**, which you can now also see in the environment. This particular data set contains some socio-economic indicators for each of 47 French-speaking provinces of Switzerland. In this assignment, you will focus on the column **Education** (the percentage education beyond primary school) and the column **Agriculture** (the percentage of males involved in agriculture as an occupation).

2.3 a) Extract the values of the two columns using the \$ sign and store them in two new variables called education and agriculture.



Hint 2.3: You can check the R environment to see the names of the objects that you have currently stored.

R code 2.3a:			

2.3 b) Use the plot() function to create a scatter plot of the two variables. Place the percentage of education beyond primary school on the x-axis and the percentage of males involved in agriculture as an occupation on the y-axis. Rename your axis labels to match the content of the figure.

R code 2.3b:		

		education beyond primary school and the percentage of males involved in agriculture as an occupation in the French-speaking provinces of Switzerland?
Ans	swe	er 2.3c:
-		
_		
2.3	d)	Give the points in your scatter plot a different color by changing the col argument after the comma. Next, improve the scatter plot further by changing the main title, the rotation of the <i>y-axis</i> labels, the shape of the points, and removing the borders.

2.3 c) Looking at the scatter plot, what can you say about the relation between the percentage of

R code 2.3d:			

Assignment 2.4: Creating and modifying a line plot in R



In this assignment, you are going to create a line plot of the daily closing prices of some major European stock indices: Germany DAX, Switzerland SMI, France CAC, and UK FTSE.

You can load this data set by running the code below:

```
data(EuStockMarkets)
stockData <- data.frame(EuStockMarkets)</pre>
```

The data is now loaded into the environment as the **EuStockMarkets** data set and immediately transformed to the **stockData** data set (this is because of technical reasons as the original data is in a time-series format). In this assignment, you will work with the **stockData** data set.

2.4 a) Create a line plot where the closing price of the DAX stock is displayed over time. Give your plot appropriate *x-axis* and *y-axis* names.

R code 2.4a:			

2.4 b) Find a function to add a separate line for the closing price of the SMI stock in red. You may try to add the other stocks as well using this function, but remember to adjust the *y-axis* accordingly so that all the lines can be viewed decently.



Hint 2.4: Do <u>not</u> use the **plot()** function to add a line to an already existing plot.

Maximum:

Assignment 2.5: Creating a box plot in R



For this next assignment, let's return to the **swiss** data set. More specifically, you will now have to focus on the values that are stored in the separate variable **agriculture**.

2.5 a)	Find out the	minimum , 1	ower quartile,	median ,	upper	quartile,	and
	maximum of	the agriculture	variable.				

2.5 b) Create a box plot of the agriculture variable.

R code 2.5b:			

2.5 c) Run the following code in R and explain the table output. How does the code work?

```
educationLevel <- rep('2.Medium', 47)
educationLevel[education <= 6] = '1.Low'
educationLevel[education >= 12] = '3.High'
table(educationLevel)
```

Answer 2.5c:		

 $2.5\,$ d) Create a box plot using the R code below and explain what you see.

poxplot(agriculture ~ educationLevel)
Answer 2.5d:

This page is intentionally left blank.