

Chapter 1

Descriptive statistics

Please read Chapter 1 (excl. section 1.3), Chapter 2, and Section 5.1-5.6 from *Learning Statistics with R* before starting these assignments.

<i>Content</i>	<i>Page</i>
Assignment 1.1: Descriptive statistics by hand	10
Assignment 1.2: Descriptive statistics of small data sets in R	12
Assignment 1.3: Descriptive statistics of larger data sets in R	14

Chapter 1: Descriptive statistics

Learning objectives of this chapter:

- Calculating the mean, mode, median, quartiles and range by hand
- Get started with R: set the working directory, inspect and load data
- Calculating the mean, mode, median and range in R

Assignment 1.1: Descriptive statistics by hand



Calculate by hand (or by the use of a pocket calculator) the descriptive statistics that are listed below:

- **mean**
- **mode**
- **median**
- **range**
- **lower and upper quartiles**
- **interquartile range**

1.1 a) 2 7 4 5 8 10 10 7 9 2 8 8 9 4 6

Answer 1.1a:

Mean:	_____	Range:	_____
Mode:	_____	Lower quartile:	_____
Median:	_____	Upper quartile:	_____
		Interquartile range:	_____

1.1 b) 7 7 6 5 2 1 3 7 5 9 9 10

Answer 1.1b:

Mean:	_____	Range:	_____
Mode:	_____	Lower quartile:	_____
Median:	_____	Upper quartile:	_____
		Interquartile range:	_____

1.1 c) Describe which question (1.1a or 1.1b) was more difficult to calculate and why.

Answer 1.1c:

1.1 d) Are these data sets **positively skewed** or **negatively skewed**? Circle the correct answer and explain why you chose this answer using the relation between the **mean**, the **median**, and the **mode**.

Answer 1.1d:

These data sets are **positively** / **negatively** skewed.

Explanation:

Assignment 1.2: Descriptive statistics of small data sets in R

This assignment assumes you opened a new script in the code editor in RStudio. Write and run your own code to answer the following questions.

In R, a one-dimensional row of numbers is represented by a **vector**.

- 1.2 a) Use the **c()** function to enter the numbers from assignment 1.1a in a new vector called **dataset1**. Next, run the following code in R and explain what you see.

```
View(dataset1)
```



*Hint 1.1: Check Part I of the R help on page 142 for more information on how to make a **vector**.*

R code 1.2a:

Answer 1.2a:

- 1.2 b) Write and run your own code in R to find the **mean**, **mode**, **median**, and **range** for the vector **dataset1**. Compare your answers with those of assignment 1.1a.



Hint 1.2: Check part IV of the R help on page 148 for descriptive statistics functions.



*Hint 1.3: There is no **mode** function in R, but you can find the mode in a frequency **table**.*

R code 1.2b:

Answer 1.2b:

Mean:	_____	Median:	_____
Mode:	_____	Range:	_____

1.2 c) Run the following code in R and explain what you see.

```
quantile(dataset1, type = 6)
```

Answer 1.2c:

1.2 d) Use the `c()` function to create a vector `dataset2` with the data from assignment 1.1b and find the **mean**, **mode**, **median**, **range**, and **quartiles** for these data. Compare your answers with the answers of assignment 1.1b.

R code 1.2d:**Answer 1.2d:**

Mean:	_____	Range:	_____
Mode:	_____	Lower quartile:	_____
Median:	_____	Upper quartile:	_____

Assignment 1.3: Descriptive statistics of larger data sets in R

For this assignment, you need the `bloodPressure.csv` data set that you can download from the online resources. This data set contains measurements of the age, blood pressure, cholesterol level, gender, and description of a random selection of people. It is normally used to look for relationships between these variables. Note that this is fake data and does not contain actual measurements.

Let's start with importing the data set (which is available in the online resources) into R.

- 1.3 a) Inspect and run the following code in R to import the blood pressure data and store it in the object `dataset3`. Explain how the code works and describe what the `bloodPressure.csv` file contains.

```
dataset3 <- read.csv(file.choose())
```

Answer 1.3a:

This method of importing data can be a lot of work if there are many files or if the script will be run many times. Faster methods exist though, for example by providing the full file path.

- 1.3 b) Describe and test ways this code can be improved to make importing a file easier.



Hint 1.4: Look at Part I of the R help at page 142 to find out more functions for importing data.

Answer 1.3b:

R code 1.3b:

The functions you used for descriptive statistics on the small data sets in assignment 1.1 can also be applied to the data set that is currently stored in `dataset3`.

- 1.3 c) Find the **mean**, **mode**, **median**, **range**, and **quartiles** for the column **Age** in `dataset3`. Describe this variable in running text using these statistics.



Hint 1.5: First find out how to extract (index) a specific column in a data frame using the `$` sign.

R code 1.3c:

Answer 1.3c:

For large data sets, it becomes a lot of work finding the **mode** in a frequency table each time. It is possible to import a package into the R session that contains a function for calculating the **mode** automatically. However, it is also possible to create a function that calculates the **mode** ourselves.

Run the following lines of R code together:

```
getMode <- function(x){  
  uniqx <- unique(x)  
  uniqx[which.max(tabulate(match(x, uniqx)))]  
}
```

You have now created your first R function and you will see it displayed separately in the R environment. This function will give you the **mode** for any **numeric** vector or column. It works by first extracting all unique values, counting their frequency, and then selecting the value with the highest frequency. Note that you can use this function, but will not be required to understand or make functions like this. However, for the interested reader, part III of the R help contains more information on how to create your own functions.

- 1.3 d) Use the new `getMode()` function to determine the **mode** for column **Age** in **dataset3** and check if it is consistent with your answer for assignment 1.3c.

R code 1.3d:

Answer 1.3d:

Mode: _____

- 1.3 e) Find the **mean**, **mode**, **median**, **range**, and **quartiles** for the column **BloodPressure** in **dataset3**. Also use the new `getMode()` function.

R code 1.3e:

Answer 1.3e:

Mean: _____ Range: _____

Mode: _____ Lower quartile: _____

Median: _____ Upper quartile: _____

- 1.3 f) Is the distribution of the values in the **BloodPressure** column **positively skewed** or **negatively skewed**? Explain your answer using the relation between the **mean**, **median**, and **mode**.

Answer 1.3f:

These data sets are **positively** / **negatively** / **not** skewed.

Explanation:

1.3 g) Determine the **variance** and **standard deviation** for the column **Cholesterol** in **dataset3** .

R code 1.3g:

Answer 1.3g:

Variance: _____

Standard deviation: _____

1.3 h) Validate the relation between the **variance** and the **standard deviation** by performing a calculation in R.

R code 1.3h:

This page is intentionally left blank.