

# What is Math?

Koji Sunami

September 20, 2026

## Contents

|   |                                                       |   |
|---|-------------------------------------------------------|---|
| 1 | Approximation Method                                  | 1 |
| 2 | Structure and Symmetry                                | 2 |
| 3 | Encountering Conjecture                               | 3 |
| 4 | Derived Algebraic Geometry From Intuitive to Nonsense | 3 |
| 5 | Grumble about Myself                                  | 4 |

In this pdf, to discuss what math really is for expert/non-expert audiences, I'll set a grand framework of math research beginning from the elementary math to the research level. To do this, I'll introduce my point of view of numerical approximation, and symmetric structural view point, and finally where is a reasonable place to start math. The goal of this paper is recall for everybody including experts and beginners why math is needed, especially Perhaps, the purposeo of this exposition is my ego, or it works for the moderate beginners trying to decide their major in college, or those who have burned out from Alice's wonderland syndrome.

## 1 Approximation Method

One says all math is meaningless beyond of counting or lower division math, because it's already sufficient to study statistics or computational algorithm, and some even claim AI will automate calculation and conducting new theorem without human being. But math is science, that is an activity of extracting law of nature, that's happening by necessity including quantum state.

- Line and Triangle  
Line and Triangle might be necessary ingredient of understanding math no doubt by including people who are not really interested in pure math. To understand the structure of the natural world, perhap we could start from triangle (point, line, triangle, and n-tetrahedron), probably the most well-known shape, and finding invariants to discuss the relationship with the real world.
- It reminds me simplicial homology. A triangle consists of 3 vertex, 3 edges, and 1 face, and any 2-dimensional polygon has a subdivision to the union of triangles, which is triangulation in other words. It calculates topological invariants, in parcticular, Euler characteristics and orientability.
- Adding cartesian product to simplicial homology makes still simplicial complex. Especially an cartesian product with a line becomes a cylinder (precisely it happens in singular homology), whose crashing one side is a cone. This operation makes triangulated category.
- Or if the line  $[0, 1]$  is replaced by an affine line  $\mathbb{A}^1$ , which corresponds to  $k[x]$  algebraic geometrically, the cartesian product  $\mathbb{A}^n$  is fiber product  $\mathbb{A}^n \otimes_{\mathbb{A}^{n-1}} \mathbb{A}^n$ , algebraically  $k[x_1, \dots, x_n] \otimes_{k[z_1, \dots, z_{n-1}]} k[y_1, \dots, y_n]$ . This creates standard resolution.

Accordingly, any geometries seem to be approximated by some simplices. Does research stop here??

- Linear Algebra

Any  $n \times n$  matrices have a diagonalization by numerical approximation, by slightly deforming the coefficients so to have distinct roots of the characteristic polynomial. so for case of numerical approximation, Jordan form doesn't seem to be necessary.

- Calculus

One of the applications of calculus is numerical approximation using Newton's method to find the roots of polynomials and more generally any differentiable functions. There is an easy generalization to the multi-variate case, in particular any differentiable functions on complex planes.

Perhaps, imagining computer vision analysis, a collection of the finite distinct points on an Euclidean space is a connected component of the image by topological data analysis. There's many metric analysis. Does research stop here??

## 2 Structure and Symmetry

Anyways, is the study of math simply the study of finding the more efficient algorithms for the sake of computations? I personally believe The pure math has a different direction, and it is a process of finding a structure, while what computation means already assuming what the structure is.

What is "structure"? When we learn structure, we are not making the world easier, or if we believe it's easier, it's simply done by our knowledge. It's just a transition of perspective (if there is any terrible experience, is this a knowledge that we can forget? Knowledge is a data that CPU can machinery interpret. Our experience isn't data, since our brain isn't CPU). By outcome of transition of perspective, solving problem and making new theorem perhaps becomes more difficult, since apparently modern math is more difficult than math in ancient Greek. If the goal of math is to solve it, what we have done is quite ridiculous. In below, I'll try to explain some examples from my experience.

- Perhaps, the journey of pure math begins with linear algebra, initially solves a system of equations, representation theory  $\rho : G \rightarrow GL_n$  maps any math objects to represent in linear algebra. In particular, if an element can be mapped to the Jordan form, computing Jordan block is a combinatorial problem (module is called module because  $k[x]$ -module computes Jordan form, where Jordan block literally counts like Lego block?), while the question is to discuss the meaning of the operations. In particular, if there's any redundancy counting as Maschke's theorem or in some geometric interpretation using GIT quotient of  $GL_n$ , or its algebraic geometry enhancement is Hilbert scheme? Nilpotency of Jordan form in abstract algebra? This kind of activity is called pure math.
- Let's consider another example. A Galois theory proved unsolvability of quintic equation by unsolvability (non-existence of proper normal subgroup) of an alternate group  $A_5$ , meaning of the non-existence of the solution by roots of unity. Adding few more interpretations that enables to solve it, However. One thing is that p-adic henselization by putting the quintic equation over the field of finite characteristic p for some prime p, assuming all the coefficients are integers. Meanwhile, another solution of the quintic is done by stereographic projection of icosahedron and its rotation, whose symmetry is  $PSL_2(\mathbb{C})$ , which generates J-invariants, whose moduli space corresponds to the moduli space of coefficients of the quintic. In summary, these two solutions depends on an infinite order p-adic abelian group and a finite order nonabelian group, and perhaps these two notions might have some relationships. This is anabelian geometry or p-adic Hodge theory.
- Another problem is geometry, but the question is what really is. homology or cohomology might be the tools of computation but can merely calculate Euler characteristic and few invariants, and missing many data that the geometry originally has. The spontaneous question is if there's any inverse relationship? If we are doing analysis, the solution space of non-linear PDE generates another geometry, which in particular is Floer or Kuranishi structure, which is often more intricate than the original geometry, and its homology can evaluate something more than the homology of the original one. Or in algebraic geometry, moduli space of some algebraic variety contains more than itself, or the further generalization is category theory: these are not necessarily making something big, the important problem is algebraic variety and manifold, and stack, orbifold, infinity-groupoid or any other generalization is rather a substitute. Meanwhile, Toda's  $\chi$ -independence conjecture claims Gopakumar-Vafa invariant is determined independently from the Euler characteristic.

### 3 Encountering Conjecture

The moment I started doing research math is seeing the millenium problems of Clay Math Institute. By seeing its problem statement by downloading the pdf, I realized understanding the problem itself isn't super hard for me in undergraduate, even though its proof might be unknown to anybody other than Poincare conjecture. I'm explaining when I started math, in the same why I study math. Here I'll leave some comments on the seven problems with the level of difficulty of understanding the problem statements.

- P=NP Problem (difficulty: ★★★)  
idk
- Yang-Mill Mass Gap (difficulty: ★★★)  
Yang-Mill Mass Gap is originated from gauge theory in math physics. The mass gap means that the total mass of quarks is different from that of proton/neutron, and this energy appears in nuclear reaction, even though total amount of quarks doesn't change by the reaction! One points out there're some attempts by lattice gauge theory or OPE of CFT.
- Hodge Conjecture (difficulty: ★★)  
An algebraic variety can be described by homology/cohomology with Hodge decomposition, which isn't difficult, but its inverse problem is difficult.
- Poincare Conjecture (difficulty: ★★)  
This involves the notion of manifold, so the level of difficulty is higher than Navier-Stokes. This is a simple extension of Hamiltonian flow, but the notion of flow itself isn't very difficult. Consider that a manifold doesn't have a shape as a topological space, as long as the metric isn't defined. Now if the Riemannian metric changes time-dependent, the shape will change; this is the notion of flow. The question is that the flow could make some singularity in the geometry, and Perelman's technique is to resolve it differentially geometrically.
- Navier Stokes Equation (difficulty: ★)  
Navier Stokes Equation is simply a multi-variate calculus, so the difficulty is equivalent to Riemann Hypothesis. Ladyzhenskaya solved 2d NSE and 2.5d NSE using functional analysis, but 3d NSE hasn't been proved. If anticipating algebra instead of functional analysis, consider CFT lower-bound derives relativistic NSE, which makes ordinary NSE. Perhaps NSE might be quantization or deformation problem.
- Riemann Hypothesis (difficulty: ★)  
If Riemann Hypothesis might be a simple extension of prime number theorem, it is probably the easiest to access, if we can understand the basic calculus, assuming any 1-dimensional complex analysis is equivalent to 1-dimensional lower-division calculus, where we can see the zeta function and gamma function. However, if we have already known algebraic geometry, Weil conjecture proves the analogue of Riemann Hypothesis.

### 4 Derived Algebraic Geometry From Intuitive to Nonsense

#### Point of View of Functional Analysis

Let's begin with theory of function. The function (any functions including non-continuous) defined on  $\mathbb{R}^1$  might be very complicated, but functions defined on  $S^1$  are Fourier series. Accordingly, the structure of function can be easier, if the domain is simplified possibly done by geometric quotient  $\mathbb{R}/\mathbb{Z} = S^1$ , and so geometric quotient might connect to the theory of functions. More generally, there will be more interesting problems in higher dimensions e.g.  $\mathbb{R}^n$ . If we are interested in complex geometry, in particular, functions defined on torus  $\mathbb{T}^2$  is holomorphic functions satisfying equivariance by torus action, so called modular form.

In the language of geometry, the function is denoted by global section of the vector bundle, and sequence of global section of vector bundles formulates cohomology e.g. de Rham cohomology, and from here, 1st chern classes  $c_1(M)$  is curvature and 2nd chern classes  $c_2(M)$  is Lagrangian density to discuss its structure. Accordingly, geometry naturally induces cohomology to study itself, while its inverse is that whether cohomology contains all information needed to reconstruct the given original geometry. part of this inverse problem is called Hodge conjecture, or Torelli theorem in the special situation.

#### Why Polynomial? Shape of Geometry

In case we are selectively study algebraic side of geometry, the next question might be why polynomial is very important in geometry. To answer this question, let's consider the whole possible Riemann surface  $\mathcal{M}$

to classify themselves by genus and marked points  $\mathcal{M}_{g,n}$ , and each  $\mathcal{M}_{g,n}$  is nontrivial. As a result, if you are interested in any geometry of fixed  $g, n$ , then polynomial geometry is enough.

The next step is the shape of geometry beyond of  $g, n$ . Naïvely there would be many weird shapes such as urchin shape, fractal, or potato shape, drawing randomly on the coordinate, but the process of understanding geometry is to find some meaning from it. Math is roughly an activity of finding invariants and making a comparison, and clarifying what is input/output in the theory, and the topologists for instance cannot distinguish coffee cup and donut, and geometers are the same. It might be Symplectic/complex/holomy structure, or perhaps hyperkähler structure.

Other than algebraic variety, quotient of algebraic variety could be no longer algebraic variety, but it is interesting. Recalling  $S^1$  defining Fourier analysis, and geometric quotient  $X/G$  deepens the structure of the original  $X$ , while many complicated shapes can be generated by quotient. For instance quotient of torus by Klein 4-group  $\mathbb{Z}/2 \times \mathbb{Z}/2$  is either Klein bottle or real projective space  $\mathbb{RP}^2$  depending on its group action.

I'd also consider the meaning of polynomial based on its connection to CFT. According to Marc Gross, polynomial in Landau-Ginzburg model defines period integral  $\int_X e^{-f(x)}$ , where  $f(x)$  is polynomial, which induces SCFT, which induces semi-infinite variation of Hodge structure, thus mirror symmetry.

## AG vs DG

Comparing derived algebraic geometry and classical algebraic geometry and even differential geometry, all of them are interesting for different reasons. In fact, algebraic geometry can be studied differentially. Hodge structure, Symplectic structure, Lie algebra, moduli problem, sheaf cohomology, singularity, derived method, and even counting problem are still domain of differential geometry, and this is for no surprise, and in fact there's tons of analysis in Fukaya category, and algebraic/differential geometry seem to be just different tools for the same problems. While to briefly list up, algebraic geometry can better treat deformation, degeneration, singularity, intersection, geometric quotient, or argument over finite characteristics etc. Derived algebraic geometry captures the information missing in classical algebraic geometry.

By the way, geometry is typically computed by homological language, and simplicial/singular homology can calculate topological data, and typically cohomology contains more information about the geometry than homology does, and cohomology is a sequence of function spaces over the geometry instead of its geometry itself. Furthermore, deformation is a generalization of tangent space, so we can make generalization of cohomology.

## 5 Grumble about Myself

### My Research Style

Math needs to be rigorous no doubt, but there is no understanding without intuition, but to understand something, it's not just about amount of knowledge, but we might need to start the new argument, and for the journey of math research, I hope it will not be chaotic system, where one mistake nullify the whole previous efforts.

Or for non-research level, the important thing is not necessarily the accuracy but to accomplish the argument. In a sentence, quantity precedes quality in the learning process so as to cultivate intellectual stamina, so we can attempt to modify the problem rather than ability to modify itself. It's similar to football, the most important thing is to run & dribble & pass not the accuracy of dribble or pass.

### My View of AI

In 2026, I personally think AI is just little more advanced model than google search with natural language interpretation. If using google search itself is already a moderate labor, I use AI to reduce the amount.