

A_∞ Enhancement of Geometry

Koji Sunami

September 21, 2026

The goal of our research is to prove homological mirror symmetry conjecture or to develop the relevant area of study. The claim of homological mirror symmetry conjecture is A_∞ -equivalence of Fukaya category $Fuk(X)$ and A_∞ -enhancement $D^b(Coh(X))$ (typically noted as $D^b(X)$) of derived category of coherent sheaves. This statement says the elaboration of mirror symmetry, originally correspondence of symplectic data on Hodge structure and that of complex in Calabi-Yau manifold, whose symplectic side we name as A-model, the complex side as B-model, mirror symmetry solves many of QFT problem as well as Landau-Ginzburg model and others, and homological mirror symmetry is generalization of mirror symmetry in ∞ -category. The proof isn't easy, but the understanding of the problem statement itself is not difficult.

Definition 0.1. A_∞ Category

An A_∞ category $\mathcal{A}$ is a quasi-category which consists of objects $X \in Ob(\mathcal{A})$ and 1-morphisms $f \in Mor^1(\mathcal{A})$, and the 1-morphism contains higher coherence $\alpha : f \Rightarrow g$, called 2-morphism, and iteratively n -morphisms.

Definition 0.2. A_∞ Functor

Definition 0.3. Fukaya Category

For a symplectic manifold X , Fukaya category $Fuk(X)$ is an A_∞ category describing Lagrangian intersection. The objects of Fukaya category $Ob(Fuk(X))$ is Lagrangian submanifolds $L \subset X$, and the morphisms are Floer chain complex $CF^*(L_0, L_1)$ for each L_0 and L_1 . This $CF^*(L_0, L_1)$ is a $\mathbb{Z}$ -graded vector space defined by $CF^*(L_0, L_1) = \bigoplus_{p \in L_0 \cap L_1} \mathbb{k}p$. Floer chain complex is generated by its intersection points, and this is equivalent with counting critical points of Morse theory.

In Fukaya category, there is a composition $m_k : CF^*(L_{k-1}, L_k) \otimes \cdots \otimes CF^*(L_1, L_2) \rightarrow CF^*(L_0, L_k)[2-k]$ satisfying Stasheff associahedron property.

A_∞ enhancement of dg-category doesn't have unique enhancement, but the uniqueness is guaranteed if dg-category is derived category $D^b(X)$.

Definition 0.4. A_∞ Derived Category

A_∞ enhancement of derived category $D^b(X)$ is an A_∞ category $\mathcal{A}$, whose homology is $H^*(\mathcal{A}(E, F)) = Ext^*(E, F)$, while its trivialization $H^0(\mathcal{A}(E, F))$ needs to become the based derived category $H^0(\mathcal{A}(E, F)) \cong Hom_{D^b(X)}(E, F)$. This A_∞ enhancement $\mathcal{A}$ can be denoted as $D^b(X)$ by abuse of notation.

Conjecture 0.1. Homological Mirror Symmetry

There is an A_∞ equivalence of categories

$$Fuk(X) \cong D^b(X)$$

What is the structure of A_∞ category? The natural question is geometric realization of A_∞ category.

Starting from A_∞ algebra as a special case of A_∞ category,
and this can be formal scheme.

One idea is identifying A_∞ algebra to L_∞ algebra through Hochschild cohomology by Kontsevich-Soibelman, and making its generalization to A_∞ category.

L_∞ algebra is a deformation of dgla, which is an analogue of Lie algebra for homological algebra (consider Lie algebra consists of vector field, dgla consists of endomorphism of chain complex $d : V \rightarrow V$ satisfying $d^2 = 0$).

<ncfp dg-manifold>

If there is any connection to the classical or derived algebraic geometry will help.

A_∞ algebra is a generalization of commutative ring. A_∞ category is its categorical generalization, especially if A_∞ category with one object is A_∞ algebra. I'll use A_∞ algebra to describe algebraic geometry. Recall a commutative ring defines an affine scheme by its ring spectrum $X = V(\mathfrak{p})$ with its structure sheaf $\mathcal{O}_X$, and that can also be described functorially as $F : CRing \rightarrow Set$. Hereafter I focus on functorial point of view.

The proof of homological mirror symmetry conjecture is discussed from multi-dimensional perspective, including counting Gromow-Witten invariant, or Balmer spectrum, while generalizing formal geometry of non-unital A_∞ -algebra to A_∞ -category.

Definition 0.5

Contents

1 2

2 Geometric Realization of A_∞ 2

1

2 Geometric Realization of A_∞

Definition 2.1. Counital Coalgebra

Coalgebra B is a graded vector space with morphism $\Delta : B \rightarrow B \otimes B$ s.t. $(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta$ called coproduct, and counital coalgebra is coalgebra B with unit $\epsilon : B \rightarrow 1$ where 1 is the unit object of $\mathcal{C}$, satisfying $(\epsilon \otimes id) \circ \Delta = (id \otimes \epsilon) \circ \Delta$. Counital coalgebra B is cofree counital coalgebra if it's given by $B = C(V) \cong T^c(V)$.

Definition 2.2. A Non-Commutative Thin Scheme

A non-commutative thin scheme X is a functor $X : Alg_{\mathcal{C}^f} \rightarrow Sets$, that is isomorphic to $X \cong Hom(-^*, B_X)$ for some coalgebra B_X , where $\mathcal{C}^f$ is artinian category of finite dimensional objects in $\mathcal{C}$, where $\mathcal{C}$ is $\mathbb{Z}$ -graded vector spaces. In particular, X is a non-commutative formal manifold, if the functor X is isomorphic to some $Spc(T(V))$ for a cofree counital tensor coalgebra $T(V) = \bigoplus_{n \geq 1} V^{\otimes n}$. The dimension of X is defined as $dim_k V$.

Definition 2.3. A Non-Commutative Thin DG-Scheme

A non-commutative thin dg-scheme is a pair (X, d) where X is a non-commutative thin scheme, and d is a vector field on X of degree +1 satisfying bracket of dgla as $[d, d] = 0$. This d is called homological vector field.

Definition 2.4. Non-Commutative Formal Pointed DG-Manifold

A non-commutative formal pointed dg-manifold is a pair $((X, x_0), d_X)$ such that (X, x_0) is a non-commutative formal pointed graded manifold, and d_X is a homological vector field on X such that $d_X|_{x_0} = 0$.

Proposition 2.1. Taylor Decomposition of Graded Vector Field

For the case of formal manifold X , let the corresponding cofree counital coalgebra be $B_X \cong T^c(V)$, and the homological vector field be coderivation $d_X : B_X \rightarrow B_X$, that can have restriction by cofreeness, thus by using projection, $\pi \circ d_X : B_X \rightarrow V[1]$. If in addition, d_X can be decomposed degreewise for its linearity, as $d_X = \sum_{n \geq 1} d_X^{(n)}$ for linear maps $d_X^{(n)} : V^{\otimes n} \rightarrow V[1]$.

Definition 2.5. Non-Unital A_∞ Algebra

A non-unital A_∞ -algebra $A = (A, m_n)$ is a graded vector space A and coherence condition $m_n : A^{\otimes n} \rightarrow A[2 - n]$ with Stasheff relation, whose geometric interpretation is the structure of a nc fp dg-manifold $((X, x_0), d_X)$ together with a coalgebra isomorphism $B_X \cong T(T_{x_0}X)$ with $x_0 : k \rightarrow B_X$.

Definition 2.6. Structure of A_∞ Algebra

The structure of A_∞ Algebra on $V \in Ob(Vect_k^{\mathbb{Z}})$ is given by coderivation d of degree $+1$ of the non-counital cofree coalgebra $T_+(V[1]) = \bigoplus_{n \geq 1} V^{\otimes n}$ s.t. $[d, d] = 0$.

Proposition 2.2. Stasheff Relation

The condition of nc fp dg-manifold $((X, x_0), d_X)$ derives Stasheff relation of the A_∞ algebra, that is $m_n := d_X^{(n)}$ for $m_n : A^{\otimes n} \rightarrow A$ s.t. $\sum_{r+s+t=n} (-1)^{r+st} m_{r+1+t}(id^{\otimes r} \otimes m_s \otimes id^{\otimes t}) = 0$, whose degree is $|m_n| = 2 - n$. The Taylor decomposition of homological vector field $\pi \circ d_X = \sum_{n \geq 1} d_X^{(n)}$ is degreewise by linearity, as $d_X^{(n)} : B_X^{(n)} \rightarrow B_X^{(1)}$.

Definition 2.7. Non-Commutative Formal n-Pointed DG-Manifold

Nc formal n -pointed dg-manifold $(\mathcal{X}, d_{\mathcal{X}}, \{pt_i\})$ is nc formal graded dg-manifold $(\mathcal{X}, d_{\mathcal{X}})$, and the points are $pt_i \in \mathcal{X}(k)$ for $1 \leq i \leq n$, and coderivation satisfies $d_{\mathcal{X}}|_{pt_i} = 0$, $d_{\mathcal{X}}|_{TX_i} : TX_i \rightarrow \bigoplus_{j,k} TX_j \oplus TX_k$ where X_i is formal neighborhood of X at pt_i .

Definition 2.8. A_∞ -Category

A_∞ -category $\mathcal{A}$ is nc formal dg-manifold $(\mathcal{X}, d_{\mathcal{X}}, \{pt_i\})$ with its coalgebra $B_{\mathcal{X}} \cong T^c(V[1])$. This V has a decomposition $V = \bigoplus_{i,j} E_{ij}$, where $E_{ij} := T(\widehat{X_i \times X_{j_{pt_i, pt_j}}})$.

Remark 2.1. T

is V is a k -linear associative algebra generated from k -linear category $\mathcal{C}$: let $Ob(\mathcal{C}) = I$ to define an algebra $V = \bigoplus Hom(i, j)$ for $i, j \in I$ is a k -linear algebra with the unit $1_V = \sum_{i \in I} id_i$, where $\pi_i = id_i \in V$ is a commuting idempotent i.e. $\pi_i^2 = \pi_i$, $\pi_i \pi_j = \pi_j \pi_i$. This makes homomorphism $k^I \rightarrow V$ of algebra with unit.

Proposition 2.3. Stasheff Relation

From non-unital A_∞ category $\mathcal{A}$, the coalgebra $d_\mathcal{X} : T^c(\oplus E_{ij}[1]) \rightarrow T^c(\oplus E_{ij}[1])$ satisfies the Stasheff relation. Consider the tensor coalgebra can be given by the grading structure $T^c(\oplus_{1 \leq i, j \leq n} E_{ij}[1]) = \oplus_{m \geq 0} \oplus_{1 \leq i_0, i_1, i_2, \dots, i_m \leq n} E_{i_0 i_1} \otimes E_{i_1 i_2} \otimes \dots \otimes E_{i_{m-1} i_m} := \oplus_{m \geq 0} B_\mathcal{X}^m$. From cofreeness of the coalgebra, the coderivation $d_\mathcal{X}$ canonically projected onto $\pi : B_\mathcal{X} \rightarrow B_\mathcal{X}^1$, and that can be decomposed linearwise, so $\pi \circ d_\mathcal{X} = \Sigma d_\mathcal{X}^n$ where $d_\mathcal{X}^n : B_\mathcal{X}^n \rightarrow B_\mathcal{X}^1$, and we have Stasheff relation from $[d_\mathcal{X}, d_\mathcal{X}] = 0$, and $m_n := d_\mathcal{X}^n$.

Remark 2.2

$d_\mathcal{X} \circ \eta = 0$ implies $d_\mathcal{X} \circ \eta(1) = 0$ or $d_\mathcal{X}(1) = 0$. This means curvature is $m_0 = 0$.

Remark 2.3

For the case of $I = \{i\}$ consists of one vertex, we write $d|_{pt} = d \circ \eta$.

For the case of infinite objects A_∞ category is its inductive limit of finite objects.

References

- [1] M. Doubek, M. Markl, and P. Zima. *Deformation Theory (lecture notes)*. 2009. arXiv: 0705.3719 [math.AG]. URL: <https://arxiv.org/abs/0705.3719>.
- [2] Fabian Haiden et al. *Towards Categorical Kähler Geometry*. 2026. arXiv: 2609.00978 [math.AG]. URL: <https://arxiv.org/abs/2609.00978>.
- [3] Robin Hartshorne. *Algebraic Geometry*. Springer, 1977.
- [4] Amanda Hirschi and Kai Hugtenburg. *Open-closed Deligne-Mumford field theories: construction*. 2026. arXiv: 2605.03521 [math.SG]. URL: <https://arxiv.org/abs/2605.03521>.
- [5] Joseph Hirsh and Joan Millès. *Curved Koszul duality theory*. 2010. arXiv: 1008.5368 [math.KT]. URL: <https://arxiv.org/abs/1008.5368>.
- [6] Isamu Iwanari. *Categorified Koszul duality of algebras*. 2026. arXiv: 2512.12263 [math.AG]. URL: <https://arxiv.org/abs/2512.12263>.
- [7] Dominic Joyce. *Kuranishi homology and Kuranishi cohomology: a User's Guide*. 2008. arXiv: 0710.5634 [math.SG]. URL: <https://arxiv.org/abs/0710.5634>.
- [8] Bernhard Keller. *On differential graded categories*. 2006. arXiv: math/0601185 [math.KT]. URL: <https://arxiv.org/abs/math/0601185>.
- [9] Maxim Kontsevich and Yan Soibelman. *Notes on A-infinity algebras, A-infinity categories and non-commutative geometry. I*. 2024. arXiv: math/0606241 [math.RA]. URL: <https://arxiv.org/abs/math/0606241>.
- [10] Tatsuki Kuwagaki. *A remark on monoidal structure and homological mirror symmetry*. 2026. arXiv: 2603.08384 [math.SG]. URL: <https://arxiv.org/abs/2603.08384>.
- [11] Siu-Cheong Lau, Junzheng Nan, and Ju Tan. *Mirror Symmetry for Quiver Algebroid Stacks*. 2025. arXiv: 2206.03028 [math.AG]. URL: <https://arxiv.org/abs/2206.03028>.
- [12] B. Toen. *The homotopy theory of dg-categories and derived Morita theory*. 2006. arXiv: math/0408337 [math.AG]. URL: <https://arxiv.org/abs/math/0408337>.
- [13] Bertrand Toën and Gabriele Vezzosi. "Homotopical Algebraic Geometry II: Geometric Stacks and Applications". In: *Memoirs of the American Mathematical Society* 193.902 (2008).