

What is Math?

Koji Sunami

September 12, 2026

In this pdf, I'll discuss numerical approximation view point of math, and symmetric structural view point of math, and finally I'll introduce where is a reasonable place to start math. The goal of paper is recall for everybody including experts and beginners why math is needed, especially those who have burned out from Alice's wonderland syndrome.

1 Approximation

One says all math is meaningless because numerical approximation. But math is the same as science, it is an activity of extracting natural law.

Line and Triangle

There's probably many people, who are wondering what math really is. It's probably an activity of extracting new structure from the natural world. To understand the structure of the natural world, perhaps we could start from triangle (point, line, triangle, and n-tetrahedron), probably the most well-known shape, and finding invariants to discuss the relationship with the real world.

One thing is simplicial homology. A triangle consists of 3 vertex, 3 edges, and 1 face, and any 2-dimensional polygon has a subdivision to the union of triangles, which is triangulation in other words. It calculates topological invariants, in particular, Euler characteristics and orientability.

Adding cartesian product to simplicial homology makes still simplicial complex. Especially an cartesian product with a line becomes a cylinder (precisely it happens in singular homology), whose crashing one side is a cone. This operation makes triangulated category.

Or if the line $[0, 1]$ is replaced by an affine line $\mathbb{A}^1$, which corresponds to $k[x]$ algebraic geometrically, the cartesian product $\mathbb{A}^n$ is fiber product $\mathbb{A}^n \otimes_{\mathbb{A}^{n-1}} \mathbb{A}^n$, algebraically $k[x]^n \otimes_{k[x]^{n-1}} k[x]^n$. This creates standard resolution.

Any geometries seem to be approximated by some simplices. Does research stop here??

Perhaps, imagining computer vision analysis, a collection of the finite distinct points on an Euclidean space is a connected component of the image by

topological data analysis.

Does research stop here??

Linear Algebra

Any $n \times n$ matrices have a diagonalization by numerical approximation, by slightly deforming the coefficients so to have distinct roots of the characteristic polynomial. so for case of numerical approximation, Jordan form doesn't seem to be necessary.

Calculus

One of the applications of calculus is numerical approximation using Newton's method to find the roots of polynomials and more generally any differentiable functions. There is an easy generalization to the multi-variate case, in particular any differentiable functions on complex planes.

2 Structure and Symmetry

Anyways, is the study of math simply the study of finding the more efficient algorithms for the sake of computations? I personally believe The pure math has a different direction, and it is a process of finding a structure, while what computation means already assuming what the structure is.

What is "structure"? When we learn structure, we are not making the world easier, or if we believe it's easier, it's simply done by our knowledge. It's just a transition of perspective (if there is any terrible experience, is this a knowledge that we can forget? Knowledge is a data that CPU can machinery interpret. Our experience isn't data, since our brain isn't CPU). By outcome of transition of perspective, solving problem and making new theorem perhaps becomes more difficult, since apparently modern math is more difficult than math in ancient Greek. If the goal of math is to solve it, what we have done is quite ridiculous. In below, I'll try to explain some examples from my experience.

Perhaps, the journey of pure math begins with linear algebra, initially solves a system of equations, representation theory $\rho : G \rightarrow GL_n$ maps any math objects to represent in linear algebra. In particular, if an element can be mapped to the Jordan form, computing Jordan block is a combinatorial problem (module is called module because $k[x]$ -module computes Jordan form, where Jordan block literally counts like Lego block?), while the question is to discuss the meaning of the operations. In particular, if there's any redundancy counting as Maschke's theorem or in some geometric interpretation using GIT quotient of GL_n , or its algebraic geometry enhancement is Hilbert scheme? Nilpotency of Jordan form in abstract algebra? This kind of activity is called pure math.

Let's consider another example. A Galois theory proved unsolvability of quintic equation by unsolvability (non-existence of proper normal subgroup) of an alternate group A5, meaning of the non-existence of the solution by roots of unity. Adding few more interpretations that enables to solve it, However.

One thing is that p-adic henselization by putting the quintic equation over the field of finite characteristic p for some prime p , assuming all the coefficients are integers. Meanwhile, another solution of the quintic is done by stereographic projection of icosahedron and its rotation, whose symmetry is $PSL_2(\mathbb{C})$, which generates J-invariants, whose moduli space corresponds to the moduli space of coefficients of the quintic. In summary, these two solutions depends on an infinite order p-adic abelian group and a finite order nonabelian group, and perhaps these two notions might have some relationships. This is anabelian geometry or p-adic Hodge theory.

Another problem is geometry, but the question is what really is. homology or cohomology might be the tools of computation but can merely calculate Euler characteristic and few invariants, and missing many data that the geometry originally has. The spontaneous question is if there's any inverse relationship? If we are doing analysis, the solution space of non-linear PDE generates another geometry, which in particular is Floer or Kuranishi structure, which is often more intricated than the original geometry, and its homology can evaluate something more than the homology of the original one. Or in algebraic geometry, moduli space of some algebraic variety contains more than itself, or the further generalization is category theory: these are not necessarily making something big, the important problem is algebraic variety and manifold, and stack, orbifold, infinity-groupoid or any other generalization is rather a substitute.

3 Millennium Problems

I started doing math from the millenium problems of Clay Math Institute in undergraduate. By seeing its problem statement by downloading the pdf, I realized understanding the problem itself isn't super hard, though its proof might be unknown to anybody. I set the level of difficulty of understanding the problem statements based on my intuition.

- P=NP Problem (difficulty: ★★★)
idk
- Yang-Mill Mass Gap (difficulty: ★★★)
Yang-Mill Mass Gap is originated from gauge theory in math physics. The mass gap means that the total mass of quarks is different from that of proton/neutron, and this energy appears in nuclear reaction, even though total amount of quarks doesn't change by the reaction! One points out there're some attempts by lattice gauge theory or OPE of CFT.
- Hodge Conjecture (difficulty: ★★)
An algebraic variety can be described by homology/cohomology with Hodge decomposition, which isn't difficult, but its inverse problem is difficult.
- Poincare Conjecture (difficulty: ★★)
This involves the notion of manifold, so the level of difficulty is higher than

Navier-Stokes. This is a simple extension of Hamiltonian flow, but the notion of flow itself isn't very difficult. Consider that a manifold doesn't have a shape as a topological space, as long as the metric isn't defined. Now if the Riemannian metric changes time-dependently, the shape will change; this is the notion of flow. The question is that the flow could make some singularity in the geometry, and Perelman's technique is to resolve it differentially geometrically.

- Navier Stokes Equation (difficulty: ★)
Navier Stokes Equation is simply a multi-variate calculus, so the difficulty is equivalent to Riemann Hypothesis. Ladychenskaya solved 2d NSE and 2.5d NSE using functional analysis, but 3d NSE hasn't been proved. If anticipating algebra instead of functional analysis, consider CFT lower-bound derives relativistic NSE, which makes ordinary NSE. Perhaps NSE might be quantization or deformation problem.
- Riemann Hypothesis (difficulty: ★)
If Riemann Hypothesis might be a simple extension of prime number theorem, it is probably the easiest to access, if we can understand the basic calculus, assuming any 1-dimensional complex analysis is equivalent to 1-dimensional lower-division calculus, where we can see the zeta function and gamma function. However, if we have already known algebraic geometry, Weil conjecture proves the analogue of Riemann Hypothesis.