

A_∞ Enhancement of Geometry

Koji Sunami

September 14, 2026

The proof of homological mirror symmetry conjecture is discussed from multi-dimensional perspective, including counting Gromow-Witten invariant, or Balmer spectrum, while generalizing formal geometry of non-unital A_∞-algebra to A_∞-category.

Definition 0.0.1 (*Counital Coalgebra*)

Coalgebra B is a graded vector space with morphism $\Delta : B \rightarrow B \otimes B$ s.t. $(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta$ called coproduct, and counital coalgebra is coalgebra B with unit $\epsilon : B \rightarrow 1$ where 1 is the unit object of $\mathcal{C}$, satisfying $(\epsilon \otimes id) \circ \Delta = (id \otimes \epsilon) \circ \Delta$. Counital coalgebra B is cofree counital coalgebra if it's given by $B = C(V) \cong T^c(V)$.

Definition 0.0.2 (*A Non-Commutative Thin Scheme*)

A non-commutative thin scheme X is a functor $X : Alg_{\mathcal{C}^f} \rightarrow Sets$, that is isomorphic to $X \cong Hom(-^*, B_X)$ for some coalgebra B_X , where $\mathcal{C}^f$ is artinian category of finite dimensional objects in $\mathcal{C}$, where $\mathcal{C}$ is $\mathbb{Z}$ -graded vector spaces. In particular, X is a non-commutative formal manifold, if the functor X is isomorphic to some $Spc(T(V))$ for a cofree counital tensor coalgebra $T(V) = \bigoplus_{n \geq 1} V^{\otimes n}$. The dimension of X is defined as $\dim_k V$.

Definition 0.0.3 (*A Non-Commutative Thin DG-Scheme*)

A non-commutative thin dg-scheme is a pair (X, d) where X is a non-commutative thin scheme, and d is a vector field on X of degree $+1$ satisfying bracket of dgla as $[d, d] = 0$. This d is called homological vector field.

Definition 0.0.4 (*Non-Commutative Formal Pointed DG-Manifold*)

A non-commutative formal pointed dg-manifold is a pair $((X, x_0), d_X)$ such that (X, x_0) is a non-commutative formal pointed graded manifold, and d_X is a homological vector field on X such that $d_X|_{x_0} = 0$.

Proposition 0.1 (*Taylor Decomposition of Graded Vector Field*)

For the case of formal manifold X , let the corresponding cofree counital coalgebra be $B_X \cong T^c(V)$, and the homological vector field be coderivation $d_X : B_X \rightarrow B_X$, that can have restriction by cofreeness, thus by using projection, $\pi \circ d_X : B_X \rightarrow V[1]$. If in addition, d_X can be decomposed degreewise for its linearity, as $d_X = \sum_{n \geq 1} d_X^{(n)}$ for linear maps $d_X^{(n)} : V^{\otimes n} \rightarrow V[1]$.

Definition 0.1.1 (Non-Unital A_∞ Algebra)

A non-unital A_∞ -algebra $A = (A, m_n)$ is a graded vector space A and coherence condition $m_n : A^{\otimes n} \rightarrow A[2-n]$ with Stasheff relation, whose geometric interpretation is the structure of a nc fp dg-manifold $((X, x_0), d_X)$ together with a coalgebra isomorphism $B_X \cong T(T_{x_0}X)$ with $x_0 : k \rightarrow B_X$.

Definition 0.1.2 (Structure of A_∞ Algebra)

The structure of A_∞ Algebra on $V \in \text{Ob}(\text{Vect}_k^{\mathbb{Z}})$ is given by coderivation d of degree $+1$ of the non-counital cofree coalgebra $T_+(V[1]) = \bigoplus_{n \geq 1} V^{\otimes n}$ s.t. $[d, d] = 0$.

Proposition 0.2 (Stasheff Relation)

The condition of nc fp dg-manifold $((X, x_0), d_X)$ derives Stasheff relation of the A_∞ algebra, that is $m_n := d_X^{(n)}$ for $m_n : A^{\otimes n} \rightarrow A$ s.t. $\sum_{r+s+t=n} (-1)^{r+st} m_{r+1+t}(id^{\otimes r} \otimes m_s \otimes id^{\otimes t}) = 0$, whose degree is $|m_n| = 2 - n$. The Taylor decomposition of homological vector field $\pi \circ d_X = \sum_{n \geq 1} d_X^{(n)}$ is degreewise by linearity, as $d_X^{(n)} : B_X^{(n)} \rightarrow B_X^{(1)}$.

Definition 0.2.1 (Non-Commutative Formal n -Pointed DG-Manifold)

Nc formal n -pointed dg-manifold $(\mathcal{X}, d_{\mathcal{X}}, \{pt_i\})$ is nc formal graded dg-manifold $(\mathcal{X}, d_{\mathcal{X}})$, and the points are $pt_i \in \mathcal{X}(k)$ for $1 \leq i \leq n$, and coderivation satisfies $d_{\mathcal{X}}|_{pt_i} = 0$, $d_{\mathcal{X}}|_{TX_i} : TX_i \rightarrow \bigoplus_{j,k} TX_j \oplus TX_k$ where X_i is formal neighborhood of X at pt_i .

Definition 0.2.2 (A_∞ -Category)

A_∞ -category $\mathcal{A}$ is nc formal dg-manifold $(\mathcal{X}, d_{\mathcal{X}}, \{pt_i\})$ with its coalgebra $B_{\mathcal{X}} \cong T^c(V[1])$. This V has a decomposition $V = \bigoplus_{i,j} E_{ij}$, where $E_{ij} := T(\widehat{X_i \times X_{j, pt_j}})$.

Remark 0.3 () This V is a k -linear associative algebra generated from k -linear category $\mathcal{C}$: let $\text{Ob}(\mathcal{C}) = I$ to define an algebra $V = \bigoplus \text{Hom}(i, j)$ for $i, j \in I$ is a k -linear algebra with the unit $1_V = \sum_{i \in I} id_i$, where $\pi_i = id_i \in V$ is a commuting idempotent i.e. $\pi_i^2 = \pi_i$, $\pi_i \pi_j = \pi_j \pi_i$. This makes homomorphism $k^I \rightarrow V$ of algebra with unit.

Proposition 0.4 (Stasheff Relation)

From non-unital A_∞ category $\mathcal{A}$, the coalgebra $d_{\mathcal{X}} : T^c(\bigoplus E_{ij}[1]) \rightarrow T^c(\bigoplus E_{ij}[1])$ satisfies the Stasheff relation. Consider the tensor coalgebra can be given by the grading structure $T^c(\bigoplus_{1 \leq i, j \leq n} E_{ij}[1]) = \bigoplus_{m \geq 0} \bigoplus_{1 \leq i_0, i_1, i_2, \dots, i_m \leq n} E_{i_0 i_1} \otimes E_{i_1 i_2} \otimes \dots \otimes E_{i_{m-1} i_m} := \bigoplus_{m \geq 0} B_{\mathcal{X}}^m$. From cofreeness of the coalgebra, the coderivation $d_{\mathcal{X}}$ canonically projected onto $\pi : B_{\mathcal{X}} \rightarrow B_{\mathcal{X}}^1$, and that can be decomposed linearwise, so $\pi \circ d_{\mathcal{X}} = \sum d_{\mathcal{X}}^n$ where $d_{\mathcal{X}}^n : B_{\mathcal{X}}^n \rightarrow B_{\mathcal{X}}^1$, and we have Stasheff relation from $[d_{\mathcal{X}}, d_{\mathcal{X}}] = 0$, and $m_n := d_{\mathcal{X}}^n$.

Remark 0.5 () $d_{\mathcal{X}} \circ \eta = 0$ implies $d_{\mathcal{X}} \circ \eta(1) = 0$ or $d_{\mathcal{X}}(1) = 0$. This means curvature is $m_0 = 0$.

Remark 0.6 () For the case of $I = \{i\}$ consists of one vertex, we write $d|_{pt} = d \circ \eta$.

For the case of infinite objects A_{∞} category is its inductive limit of finite objects.

References

- [1] M. Doubek, M. Markl, and P. Zima. *Deformation Theory (lecture notes)*. 2009. arXiv: 0705.3719 [math.AG]. URL: <https://arxiv.org/abs/0705.3719>.
- [2] Fabian Haiden et al. *Towards Categorical Kähler Geometry*. 2026. arXiv: 2609.00978 [math.AG]. URL: <https://arxiv.org/abs/2609.00978>.
- [3] Robin Hartshorne. *Algebraic Geometry*. Springer, 1977.
- [4] Amanda Hirschi and Kai Hugtenburg. *Open-closed Deligne-Mumford field theories: construction*. 2026. arXiv: 2605.03521 [math.SG]. URL: <https://arxiv.org/abs/2605.03521>.
- [5] Joseph Hirsh and Joan Millès. *Curved Koszul duality theory*. 2010. arXiv: 1008.5368 [math.KT]. URL: <https://arxiv.org/abs/1008.5368>.
- [6] Isamu Iwanari. *Categorified Koszul duality of algebras*. 2026. arXiv: 2512.12263 [math.AG]. URL: <https://arxiv.org/abs/2512.12263>.
- [7] Dominic Joyce. *Kuranishi homology and Kuranishi cohomology: a User's Guide*. 2008. arXiv: 0710.5634 [math.SG]. URL: <https://arxiv.org/abs/0710.5634>.
- [8] Bernhard Keller. *On differential graded categories*. 2006. arXiv: math/0601185 [math.KT]. URL: <https://arxiv.org/abs/math/0601185>.
- [9] Maxim Kontsevich and Yan Soibelman. *Notes on A-infinity algebras, A-infinity categories and non-commutative geometry. I*. 2024. arXiv: math/0606241 [math.RA]. URL: <https://arxiv.org/abs/math/0606241>.
- [10] Tatsuki Kuwagaki. *A remark on monoidal structure and homological mirror symmetry*. 2026. arXiv: 2603.08384 [math.SG]. URL: <https://arxiv.org/abs/2603.08384>.
- [11] B. Toën. *The homotopy theory of dg-categories and derived Morita theory*. 2006. arXiv: math/0408337 [math.AG]. URL: <https://arxiv.org/abs/math/0408337>.
- [12] Bertrand Toën and Gabriele Vezzosi. “Homotopical Algebraic Geometry II: Geometric Stacks and Applications”. In: *Memoirs of the American Mathematical Society* 193.902 (2008).