

Distality in ordered abelian groups

Koki Okura

University of Tsukuba
Doctoral Program in Mathematics

August 26th, 2026


Model Theory Summer School
and Workshop 2026
at University of Tsukuba

Slides available at Koki Okura's Home Page→




Summary

Distality is the opposite of stability (among NIP theories).



Though all ordered abelian groups (OAGs) have NIP [GS84], only a distality criterion for OAGs with finite spines was known [Asc+22].



We have

- given a distality criterion for *all* OAGs.
- shown all OAGs have distal expansions.


Definition ([Sim13])

A theory is *distal* if for any sequence $(\bar{a}_i)_{i \in \mathbb{Q}}$ and any tuple $\bar{b}$, if

- $(\bar{a}_i)_{i \in \mathbb{Q}}$ is indiscernible and
- $(\bar{a}_i)_{i \in \mathbb{Q} \setminus \{0\}}$ is indiscernible over $\bar{b}$,

then $(\bar{a}_i)_{i \in \mathbb{Q}}$ is indiscernible over $\bar{b}$.

In DLO...



Roughly speaking, distal theories are those NIP theories which do not have a stable part.

Example (Distal theories)

- DLO
- $(\mathbb{R}; +, \cdot), (\mathbb{Q}_p; +, \cdot)$
- $(\mathbb{Z}; <, +)$


Example (NIP non-distal theories)

- Stable theories (with infinite models)
- $(\mathbb{Q}_{>0}; <, \cdot)$ ($|\mathbb{Q}_{>0}/(\mathbb{Q}_{>0})^p| = \infty$ for any prime p)
- $(\mathbb{Q}_p^{\text{alg}}; +, \cdot, v_p)$

Example (Theories without a distal expansion)


- Theories interpreting an infinite field of characteristic $p > 0$ ([CS18, Corollary 6.3]).

A valued field $(\mathbb{Q}_p^{\text{alg}}; +, \cdot, v_p)$ is a combination of its value group $\mathbb{Q}$ and residue field $\mathbb{F}_p^{\text{alg}}$.



Why distality? —Zarankiewicz's problem

Fix $k \geq 2$. Let $(V_1, V_2; E)$ be a $K_{k,k}$ -free bipartite graph and $n = \max\{|V_1|, |V_2|\}$. It is well-known ([KST54]) that



$$|E| = O(n^{2-(1/k)}).$$

If V_1 is points in $\mathbb{R}^2$, V_2 is lines in $\mathbb{R}^2$, and E is the incidence relation (this graph is $K_{2,2}$ -free), it is known ([ST83]) that

$$|E| = O(n^{4/3}).$$

Chernikov, Galvin, and Starchenko [CGS20] showed that in a distal theory, for every formula $\varphi(\bar{x}, \bar{y})$ there are $t, d \in \mathbb{N}_{\geq 2}$ such that

$$\text{if } (V_1, V_2; \varphi(V_1, V_2)) \text{ is } K_{k,k}\text{-free, then } |\varphi(V_1, V_2)| = O(n^{\frac{2td-t-d}{td-1}}).$$


Why OAGs? —AKE principle

We need to study OAGs to study valued fields:

Theorem (Aschenbrenner, Chernikov, Gehret, and Ziegler [Asc+22])

Let (K, v) be a henselian valued fields. (K, v) is distal (has a distal expansion) iff


- *(K, v) is finitely ramified,*
- *The value group of K is distal (has a distal expansion) as an OAG, and*
- *The residue field of K is distal (has a distal expansion) as a pure field.*



However, only a partial result was known for OAGs. Fix an OAG G :

Theorem (ibid.)

If G has finite 'spines', then G is distal iff $|G/pG| < \infty$ for all primes p .



Theorem (O. [Oku26])

G is distal iff for every prime p , the sizes of the 'ribs' with respect to a valuation $\mathfrak{s}_p$ is uniformly bounded.

Theorem (O. [Oku26])

Every OAG has a distal expansion.

To study the theory of an OAG G , it is essential to consider definable convex subgroups of G .

Definition

For $n \geq 2$ and $a \in G \setminus nG$, let $\mathfrak{s}_n(a)$ be the largest convex subgroup H such that $a \notin H + nG$. If $a \in nG$, we set $\mathfrak{s}_n(a) = \emptyset$.

Lemma

$\mathfrak{s}_n(a)$ is definable.

Proof.

Let $\langle x \rangle^{\text{conv}}$ be the convex subgroup generated by x . Then, $x \in \mathfrak{s}_n(a) \iff \langle x \rangle^{\text{conv}} \subseteq \mathfrak{s}_n(a) \iff a \notin \langle x \rangle^{\text{conv}} + nG = [0, nx] + nG$. ■

For convex subgroups H_1, H_2 , we write $H_1 \leq H_2 \iff H_1 \subseteq H_2$.

Example

Let $G = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ with the reversed lexicographic order. The convex subgroups are $(0) \oplus (0) \oplus (0) < \mathbb{Z} \oplus (0) \oplus (0) < \mathbb{Z} \oplus \mathbb{Z} \oplus (0) < G$. For $a = (0, 1, 0)$, $a \notin \mathbb{Z} \oplus (0) \oplus (0) + 2G$ and $a \in \mathbb{Z} \oplus \mathbb{Z} \oplus (0) + 2G$. Hence, $\mathfrak{s}_2(a) = \mathbb{Z} \oplus (0) \oplus (0)$.

An important fact is that we may regard $\mathfrak{s}_n$ as a valuation on G :

Lemma

- 1 $\mathfrak{s}_n(-a) = \mathfrak{s}_n(a)$.
- 2 $\mathfrak{s}_n(a + b) \leq \max \{\mathfrak{s}_n(a), \mathfrak{s}_n(b)\}$ (*ultrametric inequality*).

Proof of (2).

For a convex subgroup H , if $H > \max \{\mathfrak{s}_n(a), \mathfrak{s}_n(b)\}$ then $a, b \in H + nG$, and hence $a + b \in H + nG$. Thus $H > \mathfrak{s}_n(a + b)$. ■

We view $(G, \mathfrak{s}_n)$ in an analogue to valued fields.

Valued field	OAG
v	$(\mathfrak{s}_n)_{n \geq 2}$
Value group	spines
Residue field	ribs

Definition

We define the *spine* $\mathcal{S}_n = \text{Im}(\mathfrak{s}_n)$.

For $G = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$,

$$\mathcal{S}_2 = \{\emptyset < (0) \oplus (0) \oplus (0) < \mathbb{Z} \oplus (0) \oplus (0) < \mathbb{Z} \oplus \mathbb{Z} \oplus (0)\}.$$

Theorem (Aschenbrenner, Chernikov, Gehret, and Ziegler [Asc+22])

Suppose that $|\mathcal{S}_p| < \infty$ for every prime p . Then G is distal iff $|G/pG| < \infty$ for all primes p .

For $G = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$, $G/pG \cong (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/p\mathbb{Z})$. Hence it is distal.

Definition

For $H \in \mathcal{S}_n \setminus \{\emptyset\}$, let $G_{\leq H}^{\mathfrak{s}_n} = \{x \in G \mid \mathfrak{s}_n(x) \leq H\}$ and $G_{< H}^{\mathfrak{s}_n}$ (these are subgroups of G). We define $R_H^{\mathfrak{s}_n} = G_{\leq H}^{\mathfrak{s}_n} / G_{< H}^{\mathfrak{s}_n}$, the *rib* at H .

For $G = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ and $H = \mathbb{Z} \oplus (0) \oplus (0)$, $G_{\leq H}^{\mathfrak{s}_2} = \mathbb{Z} \oplus \mathbb{Z} \oplus 2\mathbb{Z}$ and $G_{< H}^{\mathfrak{s}_2} = \mathbb{Z} \oplus 2\mathbb{Z} \oplus 2\mathbb{Z}$. Thus $R_H^{\mathfrak{s}_2} \cong \mathbb{Z}/2\mathbb{Z}$.

Lemma


The rib $R_H^{\mathfrak{s}_n}$ is of exponent n : For every $a + G_{< H}^{\mathfrak{s}_n} \in R_H^{\mathfrak{s}_n}$, $n(a + G_{< H}^{\mathfrak{s}_n}) = 0$.

Proof.

Since $\mathfrak{s}_n(na) = \emptyset$, we have $na \in G_{< H}^{\mathfrak{s}_n}$ and hence $n(a + G_{< H}^{\mathfrak{s}_n}) = 0$. ■

In particular, $R_H^{\mathfrak{s}_p}$ is an $\mathbb{F}_p$ -vector space.

Valued field	OAG
v	$(\mathfrak{s}_n)_{n \geq 2}$
Value group	spines
Residue field	ribs



Theorem (O. [Oku26])

G is distal iff for every prime p , there is $N_p \in \mathbb{N}$ such that $|R_H^{s_p}| < N_p$ for every $H \in \mathcal{S}_p \setminus \{\emptyset\}$.

If G is sufficiently saturated, G is distal iff for every prime p , $|R_H^{s_p}| < \infty$ for every $H \in \mathcal{S}_p \setminus \{\emptyset\}$.

$R_H^{s_p}$ is a $\mathbb{F}_p$ -vector space (hence stable). So, the distality of an OAG is determined by whether or not an infinite vector space appears as a rib.

While some OAGs are not distal, we have also shown:

Theorem (O.)

Every OAG has a distal expansion.

An infinite rib makes an OAG non-distal. Hence, our strategy is to introduce another valuation on each rib so that resulting ribs are finite. Recall that $R_H^{\mathbb{F}_p}$ is a $\mathbb{F}_p$ -vector space.

Example

Let $V = \bigoplus_{i \in I} \mathbb{F}_p$ be a $\mathbb{F}_p$ -vector space. Introduce a linear order on the index set I and define a valuation v on V as follows:
 $v((a_i)_{i \in I}) = \max \{i \in I \mid a_i \neq 0\}$. Then, every rib with respect to v is isomorphic to $\mathbb{F}_p$. Furthermore, $(V, +, v)$ is distal.

Tools for Proof

OAGs have other families of valuations:

Definition

For $n \geq 2$ and $a \in G$, let $\mathfrak{t}_n(a) = \bigcup_{H \in \mathcal{S}_n, a \notin H} H$ and $\mathfrak{t}_n^+(a) = \bigcap_{H \in \mathcal{S}_n, a \in H} H$. Let $\mathcal{T}_n = \text{Im}(\mathfrak{t}_n)$ and $\mathcal{T}_n^+ = \text{Im}(\mathfrak{t}_n^+)$.




Cluckers and Halupczok [CH11] introduced a many-sorted language $(G, (\mathcal{S}_p), (\mathcal{T}_p), (\mathcal{T}_p^+); (\mathfrak{s}_{p^r}), (\mathfrak{t}_p))$ (p is prime, $r \geq 1$), and established home-sort quantifier elimination.

Distality involves indiscernible sequences. Values of indiscernible sequences behave well.

Lemma

Let $(a_i)_{i \in I}$ be an indiscernible sequence, and let $v \in \{\mathfrak{s}_n, \mathfrak{t}_n \mid n \geq 2\}$. Then exactly one of the following holds:

- $v(a_j - a_i) < v(a_k - a_j) = v(a_i - a_k)$ for every $i < j < k$ in I .
- $v(a_k - a_j) < v(a_i - a_k) = v(a_j - a_i)$ for every $i < j < k$ in I .
- $v(a_j - a_i) = v(a_k - a_j) = v(a_i - a_k)$ for every $i < j < k$ in I .




Typically, *AKE-principles* are theorems of the following form:

AKE-Principle

Suppose that a henselian valued field has some kind of tameness (such as, equicharacteristic 0). Then, it has a property P iff both its value group and residue field have P .

Examples of P : elementary equivalence, NIP, dp-rank, NTP_2 , burden, NATP, distality, distal expansions, elimination of imaginaries, stable embeddedness, etc.




Since OAGs themselves have valuations, we may ask:


Question

What kind of AKE-principles can we establish for OAGs?

Some have already been established: elementary equivalence [Sch82, Corollary 4.7], NIP [GS84], dp-rank [Far17, Theorem 7.2], and a partial result of stable embeddedness [HLT23].



Our results may be viewed as AKE-principles of OAGs for distality. Every spine $\mathcal{S}_p$ is a linearly ordered set, which is distal. As for ribs, a vector space is distal iff it is finite. Thus, a sufficiently saturated OAG is distal iff for every prime p and $H \in \mathcal{S}_p \setminus \{\emptyset\}$, the rib at H is distal. Also, every $\mathbb{F}_p$ -vector space has a distal expansion, and so every OAG does.



- [Asc+22] Matthias Aschenbrenner, Artem Chernikov, Allen Gehret, and Martin Ziegler. “Distality in valued fields and related structures”. *Transactions of the American Mathematical Society* 375.7 (2022), pp. 4641–4710 (cit. on pp. 2, 7, 11).
- [CGS20] Artem Chernikov, David Galvin, and Sergei Starchenko. “Cutting lemma and Zarankiewicz’s problem in distal structures”. *Selecta Mathematica* 26.2, 25 (2020) (cit. on p. 6).
- [CS18] Artem Chernikov and Sergei Starchenko. “Regularity lemma for distal structures”. *Journal of the European Mathematical Society* 20.10 (2018), pp. 2437–2466 (cit. on p. 4).
- [CH11] Raf Cluckers and Immanuel Halupczok. “Quantifier elimination in ordered abelian groups”. *Confluentes Mathematici* 03.04 (2011), pp. 587–615 (cit. on p. 16).
- [Far17] Rafel Farré. *Strong ordered Abelian groups and dp-rank*. 2017. arXiv: 1706.05471 [math.LO] (cit. on p. 19).

- [GS84] Yuri Gurevich and Peter H. Schmitt. “The theory of ordered abelian groups does not have the independence property”. *Transactions of the American Mathematical Society* 284.1 (1984), pp. 171–182 (cit. on pp. 2, 19).
- [HLT23] Martin Hils, Martina Liccardo, and Pierre Touchard. *Stably embedded pairs of ordered abelian groups*. 2023. arXiv: 2308.09989 [math.LO] (cit. on p. 19).
- [KST54] T. Kvári, V. Sós, and P. Turán. “On a problem of K. Zarankiewicz”. *Colloquium Mathematicum* 3.1 (1954), pp. 50–57 (cit. on p. 6).
- [Oku26] Koki Okura. *Distality in ordered abelian groups*. 2026. arXiv: 2602.23256 [math.LO] (cit. on pp. 8, 14).
- [Sch82] Peter H. Schmitt. “Model theory of ordered abelian groups”. Habilitationsschrift. Universität Heidelberg, 1982 (cit. on p. 19).

- [Sim13] Pierre Simon. “Distal and non-distal NIP theories”. *Annals of Pure and Applied Logic* 164.3 (2013), pp. 294–318 (cit. on p. 3).
- [ST83] E. Szemerédi and W. T. Trotter. “Extremal problems in discrete geometry”. *Combinatorica* 3.3 (1983), pp. 381–392 (cit. on p. 6).