Model MEML 75:

Quark sector and axion-like dark matter from $T^2/\mathbb{Z}_3$ geometry

D. V. Yurov

2026

Abstract

We present model MEML 75, which parametrizes the quark sector of the Standard Model through the geometry of the six-dimensional orbifold $T^2/\mathbb{Z}_3$ with two independent tori. The model has 11 nominal parameters (including the ALP misalignment angle θ_i), of which one is fixed by an unphysical overall scale ($a_u = 1$): 10 effective parameters fit 9 quark observables, and adding the ALP misalignment angle θ_i and the relic density $\Omega_a h^2$ gives 11 effective parameters fitting 10 observables ($N_{\rm dof} = -1$). The model is therefore borderline overfitted, which we state explicitly.

The main testable consequence: axion-like dark matter with $f_a = 1.9801 \times 10^{11}$ GeV, $m_a = 28.407$ μeV, $\nu_a = 6.8688$ GHz, and ALP-photon coupling $g_{a\gamma\gamma} = +4.38 \times 10^{-15}$ GeV^{-1} (DFSZ) or -1.13×10^{-14} GeV^{-1} (KSVZ), within reach of future high-frequency haloscope searches such as ADMX and HAYSTAC [3, 4]. The resulting ALP mass also implies a black hole superradiance window at $M_{\rm BH} \sim 10^{25}$ kg. These are consequences of the chain $\alpha_G \to \epsilon \to f_a$, not predictions from the orbifold geometry.

Quark sector. A global fit with loss function $\chi^2 = \sum_i ((O_i - O_i^{\rm target})/\sigma_i)^2$ yields $\chi^2 < 10^{-15}$ (numerical precision limit) at $\tau_u = -0.1577944085 + 2.3851782590i$, $\tau_d = +0.3693905617 + 1.5410496057i$. The hierarchical Yukawa coefficients can be qualitatively understood via the geometric overlap of wavefunctions at the orbifold fixed points, with a universal scale $\alpha \sim 10$ and appropriate prefactors.

Lepton sector (negative result). We attempted to extend the modular construction to the leptons using 3_S contractions of $Y^{(2)}$, $Y^{(4)}$, $Y^{(6)}$ and a uniform diagonal shift. The 3_S sector with $Y^{(2)}$ and $Y^{(4)}$ alone cannot simultaneously reproduce the PMNS angles and the mass-squared differences (best joint fit $\chi^2 \approx 39$). Adding $M_{3_S}(Y^{(6)})$ gives $\chi^2 = 0$ formally, but the modulus τ_l becomes undetermined: six different values of τ_l (spread $\Delta{\rm Re}\,\tau_l = 2.25$) give equivalent fits. A viable lepton sector requires a different modular structure; we leave it for future work and do not include leptons in the main results.

Verdict: the model is a phenomenological quark-sector + ALP construction with a negative result on the lepton sector. It reproduces the 9 quark observables with high precision and makes one independently testable prediction (ALP dark matter at 6.87 GHz).

1 Introduction

Modular models of fermion masses [5, 6] build the mass hierarchy and mixing from compactification geometry. In model MEML 75, quarks are localized on two independent tori $T^2/\mathbb{Z}_3$ with moduli τ_u and τ_d. The mass matrices are built from A_4 modular forms of weights $(2, 4, 6)$. A third torus τ_l for leptons was attempted but does not yield a viable description; we report this negative result in §8.

2 Geometry of $T^2/\mathbb{Z}_3$

The orbifold $T^2/\mathbb{Z}_3$ is defined by:

$$z = x + \tau y, \quad ds^2 = dx^2 + dxdy + dy^2, \tag{2.1}$$

where $\tau = e^{i\pi/3} = \frac{1}{2} + i\frac{\sqrt{3}}{2}$. The Z_3-action $z \to \omega z$ (with $\omega = e^{2\pi i/3}$) has three fixed points, providing a topological explanation for three generations.

3 Bulk coupling α_G (postulate)

$$\alpha_G = \frac{2}{3\pi} = 0.2122065908. \tag{3.1}$$

Geometric motivation: $2/3$ is the volume fraction of the fundamental domain of the $\mathbb{Z}_3$-orbifold; $1/\pi$ is the normalization of the theta-function period.

Geometric motivation via coarse-space topology. The numerical value $2/(3\pi)$ can be read off from the topology of the coarse space of $T^2/\mathbb{Z}_3$. Under the $\mathbb{Z}_3$ action, the torus T^2 collapses onto a sphere: $T^2/\mathbb{Z}_3 \cong S^2$, whose topological Euler characteristic is

$$\chi(S^2) = 2. \tag{3.2}$$

The number of fixed points of the $\mathbb{Z}_3$-action is $N_{\text{fix}} = 3$. Combining these with the theta-function period normalization $1/\pi$ gives

$$\alpha_G = \frac{\chi(S^2)}{N_{\text{fix}}\pi} = \frac{2}{3\pi}. \tag{3.3}$$

We emphasize that this is a geometric motivation, not a derivation: the relation $\alpha_G = \chi(S^2)/(N_{\text{fix}}\pi)$ is not forced by the orbifold geometry. The status of α_G therefore remains that of a postulate.

4 Suppression factor ϵ (ansatz)

$$\epsilon = \left(\frac{\alpha_G}{4\pi}\right)^2 = \frac{1}{36\pi^4} = 2.851662 \times 10^{-4}. \tag{4.1}$$

Two-loop interpretation. The factor 4π in the denominator can be read as the standard two-loop suppression $1/(16\pi^2)$:

$$\epsilon = \left(\frac{\alpha_G}{4\pi}\right)^2 = \frac{\alpha_G^2}{16\pi^2}. \tag{4.2}$$

This provides a physical interpretation of the ansatz, but does not change its numerical value. The status of ϵ therefore remains that of an ansatz.

5 Conditionally derived quantities

$$m_{KK} = \frac{M_{Pl}^{\rm red}}{\sqrt{\mathcal{I}} e^{\pi^2}} = 2.0838 \times 10^{14}\ \text{GeV}, \tag{5.1}$$

$$M_6 = \left(3(M_{Pl}^{\rm red})^2 m_{KK}^2\right)^{1/4} = 2.9646 \times 10^{16}\ \text{GeV}, \tag{5.2}$$

at $\mathcal{I} = 0.3653$.

6 Axion-like (ALP) sector

6.1 ALP parameters

$$f_a = M_{Pl}^{\rm red} \epsilon^2 = 1.9801 \times 10^{11}\ \text{GeV}, \tag{6.1}$$

$$m_a = \frac{(\chi^{1/4})^2}{f_a} = 28.407\ \mu\text{eV}, \tag{6.2}$$

$$\nu_a = \frac{m_a c^2}{h} = 6.8688\ \text{GHz}. \tag{6.3}$$

Here $\chi^{1/4} \approx 75.0$ MeV is the fourth root of the topological susceptibility of QCD. We emphasize that no explicit $U(1)_{PQ}$ symmetry is introduced; the object is an ALP.

Consequence, not prediction. The numerical values of f_a and m_a follow from the chain $\alpha_G \to \epsilon \to f_a \to m_a$ (with $\chi^{1/4}$ taken from QCD), and not from a derivation of these quantities from the orbifold geometry.

6.2 ALP-photon coupling

$$g_{a\gamma\gamma} = \frac{\alpha_{\rm EM}}{2\pi f_a} \left(\frac{E}{N} - 1.92 \right). \tag{6.4}$$

At $f_a = 1.9801 \times 10^{11}$ GeV, $\alpha_{\rm EM}/(2\pi f_a) = 5.866 \times 10^{-15}\ \text{GeV}^{-1}$.

Table 1: Values of $g_{a\gamma\gamma}$ for the two benchmark scenarios.

Scenario	E/N	$g_{a\gamma\gamma}$ (GeV^{-1})
KSVZ	0	-1.126×10^{-14}
DFSZ	8/3	$+4.379 \times 10^{-15}$

6.3 ALP dark matter

$$\Omega_a h^2 \approx 0.12\, \theta_i^2\, f_{\rm anh}(\theta_i) \left(\frac{f_a}{3 \times 10^{11}} \right)^{7/6}, \tag{6.5}$$

where $f_{\rm anh}(\theta_i) = [\ln(e/(1-(\theta_i/\pi)^2))]^{7/6}$ is the anharmonicity factor. Numerical T-effects introduce a +12.4% correction, leading to agreement with Planck at $\theta_i = 1.1162$.

6.4 Black hole superradiance

The ultra-light ALP mass obtained from the chain above has important astrophysical consequences. In the presence of a rotating black hole, the ALP field can extract rotational energy via the superradiance mechanism. The critical black hole mass for which this process is most efficient is given by:

$$M_{\rm BH} \simeq \frac{\hbar c}{G m_a}. \tag{6.6}$$

For the resulting mass $m_a = 28.407\ \mu$eV, this yields:

$$M_{\rm BH} \approx 9.36 \times 10^{24}\ {\rm kg} \approx 4.7 \times 10^{-6} M_\odot. \tag{6.7}$$

The superradiance window is an algebraic consequence of the ALP mass m_a; it does not constitute new physics beyond the ALP sector. We note that black holes in this mass range correspond to the primordial black hole (PBH) window. Currently, no direct detections of PBHs with $M \sim 10^{-6} M_\odot$ have been confirmed by LIGO-Virgo-KAGRA. Microlensing surveys (such as OGLE) place strict upper bounds on the fraction of dark matter that can be composed of PBHs in this mass range, but these constraints do not exclude a subdominant population sufficient to test the superradiance effect.

7 Quark sector (11 nominal parameters)

7.1 Matrix structure

$$M_u = \begin{pmatrix} a_u Y_1^{(2)} & b_u Y_3^{(4)} & c_u Y_2^{(6)} \\ a_u Y_3^{(2)} & b_u Y_2^{(4)} & c_u Y_1^{(6)} \\ a_u Y_2^{(2)} & b_u Y_1^{(4)} & c_u Y_3^{(6)} \end{pmatrix}, \tag{7.1}$$

$$M_d = \begin{pmatrix} a_1 Y_1^{(6)} & b Y_3^{(2)} & c Y_2^{(2)} \\ a_1 Y_3^{(6)} & b Y_2^{(2)} & c Y_1^{(2)} \\ a_2 Y_2^{(2)} & b Y_1^{(2)} & c Y_3^{(4)} \end{pmatrix}. \tag{7.2}$$

7.2 Target values

Table 2: Target values (PDG 2024) [2].

Observable	$O_i^{\rm target}$	σ_i
m_u/m_t	$1.27 \times 10^{-3}/172.0$	$0.30 \times 10^{-3}/172.0$
m_c/m_t	$0.620/172.0$	$0.020/172.0$
m_d/m_b	$2.70 \times 10^{-3}/2.860$	$0.20 \times 10^{-3}/2.860$
m_s/m_b	$55.0 \times 10^{-3}/2.860$	$5.0 \times 10^{-3}/2.860$
m_b/m_t	$2.860/172.0$	$0.030/172.0$
V_{us}	0.22501	0.00068
V_{cb}	0.04053	0.00042
V_{ub}	0.00382	0.00020
J	3.080×10^{-5}	0.130×10^{-5}

7.3 Results

Independent optimization yields $\chi^2 < 10^{-15}$ at the moduli given in the abstract. The residual χ^2 is the machine-precision limit of the numerical fit, not a physical measurement. The coefficients are:

$$c^{\text{fit}} \approx (5.31 \times 10^{-5}, 7.19, 0.039, 2.43 \times 10^{-4}, 0.503, 1.37 \times 10^{-3}, 2.22 \times 10^{-3}). \tag{7.3}$$

Calibration of the overall scale. The overall scale of each mass matrix is unphysical: the observables depend only on mass ratios and CKM elements, which are invariant under $M_u \to kM_u$, $M_d \to kM_d$. We fix $a_u = 1$ without loss of generality.

8 Attempted lepton extension (negative result)

We attempted to extend the modular construction to the lepton sector using the same 3_S contractions of $Y^{(2)}$ and $Y^{(4)}$ that appear in the quark sector, plus a uniform diagonal shift from the singlet contractions $S_k = (Y_1^{(k)})^2 + 2Y_2^{(k)}Y_3^{(k)}$.

$$M_\nu = g_1 M_{3_S}(Y^{(2)}) + g_2 M_{3_S}(Y^{(4)}) + (g_3 S_2 + g_4 S_4 + g_5 S_6)\mathbb{I}. \tag{8.1}$$

We fitted five lepton observables — the three PMNS angles $\theta_{12}, \theta_{13}, \theta_{23}$ and the two mass-squared differences $\Delta m^2_{21}, \Delta m^2_{31}$ (NuFIT 5.2, normal ordering [7]) — and found two problems.

(i) Without $Y^{(6)}$: insufficient structure. With g_1, g_2 and the diagonal shift, the best joint fit gives $\chi^2 \approx 39$, with the residual localized in θ_{23} ($\theta_{23} \approx 46^\circ$ vs. target 49.1°). The 3_S structure with $Y^{(2)}$ and $Y^{(4)}$ alone cannot simultaneously reproduce the three PMNS angles and the two mass-squared differences.

(ii) With $Y^{(6)}$: loss of predictivity. Adding $M_{3_S}(Y^{(6)})$ gives $\chi^2 = 0$ formally, but the fit succeeds for any τ_l: six different values of τ_l (with spread $\Delta\mathrm{Re}\,\tau_l = 2.25$ across the search region) give equivalent fits with $\chi^2 < 10^{-20}$. The modulus τ_l is therefore not determined by the lepton data, and the modular structure of the lepton sector does not constrain the model.

Conclusion. We do not include the lepton sector in the main results. A viable lepton sector within the $T^2/\mathbb{Z}_3$ modular framework requires a different structure (e.g., a nonuniform diagonal shift, additional modular forms, or a different S_3 contraction). We leave this for future work.

9 Full balance

Table 3: Free parameters. Quark sector only, plus the ALP misalignment angle. Nominal count for quarks is 11 (4 moduli + 7 coefficients); fixing $a_u = 1$ removes one, leaving 10 effective quark parameters. Adding θ_i gives 11 effective parameters fitting 10 observables.

Category	Parameters	Count
Quark moduli	τ_u, τ_d (complex)	4
Quark coefficients	$a_u, b_u, c_u, a_1, a_2, b, c$	7
fixed by calibration	$a_u \equiv 1$	-1
ALP misalignment angle	θ_i	1
Total fitted		**11**

$$N_{\text{dof}} = 10 - 11 = -1. \tag{9.1}$$

Parameter classes. The 11 parameters group into three classes: (i) the four real geometric moduli of the two quark tori (τ_u, τ_d), which are the core of the model; (ii) the six Lagrangian Yukawa coefficients $(b_u, c_u, a_1, a_2, b, c)$, which are phenomenological at this stage — deriving them from holomorphic 6D $\to$ 4D couplings is the Critical item of the roadmap (§13); (iii) the ALP misalignment angle θ_i, which is a cosmological initial condition rather than a Lagrangian parameter.

Information content. The model fits 9 quark observables with 10 effective parameters, and 10 observables (9 quarks + $\Omega_a h^2$) with 11 effective parameters. In both cases $N_{\text{dof}} = -1$: the model is borderline overfitted. This is common for string-motivated models, but a reduction of parameters (e.g., deriving the Yukawa coefficients from geometry) is desirable and is the subject of the roadmap.

9.1 Information criteria

We caution that AIC and BIC are derived in the asymptotic regime $n \to \infty$ with fixed k. Here $k/n = 11/10 > 1$, beyond the asymptotic regime; the values are not meaningful and we do not report them. The residuals $< 10^{-15}$ reflect the exactly solvable structure of the quark fit, not an exceptionally good agreement beyond the model's degrees of freedom.

10 Geometric interpretation of the Yukawa coefficients

The numerical values of the coefficients exhibit a clear hierarchical pattern. In a 6D setup, the three generations can be localized at the fixed points $z_1 = 0$, $z_2 = (1+\tau)/3$, and $z_3 = 2(1+\tau)/3$. The Yukawa couplings are given by the overlap integral:

$$c_i \sim A_i e^{-\alpha d_{jk}^2}, \tag{10.1}$$

$$d_{12}^2 = d_{23}^2 = \frac{1 + 2\text{Re}\tau + |\tau|^2}{9\text{Im}\tau}, \quad d_{13}^2 = 4d_{12}^2. \tag{10.2}$$

For the fitted moduli τ_u and τ_d, these distances naturally produce the observed exponential hierarchies. In the up-sector, the extreme suppression of $a_u \sim 10^{-5}$ corresponds to the largest distance d_{13}^2, while $b_u \sim 7$ corresponds to the shorter adjacent fixed points and requires an $\mathcal{O}(1)$ prefactor A_i. While this mechanism does not analytically predict the exact values (as the spread of α across different coefficients is about a factor of 1.7), it provides a natural physical explanation for their vast hierarchy.

11 Flux stabilization of the moduli

The complex structure moduli are dynamically favored by internal fluxes. Following the GVW framework [8], the superpotential is $W(\tau) = a_0 + a_1\tau + a_2\tau^2$, where $a_i \in \mathbb{Z}$ are quantized flux numbers. Moduli stabilization requires the F-term condition:

$$D_\tau W = \partial_\tau W + (\partial_\tau K)W = 0. \tag{11.1}$$

By performing a discrete search over integer flux configurations, we find that the phenomenological values $\tau_u \approx -0.158 + 2.385i$ and $\tau_d \approx 0.369 + 1.541i$ lie near the supersymmetric points of the following fluxes:

$$\begin{aligned} \text{Up} - \text{sector}: &\quad (a_0, a_1, a_2) = (-17, 0, 1), \\ \text{Down} - \text{sector}: &\quad (a_0, a_1, a_2) = (-7, -1, 1). \end{aligned} \tag{11.2}$$

At these configurations, the flux potential yields small residuals: $|D_\tau W| \approx 0.16$ for the up-sector and 0.14 for the down-sector. While these values are not exactly zero, they indicate that the phenomenological moduli are near-supersymmetric points favored by discrete topology. The fluxes are found by a discrete search; deriving them from the topological data of the orbifold is left for future work. Exact moduli stabilization requires a full analysis of the scalar potential, including the Hessian and non-perturbative corrections, which is also left for future work.

12 Open problems

1. **Lepton sector (negative result).** The 3_S sector with $Y^{(2)}$ and $Y^{(4)}$ cannot simultaneously reproduce the PMNS angles and the mass-squared differences (best joint fit $\chi^2 \approx 39$). Adding $M_{3_S}(Y^{(6)})$ gives $\chi^2 = 0$ formally, but τ_l becomes undetermined. A viable lepton sector requires a different modular structure (§8).

2. **Borderline overfitting.** $N_{\text{dof}} = -1$: the model fits 10 observables with 11 effective parameters. A reduction of parameters is desirable and is the subject of the roadmap.

3. **Flatness of** $\text{Re}\,\tau$**.** The flatness of the real parts of the moduli remains an open problem (Appendix E).

4. **Status of** α_G **and** ϵ**.** Both remain postulates/ansätze (§3, §4).

5. **LOO instability.** The Leave-One-Out test shows extreme instability, typical for models with $N_{\text{dof}} < 0$.

6. **Hierarchy problem.** The model explains the fermion hierarchy but does not address the Higgs hierarchy. This is beyond the scope of the present work.

13 Roadmap

Table 4: Phases of development.

Phase	Task	Priority
I	Mechanism for c_i: geometric fixed-point overlap (§10)	Done (qualitative)
I	Flux stabilization of $\mathrm{Im}\,\tau$ (§11)	Done (near-SUSY)
I	Viable lepton sector (different S_3 structure)	Critical (new)
II	Reduce quark parameters (beyond $a_u = 1$ calibration)	Medium
III	ADMX, DESI, DUNE, Hyper-K	External

14 Conclusions

What has been done:

1. A quark-sector fit with 10 effective parameters fitting 9 observables, plus the ALP misalignment angle θ_i fitting $\Omega_a h^2$: 11 effective parameters, 10 observables, $N_{\mathrm{dof}} = -1$.

2. Quark sector: $\chi^2 < 10^{-15}$ (machine limit). The hierarchical Yukawa coefficients are qualitatively explained by geometric fixed-point overlap with $\alpha \sim 10$ and $\mathcal{O}(1)$ prefactors.

3. Moduli: the phenomenological values of τ_u, τ_d lie near the supersymmetric points favored by discrete GVW fluxes.

4. ALP sector: $f_a = 1.9801 \times 10^{11}$ GeV, $m_a = 28.407$ μeV, $\nu_a = 6.8688$ GHz, testable at future high-frequency haloscopes; black hole superradiance window at $M_{\mathrm{BH}} \sim 10^{25}$ kg.

5. Lepton sector: negative result. The 3_S structure with $Y^{(2)}$ and $Y^{(4)}$ is insufficient; adding $Y^{(6)}$ destroys predictivity. A different lepton sector is required (§8).

Verdict. The model is a phenomenological quark-sector + ALP construction. It reproduces the 9 quark observables with high precision, makes one independently testable prediction (ALP dark matter at 6.87 GHz), and reports a negative result on the lepton sector. The borderline overfitting ($N_{\mathrm{dof}} = -1$) is acknowledged, and parameter reduction is left for future work.

Data Availability

The datasets and the Python verification script are available in the Zenodo repository at https://doi.org/10.5281/zenodo.23575072 (Version 13.0, Version DOI; the Concept DOI https://doi.org/10.5281/zenodo.23439547 always resolves to the latest version) and in the GitHub repository at https://github.com/kolbasa1972/MEML-75-Complete. The script reproduces the numerical results of Sections 3–7 and 10. The lepton sector (§8) is a negative result and is not reproduced by the script; the flux stabilization of Section 11 is analytic and is not part of the numerical verification. The flatness scan of Appendix E is not part of the fast verification script.

Declaration of Generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work, the author used AI-assisted tools to verify the Python code, refine the theoretical text, and format the LaTeX document. The author reviewed and edited the content as needed and takes full responsibility for the content of the published article.

A A_4 modular forms: full q-expansions

Definition: $q = e^{2\pi i\tau}$.

$$Y_1^{(2)} = 1 + \sum_{n=1}^{17} C_1^{(n)} q^n, \tag{A.1}$$

$$Y_2^{(2)} = -6\, q^{1/3} \left(1 + \sum_{n=1}^{11} C_2^{(n)} q^n\right), \tag{A.2}$$

$$Y_3^{(2)} = -18\, q^{2/3} \left(1 + \sum_{n=1}^{10} C_3^{(n)} q^n\right), \tag{A.3}$$

$C_1 = [12, 36, 12, 84, 72, 36, 96, 180, 12, 216, 144, 36, 156, 96, 84, 240, 36]$; $C_2 = [1, 7, 8, 18, 14, 30, 20, 42, 30, 56, 42]$; $C_3 = [1, 2, 5, 6, 12, 8, 14, 10, 22, 16]$.

Convolutions:

$$Y_1^{(4)} = (Y_1^{(2)})^2 - Y_2^{(2)} Y_3^{(2)}, \tag{A.4}$$

$$Y_2^{(4)} = (Y_3^{(2)})^2 - Y_1^{(2)} Y_2^{(2)}, \tag{A.5}$$

$$Y_3^{(4)} = (Y_2^{(2)})^2 - Y_1^{(2)} Y_3^{(2)}, \tag{A.6}$$

$$Y_1^{(6)} = \tfrac{1}{3}\left[2Y_1^{(2)} Y_1^{(4)} - Y_2^{(2)} Y_3^{(4)} - Y_3^{(2)} Y_2^{(4)}\right], \tag{A.7}$$

$$Y_2^{(6)} = \tfrac{1}{3}\left[2Y_3^{(2)} Y_3^{(4)} - Y_1^{(2)} Y_2^{(4)} - Y_2^{(2)} Y_1^{(4)}\right], \tag{A.8}$$

$$Y_3^{(6)} = \tfrac{1}{3}\left[2Y_2^{(2)} Y_2^{(4)} - Y_3^{(2)} Y_1^{(4)} - Y_1^{(2)} Y_3^{(4)}\right]. \tag{A.9}$$

B Numerical fit procedure

The global fit was performed using `differential_evolution` (SciPy): 4 seeds (42, 43, 44, 45); maxiter= 800; popsize= 25; tol= 10^{-8}.

Quark sector. Bounds: $\mathrm{Re}\,\tau_{u,d} \in [-0.5, 0.5]$, $\mathrm{Im}\,\tau_u \in [0.87, 2.5]$, $\mathrm{Im}\,\tau_d \in [0.87, 1.60]$, $c_i \in [10^{-8}, 10^2]$.

C Reproduction of τ and χ^2

Table 5: Independent DE runs (quark sector).

Source	$\mathrm{Re}\,\tau_u$	$\mathrm{Im}\,\tau_u$	$\mathrm{Re}\,\tau_d$	$\mathrm{Im}\,\tau_d$
Run 1	−0.15779	2.38518	+0.36939	1.54105
Run 2	−0.15779	2.38518	+0.36939	1.54105
Run 3	−0.12893	2.38411	−0.37212	1.54102

The fitted moduli $\mathrm{Im}\,\tau$ are reproducible across seeds to 10^{-3}. The quantity χ^2 at the minimum is a machine zero; we quote it as $\chi^2 < 10^{-15}$.

D Proof of coefficient degeneracy

At fixed τ, multiply all c_i by k:

Table 6: Invariance of χ^2 under $c_i \to k c_i$.

k	χ^2
0.5	$< 10^{-15}$
1.0	$< 10^{-15}$
1.87	$< 10^{-15}$
10.0	$< 10^{-15}$

The residual values differ only at the level of numerical noise; the invariance is exact up to machine precision.

E Flatness of $\mathrm{Re}\,\tau$

Procedure. The flatness scan below was obtained by refitting the remaining parameters at each value of $\mathrm{Re}\,\tau$. The corresponding refit cycle is computationally expensive and is not part of the fast verification script. Scan in $\mathrm{Re}\,\tau_u$: width of $\chi^2 < 10$: $[-0.40, +0.00]$ (0.40). Scan in $\mathrm{Re}\,\tau_d$: width of $\chi^2 < 10$: $[+0.30, +0.50]$ (0.20).

F LOO test

For each $j \in \{1, \ldots, 9\}$: remove the j-th observable, predict the j-th. $N_{\mathrm{dof}} = +1$. Results: mean $|V_{ub}|$ +16728%, std 11819%.

G Systematic search for α_G

Tested $N = 392$ combinations. 2 coincidences below 0.1%: $|Y_1^{(6)}|/|Y_3^{(4)}| = 0.212125$ (-0.0385%); $|Y_2^{(6)}|/\pi = 0.212411$ ($+0.0965\%$). Poisson: $\langle n \rangle = 0.78$, $n_{\mathrm{obs}} = 2$; $P(n \geq 2) = 0.185$.

References

[1] Planck Collaboration, *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. **641**, A6 (2020), arXiv:1807.06209, DOI: 10.1051/0004-6361/201833910.

[2] R. L. Workman et al. (Particle Data Group), *Review of Particle Physics*, Prog. Theor. Exp. Phys. **2022**, 083C01 (2022), DOI: 10.1093/ptep/ptac097.

[3] ADMX Collaboration, Phys. Rev. Lett. **120**, 151301 (2018), arXiv:1804.05750.

[4] HAYSTAC Collaboration, *A quantum-enhanced search for dark matter axions*, Nature **590**, 238 (2021), arXiv:2008.01853.

[5] F. Feruglio, *Are neutrino masses modular forms?*, Eur. Phys. J. C **77**, 713 (2017), arXiv:1706.08749.

[6] T. Kobayashi, Y. Shimizu, K. Takagi, M. Tanimoto, T. H. Tatsuishi, *Modular S_3-invariant flavor model in SU(5) grand unified theory*, PTEP **2020**, 053B05 (2020), arXiv:1906.10341.

[7] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, *The fate of hints: updated global analysis of three-flavor neutrino oscillations*, JHEP **09** (2020) 178, arXiv:2007.14792 (NuFIT 5.2, 2022).

[8] S. Gukov, C. Vafa, E. Witten, *CFT's from Calabi-Yau four-folds*, Nucl. Phys. B **584** (2000) 69, arXiv:hep-th/9906070.