Model MEML 75 (v16.0):

Quark and lepton mass matrices from mixed
$T^2/\mathbb{Z}_3 \times T^2/\mathbb{Z}_4$ geometry
with axion-like dark matter and leptogenesis

D. V. Yurov

2026

Abstract

We present model MEML 75 (v16.0), which parametrizes the fermion sector of the Standard Model through the geometry of mixed six-dimensional orbifolds. The model has **18 effective fitted parameters** — 17 Lagrangian parameters plus the ALP misalignment angle θ_i, which is a cosmological initial condition — fitting **16 observables** (9 quarks + 5 lepton observables + $\Omega_a h^2$ + η_B), yielding $N_{\rm dof}^{\rm fit} = -2$. We explore a phenomenological reduction $c_2 + c_3 = 0$ in the lepton sector, which yields a slightly improved balance, though it is not derived from first principles. Without it, the balance would be $N_{\rm dof}^{\rm fit} = -3$.

New in v16.0: We evaluate the electron electric dipole moment (EDM) and lepton flavor violation (LFV) in the Type-I See-Saw framework. We find $|d_e| \simeq 2.6 \times 10^{-54}\ e \cdot$cm and BR$(\mu \to e\gamma) \sim 10^{-61}$, both safely below current experimental limits. We also report a *negative result* on the geometric derivation of the lepton Yukawa coefficients from zero-mode overlap integrals with magnetic flux.

The main testable consequences: axion-like dark matter with $f_a = 1.9801 \times 10^{11}$ GeV, $m_a = 28.407\ \mu$eV, testable at future high-frequency haloscope upgrades (e.g., ADMX-EFR or HAYSTAC Phase III) [3, 4]. The model makes 3 conditional predictions: $\sum m_i \approx 0.068$ eV, $m_{\beta\beta} \approx 8.37$ meV (assuming vanishing Majorana phases; for arbitrary phases, $m_{\beta\beta}$ lies in the range 0–8.7 meV), and $\delta_{CP} \approx 23^\circ$ or 157°.

It also selects the upper octant of θ_{23}: this is consistent with NuFIT 5.0 ($\chi^2 = 0.28$) and with NuFIT 5.2 without SK atmospheric data ($\chi^2 = 0.39$), but in tension at high confidence with NuFIT 5.2 including SK atmospheric data ($\chi^2 \approx 58$). This tension is a genuine testable consequence that will be settled by DUNE and Hyper-Kamiokande.

1 Introduction

Modular models of fermion masses [5, 6, 7, 8] build the mass hierarchy and mixing from compactification geometry. In model MEML 75, we explore a mixed geometric setup: quarks are localized on a $T^2/\mathbb{Z}_3$ torus, while leptons are localized on a $T^2/\mathbb{Z}_4$ torus. This dual-orbifold approach allows us to leverage the best features of two different modular groups: A_4 for the hierarchical quark masses, and S_4' for the rich leptonic mixing pattern.

2 Geometry of $T^2/\mathbb{Z}_3$ and $T^2/\mathbb{Z}_4$

The orbifold $T^2/\mathbb{Z}_3$ is defined by $z = x + \tau y$ with $\tau = e^{i\pi/3}$. The quark mass matrices are built from A_4 modular forms of weights $(2, 4, 6)$. The mixed geometry naturally arises in Type IIB orientifold compactifications. The lepton sector uses the $T^2/\mathbb{Z}_4$ orbifold, whose modular symmetry is the double cover group S_4' [7, 8].

3 Bulk coupling α_G (postulate)

$$\alpha_G = \frac{2}{3\pi} = 0.2122065908. \tag{3.1}$$

Geometric motivation: The orbifold $T^2/\mathbb{Z}_3$ has $N_{\text{fix}} = 3$ fixed points. The factor

$$\frac{2}{3} = \frac{N_{\text{fix}} - 1}{N_{\text{fix}}} = 1 - \frac{1}{N_{\text{fix}}} \tag{3.2}$$

reflects the combinatorics of the three fixed points: a given fixed point accounts for $1/3$ of the fixed-point set, leaving $2/3$. The factor $1/\pi$ is the normalization of the theta-function period. We emphasize that this is a geometric motivation, not a rigorous derivation. The status of α_G remains that of a postulate.

4 Suppression factor ε (ansatz)

$$\varepsilon = \left(\frac{\alpha_G}{4\pi}\right)^2 = \frac{1}{36\pi^4} = 2.851662 \times 10^{-4}. \tag{4.1}$$

We have searched for a geometric derivation of ε. The natural candidate, a KK-mode suppression $\varepsilon \sim (g_s/4\pi)^2 (M_{KK}/M_6)^2$, fails by three orders of magnitude: with $M_{KK}/M_6 = 7.03 \times 10^{-3}$ and $g_s \sim 1$, it gives $\varepsilon \sim 3 \times 10^{-7}$, not 2.85×10^{-4}. We do not claim a KK origin. We do note a numerical coincidence at the 11% level:

$$\alpha_G^2 \frac{M_{KK}}{M_6} = 3.17 \times 10^{-4} \quad \text{vs.} \quad \varepsilon = 2.85 \times 10^{-4}, \qquad \frac{\alpha_G^2 \, (M_{KK}/M_6)}{\varepsilon} = 1.11. \tag{4.2}$$

Equivalently, $M_{KK}/M_6 \approx 1/(4\pi)^2$ to within 11%. Since M_{KK}/M_6 is a geometric ratio and $(4\pi)^2$ is a loop factor, the coincidence is not a derivation: no principle in the model relates them. We keep ε as an ansatz, and we flag this numerical coincidence as a possible hint of a deeper structure that we cannot establish here.

5 Conditionally derived quantities

$$m_{KK} = \frac{M_{Pl}^{\rm red}}{\sqrt{\mathcal{I}}\, e^{\pi^2}} = 2.0838 \times 10^{14}\ \text{GeV}, \tag{5.1}$$

$$M_6 = \left(3(M_{Pl}^{\rm red})^2 m_{KK}^2\right)^{1/4} = 2.9646 \times 10^{16}\ \text{GeV}, \tag{5.2}$$

at $\mathcal{I} = 0.3653$.[1]

6 Axion-like (ALP) sector

$$f_a = M_{Pl}^{\rm red}\, \varepsilon^2 = 1.9801 \times 10^{11}\ \text{GeV}, \tag{6.1}$$

$$m_a = \frac{(\chi^{1/4})^2}{f_a} = 28.407\ \mu\text{eV}, \tag{6.2}$$

$$\nu_a = \frac{m_a c^2}{h} = 6.8688\ \text{GHz}. \tag{6.3}$$

Here $\chi^{1/4} \approx 75.0$ MeV. The relic density is:

$$\Omega_a h^2 \approx 0.12\, \theta_i^2\, f_{\rm anh}(\theta_i) \left(\frac{f_a}{3 \times 10^{11}}\right)^{7/6}, \tag{6.4}$$

where $f_{\rm anh}(\theta_i) = [\ln(e/(1-(\theta_i/\pi)^2))]^{7/6}$. We fit θ_i to the Planck value $\Omega_a h^2 = 0.12$ [1], obtaining $\theta_i \approx 1.174$. This is a cosmological initial condition, not a Lagrangian parameter; it is fitted to $\Omega_a h^2$ and is counted in the effective parameter balance of Section 13.

At $f_a = 1.9801 \times 10^{11}$ GeV: $g_{a\gamma\gamma} = +4.38 \times 10^{-15}$ GeV^{-1} (DFSZ) or -1.13×10^{-14} GeV^{-1} (KSVZ). The ALP mass implies a black hole superradiance window at $M_{\rm BH} \simeq \hbar c/(G m_a) \approx 9.36 \times 10^{24}$ kg.

7 Quark sector ($T^2/\mathbb{Z}_3$, A_4)

7.1 Matrix structure

$$M_u = \begin{pmatrix} a_u Y_1^{(2)} & b_u Y_3^{(4)} & c_u Y_2^{(6)} \\ a_u Y_3^{(2)} & b_u Y_2^{(4)} & c_u Y_1^{(6)} \\ a_u Y_2^{(2)} & b_u Y_1^{(4)} & c_u Y_3^{(6)} \end{pmatrix}, \tag{7.1}$$

$$M_d = \begin{pmatrix} a_1 Y_1^{(6)} & b Y_3^{(2)} & c Y_2^{(2)} \\ a_1 Y_3^{(6)} & b Y_2^{(2)} & c Y_1^{(2)} \\ a_2 Y_2^{(2)} & b Y_1^{(2)} & c Y_3^{(4)} \end{pmatrix}. \tag{7.2}$$

[1]The geometric factor $\mathcal{I}$ and the neutrino Yukawa coupling $y_\nu = 0.934$ are *not* derived from first principles in this work. To be conservative, we count them as free parameters in the balance of Section 13. The value $\mathcal{I} = 0.3653$ is numerically close to $N_3 V_3 - N_4 V_4 = (\sqrt{3}-1)/2 \approx 0.3660$, evaluated at the symmetric orbifold point $\tau_3 = e^{i\pi/3}$, $\tau_4 = i$, where $V_3 = \text{Im}\,\tau_3/3$, $V_4 = \text{Im}\,\tau_4/4$, and $N_3 = 3$, $N_4 = 2$ are the numbers of fixed points of $\mathbb{Z}_3$ and of order-4 $\mathbb{Z}_4$, respectively. We do *not* claim a derivation: (i) the fitted moduli are not at the symmetric point; (ii) the number of $\mathbb{Z}_4$ fixed points is convention-dependent (2, 3, or 4); (iii) there is no principle selecting subtraction over addition. A refit with $\mathcal{I} = (\sqrt{3}-1)/2$ fixed leaves all predictions unchanged at the 0.1% level (Appendix G), so the data are consistent with the geometric value, but do not prove it. We therefore keep $\mathcal{I}$ as a free parameter.

Independent optimization yields $\chi^2 < 10^{-19}$ at $\tau_u = -0.1578 + 2.3852i$ and $\tau_d = 0.3694 + 1.5410i$ [2]. The overall scale is unphysical; we fix $a_u = 1$ without loss of generality. We stress that the quark sector has 10 parameters for 9 observables ($\chi^2 < 10^{-19}$ by construction); the fit quality is not a prediction, and the geometric overlap discussed below provides only a qualitative explanation of the hierarchy.

8 Lepton sector ($T^2/\mathbb{Z}_4$, S_4')

The neutrino mass matrix is built from the contractions of the **2** and **3′** representations of S_4':

$$M_\nu = c_1 M_{\mathbf{2}}(\tau_l) + c_2 M_{\mathbf{3'}}(\tau_l) + c_3 \mathbb{I}. \tag{8.1}$$

We fit the model to NuFIT 5.0 [10], yielding $\chi^2 \approx 0.089$ at $\tau_l = 1.5398 + 1.5511\,i$, $c_1 = 6.371$, $c_2 = 17.460$, $c_3 = -17.199$ (the earlier value $\chi^2 \approx 0.23$ in earlier drafts was obtained with a less-converged differential-evolution fit; the refined global scan gives 0.089).

Phenomenological relation $c_2 + c_3 = 0$. The unconstrained fit gives $c_2 + c_3 = 0.261$, which is 1.5% of the individual coefficients $|c_2| \approx |c_3| \approx 17.4$. We therefore consider the reduced model obtained by imposing $c_2 + c_3 = 0$. This is a *phenomenological reduction*, not a derivation: it replaces the combination $M_{\mathbf{3'}} + (c_3/c_2)\mathbb{I}$ by $M_{\mathbf{3'}} - \mathbb{I}$. The constrained fit gives $\chi^2 \approx 0.309$ at $\tau_l = 1.584 + 1.552\,i$, $c_1 = -6.591$, $c_2 = -17.400$ (with $c_3 = -c_2 = +17.400$), compared with $\chi^2 \approx 0.089$ unconstrained, i.e. $\Delta\chi^2 = 0.22$ for one fewer parameter ($p \approx 0.64$). In the lepton sector ($k = 3$ full, $k = 2$ reduced, $n = 5$), AIC and BIC give $\Delta\mathrm{AIC} = -1.78$ and $\Delta\mathrm{BIC}(N{=}5) = -1.39$, both weakly favouring the reduction. Using the global $N = 16$ in the BIC penalty gives $\Delta\mathrm{BIC}(N{=}16) = -2.55$. Since the fit does not strongly distinguish the two scenarios (and $k/n > 1$ places AIC/BIC outside their asymptotic regime), we adopt $c_2 + c_3 = 0$ as a phenomenological simplification for the remainder of this work.

We tested the complete set of linear relations among c_1, c_2, c_3 with coefficients in $\{-1, 0, +1\}$ and pairwise equalities (seven in total); the six alternatives are excluded at $\chi^2 > 38$, so even with a multiple-comparisons penalty for seven tests the selection is not marginal; nevertheless, it remains a post-hoc model choice, not a derivation. A derivation from a $\mathbb{Z}_2$ symmetry of S_4' is not possible in the minimal setup: the identity term $c_3\mathbb{I}$ is a trivial singlet and is invariant under any subgroup, so a genuine derivation would require additional structure in the heavy-neutrino sector. We therefore treat $c_2 + c_3 = 0$ as a phenomenological reduction, not a symmetry consequence. Because the reduction is not derived, we report both balances in Section 13: $N_{\text{eff}} = 19$ without the reduction, and $N_{\text{eff}} = 18$ with it. Our main balance uses the latter, justified by the model-selection argument above.

With the reduction $c_2 + c_3 = 0$ imposed, the PMNS angles and mass-squared differences are

$$\theta_{12} = 33.64^\circ, \quad \theta_{13} = 8.53^\circ, \quad \theta_{23} = 49.04^\circ, \tag{8.2}$$

$$\Delta m^2_{21} = 7.48 \times 10^{-5}\ \text{eV}^2, \quad \Delta m^2_{31} = 2.51 \times 10^{-3}\ \text{eV}^2. \tag{8.3}$$

Using Type-I See-Saw, $\Lambda_\nu = y_\nu^2 v^2 / M_6 \approx 1.78$ meV, we find the physical masses $m_1 \approx 6.39$ meV, $m_2 \approx 10.75$ meV, $m_3 \approx 50.55$ meV. This gives $\sum m_i \approx 0.068$ eV. For the effective Majorana mass, assuming vanishing Majorana phases, we find $m_{\beta\beta} \approx 8.37$ meV, which is below the current KamLAND-Zen limit [11]; for arbitrary Majorana phases,

$m_{\beta\beta}$ lies in the range 0–8.7 meV, entirely below the current KamLAND-Zen limit, and this is a testable prediction for next-generation experiments (nEXO, LEGEND-1000). We note that the condition for $m_{\beta\beta} = 0$ is satisfied only marginally ($A = 4.33$ meV vs $B + C = 4.34$ meV). Small shifts in the input values could result in a non-zero lower bound. Therefore, the robust prediction is the upper bound: $m_{\beta\beta} < 8.7$ meV.

The CP phase is $\delta_{CP} \approx 23°$. The Jarlskog invariant determines only $|\sin\delta_{CP}|$, allowing a second solution $\delta_{CP} \approx 157°$; d_e and η_B depend on δ_{CP} through $\sin\delta_{CP}$, while $m_{\beta\beta}$ depends on δ_{CP} through $\cos(2\delta_{CP})$; since $\sin(\pi - \delta) = \sin\delta$ and $\cos(2(\pi - \delta)) = \cos(2\delta)$, all three predictions are identical for both branches.

Octant ambiguity in θ_{23}. The fit gives $\theta_{23} = 49.04°$, in the upper octant. This is consistent with NuFIT 5.0 (IC19+SK atm, $\chi^2 = 0.28$) and with NuFIT 5.2 without SK atmospheric data ($\chi^2 = 0.39$). However, NuFIT 5.2 with SK atmospheric data prefers the lower octant ($\theta_{23} \approx 42.2°$); against this dataset our prediction gives $\chi^2 \approx 58$, which is in tension at high confidence. We emphasize that the octant of θ_{23} is not yet settled: T2K and NOvA show a mild preference for the upper octant, while SK atmospheric data prefer the lower one. The model predicts the upper octant as a consequence of the S_4' structure and the fitted value of τ_l; this is a genuine testable consequence that will be settled by DUNE and Hyper-Kamiokande. If future data confirm the lower octant, the model would need a different S_4' representation assignment.

9 Baryon asymmetry via leptogenesis

The heavy neutrinos with $M_R \sim M_6 \approx 3 \times 10^{16}$ GeV generate the baryon asymmetry [9]. Using the Davidson-Ibarra formula [12], we find $\varepsilon_{CP} \approx 0.57$ (consistent with the Davidson-Ibarra upper bound). The resulting baryon asymmetry before dilution is

$$\eta_B^{\rm pre} \approx \frac{28}{79}\,\varepsilon_{CP}\,\kappa\,g_*^{-1} \approx 1.9 \times 10^{-3}, \tag{9.1}$$

where $\kappa \approx 1$ is the washout factor in the weak-washout regime ($K \approx 0.83 < 1$) and $g_* = 106.75$ is the effective number of relativistic degrees of freedom. This overproduction is resolved by entropy dilution from the late decay of moduli [13], with a dilution factor $\Delta \approx 3.13 \times 10^6$, yielding the observed $\eta_B^{\rm obs} \approx 6.1 \times 10^{-10}$. We stress that η_B is not an independent prediction: it is obtained only after fitting Δ.

Status of the dilution factor Δ. Using the standard moduli-decay formula

$$\Delta = \frac{\frac{1}{2}m_\phi^2\phi_0^2}{(\pi^2/30)\,g_*\,T_{\rm RH}\,T_i^3}, \qquad T_{\rm RH} \approx \left(\frac{90}{\pi^2 g_*}\right)^{1/4}\sqrt{\frac{m_\phi^3}{8\pi M_{Pl}}}, \tag{9.2}$$

we find that $\Delta \sim 10^6$ requires either $\phi_0 \sim 10^{12}$ GeV (with $m_\phi \sim 3 \times 10^6$ GeV, $T_i \sim 10^{10}$ GeV) or a specific combination $\phi_0 \sim 10^{16}$–10^{17} GeV with $T_i \sim 10^{13}$ GeV (with $m_\phi \sim 10^9$ GeV). Neither is fixed by the model: m_ϕ depends on moduli stabilization (Section 11), ϕ_0 on the inflationary sector, and T_i on the leptogenesis scale. We therefore keep Δ as a free parameter; a full derivation requires a UV completion of moduli stabilization and reheating, which is beyond the scope of this work.

10 Lepton Flavor Violation and Electric Dipole Moments

Lepton Flavor Violation. In the Type-I See-Saw mechanism, the branching ratio for $\mu \to e\gamma$ is:

$$\mathrm{BR}(\mu \to e\gamma) \approx \frac{3\alpha_{\mathrm{EM}}}{32\pi} \left| \sum_{i=1}^{3} (Y_\nu)_{\mu i} (Y_\nu^\dagger)_{ie} \frac{M_W^2}{M_i^2} \right|^2 . \tag{10.1}$$

Using the predicted PMNS matrix and neutrino masses, we find $\mathrm{BR}(\mu \to e\gamma) \sim 10^{-61}$. This is many orders of magnitude below the current experimental limit (4.2×10^{-13} from MEG II).

Electron Electric Dipole Moment. The CP-violating phase $\delta_{CP} \approx 23^\circ$ contributes to the electron EDM via two-loop Barr-Zee diagrams:

$$|d_e| \sim \frac{\alpha_{\mathrm{EM}}}{4\pi} \frac{m_e}{M_6^2} \sin \delta_{CP}. \tag{10.2}$$

With $M_6 \approx 2.96 \times 10^{16}$ GeV, this yields $|d_e| \simeq 2.6 \times 10^{-54}\ e \cdot \mathrm{cm}$. This is safely below the ACME III limit ($< 4.1 \times 10^{-30}\ e \cdot \mathrm{cm}$).

11 Attempted geometric derivation of the Yukawa coefficients

We attempted to derive c_1, c_2, c_3 from the overlap integrals of zero-mode wavefunctions on $T^2/\mathbb{Z}_4$ with a uniform magnetic flux M. Using a numerical grid of $N_{\mathrm{grid}} = 50$ and $N_{\max} = 30$, we find that for flux $M = 2$, the diagonal contraction strongly dominates over the off-diagonal ones, with a ratio of order $Y_{(0,0,0)}/Y_{(0,1,1)} \sim 65$. However, the phenomenological fit requires $c_1/c_2 \approx 0.36$. Thus, a uniform magnetic flux cannot reproduce the required coefficients.

12 Flux stabilization of the moduli

Following the GVW framework [14], $W(\tau) = a_0 + a_1\tau + a_2\tau^2$. The values $\tau_u \approx -0.158 + 2.385i$ and $\tau_d \approx 0.369 + 1.541i$ lie near the supersymmetric points of the fluxes $(-17, 0, 1)$ and $(-7, -1, 1)$, yielding small residuals $|D_\tau W| \approx 0.16$ and 0.14. Exact stabilization requires non-perturbative terms (KKLT).

13 Full balance

Table 1: Free parameters. The model has 18 effective fitted parameters (17 Lagrangian parameters plus the ALP misalignment angle θ_i, a cosmological initial condition) fitting 16 observables. The reduction from 19 to 18 effective parameters follows from the phenomenological relation $c_2 + c_3 = 0$ (Section 8), which removes one parameter from the lepton sector. The alternative balance without this reduction (variant C) is $N_{\rm eff} = 19$, $N_{\rm dof}^{\rm fit} = -3$, $N_{\rm dof}^{\rm total} = 0$. θ_i is fitted to $\Omega_a h^2$; it is counted in $N_{\rm eff}$ but is not a Lagrangian parameter.

Category	Parameters	Count
Quark moduli	τ_u, τ_d	4
Quark coefficients	$a_u, b_u, c_u, a_1, a_2, b, c$	7
fixed by calibration	$a_u \equiv 1$	-1
Lepton modulus	τ_l	2
Lepton coefficients	c_1, c_2 (with $c_3 = -c_2$)	2
Entropy dilution factor	Δ	1
See-Saw Yukawa coupling	y_ν	1
See-Saw geometric factor	$\mathcal{I}$	1
Total fitted (Lagrangian)		**17**
ALP misalignment angle (cosmological IC)	θ_i	1
Total effective fitted		**18**

[2]

Note that among the 16 observables, η_B is a postdiction (obtained via the free parameter Δ), while the 3 quantities (δ_{CP}, $\sum m_i$, $m_{\beta\beta}$) are genuine conditional predictions.

The model fits 16 observables with 18 effective parameters ($N_{\rm dof}^{\rm fit} = -2$) and makes 3 conditional predictions that can be cross-checked against external data not used in the fit: $\delta_{CP} \approx 23^\circ$ or 157°, $\sum m_i \approx 0.068$ eV, and $m_{\beta\beta} \approx 8.37$ meV (the latter assuming vanishing Majorana phases; for arbitrary phases, $m_{\beta\beta}$ lies in the range 0–8.7 meV). If these three predictions are treated as additional data points (appropriate once they are measured), the counting gives $N_{\rm dof}^{\rm total} = (16 + 3) - 18 = +1$ (conditional). *Without* the phenomenological reduction $c_2 + c_3 = 0$, the balance would be $N_{\rm eff} = 19$, $N_{\rm dof}^{\rm fit} = -3$, $N_{\rm dof}^{\rm total} = 0$ (variant C); we have adopted the reduced balance (18, -2, $+1$) on the basis of the model-selection argument of Section 8. Separately, we note that the model selects the upper octant of θ_{23} (Section 8); since this is already used in the fit through NuFIT 5.0, it is a *testable consequence* rather than an independent prediction. A UV completion may fix y_ν and $\mathcal{I}$, but we do not claim it in this work.

[2]One might attempt to fix y_ν via the SO(10) relation $y_\nu = y_u$ at the GUT scale. Running y_t from M_Z to $M_{\rm GUT} \approx 2 \times 10^{16}$ GeV with 1-loop MSSM RGEs gives $y_t(M_{\rm GUT}) \approx 0.48$ (the Pendleton–Ross quasi-fixed point [15]), while $y_\nu = 0.934$ would require a Clebsch–Gordan factor $f \approx 1.95$ from $10_H + 126_H$ representations. Since y_{10} and y_{126} are then independent parameters, this does not reduce the parameter count. At $\tan\beta = 2$, where $f \approx 1.1$, the value is marginally compatible, but this regime is excluded by the Higgs mass and by t-b-τ unification. We therefore keep y_ν as a free parameter.

14 Conclusions

What has been done:

1. A full model with 18 effective parameters fitting 16 observables ($N_{\text{dof}}^{\text{fit}} = -2$), making 3 conditional predictions (δ_{CP}, $\sum m_i$, $m_{\beta\beta}$), with the upper octant of θ_{23} as an additional testable consequence (Section 8). The reduction from 19 to 18 parameters (from three to two lepton coefficients) follows from the phenomenological relation $c_2 + c_3 = 0$, justified by the model-selection argument of Section 8 ($\Delta\chi^2 = 0.22$, $p = 0.64$, $\Delta\text{AIC} = -1.78$, and $\Delta\text{BIC}(N{=}16) = -2.55$ as an asymptotic heuristic).

2. Quark sector: $\chi^2 < 10^{-19}$ by construction (10 parameters, 9 observables). The hierarchical Yukawa coefficients are qualitatively reproduced by geometric fixed-point overlap with $\alpha \sim 10$ and $\mathcal{O}(1)$ prefactors; the fit quality is not a prediction.

3. Lepton sector (S_4', $T^2/\mathbb{Z}_4$): $\chi^2 \approx 0.309$ at $\tau_l = 1.584 + 1.552i$, $c_1 = -6.591$, $c_2 = -17.400$, $c_3 = -c_2 = +17.400$; PMNS angles and mass-squared differences reproduced.

4. Moduli: the phenomenological values of τ_u, τ_d lie near the supersymmetric points favored by discrete GVW fluxes.

5. ALP sector: $f_a = 1.9801 \times 10^{11}$ GeV, $m_a = 28.407$ μeV, $\nu_a = 6.8688$ GHz, testable at future high-frequency haloscopes; black hole superradiance window at $M_{\text{BH}} \sim 10^{25}$ kg.

6. Leptogenesis: $\eta_B^{\text{pre}} \approx 1.9 \times 10^{-3}$, diluted by $\Delta \approx 3.13 \times 10^6$ to the observed value; Δ is treated as a free parameter, with the moduli-decay formula discussed in Section 9.

7. LFV and EDM: $\text{BR}(\mu \to e\gamma) \sim 10^{-61}$, $d_e \sim 2.6 \times 10^{-54}$ $e \cdot$ cm, safely below current experimental limits.

8. Negative results: (i) $y_\nu = 0.934$ cannot be derived from SO(10)-like GUT unification without introducing new parameters; (ii) $\mathcal{I} = 0.3653$ is numerically close to $(\sqrt{3} - 1)/2$ but not derived (Appendix G); (iii) lepton Yukawa coefficients cannot be derived from uniform magnetic flux overlaps; (iv) ε does not arise from a simple KK-mode suppression (Section 4); (v) the dilution factor Δ cannot be derived from moduli dynamics without a UV completion (Section 9).

While the model contains phenomenological parameters, it is not a mere polynomial fit. It arises from a specific geometric structure ($T^2/\mathbb{Z}_3 \times T^2/\mathbb{Z}_4$), which correlates the quark, lepton, and ALP sectors. Crucially, it predicts the exact frequency of the ALP signal (6.87 GHz) and the upper bound on $m_{\beta\beta}$, which will be tested by upcoming experiments. This predictive power distinguishes it from generic parameterizations.

Verdict. The model is a phenomenological construction with 18 effective parameters (17 Lagrangian + θ_i), $N_{\text{dof}}^{\text{fit}} = -2$, three conditional predictions (δ_{CP}, $\sum m_i$, $m_{\beta\beta}$), one independently testable ALP signal at 6.87 GHz, and one testable consequence — the upper octant of θ_{23} — that will be settled by DUNE and Hyper-Kamiokande. The overfitting is acknowledged, as are the negative results on y_ν, $\mathcal{I}$, ε, Δ, and the flux derivation of c_i. The relation $c_2 + c_3 = 0$ is a phenomenological reduction, not a derivation; without it, the balance would be (19, −3, 0). Further parameter reduction requires a UV completion.

Data Availability

The datasets and the Python verification script (v16.0) are available in the Zenodo repository at https://doi.org/10.5281/zenodo.23638606 (Concept DOI: https://doi.org/10.5281/zenodo.23439547) and in the GitHub repository at https://github.com/kolbasa1972/MEML-75-Mixed-Geometry. The script `meml75_v16.0_master.py` reproduces the numerical results of Sections 3–12, implements the consistency check of Appendix G, the lepton-modulus coarse scan discussed in Section 8, the GUT test quoted in Section 13, and the NuFIT 5.0 versus 5.2 comparison discussed in Section 8. The test of the seven linear relations in the lepton sector (Section 8) is implemented in `meml75_constraint_test_extended.py`. The global scans of the lepton modulus are implemented in `meml75_tau_l_global_scan_v2.py` (unconstrained) and `meml75_constrained_global_scan.py` (constrained).

Declaration of Generative AI and AI-assisted technologies

During the preparation of this work, the author used AI-assisted tools to verify the Python code, refine the theoretical text, and format the LaTeX document. The author reviewed and edited the content as needed and takes full responsibility for the content of the published article.

A A_4 modular forms: full q-expansions

Definition: $q = e^{2\pi i\tau}$.

$$Y_1^{(2)} = 1 + \sum_{n=1}^{17} C_1^{(n)} q^n, \tag{A.1}$$

$$Y_2^{(2)} = -6\, q^{1/3} \left(1 + \sum_{n=1}^{11} C_2^{(n)} q^n\right), \tag{A.2}$$

$$Y_3^{(2)} = -18\, q^{2/3} \left(1 + \sum_{n=1}^{10} C_3^{(n)} q^n\right), \tag{A.3}$$

$C_1 = [12, 36, 12, 84, 72, 36, 96, 180, 12, 216, 144, 36, 156, 96, 84, 240, 36]$; $C_2 = [1, 7, 8, 18, 14, 30, 20, 42, 30, 56, 42]$; $C_3 = [1, 2, 5, 6, 12, 8, 14, 10, 22, 16]$.

Convolutions:

$$Y_1^{(4)} = (Y_1^{(2)})^2 - Y_2^{(2)} Y_3^{(2)}, \tag{A.4}$$

$$Y_2^{(4)} = (Y_3^{(2)})^2 - Y_1^{(2)} Y_2^{(2)}, \tag{A.5}$$

$$Y_3^{(4)} = (Y_2^{(2)})^2 - Y_1^{(2)} Y_3^{(2)}, \tag{A.6}$$

$$Y_1^{(6)} = \tfrac{1}{3}\left[2Y_1^{(2)} Y_1^{(4)} - Y_2^{(2)} Y_3^{(4)} - Y_3^{(2)} Y_2^{(4)}\right], \tag{A.7}$$

$$Y_2^{(6)} = \tfrac{1}{3}\left[2Y_3^{(2)} Y_3^{(4)} - Y_1^{(2)} Y_2^{(4)} - Y_2^{(2)} Y_1^{(4)}\right], \tag{A.8}$$

$$Y_3^{(6)} = \tfrac{1}{3}\left[2Y_2^{(2)} Y_2^{(4)} - Y_3^{(2)} Y_1^{(4)} - Y_1^{(2)} Y_3^{(4)}\right]. \tag{A.9}$$

B Numerical fit procedure

The global fit was performed using `differential_evolution` (SciPy): 4 seeds (42, 43, 44, 45); maxiter= 800; popsize= 25; tol= 10^{-8}.

Quark sector. Bounds: $\mathrm{Re}\,\tau_{u,d} \in [-0.5, 0.5]$, $\mathrm{Im}\,\tau_u \in [0.87, 2.5]$, $\mathrm{Im}\,\tau_d \in [0.87, 1.60]$, $c_i \in [10^{-8}, 10^2]$.

C Reproduction of τ and χ^2

Table 2: Independent DE runs (quark sector).

Source	$\mathrm{Re}\,\tau_u$	$\mathrm{Im}\,\tau_u$	$\mathrm{Re}\,\tau_d$	$\mathrm{Im}\,\tau_d$
Run 1	−0.15779	2.38518	+0.36939	1.54105
Run 2	−0.15779	2.38518	+0.36939	1.54105
Run 3	−0.12893	2.38411	−0.37212	1.54102

The fitted moduli $\mathrm{Im}\,\tau$ are reproducible across seeds to 10^{-3}. Run 3 converges to a second minimum at $\mathrm{Re}\,\tau_d \approx -0.372$, related to the first by the residual symmetry

$\tau \to -\tau$ of the modular forms; both minima give identical observables with $\chi^2 < 10^{-19}$. The fit does not distinguish the two minima; we adopt the minimum with $\mathrm{Re}\,\tau_d > 0$ by convention. The quantity χ^2 at the minimum is below double-precision roundoff; we quote it as $\chi^2 < 10^{-19}$.

D Proof of coefficient degeneracy

At fixed τ, multiply all c_i by k:

Table 3: Invariance of χ^2 under $c_i \to k c_i$.

k	χ^2
0.5	$< 10^{-19}$
1.0	$< 10^{-19}$
1.87	$< 10^{-19}$
10.0	$< 10^{-19}$

The residual values differ only at the level of numerical noise; the invariance is exact up to machine precision.

E Flatness of $\mathrm{Re}\,\tau$

Procedure. The flatness scan below was obtained by refitting the remaining parameters at each value of $\mathrm{Re}\,\tau$. The corresponding refit cycle is computationally expensive and is not part of the fast verification script. Scan in $\mathrm{Re}\,\tau_u$: width of $\chi^2 < 10$: $[-0.40, +0.00]$ (0.40). Scan in $\mathrm{Re}\,\tau_d$: width of $\chi^2 < 10$: $[+0.30, +0.50]$ (0.20).

F LOO test

The Leave-One-Out test is not informative for $N_{\mathrm{dof}} \leq 0$: with as many parameters as observables, removing one observable necessarily produces a large shift in the fitted parameters. We report the LOO numbers for completeness, but they should not be interpreted as a measure of predictive stability. For each $j \in \{1, \ldots, 9\}$: remove the j-th observable, predict the j-th. Results: mean $|V_{ub}|$ +16728%, std 11819%.

G Consistency check for the geometric value of $\mathcal{I}$

We test whether the data can distinguish the fitted value $\mathcal{I} = 0.3653$ from the geometric value $\mathcal{I}_{\mathrm{geom}} = (\sqrt{3} - 1)/2 \approx 0.3660$. We fix $\mathcal{I} = \mathcal{I}_{\mathrm{geom}}$ and refit the neutrino Yukawa coupling y_ν to reproduce the observed Δm^2_{31}. The results are summarized in Table 4.

Table 4: Comparison between the baseline fit ($\mathcal{I} = 0.3653$) and the geometric value ($\mathcal{I} = (\sqrt{3}-1)/2$). The baseline values in this table are computed with y_ν refit to reproduce Δm^2_{31} at the same target as the geometric case, following the procedure of this appendix; they differ from the values quoted in Section 8 at the 0.1% level. The maximum relative shift across all predictions is 0.0992%.

Quantity	Baseline	Geometric	Relative change
y_ν	0.93432	0.93409	-0.025%
M_6 [GeV]	2.9646×10^{16}	2.9631×10^{16}	-0.050%
$\sum m_i$ [eV]	0.06763	0.06763	0.000%
$m_{\beta\beta}$ [meV]	8.361	8.361	0.000%
$\mathrm{BR}(\mu \to e\gamma)$	1.0020×10^{-61}	1.0030×10^{-61}	$+0.099\%$
d_e [$e \cdot$ cm]	2.5925×10^{-54}	2.5951×10^{-54}	$+0.099\%$
η_B^{pre}	1.9075×10^{-3}	1.9056×10^{-3}	-0.099%
Δ	3.1270×10^{6}	3.1239×10^{6}	-0.099%

The fit quality is unchanged, and all predictions shift by less than 0.1%. This demonstrates that the data are fully consistent with the geometric value of $\mathcal{I}$. However, this is a consistency check, not a derivation: the fitted moduli are not exactly at the symmetric point, the number of $\mathbb{Z}_4$ fixed points is convention-dependent, and there is no dynamical principle that selects the weighted difference $N_3V_3 - N_4V_4$ over other combinations. We therefore conservatively keep $\mathcal{I}$ as a free parameter in the statistical balance of Section 13.

References

[1] Planck Collaboration, Astron. Astrophys. **641**, A6 (2020).

[2] S. Navas et al. (Particle Data Group), *Review of Particle Physics*, Phys. Rev. D **110**, 030001 (2024).

[3] ADMX Collaboration, Phys. Rev. Lett. **120**, 151301 (2018).

[4] HAYSTAC Collaboration, Nature **590**, 238 (2021).

[5] F. Feruglio, Eur. Phys. J. C **77**, 713 (2017).

[6] T. Kobayashi et al., PTEP **2020**, 053B05 (2020).

[7] P. P. Novichkov et al., JHEP **04** (2019) 005.

[8] G. J. Ding, S. F. King, C. Y. Yao, Phys. Rev. D **104**, 055034 (2021).

[9] M. Fukugita, T. Yanagida, Phys. Lett. B **174** (1986) 45.

[10] I. Esteban et al., JHEP **09** (2020) 178.

[11] KamLAND-Zen Collaboration, Phys. Rev. Lett. **130**, 051801 (2023).

[12] S. Davidson, A. Ibarra, Phys. Lett. B **535** (2002) 25.

[13] S. Kachru et al., Phys. Rev. D **68**, 046005 (2003).

[14] S. Gukov, C. Vafa, E. Witten, Nucl. Phys. B **584** (2000) 69.

[15] B. Pendleton, G. G. Ross, Phys. Lett. B **98**, 291 (1981).