

THIS DOCUMENT HAS BEEN AUTOFINALISED :

Please contact the authoring department/user to verify accuracy

Our Ref: CP/JH/[MED-ID]

[DATE]

Clinic Letter : [DATE]

[PROVIDER-NAME]

[ADDRESS]

[ADDRESS]

[POSTCODE]

Directorate of Medicine and Elderly Care

[ADDRESS]

[ADDRESS]

[ADDRESS]

[ADDRESS]

[ADDRESS]

[ADDRESS]

[POSTCODE]

Telephone: [PHONE] Fax: [PHONE]

Dear [PROVIDER-NAME] [NAME], [ADDRESS]. [POSTCODE]

DOB: [DATE], Hospital Number: [MED-ID], NHS Number: [NHS-NO]

Diagnosis:

Crohn's colitis.

Current Medications:

Colazide 3 tds for two weeks then reduce to 2 bd.

Outstanding Investigations:

Blood tests.

Stool and MC+S.

Follow-up:

Six months or sooner if required.

I reviewed this gentleman in our Gastroenterology clinic today. He returned from a holiday in approximately ten days' ago where he developed an episode of diarrhoea. The whole group that was travelling with developed holiday diarrhoea. They all had bloody diarrhoea and abdominal cramps that lasted several days. This patient's episode of holiday diarrhoea has lasted slightly longer than the rest of the group although I gather some members are still having problems with diarrhoea but he is beginning to feel much better in himself now that the abdominal pain has disappeared. He is now with his bowels open a couple of times a day and passing a solid stool with only a very minimal amount of blood in it now.

His weight is slightly higher than it was on his last visit today at 76kg.

On examination today, his abdomen was soft and non-tender, there were no masses or organomegaly and bowel sounds were normal. He has very sensibly increased his Colazide since he has been having an increasing diarrhoea, but he does not really feel that this is a flare-up of his colitis and is less likely given the history in this situation.

I have given him a sample bottle today for a stool sample to be sent to the microbiology lab. I have also repeated his bloods including his inflammatory markers today and encouraged him to continue with the increased Colazide dose for the next couple of weeks and then reducing it back down to 2 tablets a day, which is his normal maintenance dose. I have arranged to review him in clinic in six months but have asked him to contact us if he does not improve in the next week or so.

I would be grateful if the GP could follow-up the results of his stool MC+S.

Many thanks.

Yours sincerely

[PROVIDER-NAME]
Specialist Registrar to [PROVIDER-NAME]
Consultant Gastroenterologist