

Model Theory of Valued Fields: Midterm

Li Ling Ko
lko@nd.edu

October 15, 2017

Exercise 12: Show that for local rings $R_0 \subseteq R$ we have $\mathfrak{m}_0 = \mathfrak{m} \cap R_0$.

Proof. We are assuming that R_0 is a subring of R .

$\mathfrak{m}_0 \subseteq \mathfrak{m} \cap R_0$ follows by definition of $R_0 \subseteq R$. For the reverse containment, since R_0 is local, it suffices to show that $\mathfrak{m} \cap R_0$ is an ideal of R_0 , and that $\mathfrak{m} \cap R_0$ is not equal to R_0 . To prove the second claim, observe that R_0 cannot be contained in \mathfrak{m} , because as a subring of R , R_0 must contain the identity of R , which does not lie in \mathfrak{m} since \mathfrak{m} is a non-trivial ideal of R . Thus $\mathfrak{m} \cap R_0 \neq R_0$.

To prove the first claim, observe that as rings, both \mathfrak{m} and R_0 are closed under subtraction. As an ideal of R , \mathfrak{m} is closed under multiplication with elements of R_0 . R_0 is also clearly closed under multiplication with elements of R_0 . Thus $\mathfrak{m} \cap R_0$ is an ideal of R_0 . \square