A fixed-target approach to the sharp iid Berry–Esseen inequality*

Bartosz Kołodziejek Kacper Rodziewicz

28 October 2026

Abstract

The conjectured optimal constant in the iid Berry–Esseen inequality is $C_{\mathrm{E}} = (3 + \sqrt{10})/(6\sqrt{2\pi})$. We study this upper bound by maximizing the excess over the fixed target $C_{\mathrm{E}}\mathbb{E}|X|^3$ at a fixed sample size. This choice gives compact-support contact and an exact conservation identity in which excess, moment defect and leakage share one nonnegative budget. An ordered Fourier calculation and elementary moment inequalities then restrict a possible counterexample at $n \geq 42$, under the stated predecessor bounds, to a neighbourhood of a two-point law. We give the analytical steps of this approach, including the bounds and coverage conditions needed by its rational certificates. A continuous contact argument proves half-amplitude confinement for small conditional noise on the acquired probability range $[.40, .53]$: both support rays reduce to a single quintic and elementary derivative signs. A signed resource argument also completes the conditional confinement step on the panel $[.41, .42]$ for its full acquired noise range. The exact conditional-count representation describes the comparison still needed after confinement. A sharp finite base through 41 and a complete local exclusion at every later index remain unproved here. The paper is a self-contained account of the stated reductions, not a proof of the full conjecture.

*This paper was written with substantial assistance from AI systems; see the end of Section 9.

Contents

1 The problem and the approach **3**

2 Fixed-target extremizers and the conserved resource **7**

3 The two-point skeleton and its moment defect **12**

4 From fixed-target contact to a coarse acquisition **16**

5 Fixed-target moments and signed phase resources **29**

6 Conditional confinement in Bernoulli coordinates **34**

7 Conditional counts and discrete curvature **44**

8 Certificates and the remaining global implication **50**

9 Discussion **55**

A A uniform Fourier enclosure with explicit input hypotheses **55**

B Conditional-count estimates **60**

C Binomial normalization and elementary constants **70**

D Signed refinements of the local comparison **72**

E Closed-half-line smoothing **78**

F An original-law reduction for four summands **82**

1 The problem and the approach

The central limit theorem describes the shape of a long sum. The Berry–Esseen inequality asks how far that description can be from the truth at a specified sample size. Even when the summands have the same distribution, the best universal constant is a delicate question: the largest errors come from the interaction of skewness with the jumps of a lattice distribution. A proof of the sharp inequality must control both effects uniformly in the law and in the sample size.

Let $\mathcal{P}$ denote the probability laws on $\mathbb{R}$ satisfying $\mathbb{E}X = 0$, $\mathbb{E}X^2 = 1$, and $\rho(P) := \mathbb{E}|X|^3 < \infty$. For independent $X_1, \dots, X_n$ with common law P, write $S_n = X_1 + \cdots + X_n$ and

$$\Delta_n(P) = \sup_{H \in \mathbb{R}} |\mathbb{P}_P(S_n \le H) - \Phi(H/\sqrt{n})|,$$
$$R_n(P) = \frac{\sqrt{n}\,\Delta_n(P)}{\rho(P)}, \qquad C_n = \sup_{P \in \mathcal{P}} R_n(P), \qquad C^* = \sup_{n \ge 1} C_n.$$

Here Φ and ϕ are the standard normal distribution function and density. The conjecture is

$$C^* = C_{\mathrm{E}}, \qquad C_{\mathrm{E}} = \frac{3 + \sqrt{10}}{6\sqrt{2\pi}} = 0.409732\ldots. \tag{1}$$

Throughout, H is a threshold for the *raw sum* S_n and $H/\sqrt{n}$ its standardized version. All terminating decimals in numerical inequalities mean exact rational numbers.

The normalization costs no generality. If Z has mean μ, variance $\varsigma^2 > 0$, and finite third absolute central moment β_3, apply the definitions to $X = (Z - \mu)/\varsigma$. Then $\rho = \beta_3/\varsigma^3$ and

$$\mathbb{P}\left(\frac{\sum_{i=1}^n Z_i - n\mu}{\varsigma\sqrt{n}} \le x \right) = \mathbb{P}(S_n \le \sqrt{n}\,x).$$

Thus $C_n \le C_{\mathrm{E}}$ is precisely the usual sharp inequality at index n.

1.1 From Berry and Esseen to the sharp-constant problem

Berry [8] and Esseen [9, 10] established quantitative normal approximation of order $n^{-1/2}$ under a finite third absolute moment. Their results initiated the search for the smallest constant in a bound that is uniform over the summand law. Esseen's later moment analysis [11] identified the lower constant in (1). The conjecture that this lower constant is also the universal upper constant is associated with Zolotarev's work of 1966–1967 [31, 32].

The two-point example explains the value of the constant. Put

$$p_* = (4 - \sqrt{10})/2, \qquad q_* = 1 - p_*, \qquad X = (B - p_*)/\sqrt{p_* q_*}, \quad B \sim \mathrm{Bernoulli}(p_*).$$

For this fixed law, $\rho = (p_*^2 + q_*^2)/\sqrt{p_* q_*}$ and the classical lattice asymptotic gives

$$\lim_{n \to \infty} R_n(P) = \frac{3 + q_* - p_*}{6\sqrt{2\pi}(p_*^2 + q_*^2)} = C_{\mathrm{E}}. \tag{2}$$

Indeed $p_*^2 + q_*^2 = 10 - 3\sqrt{10}$ and $(3 + q_* - p_*)/(p_*^2 + q_*^2) = 3 + \sqrt{10}$. Consequently every constant smaller than C_{E} fails for a sufficiently large sum of this one fixed Bernoulli law. This proves $C^* \ge C_{\mathrm{E}}$; it supplies no upper bound for other laws. Hipp and Mattner settled the symmetric binomial case [14]; Schulz's dissertation [19] proves the sharp upper bound for the entire binomial family at every sample size. We use Schulz's theorem for the noiseless case in Section 7.

Upper estimates developed through increasingly precise smoothing and characteristic-function inequalities. Zolotarev's work and van Beek's Fourier analysis [28, 32] are early milestones; Prawitz's one-sided smoothing kernel [17] remains useful in modern computations. Shiganov obtained the iid bound 0.7655 in work originally published in 1982 [24, 27]. Subsequent improvements include Shevtsova's bound 0.7056 [22], the bound 0.5129 of Korolev and Shevtsova [15], Tyurin's bound 0.4785 obtained using zero-bias estimates and characteristic-function bounds [26], and Shevtsova's bounds 0.4748 and 0.4690 [20, 23]. Their structural forms are more informative than the constants alone. For example, the inequalities

$$\sqrt{n}\,\Delta_n(P) \le \min\{0.4690\rho,\ 0.3322(\rho+0.429),\ 0.3031(\rho+0.646)\} \tag{3}$$

hold for every $P \in \mathcal{P}$ and $n \ge 1$ [23]. If $R_n(P)$ is near C_{E}, they bound ρ from above. This supplies the first quantitative moment cap for the support calculation.

1.2 Recent manuscripts, repositories, and formalization

Several recent projects combine analytic inequalities with extensive computation. Xiao and Li report a global iid upper bound of 0.4395; Lin reports 0.423 using a refinement of their analytic framework [16, 29]. Both supply Lean developments and exact finite certificates. Tao and contributors' *Optimization Constants in Mathematics*, especially Problem 19a, records this work and its verification status [25].

He and Cheng's preprint [13] reports the sharp inequality for every

$$n \ge 2\lceil \exp(10^{17}) \rceil,$$

with a threshold independent of the summand law. This uniform quantifier is essential. A theorem for each fixed law, with a law-dependent threshold, would not exclude a sequence of counterexamples whose laws change with n. A finite but astronomically large range of n remains.

Bell's collection treats independent, not necessarily identically distributed, summands. It reports a universal upper bound below 0.44988, results for finite arrays, and sharp bounds when the Lyapunov ratio is small [3–5]; its structural papers study extremizers in that setting [2, 6].

In Lean, Yan's *ProbabilityApproximation* formalizes classical nonuniform and multivariate normal-approximation theorems [30], and the *BerryEsseenLean* project formalizes an eventual-sharpness framework accompanying an earlier version of He and Cheng, relative to explicitly listed external mathematical inputs [1].

1.3 The present approach and its scope

Rather than first finding the largest normalized ratio C_n, fix $c = C_{\mathrm{E}}$ and maximize the excess

$$a(P,H) = \sqrt{n}\{\mathbb{P}_P(S_n \le H) - \Phi(H/\sqrt{n})\} - c\rho(P). \tag{4}$$

The target coefficient stays fixed when the law is varied. If the sharp bound fails at index n, this problem has a maximizing pair with positive excess. The equality case also has an attained witness when $C_n = c$. The negative third-moment penalty is useful: it prevents loss of third moment in a maximizing limit, without assuming uniform integrability of third moments. The contact inequality and this compactness argument are proved in Section 2.

Integrating contact against the zero-bias density connects the excess to a Fourier remainder and the nonnegative defect

$$D = (3+\sqrt{10})\rho - \gamma - 3J.$$

Here $\gamma = \mathbb{E}X^3$ and J measures the actual support gaps, as defined in Section 3. The identity retains the whole term $(n + H^2)a$. It therefore gives one budget for the error excess, the defect and leakage, rather than separate bounds whose best cases need not occur for the same law. The zero-bias transform was introduced by Goldstein and Reinert [12]; related normal-approximation methods appear in Tyurin and, for independent summands, in Bell's transport work [7, 26].

The main quantitative advances concern what happens before the final count comparison. The selected coarse entry uses two quadratic updates and four signed updates. Its Fourier bounds are proved in Appendix A; the finite calculation keeps the sign of the third moment and feeds the defect back into the first moment. Elementary inequalities then give simultaneous moment, gap and noise bounds. For small noise, contact excludes both exterior support rays by one polynomial argument, uniformly in the raw threshold and sample size. For larger noise on $p \in [.41, .42]$, a signed resource calculation completes the same confinement step. The different scopes of these results are stated in Sections 5 and 6.

Confinement is not the final comparison. Conditioning on the number of positive summands gives an exact binomial mixture, and the remaining error must be bounded proportionally to its conditional noise variance, even as that variance tends to zero. Section 7 and Appendix B give the identities and sufficient bounds. They are not accompanied here by a complete acceptance over the remaining domain. The finite base $C_j \le C_{\mathrm{E}}$, $1 \le j \le 41$, is another independent obligation. Section 8 states exactly how these missing pieces would complete the proof and separates mathematical results from arithmetic checks.

An earlier AI-assisted working draft is available on Hexagon [18]. The present paper replaces its main proof organization by the fixed-target argument above. None of the bounds reported in the recent projects cited above is used as a premise of this argument.

1.4 A guide to the proof

A possible counterexample must satisfy two very different demands. Its moments and support must make the error of normal approximation large; its repeated iid convolution must also put enough mass on one side of a threshold. The first demand brings the marginal close to two points. The second is a discrete comparison with a binomial law. The argument below keeps those two tasks separate, and uses contact to pass between them.

Fix the value we want to prove.

At a fixed index, maximize (4) over standardized laws and raw thresholds. The target $c = C_{\mathrm{E}}$ remains fixed. At a first failure above an independently established base through 41, the predecessor bound is $C_{n-1} \le c$. Thus the deleted sum can be estimated with the same coefficient that appears in the extremal problem. No estimate of an unknown maximizing ratio is needed at this stage.

A maximizing limit needs care because third moments need not be uniformly integrable. The moment penalty in the objective supplies the missing inequality. Thresholds are bounded along a maximizing sequence with nonnegative limiting excess, and the closed-event Portmanteau inequality handles moving thresholds and atoms. Contamination by a new atom, followed by centering and scaling, gives contact on the entire closed support. Its cubic term proves compact support before higher moments are used.

Find one budget for the departures from two points.

Write $p = \mathbb{P}(X > 0)$, $q = 1 - p$ and $m = \mathbb{E}X_+ = \mathbb{E}X_-$. The sign decomposition is exactly

$$X = \ell(B_0 + U - p), \qquad B_0 = \mathbf{1}_{\{X>0\}}, \qquad \mathbb{E}(U \mid B_0) = 0, \qquad \ell = m/(pq).$$

The variance of the physical noise $W = \ell U$ is θ; the variance of the noise in a sum of n Bernoulli-coordinate observations is $\lambda = n\mathbb{E}U^2$. These are different quantities.

The gap defect D measures both moment asymmetry and the loss of a single large support gap. Integrating contact against the zero-bias density gives the exact identity in (10). After a raw bound licenses the phase conversion, the resulting nonnegative resource is

$$\mathcal{U} = (n + H^2)a + \frac{n\phi(0)}{6}D + n^{3/2}L.$$

The leakage L is defined using half-open intervals, so an atom at a contact endpoint has its proper weight. The identity must be applied to the same marginal throughout; separate maxima for its costs cannot be added as though they had independent budgets.

Acquire the moment and support bounds in order.

A first support bound gives finite fourth and sixth moments. Two quadratic updates and four signed updates then bound the Fourier remainder and contract the possible moments. A lower bound for m follows from $D \geq \sqrt{10}\rho - \gamma - 6m$; combining it with $4m^2 + \beta^2 \leq 1$ makes the signed updates effective. Each pass uses only bounds already proved. In particular, a density floor comes from the preceding raw ceiling, and a new phase price is used only after its own raw ceiling has been established.

The selected rational calculation retains every actual-law cell and every closed probability overlap. It gives $p \in [.24, .54]$; an elementary moment contraction then gives $p \in [.25, .53]$. These intervals are necessary conditions for a witness, not descriptions of all feasible laws. The Fourier proof and the finite arithmetic have distinct roles: the first explains the implication, the second verifies all of its specified coefficient and coverage gates.

Use contact again to confine the conditional noise.

On the acquired range $p \in [.40, .53]$, the initial strip is $|U| < .6$. If $\lambda \leq .075$, elementary moment bounds give $\theta < .0074$, $\ell > 1.992$ and $\rho < 1.105$. For the negative support ray set $z = p$; for the positive ray set $z = 1 - p$. Both contact bounds reduce to the same quintic on $u = 1.992(z + 1/2) \in [1.7928, 2.1912]$. Its third derivative is negative, its first derivative is concave with positive endpoint values, and its right endpoint is negative. Separate radial derivative signs extend the boundary contradiction along the whole rays. This proves $|U| < 1/2$ without subdividing this probability interval. The two rays are rays of one physical law, not a reflection argument for its CDF.

On $p \in [.41, .42]$, the complementary noise range can also be handled. The sign of H is retained in the phase resource. Noise and endpoint motion consume that same resource before the deletion errors are bounded. Antichain concentration and signed deletion then give the two required ray inequalities. The finite closed cover has 128 sign, threshold and noise boxes: the low-noise lemma handles 16, necessary conditions exclude 84, and both ray tests pass on the remaining 28. This completes confinement on that one panel; it does not complete the count comparison.

Compare the actual convolution with its binomial skeleton.

At $np + H/\ell = k + u$, conditioning on the sign count gives

$$\mathbb{P}(S_n \leq H) = \sum_{j=0}^{n} \binom{n}{j} p^j q^{n-j} \mathbb{P}(V_j \leq k + u - j),$$

where V_j is the sum of the actual conditional noises. Section 7 rewrites the excess over the fixed target exactly as outgoing count mass, incoming count mass and a Gaussian/moment

repair. A sufficiently negative upper bound contradicts nonnegative excess. The needed error bounds must be proportional to λ uniformly down to zero. The marked-crossing and curvature calculations are given in full in Appendix B. Their positivity gates and complete cover remain to be supplied on the new acquired domain. At zero noise the law is Bernoulli and is treated directly.

1.5 Notation and reading order

Symbol	Meaning
$c = C_{\mathrm{E}}$, a	Fixed target and unnormalized excess over that target.
H, $H/\sqrt{n}$	Raw and standardized thresholds.
ρ, γ, β	$\mathbb{E}\lvert X\rvert^3$, $\mathbb{E}X^3$, $\mathbb{E}[X\lvert X\rvert]$.
p, q, m	Positive and nonpositive probabilities; common one-sided first moment.
ℓ, W, U	Conditional-mean span, physical noise and $U = W/\ell$.
θ, s, λ	$\mathbb{E}W^2$, $\mathbb{E}U^2$, ns; $\theta = s/(pq+s)$.
D, L, $\mathcal{U}$	Physical gap defect, leakage, and their joint resource with excess.
$d = d_Y$	Inward displacement in Bernoulli coordinates.
k, u	Integer count and phase, $np + H/\ell = k + u$, $\lvert u\rvert \le 1/2$.
$\zeta = 1 - d_0$	A lower count increment used only in the count/curvature formulas.
$\sigma = \sqrt{10}$	Constant in the moment defect.

The main text follows this order from fixed-target extremality to confinement and counts. The appendices give the selected Fourier proof, the detailed count estimates and their refinements, and a separate original-law reduction for four summands. The latter concerns the missing finite base; it is not an input to the confinement theorem. Historical acquisition schedules are preserved in the repository but are not silently included among the premises of this argument.

2 Fixed-target extremizers and the conserved resource

We optimize the excess above a fixed candidate constant. This gives a variational problem whose coefficient is known before any acquisition step. The positive excess is retained through the contact calculation and the conserved identity; it becomes a nonnegative term in the final resource. No upper bound for the maximizing normalized score is needed. Throughout, sums have independent identically distributed summands with mean zero, variance one and finite third absolute moment ρ. Fix $n \ge 2$, $c > 0$, and a real raw sum threshold H. Set

$$z = H/\sqrt{n}, \quad F = \mathbb{P}(S_n \le H), \quad a = \sqrt{n}\{F - \Phi(z)\} - c\rho,$$
$$\varphi = \phi(z), \quad \beta = \mathbb{E}[X\lvert X\rvert], \quad \gamma = \mathbb{E}X^3.$$

In particular a is not divided by ρ.

2.1 Attainment, contamination and the entire closed support

Proposition 2.1. *For each fixed n and $c > 0$, the supremum of a over these laws and thresholds is attained, and the closed-CDF discrepancy at every attaining pair is positive. At every maximizing pair,*

$$I(y) = n^{3/2}\{h(y) - F\} + ny\varphi + \tfrac{1}{2}H\varphi(y^2-1) + \tfrac{3}{2}c\rho(y^2-1) - c\{\lvert y\rvert^3 - \rho - 3y\beta\} \le 0, \qquad h(y) = \mathbb{P}(S_{n-1} \le H - y), \tag{5}$$

with equality on the whole closed support. That support is compact. Neither statement assumes $a \ge 0$ or $H \ge 0$.

Proof. Write A for the supremum. Since $\rho \geq 1$ and the discrepancy is at most one, $A \leq \sqrt{n}-c$. Symmetric Bernoulli evaluated at $H = n$ gives $A \geq \sqrt{n}(1-\Phi(\sqrt{n}))-c > -c$. For a maximizing sequence, $a_j \to A$ and $c\rho_j \leq \sqrt{n} - a_j$ give a common third-moment ceiling. The discrepancies satisfy $\sqrt{n}(F_j - \Phi(H_j/\sqrt{n})) = a_j + c\rho_j \geq a_j + c > \eta$ eventually for some $\eta > 0$. Chebyshev bounds the negative threshold tail by n/H_j^2; the positive discrepancy is at most the normal tail, also at most n/H_j^2. Thus H_j is bounded.

Tightness gives a subsequence $P_j \Rightarrow P$, $H_j \to H$ and $\rho_j \to b$. The common third-moment ceiling M gives tail bounds $\int_{|x|>K} x^2 dP_j \leq M/K$ and $\int_{|x|>K} |x| dP_j \leq M/K^2$. Bounded continuous truncations therefore preserve mean and variance. Lower semicontinuity gives $1 \leq \rho(P) \leq b$. For fixed n, the sum laws converge weakly. Applying the closed-set inequality first at $H + t$ and then letting $t \downarrow 0$ gives $\limsup F_j(H_j) \leq F_P(H)$, including an atom at H. Consequently

$$a(P, H) \geq A + c\{b - \rho(P)\} \geq A.$$

Maximality forces equality and $\rho(P) = b$. Convergence of third moments is a conclusion here; it was not inferred from bounded third moments. The discrepancy remains positive since $A + c\rho \geq A + c > 0$.

For contact, contaminate the raw marginal by $(1 - \epsilon)P + \epsilon\delta_y$. Its mean is ϵy, variance $v_\epsilon = 1 + \epsilon(y^2 - 1) - \epsilon^2 y^2$, and centered third absolute moment has derivative $|y|^3 - \rho - 3y\beta$ at zero. The last assertion follows by dominated differentiation of $|x - \epsilon y|^3$, with bound $3|y|(|x| + |y|)^2$. Evaluate the standardized law at the standardized image of the fixed raw event $S_n \leq H$. The raw event probability is a polynomial in ϵ, with derivative $n(h(y) - F)$ even at an atom collision. The normal argument is $(H - n\epsilon y)/\sqrt{nv_\epsilon}$, whose derivative is $-\sqrt{n}y - H(y^2 - 1)/(2\sqrt{n})$. The one-sided derivative of the excess is exactly (5); maximality makes it nonpositive.

Its integral against P is zero. All terms are integrable. The closed-event map h is upper semicontinuous; at an atom it takes the larger one-sided value. Thus I is upper semicontinuous. Strict negativity at a closed-support point would hold with a uniform negative margin on a neighborhood of positive P-mass, contradicting its zero integral. Finally $I(y) \leq -c|y|^3 + C_2 y^2 + C_1|y| + C_0$ for constants depending on this pair. It is strictly negative on both sufficiently distant rays, proving compact support. No higher moments were used before this step. □

2.2 Reflection, equality, and the first unresolved index

The sign of the attained maximum detects failure of a strict row bound, including equality. If C_n denotes the supremum of the normalized absolute discrepancy, reflection gives, including an atom at H,

$$\mathbb{P}(-S_n \leq -H) - \Phi(-z) = \Phi(z) - \mathbb{P}(S_n < H) \geq \Phi(z) - F.$$

Thus each negative closed-CDF error is bounded by a positive closed-CDF error of the reflected law; equality at an atom is not assumed. Consequently C_n is also the supremum of positive closed scores r. Since $a = \rho(r - c)$, $C_n < c$ gives $A \leq C_n - c < 0$, and $C_n > c$ gives $A > 0$. If $C_n = c$, choose $r_j \uparrow c$ with $r_j \geq c/2$. Then $\rho_j \leq 2\sqrt{n}/c$, so $\rho_j(r_j - c) \to 0$ and $A = 0$. Hence $C_n \geq c$ supplies an attained pair with $a \geq 0$.

Corollary 2.2 (First failure of a strict bound)**.** *Assume a genuine finite base $C_j \leq c$ for $1 \leq j < N$. If a strict bound $C_n < c$ fails at any index $n \geq N$, there is a least such index, an attained fixed-target pair with $a \geq 0$, and true predecessor bounds $C_j \leq c$ for every $j < n$. The bounds are strict for $N \leq j < n$.*

Proof. The nonempty set of failing indices at least N has a least member. The preceding equivalence supplies its maximizing pair. Indices below N use the finite base; the other earlier indices use minimality. □

Thus an argument excluding every such pair for $n \geq N$ would give strict bounds above the finite base, including exclusion of equality. The subsequent entry argument uses only $C_{n-1} \leq c$; it does not use an upper bound for the actual score $c + a/\rho$. The finite base and the eventual exclusion remain separate obligations.

Equivalently, $\Delta = Q - h \geq 0$ everywhere and $\Delta = 0$ on the closed support, where

$$Q(y) = F + \frac{H\varphi + c\rho}{2n^{3/2}} - \frac{n\varphi + 3c\beta}{n^{3/2}}y - \frac{H\varphi + 3c\rho}{2n^{3/2}}y^2 + \frac{c}{n^{3/2}}|y|^3. \tag{6}$$

2.3 Zero-bias integration and the exact identity

We now allow an arbitrary compactly supported standardized law, retaining the definitions of a, Q, h, Δ above without assuming contact. Put $\nu = \mathcal{L}(H - S_{n-1})$ and $\tau(t) = \mathbb{E}[X\mathbf{1}_{X \geq t}]$. Then $\tau \geq 0$, $\int \tau = 1$. Over the bounded complementary intervals (u, v) of the closed support K, define

$$J = \sum \tau_{u,v}(v-u)^2, \qquad J_1 = \sum \tau_{u,v} \int_u^v (t-u)t\, dt,$$
$$J_2 = \sum \tau_{u,v} \int_u^v (t-u)t|t|\, dt, \qquad L = \sum \tau_{u,v} \int_{[u,v)} (t-u)\, d\nu(t).$$

Here $\tau_{u,v}$ is the constant interior value. Introduce two contact residuals:

$$\xi = \mathbb{E}[X\Delta(X)], \qquad B = \int_K \tau(t)\Delta(t)\, dt + \sum_{(u,v)} \tau_{u,v}(v-u)\Delta(v). \tag{7}$$

The integral over K is a Lebesgue integral, not an expectation under P. It must be retained when the closed support has positive Lebesgue measure. If $K \subset [-R, R]$, then $J \leq 2R$, $|J_1| \leq RJ/2$, $|J_2| \leq R^2 J/2$, and $0 \leq L \leq 2R$. Also Δ is bounded on the convex hull, so the sum in B is absolutely convergent because $\sum \tau_{u,v}(v-u) \leq 1$.

Let $M_4 = \mathbb{E}X^4$, $M_5 = \mathbb{E}|X|^5$, and put

$$K_q = M_5/4 - \rho M_4/2 - 3\beta\gamma/2 + \rho/2, \qquad K_1 = \mathbb{E}[X|X|^3] - 3\beta - 3\rho\gamma/2,$$

$$\Omega = J_1 + (M_4 - 3)/6, \quad \varepsilon = \mathbb{E}q_z(S_n/\sqrt{n}) - \mathbb{E}q_z(Z_0) - 2\gamma\varphi/(3\sqrt{n}), \quad q_z(x) = (x-z)|x-z|,$$

where Z_0 is standard normal. Define $E_n = -n^{3/2}\varepsilon/2$ and

$$\mathcal{R}_c = c(3\beta J/2 + 3\rho J_1 - 3J_2 - K_q) + H(cK_1 + \varphi\Omega) - H^2(c\rho + \gamma\varphi/2) + E_n. \tag{8}$$

Theorem 2.3 (Conservation with contact residuals)**.** *For every such law, every $H \in \mathbb{R}$ and every $c > 0$,*

$$(n + H^2)a + nc\rho - \frac{n\varphi}{6}(\gamma + 3J) + n^{3/2}L = \mathcal{R}_c + n^{3/2}(B - H\xi). \tag{9}$$

At a fixed-target maximizing pair, $B = \xi = 0$. In particular,

$$(n + H^2)a + nc\rho - \frac{n\varphi}{6}(\gamma + 3J) + n^{3/2}L = \mathcal{R}_c. \tag{10}$$

The coefficient of the unnormalized excess is $n + H^2$.

Proof. There are two direct evaluations of $\mathcal{A} = \int \tau\Delta$. Since $h(t) = \nu([t,\infty))$, one has $h(t) = h(v) + \nu([t,v))$ inside a gap. Thus bounded Fubini gives

$$\int_u^v \Delta(t)dt = (v-u)\Delta(v) + \int_u^v (t-u)(-Q'(t))dt - \int_{[u,v)} (t-u)d\nu(t).$$

An atom at v belongs to $h(v)$ and is not in the last integral; an atom at u is included but multiplied by zero. Summing, including the support integral in (7), gives

$$\mathcal{A} = \frac{\varphi J}{2\sqrt{n}} + \frac{3c\beta J/2 + (H\varphi + 3c\rho)J_1 - 3cJ_2}{n^{3/2}} - L + B. \tag{11}$$

For the second evaluation, signwise Fubini proves $\int \tau g' = \mathbb{E}[Xg(X)]$ for absolutely continuous g with bounded derivative on the relevant compact range. In particular

$$\int t\tau = \gamma/2, \quad \int t^2\tau = M_4/3, \quad \int |t|^3\tau = M_5/4.$$

Apply that same identity conditionally to $g_s(x) = \min(s+x, H)$ and then use exchangeability. The choice of derivative at its single corner is immaterial under the density τ. It follows directly that

$$\int \tau(t)h(t)dt = \mathbb{E}[X_n \min(S_n, H)] = \frac{1}{n}\mathbb{E}[S_n \min(S_n, H)] = T,$$

where $T = \mathbb{E}[(Z^2 - zZ)\mathbf{1}_{Z\le z}]$, $Z = S_n/\sqrt{n}$. The last expression vanishes at an endpoint atom. Integrating (6) against τ therefore yields

$$\mathcal{A} = F - \frac{\gamma\varphi}{2\sqrt{n}} + \frac{H\varphi(3-M_4)/6 + cK_q}{n^{3/2}} - T. \tag{12}$$

Set $M = \mathbb{E}[Z\mathbf{1}_{Z\le z}]$. Since $\mathbb{E}[Xh(X)] = M/\sqrt{n}$, integration of $XQ(X)$ against P gives

$$M + \varphi = -\frac{\gamma H\varphi}{2n} + \frac{cK_1}{n} - \sqrt{n}\xi. \tag{13}$$

The pointwise identity $q_z(x) = (x-z)^2 - 2(x-z)^2\mathbf{1}_{x\le z}$, with the same identity for a normal variable, gives

$$T - \Phi(z) = -\varepsilon/2 - \gamma\varphi/(3\sqrt{n}) + z(M+\varphi) - z^2(F - \Phi(z)).$$

Inserting (13) and $\sqrt{n}(F-\Phi(z)) = c\rho + a$ into (12) gives

$$\begin{aligned}\sqrt{n}\mathcal{A} &= c\rho + a - \gamma\varphi/6 + P_0/n + \sqrt{n}\varepsilon/2 + H\sqrt{n}\xi,\\ P_0 &= H^2(c\rho + a + \gamma\varphi/2) - H\{cK_1 + \varphi(M_4-3)/6\} + cK_q.\end{aligned}$$

Equate this with (11) multiplied by $\sqrt{n}$, then multiply by n and collect terms. This is (9). Under closed-support contact, every term defining B and ξ vanishes. □

The compensated Fourier representation of this same E_n is proved in Appendix A. Its comparator has transform $e^{-nt^2/2}(1 - in\gamma t^3/6)$. The residual identity also explains why applying the residual-free formula to arbitrary laws would be invalid. For example, let X be uniform on $[-\sqrt{3}, \sqrt{3}]$ and take $H = 0$. There are no support gaps, $F = T = 1/2$, $\beta = \gamma = 0$, and $K_q = 3\sqrt{3}/40$. Therefore $B = \int_K \tau\Delta = cK_q/n^{3/2} > 0$. Omitting the support integral would give a false identity, even though the gap sums are all empty. This is an off-contact negative control, not a counterexample to the contact theorem.

2.4 A positive resource and its ordered phase price

Put $m = \mathbb{E}X_+ = \mathbb{E}X_- > 0$ and $\sigma = \sqrt{10}$. Signwise Cauchy gives $a_\pm^2 \le m b_\pm$ and $m^2 \le a_+ a_-$, where $a_\pm = \mathbb{E}X_\pm^2$ and $b_\pm = \mathbb{E}X_\pm^3$. Therefore $b_+ b_- \ge m^2$, or $\gamma^2 + 4m^2 \le \rho^2$. A gap on one side of zero has $(v-u)^2 \le 2\int_u^v |t|dt$. If a central gap $(-A, B)$ exists, its extra contribution is $2mAB$, and $0 \le \mathbb{E}[(X+A)(X-B)] = 1 - AB$. Signwise Tonelli then gives $J \le \rho + 2m$. It follows that

$$D = (3+\sigma)\rho - \gamma - 3J = 3(\rho + 2m - J) + (\sigma\rho - \gamma - 6m) \ge 0. \tag{14}$$

The last term is nonnegative by Cauchy in two dimensions. These statements hold with or without an atom at zero.

Specialize to $c = C_{\mathrm{E}} = (3+\sigma)\phi(0)/6$ and contact. With $w = H^2/(2n)$, $e = e^{-w}$, the identity becomes

$$\mathcal{R}_c = (n + H^2)a + nc\rho(1-e) + \frac{n\phi(0)e}{6}D + n^{3/2}L. \tag{15}$$

If $a \ge 0$, every term is nonnegative. This specialization to C_{E} is essential for this definition of D.

Corollary 2.4 (Raw bound, density, and phase price)**.** *At such a contact pair with $a \ge 0$, assume $n \ge N \ge 2$, $\rho \ge l > 0$, $0 < c_- \le c$, and a proved raw bound $\mathcal{R}_c \le T < Nc_-l$. A negative T excludes the pair. Otherwise*

$$\gamma + 3J > 0, \qquad e \ge 1 - T/(Nc_-l) > 0,$$
$$\mathcal{U} := (n+H^2)a + n\phi(0)D/6 + n^{3/2}L \le \mathcal{R}_c,$$
$$\mathcal{U} \le \mathcal{R}_c - \tfrac{1}{2}(c_-l - T/N)H^2.$$

No condition $D < 2$ or own-score upper bound is required.

Proof. Set $M_c = c\rho - \phi(0)D/6 = \phi(0)(\gamma + 3J)/6$. The general master identity gives

$$eM_c = (1 + H^2/n)a + c\rho + \sqrt{n}L - \mathcal{R}_c/n \ge c\rho - T/n > 0.$$

Thus $M_c > 0$ and, since $D \ge 0$, $M_c \le c\rho$. The density bound follows. Regrouping (15) gives $\mathcal{R}_c = \mathcal{U} + n(1-e)M_c$, proving the unweighted bound. Finally

$$n(1-e)M_c = n(e^w - 1)eM_c \ge \frac{H^2}{2}(c\rho - T/n) \ge \frac{H^2}{2}(c_-l - T/N).$$

This prices one term once. It is a bound on $\mathcal{U}$, not an improved bound on the raw remainder. □

Every conclusion holds at each actual index and raw threshold. Common bounds on a closed moment cell make the result uniform even when the maximizing law varies with n. There is no exchange of a limit with contact, support gaps, moments, or an integral. In particular, the order of use is essential: a raw ceiling first proves $M_c > 0$ and a density floor; only then may the Gaussian phase term be subtracted to improve $\mathcal{U}$. A smaller resource ceiling is not a replacement for the raw ceiling in that licensing step.

2.5 Two elementary moment domains

The moment inequalities below are used before any finite-dimensional relaxation. They hold for the actual marginal, without a bounded-support assumption. In particular, they are unaffected by adding a very small atom far from the origin.

Theorem 2.5 (Signed and absolute third moments)**.** *If* $\mathbb{E}X = 0$, $\mathbb{E}X^2 = 1$, $\rho = \mathbb{E}|X|^3 < \infty$, *and* $\gamma = \mathbb{E}X^3$, *then*

$$\rho \geq \frac{\gamma^2 + 2}{\sqrt{\gamma^2 + 4}}.$$

This is the sharp moment relation of Shevtsova [21]; in the cited arXiv version its equivalent form is Theorem 2. We include a factorization proof to specify the domain used by the finite certificates.

Proof. Set $s = \sqrt{\gamma^2 + 4}$, $a = (s - \gamma)/2$, $b = (s + \gamma)/2$ and $c = \gamma(\gamma^2 + 6)/s^3$. Then $a, b > 0$, $ab = 1$ and $a + b = s$. Define

$$w(y) = |y|^3 - cy^3 - \frac{12y^2}{s^3} + \frac{6\gamma y}{s^3} + \frac{4}{s^3}.$$

For $y, t \geq 0$, direct multiplication gives

$$w(y) = \frac{2a^2}{s^3}(y - b)^2\{2 + (a + 3b)y\}, \qquad w(-t) = \frac{2b^2}{s^3}(t - a)^2\{2 + (3a + b)t\}.$$

Thus $w \geq 0$ on the whole real line. Taking expectations and using the first three signed moments gives

$$0 \leq \mathbb{E}w(X) = \rho - c\gamma - 8/s^3 = \rho - (\gamma^2 + 2)/s.$$

All terms are integrable under the stated third-moment hypothesis. □

Lemma 2.6 (The signed second moment)**.** *With the same normalization and* $\beta = \mathbb{E}[X|X|]$,

$$|\beta| < 1, \qquad \rho \geq \frac{1 + \beta^2}{\sqrt{1 - \beta^2}}.$$

Proof. Both signs have positive probability. Put $m = \mathbb{E}X_+ = \mathbb{E}X_- > 0$ and $a_\pm = \mathbb{E}X_\pm^2 = (1 \pm \beta)/2 > 0$. If $p_\pm = \mathbb{P}(\pm X > 0)$, Cauchy's inequality gives $m^2 \leq p_\pm a_\pm$. Since $p_+ + p_- \leq 1$, this implies

$$m^2(1/a_+ + 1/a_-) \leq 1, \qquad 2m \leq \sqrt{1 - \beta^2}.$$

Another application of Cauchy's inequality, to $X_\pm^{1/2}$ and $X_\pm^{3/2}$, gives $m\mathbb{E}X_\pm^3 \geq a_\pm^2$. Summing yields $m\rho \geq (1 + \beta^2)/2$, which proves the claim. A possible zero atom enters only through $p_+ + p_- \leq 1$ and causes no exception. □

3 The two-point skeleton and its moment defect

Every centered law can be separated into its positive and nonpositive parts. Replacing each part by its conditional mean gives a two-point law; the remaining conditional fluctuations measure its departure from that law. We first make this decomposition exact, then identify a moment defect that vanishes at Esseen's extremal distribution.

3.1 The sign split and its units

Fix $P \in \mathcal{P}$ and put

$$p = \mathbb{P}(X > 0), \quad q = 1 - p, \quad v = pq, \quad m = \mathbb{E}X_+ = \mathbb{E}X_- > 0, \quad B_0 = \mathbf{1}_{\{X>0\}}.$$

Both p, q are positive; a possible atom at zero is assigned to the nonpositive class. Define $V = \mathbb{E}[X \mid B_0]$ and $W = X - V$, so that

$$V = -m/q \quad (B_0 = 0), \qquad V = m/p \quad (B_0 = 1), \qquad \theta = \mathbb{E}W^2 = 1 - m^2/v < 1.$$

Write $\theta_- = \mathbb{E}[W^2; B_0 = 0]$ and $\theta_+ = \mathbb{E}[W^2; B_0 = 1]$. The distance between the two conditional means is

$$\ell = m/v = \sqrt{(1-\theta)/v} > 0.$$

Set $U = W/\ell$ and $Y = B_0 + U$. Then

$$X = \ell(Y - p), \qquad \mathbb{E}[U \mid B_0] = 0. \tag{16}$$

Let U_0, U_1 have the two conditional laws, and put

$$w_i = \mathbb{E}U_i^2, \quad \mu_i = \mathbb{E}U_i^3, \quad s = qw_0 + pw_1 = \frac{v\theta}{1-\theta}, \quad \lambda = ns, \quad \ell = (v+s)^{-1/2}.$$

The moments μ_i are finite because $\rho < \infty$. Higher moments are used only for extremal laws, which have compact support by Proposition 2.1.

If $s = 0$, both nonnegative conditional variances vanish, so $U_0 = U_1 = 0$ almost surely. This is exactly the standardized Bernoulli law. If $s > 0$, the parameter $\alpha = pw_1/s$ lies in $[0, 1]$ and gives $w_0 = s(1-\alpha)/q$, $w_1 = s\alpha/p$.

The signs of X are still those defining the two classes, so expansion of the actual conditional moments gives

$$\begin{aligned}
\rho/\ell^3 &= v(1-2v) + 3v(w_0 + w_1) - q\mu_0 + p\mu_1, \\
\gamma/\ell^3 &= v(q-p) + 3v(w_1 - w_0) + q\mu_0 + p\mu_1, \\
\beta/\ell^2 &= v(q-p) + pw_1 - qw_0.
\end{aligned} \tag{17}$$

For example, on the nonpositive class $|X|^3 = \ell^3(p - U_0)^3$; on the positive class it is $\ell^3(q + U_1)^3$. Conditional centering removes the linear terms and gives the first identity; the other two follow in the same way. The three identities share the same pair (μ_0, μ_1).

3.2 A nonnegative moment defect

The constant $3 + \sqrt{10}$ has a simple geometric explanation. The signed third moment and the first absolute moment satisfy a circle inequality; the lengths of the support gaps are controlled by the same first absolute moment. Neither fact requires bounded support.

Define

$$\tau(t) = \mathbb{E}[X\mathbf{1}_{\{X \geq t\}}], \qquad \sigma = \sqrt{10}.$$

For $t < 0$, centering gives $\tau(t) = -\mathbb{E}[X\mathbf{1}_{\{X<t\}}] \geq 0$; for $t \geq 0$ nonnegativity is immediate. Signwise Tonelli integration gives

$$\int_{\mathbb{R}} \tau(t)\, dt = 1, \qquad 2\int_{\mathbb{R}} |t|\tau(t)\, dt = \rho. \tag{18}$$

On each bounded complementary interval (a,b) of the closed support, τ is constant in the interior, with value $\tau_{a,b}$. Put

$$J=\sum_{(a,b)}\tau_{a,b}(b-a)^2,\qquad D=(3+\sigma)\rho-\gamma-3J. \tag{19}$$

There are at most countably many gaps; the next proof shows that the nonnegative sum is finite. Endpoint atoms do not affect its integrals.

Lemma 3.1 (A moment circle and the gap inequality)**.** *For every $P\in\mathcal{P}$,*

$$\gamma^2+(2m)^2\le\rho^2,\qquad J\le\rho+2m,\qquad \gamma+tJ\le\bigl(t+\sqrt{1+t^2}\bigr)\rho\quad(t>0). \tag{20}$$

The last constant is sharp. Equality holds exactly at the standardized Bernoulli law with positive mass $p_t=(1+t-\sqrt{1+t^2})/2$. In particular, $D\ge 0$. If zero belongs to the support, then $D\ge(\sigma-1)\rho$; hence $D<(\sigma-1)\rho$ implies a central support gap $(-A,B)$ with $A,B>0$.

Proof. Write $a_\pm=\mathbb{E}X_\pm^2$ and $b_\pm=\mathbb{E}X_\pm^3$, so $a_++a_-=1$, $b_++b_-=\rho$ and $b_+-b_-=\gamma$. Cauchy–Schwarz on each sign class gives $a_\pm^2\le mb_\pm$. It also gives $m^2\le pa_+$ and $m^2\le qa_-$, where an atom at zero is included in q. Thus $m^2(1/a_++1/a_-)\le 1$, or $m^2\le a_+a_-$. Consequently $b_+b_-\ge(a_+a_-)^2/m^2\ge m^2$, proving the circle inequality since $\rho^2-\gamma^2=4b_+b_-$.

Every gap on one side of zero satisfies $(b-a)^2\le 2\int_a^b|x|\,dx$. If a central gap $(-A,B)$ exists, its weight is m and its excess over this integral is $2mAB$. Moreover $0\le\mathbb{E}[(X+A)(X-B)]=1-AB$. Summing the gap bounds by Tonelli proves $J\le\rho+2m$. If zero is in the support there is no central gap, so $J\le\rho$. Finally Cauchy–Schwarz in $\mathbb{R}^2$ gives $\gamma+2tm\le\sqrt{1+t^2}\rho$, proving the family and the assertions about D.

Equality in the circle forces equality in the two signwise Cauchy bounds: the positive and negative parts are each concentrated at one nonzero value. Equality in $m^2\le a_+a_-$ excludes mass at zero. For such a law, with positive mass p and $q=1-p$, one has $J=\rho+2m$. Equality in the last Cauchy bound requires $\gamma=2m/t$, equivalently $t(1-2p)=2p(1-p)$. Its unique solution in $(0,1)$ is the stated p_t. □

Corollary 3.2 (The zero-defect law)**.** *For $P\in\mathcal{P}$, $D=0$ if and only if*

$$X=(B_0-p_*)/\sqrt{p_*q_*},\qquad p_*=(4-\sqrt{10})/2,\quad q_*=1-p_*,\quad B_0\sim\text{Bernoulli}(p_*).$$

Proof. Apply the equality case of Lemma 3.1 with $t=3$. □

In particular the defect separates two losses:

$$D=\underbrace{\sigma\rho-\gamma-6m}_{\text{moment loss}}+3\underbrace{(\rho+2m-J)}_{\text{gap loss}},$$

each nonnegative. For quantitative estimates we need more information than their signs. Suppose a central gap $(-A,B)$ has been established, and write

$$T_+=\mathbb{E}X_+^3-mB^2,\quad T_-=\mathbb{E}X_-^3-mA^2,\qquad R_+=\mathbb{E}X_+^2-mB,\quad R_-=\mathbb{E}X_-^2-mA.$$

These four quantities are nonnegative. For the outer gaps put $J_o=J-m(A+B)^2$. A positive outer gap satisfies $(b-a)^2\le 2\int_a^b(t-B)\,dt$, so Tonelli gives

$$J_o^+\le 2\int_B^\infty(t-B)\tau(t)\,dt=\mathbb{E}[X_+(X_+-B)^2]=T_+-2BR_+\le T_+.$$

Similarly $J_o^- \le T_- - 2AR_- \le T_-$. Expanding D, using $\sqrt{\sigma+1}\sqrt{\sigma-1} = 3$, gives

$$D = m\{\sqrt{\sigma+1}A - \sqrt{\sigma-1}B\}^2 + (\sigma-1)T_+ + (\sigma+1)T_- + 3(T_+ + T_- - J_o). \tag{21}$$

Every term is nonnegative. Cauchy–Schwarz also gives $1/(2\rho) \le m \le 1/2$: indeed $1 \le \rho\mathbb{E}|X| = 2m\rho$ and $\mathbb{E}|X| \le 1$. Since $\sigma - 1 > 2$, the decomposition yields $T_+ + T_- \le D/2$. If $D < 2$, then

$$m(A^2 + B^2) = \rho - T_+ - T_- \ge 1 - D/2, \qquad A + B \ge \sqrt{2 - D}.$$

3.3 One joint budget for endpoint motion and noise

Let $\bar{A} = m/q$, $\bar{B} = m/p$ be the conditional mean magnitudes, and let $\Delta_A = \bar{A} - A$, $\Delta_B = \bar{B} - B$. They are nonnegative.

Lemma 3.3 (Simultaneous gap inequality)**.** *For this central gap,*

$$D \ge \frac{(6+2\sigma)(1-\theta)^{3/2}}{\sqrt{pq}}(p - p_*)^2 + 6m\{(A+B)(\Delta_A + \Delta_B) + \Delta_A\Delta_B\} + 3(\sigma+1)B\theta_+ + 3(\sigma+3)A\theta_-. \tag{22}$$

Proof. The stronger outer-gap bound in the preceding proof gives

$$D \ge (\sigma-1)\mathbb{E}X_+^3 + (\sigma+1)\mathbb{E}X_-^3 - 6mAB + 6BR_+ + 6AR_-.$$

Conditional cubic expansion gives

$$\mathbb{E}X_+^3 = m\bar{B}^2 + \mathbb{E}[(X + 2\bar{B})(X - \bar{B})^2; X > 0] \ge m\bar{B}^2 + 3B\theta_+,$$

and similarly $\mathbb{E}X_-^3 \ge m\bar{A}^2 + 3A\theta_-$. Moreover $R_+ = \theta_+ + m\Delta_B$ and $R_- = \theta_- + m\Delta_A$. Substitution produces the last two terms of (22) and the deterministic expression

$$m\{(\sigma-1)\bar{B}^2 + (\sigma+1)\bar{A}^2 - 6\bar{A}\bar{B}\} + 6m\{\bar{A}\bar{B} - AB + B\Delta_B + A\Delta_A\}.$$

The second brace equals $(A+B)(\Delta_A + \Delta_B) + \Delta_A\Delta_B$. For the first, use $m^2 = (1-\theta)pq$ and the exact identity

$$(\sigma+1)p^2 + (\sigma-1)q^2 - 6pq = (6+2\sigma)(p - p_*)^2.$$

This proves the displayed inequality. □

In count coordinates the inward displacements are

$$a_0 = p - A/\ell = \Delta_A/\ell, \quad a_1 = q - B/\ell = \Delta_B/\ell, \quad d_Y = a_0 + a_1 = (\Delta_A + \Delta_B)/\ell.$$

Thus $U_0 \le a_0$ and $U_1 \ge -a_1$. The physical displacement $\Delta_A + \Delta_B$ and the count displacement d_Y differ by ℓ.

Theorem 3.4 (Weighted joint resource)**.** *Suppose $n \ge N > 0$ and*

$$nD \le Nd_D, \quad \theta \le \Theta < 1, \quad pq \le v_*, \quad D \ge b + 6mS_*(\Delta_A + \Delta_B) + K\theta,$$

where $b \ge 0$, $S_* > 0$, $K > 0$ and $v_* > 0$. Put $r = d_D - b$. If $r < 0$ there is no such law. If $r = 0$, then $d_Y = \lambda = 0$. If $r > 0$, define

$$d_* = \frac{r}{6(1-\Theta)S_*}, \qquad \Lambda = \frac{Nv_*r}{K(1-\Theta)}.$$

Then

$$\frac{d_Y}{d_*} + \frac{\lambda}{\Lambda} \le 1. \tag{23}$$

Proof. Since $m\ell = 1-\theta$ and $n\theta = (1-\theta)\lambda/(pq)$,

$$\begin{aligned} Nd_D \ge nD &\ge nb + 6n(1-\theta)S_* d_Y + K(1-\theta)\lambda/(pq) \\ &\ge Nb + 6N(1-\Theta)S_* d_Y + K(1-\Theta)\lambda/v_*. \end{aligned}$$

All weakened coefficients multiply nonnegative quantities. Subtracting Nb proves the impossible and zero cases without division. For $r > 0$, division by Nr proves (23). □

For instance, on a closed probability interval I, established endpoint floors $A \ge A_l > 0$, $B \ge B_l > 0$ and $\theta \le \Theta$ supply

$$S_* = A_l + B_l, \qquad K = \min\{3(\sigma+1)B_l, 3(\sigma+3)A_l\},$$

$$b = \frac{(6+2\sigma)(1-\Theta)^{3/2}}{\sqrt{v_*}}\operatorname{dist}(p_*, I)^2, \qquad v_* \ge \max_{p\in I} p(1-p).$$

These choices follow from Lemma 3.3 by dropping the nonnegative product $6m\Delta_A\Delta_B$. Rational lower enclosures for b, K, S_* and an upper enclosure for v_* preserve the implication.

On a band $L \le \lambda \le U \le \Lambda$, the joint plane gives $d_Y \le d_*(1 - L/\Lambda)$. Either intercept may be replaced by a larger value.

4 From fixed-target contact to a coarse acquisition

We now derive quantitative support and moment bounds, then explain the finite enclosure that propagates them. The selected calculation has two coarse quadratic passes followed by four signed-moment passes. Every improvement is used only after its hypotheses have been established for the same attaining law. The final output is an acquisition bound, not a local exclusion or a proof of the conjecture. Let P be a centered, unit-variance law, S_j a sum of j independent copies, and set

$$\rho = \mathbb{E}|X|^3, \quad \beta = \mathbb{E}[X|X|], \quad \gamma = \mathbb{E}X^3, \quad c = C_{\mathrm{E}} = \frac{3+\sqrt{10}}{6\sqrt{2\pi}}.$$

The threshold H always has the units of the raw sum. A fixed-target maximizer maximizes

$$a(P,H) = \sqrt{n}\{\mathbb{P}(S_n \le H) - \Phi(H/\sqrt{n})\} - c\rho.$$

We use Proposition 2.1 and its closed-support contact formula (5). In particular its closed-support contact equality, not ratio optimality, is the variational premise here. We consider a maximizer with $a \ge 0$. The actual normalized score is $c + a/\rho$; no upper bound on it is assumed.

4.1 The first support bound

Theorem 4.1 (Initial moment and support entry)**.** *Assume $n \ge 42$, $C_{n-1} \le c$, and the universal structural inequality*

$$\sqrt{n}\,\Delta_n(P) \le \frac{3031}{10000}\left(\rho + \frac{323}{500}\right) \tag{24}$$

for standardized laws with finite third moment. Then a fixed-target maximizer with $a \ge 0$ satisfies

$$1 \le \rho < \frac{7361}{4000} < \frac{1841}{1000}, \qquad |\beta| < \frac{321}{500}, \qquad \operatorname{supp} P \subset (-96/25, 96/25). \tag{25}$$

The theorem is uniform in all real H and in both signs of a support point. It requires neither C_{n-2} nor an excess bound of order $1/n$.

The structural inequality (24) is the external iid theorem of Shevtsova [23]. Only this affine bound is needed for the displayed moment radius; it is not a consequence of the finite arithmetic in this section.

Proof. Since $c\rho \le \sqrt{n}\Delta_n(P)$ and $c > 819/2000$, elementary division gives

$$\rho < \frac{(3031/10000)(323/500)}{819/2000 - 3031/10000} = \frac{7361}{4000}.$$

The other affine bound $0.3322(\rho + 0.429)$ alone would give the larger number $712569/386500$; it must not be used to claim $7361/4000$. Put $m = \mathbb{E}X_+ = \mathbb{E}X_- > 0$ and $a_\pm = \mathbb{E}X_\pm^2 = (1 \pm \beta)/2 > 0$. Cauchy on each sign class gives $m\rho \ge a_+^2 + a_-^2$ and $m^2 \le a_+a_-$. Hence

$$\rho \ge \frac{1+\beta^2}{\sqrt{1-\beta^2}}.$$

This function increases with $|\beta|$. For $r = 1841/1000$, $b = 321/500$,

$$(1+b^2)^2 - r^2(1-b^2) = \frac{93288849}{50000000000} > 0,$$

which proves the second assertion without assuming bounded support.

Write $v = n-1$, $s = \sqrt{n/v}$, $F_v(x) = \Phi(x/\sqrt{v})$ and $e_n = n(s-1) - 1/2$. Closed-support contact, the upper predecessor bound $F_{n-1,P}(H-t) \le F_v(H-t) + c\rho/\sqrt{v}$, and $\sqrt{n}(F_{n,P}(H) - \Phi(H/\sqrt{n})) = c\rho + a$ give

$$0 \le H_n(H,t) + c\{-|t|^3 + \tfrac{3}{2}\rho t^2 + 3\beta t + \rho e_n\} - na \quad (t \in \operatorname{supp} P), \tag{26}$$

where exactly

$$\frac{H_n(H,t)}{n^{3/2}} = F_v(H-t) - F_{v+1}(H) + tF'_{v+1}(H) - \frac{t^2-1}{2}F''_{v+1}(H).$$

The extra $-na$ can be discarded in this upper bound. This is the precise reason the predecessor assumption is $C_{n-1} \le c$, not a comparison with the actual score. No continuity assumption on the distribution function at $H - t$ is inserted.

Rationalization gives

$$ne_n = \frac{s^2(2s+1)}{2(s+1)^2}, \qquad s < 81/80 =: S, \qquad ne_n < \frac{793881}{2073680} < \frac{48}{125}.$$

The first expression increases for $s \ge 1$, as its logarithmic derivative is $2/s + 2/(2s+1) - 2/(s+1) > 0$. Thus $e_n < 8/875$.

For completeness we rederive the Gaussian estimate used at the radius. The heat identity $\partial_v F_v = F_v''/2$ permits spatial expansion of $F_v(H-t)$ to order four, variance expansion of F_{v+1} to order two, and variance expansion of F'_{v+1}, F''_{v+1} to order one. The cubic coefficient is $(-t^3+3t)/6$, and the quartic coefficient is $(t^4-6t^2+3)/24$. Therefore, for $t = \pm u$, $z_v = H/\sqrt{v}$,

$$H_n(H, \pm u)/s^3 = \pm A(1-z_v^2)\phi(z_v) + B\phi'''(z_v) + \mathcal{E}, \tag{27}$$

where $A = (u^3 - 3u)/6$, $H_4 = u^4 - 6u^2 + 3$ and $B = H_4/(24\sqrt{v})$. The four error bounds after division by s^3 are respectively

$$\frac{u^5\|\phi^{(4)}\|_\infty}{120v}, \quad \frac{\|\phi^{(5)}\|_\infty}{48v^{3/2}}, \quad \frac{u\|\phi^{(4)}\|_\infty}{8v}, \quad \frac{|u^2-1|\|\phi^{(5)}\|_\infty}{16v^{3/2}}.$$

Gaussian Fourier inversion gives norms at most $3\phi(0) < 6/5$ and $8/\pi < 51/20$. Since $v \geq 41$ and $\sqrt{v} > 6$, for $u \geq 1$ this gives

$$|\mathcal{E}| < E(u) := u^5/4100 + 3u/820 + 17(3u^2 - 2)/78720.$$

These errors are uniform in H; they use no higher moment of P.

If $A, B > 0$ and $B \leq A/4$, the leading part of (27) is at most $\phi(0)(A + 9B^2/(4A))$ for both signs. Indeed, for the positive sign and $|z_v| \leq 1$, it is at most $\phi(0)(A - Az_v^2 + 3B|z_v|)$, and complete the square. Outside that interval the first term is nonpositive. For the negative sign the first term is at most $2e^{-3/2}A\phi(0) < A\phi(0)/2$. In both cases use $\|\phi'''\|_\infty < 2\phi(0)$: on $[0, \sqrt{3}]$ the polynomial $3x - x^3$ is at most two, and beyond $\sqrt{3}$ use $x^3 e^{-x^2/2} \leq 3\sqrt{3}e^{-3/2} < 2$.

For frozen r, b above define

$$Q(u) = -u^3 + \tfrac{3}{2}ru^2 + 3bu + (8/875)r,$$
$$P(u) = \tfrac{2}{5}S^3\left(A^2 + \frac{H_4^2}{256 \cdot 41}\right) + S^3E(u)A + \frac{819}{2000}Q(u)A.$$

At $R = 96/25 < 6$ we have $A > 0$, $0 < H_4 < 36A$, hence $B < A/4$. The sign $Q(R) < 0$ is checked before replacing c by its lower endpoint. The upper bound in (26), multiplied by A, is then strictly below $P(R)$, and exact arithmetic gives

$$-P(R) = \frac{2214258283536204102723}{10250000000000000000000} > \frac{1}{5}.$$

To cover each entire exterior ray, differentiation of the exact Gaussian expression gives

$$|\partial_t H_n(H, t)| < \tfrac{2}{5}(t^2/2 + 1/2 + e_n).$$

For verification put $y = (H - t)/\sqrt{n}$, $z = H/\sqrt{n}$; the derivative is $n\phi(z) + Ht\phi(z) - ns\phi(sy)$. The second-order Taylor error between $\phi(y)$ and $\phi(z)$ costs $\phi(0)t^2/2$, while $\partial_s[s\phi(sy)] = -\phi''(sy)$ costs $\phi(0)n(s-1)$. Define $L(u) = \frac{2}{5}(u^2/2 + 1/2 + 8/875) + (819/2000)Q'(u)$. At R, the six numbers Q, Q', Q'', L, L', L'' are strictly negative. Since $Q''' = -6$ and $L'' = 2/5 - 6(819/2000) < 0$, all their required signs persist for $u \geq R$. Thus both radial upper bounds decrease from a strictly negative boundary value. The contact inequality excludes every support point with $|t| \geq R$. □

4.2 Higher moments and an actual Fourier event

Higher moments are used only after Theorem 4.1. For $x = |X|$ the nonnegative measure $d\mu_0 = (x-1)^2(x+1/2)\,dP$ has mass $\rho - 1$. Direct expansion gives

$$M_4 - 1 = \int \frac{(x+1)^2}{x+1/2}\,d\mu_0,$$
$$M_5 - 1 = \int \frac{x^3 + 2x^2 + 3x + 3/2}{x+1/2}\,d\mu_0,$$

where $M_j = \mathbb{E}|X|^j$. Both ratios increase for $x \geq 0$: they equal $x + 3/2 + 1/[4(x+1/2)]$ and $x^2 + 3x/2 + 9/4 + 3/[8(x+1/2)]$, whose derivatives are nonnegative. Writing their values at $R = 96/25$ as $k_4(R), k_5(R)$, we obtain

$$M_4 \leq \min(R\rho, 1 + k_4(R)(\rho - 1)), \quad M_5 \leq \min(RM_4, 1 + k_5(R)(\rho - 1)), \quad M_6 \leq RM_5. \tag{28}$$

Also $\gamma^2 \leq M_4 - 1$, by expanding $\mathbb{E}(X^2 - 1 - \gamma X)^2 \geq 0$.

There is no need to assume a central support gap before the first Fourier bound. For independent copies let $E = \{(X - X')^2 \geq 9/25\}$ and let M be any valid fourth-moment ceiling. Then

$$\mathbb{P}(E) \geq \alpha := \frac{(2 - 9/25)^2}{2M + 6} > 0, \quad |X - X'| \geq 3/5 \text{ on } E, \quad \mathbb{E}[(X - X')^2 \mid E] \leq 2/\alpha. \tag{29}$$

Indeed $\mathbb{E}[(X - X')^2; E] \geq 2 - 9/25$ and $\mathbb{E}(X - X')^4 = 2M_4 + 6 \leq 2M + 6$, so Cauchy proves the mass bound. The conditional second-moment bound follows from its unconditional value two. This is an event under one actual iid marginal, not a replacement characteristic function.

The same support proof will be used again on a smaller closed moment cell. Freeze upper bounds r, b for $\rho, |\beta|$ in its polynomial Q, and test a proposed radius before changing any moment bound. For $R \in [5/2, 96/25]$, the gates $A > 0$, $0 < H_4 < 36A$, $Q(R) < 0$, $P(R) < 0$, and the six negative ray signs suffice. Their strictness excludes the boundary atoms as well as the open exterior rays. A radius that fails a gate is not used. The selected finite search is $R = 2.50, 2.51, \ldots, 3.84$; it only chooses among proved implications.

4.3 The Fourier enclosure used by every pass

Let $f(t) = \mathbb{E}e^{itX}$ and $e_n(t) = e^{-nt^2/2}(1 - in\gamma t^3/6)$. For the remainder in (8), the compensated inversion proved in Appendix A gives

$$|E_n| \leq \frac{2\sqrt{n}}{\pi} \left\{ \int_0^{1/2} \frac{|\Im[e^{-itH}(f(t)^n - e_n(t))]|}{t^3} dt + \int_{1/2}^{\infty} \frac{|f(t)^n - e_n(t)|}{t^3} dt \right\}.$$

The comparator is signed, has the correct mass and first moment, and its entire high-frequency tail is included. This formula applies at atoms and for arbitrary real H.

For clarity, the exact input and output of the selected Fourier step are as follows. Acquire $|X| \leq R$ first, and choose common ceilings $M \leq 11.1$, $g \leq 1.85$, M_6 for $M_4, |\gamma|, \mathbb{E}X^6$, together with

$$|(3 - M_4)/24| \leq A_4, \qquad |(10\gamma - \mathbb{E}X^5)/120| \leq Z_5.$$

Put

$$\begin{gathered} E_6 = M_6/720 + g^2/72 + (1/2 + g^2/144)^3/5, \\ B_6 = A_4 + Z_5/2 + E_6/4, \\ a_0 = (N - 1)/(2N), \quad b_0 = (45 - M)(1 - 2/N)/96, \\ c_0 \geq \sqrt{\pi/2}(N/(N - 1))^{3/2}, \qquad c_1 \geq c_0/2. \end{gathered}$$

A complete admissible low-frequency triple is

$$\begin{aligned} L_0 &= c_0 Z_5 + c_1 A_4 g + 251 g^3/17280, \\ L_H &= c_0 A_4 + 251 g^2/4800, \\ L_2 &= E_6/(2a_0^2) + B_6^2/(2b_0^3). \end{aligned}$$

One may take $A_4 = \max(2, M - 3)/24$ and $M_6 = RM_5$. The sharper fifth-order coefficient used by the finite cover is proved in Appendix A; it follows from the exact identity for $X^2 - \gamma X - 1$, not from an assumed fifth moment.

The high-frequency input is an actual event for two iid copies:

$$\mathbb{P}(\mathcal{E}) \geq \alpha > 0, \qquad |X - X'| \geq d_E > 0 \text{ on } \mathcal{E}, \qquad \mathbb{E}[(X - X')^2 \mid \mathcal{E}] \leq M_E.$$

Proposition A.3 supplies a complete six-term bound K_H at $N = 42$, subject to its explicit positive damping and all-index gates. The first event is (29); an opposite-sign event is used only after an actual central gap is acquired below. Different valid windows may be minimized only after all six terms, including the comparator, are present. No characteristic function of an artificial binary law is substituted for f.

The exact scalar output is

$$|E_n| \le K_F + s_F|H|, \qquad K_F = \frac{16}{25}(L_0 + K_H + L_2/\sqrt{N}), \quad s_F = \frac{16}{25}L_H. \tag{30}$$

All square roots and exponential bounds in the finite evaluation are directed outward. The proof in the appendix applies to every actual $n \ge N$ and allows the law and threshold to vary with n. Thus no fixed-law limit is being used as a uniform bound.

4.4 Coarse coefficients before a central gap is known

Write the non-Fourier constant charge in (8) as

$$A_c^{\mathrm{law}} = \frac{3}{2}\beta(J+\gamma) + 3(\rho J_1 - J_2) + \frac{\rho}{2}M_4 - \frac{1}{4}M_5 - \frac{\rho}{2}.$$

Here A_c^{law} does not depend on c; the subscript records its role as the coefficient of c. We will bound it from above, retaining all signs in the later passes.

The following gap inequalities need no gap around zero:

$$-M_4/6 \le J_1 \le M_4/2, \qquad -3J_2 - M_5/4 \le 0, \qquad J \le \rho + 1. \tag{31}$$

For a gap (u, v) of length h, the first pair follows from

$$\int_u^v (t-u)t\,dt + \frac{1}{2}\int_u^v t^2 dt = hv^2/2, \quad \frac{3}{2}\int_u^v t^2 dt - \int_u^v (t-u)t\,dt = h(2u+v)^2/6.$$

Sum with the nonnegative gap weights and use $\int t^2\tau = M_4/3$. On a negative gap, reflection and

$$\frac{1}{3}\int_u^v x^3 dx - \int_u^v (v-x)x^2 dx = u^3(v-u)/3 \ge 0 \quad (0 \le u \le v)$$

prove the second assertion; the positive portion only improves it. The last bound follows from $J \le \rho + 2m$ and $2m \le 1$.

For $1 \le r \le \rho$ and $R \le 6$, choose $\Gamma(R, r) \ge 0$ with

$$\Gamma(R,r) \ge \max_{r \le v \le R} \left\{ -\frac{v^2}{4} + \frac{(R+r)v}{3} - \frac{Rr}{2} + \frac{r^3(2R-r)}{12v^2} \right\}_+ . \tag{32}$$

Then

$$3(\rho J_1 - J_2) \le \frac{3}{2}\Gamma(R,r)(\rho - \gamma) + \frac{3}{8}\rho^2 J. \tag{33}$$

On the negative portion of a gap discard $|t| < \rho$, where its charge is nonpositive. Extending the remaining positive integral gives $\int_r^v (R-x)x(x-r)\,dx$. Its quotient by v^2 is the expression in (32). Signwise Tonelli charges its upper bound against $\mathbb{E}X_-^3 = (\rho - \gamma)/2$. On the positive portion, $t(\rho - t) \le \rho^2/4$ and $\sum \tau_{u,v} \int_u^v (t-u)dt = J/2$ give the other term. This argument includes a gap crossing zero.

The finite bound for Γ partitions $[r, R]$ into 64 closed intervals. On each, maximize the concave quadratic at its clipped vertex $2(R+r)/3$ and its nonnegative reciprocal term at the left endpoint. The alternative ceiling $5/2$ follows at $r = 1, R = 6$ from $3v^4 - 28v^3 + 66v^2 - 11 > 0$ on

$[1,6]$: the endpoint values are $30, 205$ and the two interior critical values are $(85 \mp 35\sqrt{5})/2 > 0$. The positive integral representation is increasing in R and decreasing in r, so that endpoint choice encloses the present domain.

The elementary square

$$V = \mathbb{E}[|X|^3(|X| - \rho)^2] = M_5 - 2\rho M_4 + \rho^3 \geq 0$$

gives $\rho M_4/2 - M_5/4 \leq \rho^3/4$. Thus on $\rho \in [l, u]$, $|\beta| \leq b$, $M_4 \leq M$, and with $0 \leq \rho - \gamma \leq \Delta$, two complete constant bounds are

$$\begin{aligned} A_1(\Delta) &= \tfrac{3}{2}b(2u+1) + \tfrac{3}{2}\Gamma(R,l)\Delta + \tfrac{3}{8}u^2(u+1) + u^3/4 - l/2, \\ A_2 &= \tfrac{3}{2}b(2u+1) + 2uM - l/2. \end{aligned} \tag{34}$$

Use $A = \min(A_1, A_2)$. The first uses (33); the second uses (31). They are alternative complete bounds, not partial costs to be mixed.

For the linear term K_1, the following three bounds are valid:

$$\begin{aligned} K_{\text{rad}} &= 3ub + \frac{R(R+2-3l/2)}{R+1/2}(u-1), \\ K_{\text{univ}} &= 6u - 3u^2/2 + 3b, \\ K_R &= Ru/2 + \max_{r \in \{l,u\}} |R/2 - 3r/2|g + 3b. \end{aligned}$$

For the first use the positive measure $d\mu_0 = (|X| - 1)^2(|X| + 1/2)dP$, of mass $\rho - 1$, and the exact identity

$$K_1 + 3\rho\beta = \int \operatorname{sgn}(X) \frac{x(x+2-3\rho/2)}{x+1/2}\, d\mu_0, \qquad x = |X|.$$

The ratio is convex in $x \geq 0$ and decreases in ρ. Its negative part is at most $31/40$ on $\rho \leq 1.85$; require its displayed endpoint ceiling to exceed that number. The second bound uses $|\mathbb{E}[X|X|^3] - 3\gamma| \leq 3\rho$ for $R \leq 6$, and $\rho \leq 2$. For the third use the signwise fourth-moment bounds $0 \leq \mathbb{E}X_\pm^4 \leq R(\rho \pm \gamma)/2$. Finally (31) gives

$$|\varphi\Omega| \leq v_H := \frac{2}{5}\max\{1/2, 2M/3 - 1/2\}, \qquad \Omega = J_1 + (M_4 - 3)/6.$$

Lemma 4.2 (A stronger signed fourth-moment charge)**.** *Let $|X| \leq R$, $\rho \in [l, u]$, $|\beta| \leq b < 1$. Put $h_* = 1 + b$ and $a(r) = (3r - R)/2$. A valid bound for $|K_1|$ is the maximum on $[l, u]$ of*

$$B_R(r, b) = Rr - \tfrac{3}{2}r^2 + 3b + \tfrac{1}{2}(1+b)a(r)^2.$$

If the three gates

$$h_* \geq a(u), \qquad 3 + 2a(l)h_* - \tfrac{3}{2}h_*^2 \geq 0, \qquad R - 3u + \tfrac{3}{2}h_*^2 \geq 0$$

hold, it may be intersected with the smaller bound

$$Ru - \tfrac{3}{2}u^2 - 3 + 3h_* + a(u)h_*^2 - h_*^3/2.$$

Proof. For $\epsilon = \pm 1$ set $h = 1 - \epsilon\beta > 0$ and $\delta = \rho - \epsilon\gamma$. Signwise Cauchy and $m \leq 1/2$ give $\delta \geq h^2$. The fourth moments on the two signs give

$$\epsilon K_1 \leq R\rho - \tfrac{3}{2}\rho^2 - 3 + 3h + a(\rho)\delta - \delta^2/(2h).$$

Completing the square in unrestricted δ and using $h \le h_*$ proves the first bound. With $\delta \ge h^2$, the last three terms maximize to

$$\psi_a(h) = \begin{cases} h(3 + a^2/2), & 0 < h \le a, \\ 3h + ah^2 - h^3/2, & h \ge a. \end{cases}$$

The first branch is increasing. The second has derivative $3 + 2ah - 3h^2/2$, a concave quadratic whose endpoint values on $[0, h_*]$ are nonnegative under the second gate. The branches agree at $h = a$; the first gate places h_* in the second branch. The remaining derivative in ρ is $R - 3\rho + 3h_*^2/2$, nonnegative by the third gate. This proves the claim, including all endpoints. The quadratic maximum in the first bound is evaluated at its endpoints and, when present, its concave vertex. □

Take the minimum of the applicable bounds for $|K_1|$ and call it K. No assertion about the signs of $\beta\gamma$ or $K_1\gamma$ is used.

4.5 Two quadratic passes and the first actual gap

Use the 24 closed cells with endpoints

$$r_i = 1 + 3361i/96000, \quad i = 0, \dots, 24, \qquad N = 42,$$

and directed constants

$$c_- = 0.4097321837023 < C_{\mathrm{E}} < c_+ = 0.4097321837025,$$
$$\phi_l = 997/2500 < \phi(0), \quad \sigma_l = 3.162277660 < \sqrt{10}.$$

The Gaussian constants follow from the rational Machin estimates in Lemma C.2 and Appendix A; the tighter c enclosure is checked by the same alternating expansions and rational squares. The interval encloses the fixed coefficient c, not the actual normalized score.

On a cell $[l, u]$, form the moments and the universal actual event above, and compute the complete Fourier pair K_F, s_F. In pass 1 use $\Delta = 2u$, $A = \min(A_1(2u), A_2)$, and, for $t \in [c_-, c_+]$,

$$f(t) = tK + v_H + s_F, \qquad q(t) = tl - g/5.$$

Require $q > 0$. The bound $\mathcal{R}_c \le K_F + cA + f(c)|H| - q(c)H^2$ is affine in c at each fixed $|H|$, so its maximum is at an endpoint before completing squares. Let an overline denote upward rounding to 10^{-10}. Compute

$$\begin{aligned} T_1 &= \overline{\max_{t=c_-,c_+} \{K_F + tA + f(t)^2/(4q(t))\}}, \\ \eta_1 &= (c_- l - T_1/N)/2, \\ U_1 &= \min\left\{T_1, \overline{\max_{t=c_-,c_+} \{K_F + tA + f(t)^2/[4(q(t) + \eta_1)]\}}\right\}, \\ d_1 &= \overline{6U_1/(N\phi_l)}. \end{aligned} \tag{35}$$

The strict gate $0 \le T_1 < Nc_- l$ is checked before η_1 or the resource bound is used. Corollary 2.4 then proves $\mathcal{R}_c \le T_1$, $\mathcal{U} \le U_1$, and $nD \le Nd_1$. The scalar bounds hold at every actual $n \ge N$.

For the second pass the circle in Lemma 3.1 gives

$$D \ge \sigma\rho - \gamma - 3\sqrt{\rho^2 - \gamma^2}, \qquad \sigma = \sqrt{10}.$$

Put $a_0 = \sigma_l - d_1/l$. A valid lower bound for γ/ρ is

$$Y(a_0) = \begin{cases} -1, & a_0 \le -1, \\ (a_0 - 3\sqrt{10 - a_0^2})/10, & -1 < a_0 \le \sqrt{10}. \end{cases} \tag{36}$$

Indeed $y + 3\sqrt{1-y^2}$ increases from -1 to $\sqrt{10}$ on $[-1, 1/\sqrt{10}]$. Solving its level equation on that branch gives the smaller quadratic root in (36); it has $a_0 - y \ge 0$, so squaring introduced no false branch. Every y below it violates the required lower level. A downward rational bound y_l is obtained by rounding the square root upward. Thus

$$g_l = \max\{-g, \sigma_l l - 3 - d_1, \min(l y_l, u y_l)\} \le \gamma,$$
$$\Delta = \min\{2u, (1-\sigma_l)l + 3 + d_1, u - g_l\}, \qquad \phi_- = \phi_l(1 - T_1/(Nc_- l)) > 0,$$
$$A = \min\{A_1(2u), A_1(\Delta), A_2\},$$
$$q(t) = \max\{tl - g/5, tl + \tfrac{1}{2}\min(\phi_- g_l, (2/5) g_l)\}.$$

The first two bounds also use $2m \le 1$. Each value of $q(t)$ is affine in t, since its two alternatives have the same slope l. Repeat the raw bound first, intersect it with T_1, then price phase using this new raw bound, intersect the resource with U_1, and finally retain $d_2 \le d_1$. A resource bound is never substituted for a raw bound in the density or phase license.

Certificate condition 4.3 (The two coarse passes)**.** The preceding coefficient domains, Fourier gates, endpoint enclosures and strict raw gates hold on every one of the 24 closed cells. Outward evaluation gives the following ceilings, and in every second-pass cell $d_2 < (\sigma_l - 1)l$:

pass	$\max T$	$\max U$	$\max d$	strict gap gates
1	13.2818901598	10.9330610254	3.9164138937	23
2	11.5164967977	9.9790315907	3.5746638454	24.

The minimum second-pass gap margin is at least $526024602543/1600000000000 > 0$.

The stated condition is a finite exact arithmetic obligation. The named two-pass proof object records and rechecks these inequalities; it is kept distinct from the analytic implications just proved. Although its implementation retains additional previously valid minima, the fresh quadratic ceilings attain all the displayed minima in this cover, so no earlier resource-refinement schedule is a mathematical premise here.

If zero belonged to the closed support, every gap would be on one side of zero and $J \le \rho$. Hence $D \ge (\sigma - 1)\rho$, contradicting the strict second-pass test. Centering and positive variance therefore give an actual central gap $(-A, B)$, with positive endpoints. This acquires an event, rather than supposing an artificial two-point kernel.

4.6 Signed moments and physical gap costs

Write $\mu_4 = M_4$, $\eta_5 = M_5$, $\sigma = \sqrt{10}$. Put $a_\pm = (1 \pm \beta)/2$, $b_\pm = (\rho \pm \gamma)/2$. Both signs have positive probability and positive first moment, since a centered variance-one law cannot be supported on one half-line. Thus $|\beta| < 1$ and the signwise denominators below are positive whenever used; degenerate correction denominators are handled separately. Cauchy gives $m^2 \le a_+ a_-$ and $b_\pm m \ge a_\pm^2$. Consequently

$$\frac{(1+\beta)^{3/2}}{\sqrt{1-\beta}} - \rho \le \gamma \le \rho - \frac{(1-\beta)^{3/2}}{\sqrt{1+\beta}}, \qquad m \ge \max\left\{\frac{1+\beta^2}{2\rho}, \frac{a_+^2}{b_+}, \frac{a_-^2}{b_-}\right\}.$$

These statements also hold with a zero atom. Projection of $X|X|$ onto the span of X and $|X|$ gives

$$\mu_4 \geq \rho^2 + \frac{(\gamma - \rho\beta)^2}{1 - \beta^2}.$$

For completeness, this follows by minimizing the nonnegative expectation $\mathbb{E}(X|X| - uX - v|X|)^2$ over u, v; its Gram matrix has diagonal entries one and off-diagonal entry β, and is positive definite.

Two support-localizing moment matrices give the additional upper bounds as follows. If $D_\pm = Ra_\pm - b_\pm > 0$, then

$$2m + \sum_{\pm} \frac{(Rm - a_\pm)^2}{D_\pm} \leq R, \qquad \mu_4 \leq R\rho - \sum_{\pm} \frac{D_\pm^2}{Rm - a_\pm}.$$

Indeed the first follows by summing the determinant inequalities for

$$\mathbb{E}[(R - |X|)(1, |X|)^{\mathsf{T}}(1, |X|); \pm X > 0].$$

The second follows from the corresponding matrices with an additional factor $|X|$. One more factor $|X|$ yields, sign by sign,

$$\mathbb{E}[X_\pm^5] \leq R\mathbb{E}[X_\pm^4] - \frac{(Rb_\pm - \mathbb{E}[X_\pm^4])^2}{D_\pm}.$$

Zero denominators are treated by dropping their nonnegative correction; they never imply exclusion without a separate contradictory numerator. The common m is retained throughout.

A second useful upper bound for m follows by integrating $(x - a)^2(R - x) \geq 0$ for $x = |X| \in [0, R]$. For every fixed $a > 0$,

$$2m \leq \frac{R - \rho + 2a + Ra^2}{a(2R + a)}.$$

The cover uses only the eleven rational choices $a = 2/10, \ldots, 12/10$.

Write $\delta = \gamma - \rho\beta$. Cauchy and a square completion on each sign give the signed constant bound

$$\frac{\rho}{2}\mu_4 - \frac{1}{4}\eta_5 \leq \frac{\rho^3}{4} - \frac{(\rho + \gamma)(\delta_+)^2}{8(1 + \beta)^2} - \frac{(\rho - \gamma)((-\delta)_+)^2}{8(1 - \beta)^2}.$$

Indeed on either sign, write $a = \mathbb{E}X_s^2$, $b = \mathbb{E}X_s^3$, $c = \mathbb{E}X_s^4$, $f = \mathbb{E}X_s^5$. Cauchy gives $c \geq b^2/a$, $f \geq c^2/b$, and $\rho c/2 - f/4 \leq \rho^2 b/4 - (c - \rho b)^2/(4b)$. Distance to the half-line $c \geq b^2/a$ proves the stated penalty on summing both signs. Independently, $\mu_4^2 \leq \rho\eta_5$ and the projection bound above give

$$\frac{\rho}{2}\mu_4 - \frac{1}{4}\eta_5 \leq \frac{\rho^3}{4} - \frac{(\gamma - \rho\beta)^4}{4\rho(1 - \beta^2)^2}.$$

Each is a complete deduction from the same square. There is a third independent deduction which is often stronger. For $Y = |X|$ and $\delta = \gamma - \rho\beta$, put

$$V = \mathbb{E}[Y(Y^2 - \rho Y)^2] = \eta_5 - 2\rho\mu_4 + \rho^3.$$

Since $\delta = \mathbb{E}[Y \operatorname{sgn}(X)(Y^2 - \rho Y)]$, weighted Cauchy gives $\delta^2 \leq 2mV$. Hence

$$\frac{\rho}{2}\mu_4 - \frac{1}{4}\eta_5 \leq \frac{\rho^3}{4} - \frac{\delta^2}{8m}. \tag{37}$$

The maximum of these three separately valid losses can be subtracted. Their sum cannot. A zero atom contributes zero to the weighted measure.

An actual central gap without assuming $D < 2$

The physical decomposition (21) shows that $D < (\sigma-1)\rho$ implies an actual central gap. Use its same nonnegative terms, writing $T = T_+ + T_-$ and $Q = (\sqrt{\sigma+1}A - \sqrt{\sigma-1}B)^2$. Hence $T \le D/(\sigma-1)$. The additional inequality $J \le \rho + 2mAB$ gives $6mAB \ge (\sigma-1)\rho - D$. Consequently, if $\rho \ge l$, $D \le d < (\sigma_l - 1)l$ and $m \le m_u$,

$$(A+B)^2 \ge \frac{l - d/(\sigma_l - 1) + \{(\sigma_l-1)l - d\}/3}{m_u} > 0. \tag{38}$$

This is the distance lower bound used in the complete high-frequency event. It uses the actual moment m and the actual support endpoints.

Moreover $A \le a_-/m$, $B \le a_+/m$ and $J \le \rho + 2m$. The last bound gives the compulsory defect

$$D \ge \sigma\rho - \gamma - 6m.$$

The inequalities $T - J_o \ge 2BR_+ + 2AR_-$ give the useful additional endpoint constraint

$$D \ge (\sigma-1)b_+ + 6a_+B - (\sigma+5)mB^2.$$

If $D < (\sigma-1)b_+$, this downward quadratic forces B to be at least its positive root. The reflected constraint has a_-, b_-, A and the coefficients $\sigma+1, \sigma+7$. In a closed moment box, positive coefficients are rounded down, negative coefficients up, and the lower square-root endpoint is used in the resulting root bound.

The actual opposite-sign event has probability $2pq$. There is no atom at zero. Cauchy gives the necessary probability interval $2m_-^2/(1+\beta_+) \le p \le 1 - 2m_-^2/(1-\beta_-)$ on a moment cell. In particular, $2pq \ge (1+\beta^2)^2/(2\rho^2)$, using $m \ge (1+\beta^2)/(2\rho)$ and $pq \ge m^2$. The minimum of $p(1-p)$ on a closed interval is attained at an endpoint.

For independent copies one has exactly

$$\mathbb{E}[(X-X')^2 \mid XX' < 0] = \frac{1+\beta(1-2p)+4m^2}{2pq} = \frac{1+4m^2+\beta}{2p} + \frac{1+4m^2-\beta}{2q}.$$

For an upper bound, use the upper m endpoint and both endpoints of p and β. The expression is convex in p and affine in β. This improves the former $2/\mathbb{P}(XX' < 0)$ bound inside the complete high-frequency estimate. Both copies still have the same actual marginal.

Preserving central and outer integral costs

Let

$$P_1(A,B) = B^3/3 + AB^2/2 - A^3/6, \quad P_2(A,B) = B^4/4 + AB^3/3 - A^4/12, \quad G = \rho P_1 - P_2.$$

Then $J_{1,c} = mP_1$, $J_{2,c} = mP_2$. Since $\partial_B G = B(A+B)(\rho - B)$,

$$G(A,B;\rho) \le G(A,\rho;\rho) = \rho^4/12 + \rho^3 A/6 - \rho A^3/6 + A^4/12.$$

The last expression increases with A because its derivative is $(\rho - A)^2(\rho + 2A)/6$, and is convex in ρ because its second ρ derivative is $\rho^2 + \rho A$. Its closed-rectangle maximum is therefore bounded using the upper A endpoint and both ρ endpoints.

Positive outer gaps satisfy $(t-a)(\rho - t) \le (\rho - B)_+^2/4$. On negative outer gaps the positive part is bounded by $(R-\rho)^2/4$ after extracting $|t|$. Integration gives

$$3(\rho J_1 - J_2)_o \le \frac{3}{8}\{(\rho - B)_+^2 T_+ + (R-\rho)^2 T_-\}.$$

The available weighted outer budget is $D - mQ$, not the entire D. The cover intersects this budget with the individual bounds $T_+ \le b_{+,u} - m_l B_l^2$ and $T_- \le b_{-,u} - m_l A_l^2$.

For the signed phase term $\Omega = J_1 + (\mu_4 - 3)/6$, retain

$$-RT_-/2 \le J_{1,o} \le RT_+/2.$$

The central minimum of P_1 occurs at B_l and an A endpoint; its maximum occurs at B_u and A clipped to the interval at $A = B_u$. A nonnegative polynomial bound is multiplied by m_u for an upper bound and m_l for a lower bound; a negative one uses the opposite endpoint. For example, the binary endpoints $A = 3/2$, $B = 2/3$ make this maximum negative; the sign switch is necessary even when $D < 2$. The fourth-moment sign intervals also sharpen $K_1 = \mathbb{E}[X|X|^3] - 3\beta - 3\rho\gamma/2$. The complete K_1, Ω, φ rectangles are used in the signed expression $|cK_1 + \varphi\Omega|$, without replacing it by two separate absolute values.

A compulsory first moment before each signed evaluation.

The physical inequality $D \ge \sigma\rho - \gamma - 6m$ is especially useful inside a cell. If its prior ceiling is $D \le d$, and $\rho \ge l$, $\gamma \le \gamma_+$, then

$$m \ge m_* := \max\{0, (\sigma_l l - \gamma_+ - d)/6\}. \tag{39}$$

If $m_* > 1/2$, the cell is empty. Otherwise the common sign-moment inequality $4m^2 + \beta^2 \le 1$ forces $|\beta| \le \sqrt{1 - 4m_*^2}$. This restriction must be intersected with the beta interval for this gamma cell, not carried from a different gamma cell. Recompute all dependent sign-moment ranges after the intersection. These are necessary conditions on one law; no sufficiency or moment-realizability assertion is made.

For completeness, the event distance and conditional moment refinements fit the Fourier input exactly. On the acquired actual gap, use $\mathcal{E} = \{XX' < 0\}$, a downward square root of (38), and the larger of the two valid probability lower bounds

$$\alpha = \max\left\{\frac{(1 + \min|\beta|^2)^2}{2u^2}, \quad 2\min_{p=p_-,p_+} p(1-p)\right\}.$$

The conditional second-moment ceiling is the smaller of $2/\alpha$ and the corner maximum of the exact expression displayed above. Its numerators are positive, it increases with m, is affine in beta, and is convex in p, so these corner choices are valid. Each event option is inserted into all six HIGH terms. Taking the minimum with the complete universal-event bound preserves the same actual Fourier integral. None of these statements follows merely from declaring a moment rectangle feasible.

4.7 The signed enclosure and its ordered update

On one closed moment cell retain bounds for (ρ, β, γ, m), an already proved support radius R, an incoming raw ceiling T, and the weighted defect invariant $nD \le Nd$. Thus $D \le d$ for every actual $n \ge N$. Intersect all necessary moment conditions just proved, with directed endpoint operations. A positive numerator with a zero denominator proves incompatibility only through its original positive-semidefinite inequality; a zero-over-zero correction is dropped. No division is made before this check. Bisection of the monotone first-moment necessary inequality keeps its upper endpoint, so its rounding cannot remove a law.

Here is an explicit description of the coefficient enclosures. Let M be the resulting upper fourth moment and let $[J_-, J_+]$ contain J; one can use

$$J_- = \max\{0, ((3 + \sigma_l)l - \gamma_+ - d)/3\}, \quad J_+ = \min\{u + 2m_+, ((3 + \sigma_u)u - \gamma_-)/3\},$$

with a rational $\sigma_u > \sqrt{10}$. For the beta term take the full corner maximum of $3\beta(J+\gamma)/2$. For the gap charge take the minimum of the complete coarse charge (33) and the central-plus-outer charge just proved. Write this bound as G_* and the beta-term maximum as B_*. Let $P_* \geq 0$ be any lower bound for the maximum of the three independent fifth-moment penalties. Then a valid complete constant bound is

$$A = \min\{B_* + G_* + u^3/4 - l/2 - P_*, \quad B_* + 2uM - l/2\}. \tag{40}$$

The fourth semimoment inequalities and the signed product rectangle for $\rho\gamma$ give an interval $[k_-, k_+]$ containing $K_1 = \mathbb{E}X_+^4 - \mathbb{E}X_-^4 - 3\beta - 3\rho\gamma/2$. The central polynomial mP_1 and the signed outer bounds give an interval $[\omega_-, \omega_+]$ containing $\Omega = J_1 + (M_4 - 3)/6$. Intersect them with every previous valid interval. When multiplying by a moment endpoint, the sign of the coefficient is retained. These constructions explain every coefficient in the next quadratic, without assuming that unrelated corner maxima are attained by one common law.

Require the incoming raw gate $0 \leq T < Nc_l$. By Corollary 2.4,

$$\varphi = \phi(H/\sqrt{n}) \in [\phi_-, 2/5], \qquad \phi_- = \phi_l(1 - T/(Nc_l)) > 0.$$

For a formal coefficient $v \in [c_-, c_+]$ set

$$\begin{aligned} B_v &= K_F + vA, \\ F_v &= \max_{k\in\{k_-,k_+\},\ \omega\in\{\omega_-,\omega_+\},\ \psi\in\{\phi_-,2/5\}} |vk + \psi\omega| + s_F, \\ Q_v &= vl + \tfrac{1}{2}\min\{\phi_-\gamma_-, (2/5)\gamma_-\}. \end{aligned} \tag{41}$$

The actual raw remainder is at most $B_c + F_c|H| - Q_cH^2$. The maximum over the signed rectangle is valid by successive affine maximization, and the absolute value preserves convexity. For fixed $|H|$, this expression is convex in c: the constant and curvature parts are affine and F_c is convex. Thus the two endpoints c-,c+ suffice, even if A is negative. A negative lower gamma must be priced with the upper density endpoint; it cannot be reflected inside the contact law.

Lemma 4.4 (One ordered signed step)**.** *Suppose all preceding cell bounds hold and $Q_v > 0$ at both endpoints. Let upward rounding be applied to every proposed numerical ceiling. Set*

$$\begin{aligned} T' &= \min\{T, \max_{v=c_-,c_+}(B_v + F_v^2/(4Q_v))\}, \\ \eta &= (c_l - T'/N)/2, \\ U' &= \max_{v=c_-,c_+}(B_v + F_v^2/[4(Q_v+\eta)]). \end{aligned}$$

A negative T' or U' excludes the cell. Otherwise $\mathcal{R}_c \leq T'$, $\mathcal{U} \leq U'$, and the new weighted defect bound is

$$nD \leq Nd', \qquad d' = \min\{d, 6U'/(N\phi_l)\}.$$

A valid raw-threshold ceiling is

$$|H| \leq \max_{v=c_-,c_+} \frac{F_v + \sqrt{F_v^2 + 4(Q_v+\eta)\max(0, B_v)}}{2(Q_v+\eta)}. \tag{42}$$

Proof. Complete the two raw squares first. The incoming strict gate persists under $T' \leq T$, and gives $\eta > 0$. Only now apply Corollary 2.4 and complete the two resource squares with curvature $Q_v + \eta$. A negative complete ceiling is incompatible with the nonnegative raw or resource expression. The resource contains $n\phi(0)D/6$, so the displayed weighted bound follows without setting the actual sample size equal to N. If the maximum of both resource quadratics at a phase were negative, no witness could have that phase. Replacing B by its positive part enlarges each positive root and gives (42). Previous valid resource or phase bounds may also be intersected, but an unlicensed resource bound cannot be fed back as a raw ceiling. □

4.8 Closed coverage and the selected acquisition theorem

For each of the original 24 rho cells, acquire a support radius with the independent heat/ray test before recomputing higher moments. Each signed pass then partitions a full necessary beta enclosure into eight closed intervals, each resulting necessary gamma enclosure into eight closed intervals, and each necessary common-m enclosure into two closed intervals. The original beta interval is restored for every gamma slice before applying (39). Discarding an outer interval counts all its descendants, and requires one of the proved necessary conditions to be inconsistent. In every processed rho cell the active and excluded counts sum to 128.

Each surviving actual law satisfies the cell update of Lemma 4.4. Pool upper bounds by maxima and lower bounds by minima before the next pass. This may enlarge the possible set, but cannot lose an actual witness. A rho cell disappears from later passes only when all of its descendants have been excluded. Degenerate closed intervals may duplicate boundary work; they do not leave holes.

For export, intersect each surviving necessary probability interval with all 100 closed panels $[j/100, (j+1)/100]$, $j = 0, \dots, 99$. A cell meeting a shared endpoint contributes to both panels. From $\theta = 1 - m^2/[p(1-p)]$ obtain a panel upper bound using its lower m endpoint and the maximum of $p(1-p)$ on the intersection, including $1/4$ if it crosses $1/2$. Pool this simultaneously with the raw, weighted defect, rho, beta, and phase bounds of the same cells.

Certificate condition 4.5 (The four signed passes)**.** Starting with Condition 4.3, all support, moment, actual-event, Fourier, denominator, signed-curvature and ordered raw-license gates hold on the complete four-pass cover just described. Its exact rational evaluation contains 11264 nominal signed boxes in total. The complete 100-panel export satisfies, on every nonempty panel,

$$\begin{aligned} \mathcal{R}_c &\le 2.1995174, & nD &\le 32.20665378, & D &\le 0.76682509, \\ \rho &\le 71527/48000, & |\beta| &\le 0.50549963, & |H| &\le 1.83605680. \end{aligned}$$

Fourteen rho cells survive, and thirty probability panels are nonempty. The largest support radius among the surviving final rows is $157/50$; thus the export also supplies

$$\operatorname{supp} P \subset (-157/50, 157/50). \tag{43}$$

This is the named fixed-target coarse proof object, with its own targets and complete incidence requirements. It has a fresh exact acceptance record; numerical acceptance remains distinct from the analytic proof that its leaves enclose all possible witnesses. The preceding statements supply that reduction and explicitly identify the finite predicates used. They do not change an older certificate's hypotheses or its target.

Theorem 4.6 (Selected fixed-target coarse entry)**.** *Let $n \ge 42$ be an actual index and let an attained fixed-target pair have $a \ge 0$. Assume the true sharp predecessor $C_{n-1} \le C_{\mathrm{E}}$ and the external structural inequality* (24)*. Under Conditions 4.3 and 4.5, the stated complete probability cover and all the displayed coarse bounds hold for that pair.*

Proof. Theorem 4.1 supplies the actual support before higher moments are used. Appendix A and the actual-event constructions prove the complete Fourier bounds on their specified domains. The coarse coefficient inequalities and the two ordered quadratic steps establish the first actual gap. Every signed cell thereafter uses the moment and physical-gap bounds of this same law, followed by Lemma 4.4. Induction over the four passes preserves the weighted invariants. The complete closed cover and outward pooling prove that every witness reaches an incident output panel with the stated bounds. □

The argument contains no upper bound for the witness's actual score, no scalar score recurrence, and no second deleted-score assumption. It uses only the simple quadratic phase price. A further confinement and count comparison is required on every retained probability panel; the finite base required to obtain a first failure is a separate input. Neither a small defect nor a successful coarse acquisition proves that final comparison.

5 Fixed-target moments and signed phase resources

The next two sections keep the actual law, index and threshold fixed. Throughout, $n \geq N = 42$, $c = C_{\mathrm{E}}$, and the attained fixed-target excess a is nonnegative. We assume the sharp predecessor $C_{n-1} \leq c$ and the contact and conservation formulas (5) and (10). The entry of Section 4.8 supplies a complete closed cover, an actual central gap $(-A, B)$, and bounds for the same law's $\rho, \beta, \gamma, m, D$ and raw threshold H. Here $D = (3+\sqrt{10})\rho - \gamma - 3J$, with J the actual weighted gap sum. Write $\mathcal{L} = L_{\mathrm{core}}$ for the nonnegative leakage denoted L in the core conservation formula; this separates it from slope bounds used below. The physical noise remains $W = X - \mathbb{E}[X \mid B_0]$ and the count noise is $U = W/\ell$. Below, the different symbol Ω denotes a signed Fourier moment coefficient.

Certificate condition 5.1 (Panel input and finite contraction)**.** Use the nonempty probability panels indexed 24 through 53 in the selected entry, with closed endpoints $[i/100, (i+1)/100]$. On each panel I retain its actual-law bounds, including $nD \leq ND_I$ and an initial noise ceiling $\theta \leq \Theta_I$. Every law allowed by the entry must belong to at least one such panel. The support radius $R = 157/50$ is the largest already acquired radius of the surviving fourth-pass rho rows. All arithmetic in the ordered iteration below is outward: lower roots are rounded down, upper roots up, and only a strictly negative residual excludes a panel. Perform eight rounds; check every denominator is positive before division and every asserted root has nonnegative input.

The numerical input is a finite obligation, not a replacement for its entry hypotheses. The formulas below explain the implication of each recorded inequality and permit a check of each entire panel.

Write $p = \mathbb{P}(X > 0)$, $q = 1 - p$, $m = \mathbb{E}X_+ = \mathbb{E}X_- > 0$. The conditional means are $\bar{B} = m/p$, $\bar{A} = m/q$, and

$$\theta = 1 - \frac{m^2}{pq}, \qquad \ell = \frac{m}{pq} = \sqrt{\frac{1-\theta}{pq}}, \qquad d = 1 - \frac{A+B}{\ell}, \qquad \lambda = \frac{n\theta}{\ell^2}.$$

Thus d is a count-coordinate inward displacement, not D. Let $V = \bar{B}$ on the positive sign and $V = -\bar{A}$ on the negative sign; write $W = X - V$. Define the two measures on nonnegative magnitudes by $\int f(x)d\mu_{\pm}(x) = \mathbb{E}[W^2 \mathbf{1}_{\{\pm X > 0\}} f(|X|)]$. Their masses sum to θ; the central gap ensures there is no atom at zero. Also $A \leq \bar{A}$, $B \leq \bar{B}$.

5.1 A common defect budget

We use the actual-law outer-gap inequality

$$D \geq (\sigma - 1)\mathbb{E}X_+^3 + (\sigma + 1)\mathbb{E}X_-^3 - 6mAB + 6BR_+ + 6AR_-,$$
$$\sigma = \sqrt{10}, \qquad R_+ = \mathbb{E}X_+^2 - mB, \qquad R_- = \mathbb{E}X_-^2 - mA.$$

Here is the gap argument behind this input. On the positive side, summing over outer support gaps and comparing each squared gap length with $2\int(t-B)\,dt$ gives

$$J_o^+ \leq 2\int_B^\infty (t-B)\tau(t)\,dt = \mathbb{E}[X_+(X_+ - B)^2] = T_+ - 2BR_+, \quad T_+ = \mathbb{E}X_+^3 - mB^2.$$

The negative side gives $J_o^- \le T_- - 2AR_-$ by the same magnitude calculation. Adding the central term $m(A+B)^2$ and substituting in D proves the displayed inequality. Endpoint atoms are included in the tail integrals; changing their values at integration endpoints does not change an integral. Substituting $\mathbb{E}X_+^3 = m\bar{B}^2 + \int (x+2\bar{B})d\mu_+$ and $R_+ = \mu_+(\mathbb{R}) + m(\bar{B}-B)$, and their negative-sign counterparts, gives

$$D \ge \mathfrak{b}_0 + E_0 + \int k_+ d\mu_+ + \int k_- d\mu_-, \tag{44}$$

where, with $p_* = (4-\sqrt{10})/2$,

$$\begin{aligned}
\mathfrak{b}_0 &= \frac{(6+2\sigma)(1-\theta)^{3/2}}{\sqrt{pq}}(p-p_*)^2, \\
E_0 &= 6m\{(A+B)(\bar{A}-A+\bar{B}-B) + (\bar{A}-A)(\bar{B}-B)\}, \\
k_+(x) &= (\sigma-1)(x+2\bar{B}) + 6B, \qquad k_-(x) = (\sigma+1)(x+2\bar{A}) + 6A.
\end{aligned}$$

The deterministic identity is $(\sigma+1)p^2 + (\sigma-1)q^2 - 6pq = (6+2\sigma)(p-p_*)^2$. Every displayed cost is nonnegative.

5.2 Ordered contractions on closed probability panels

For $I = [l,u]$, set $v_- = \min_{p\in I} pq$, $v_+ = \max_{p\in I} pq$. The maximum includes $1/4$ if I crosses $1/2$. Use $\sigma_l = 3.162277660 < \sigma < 3.162277661 = \sigma_u$. Initially $D \le D_I < \sigma_l - 1$, and the gap bound $J \le \rho + 2mAB$ gives

$$AB \ge z := (\sigma_l - 1 - D_I)/3 > 0.$$

Indeed $\rho \ge 1, \gamma \le \rho, m \le 1/2$ imply this inequality. With $\bar{A}_u = \sqrt{u/(1-u)}$, $\bar{B}_u = \sqrt{(1-l)/l}$, obtain

$$A \ge A_l = z/\bar{B}_u, \quad B \ge B_l = z/\bar{A}_u, \quad (A+B)/\ell \ge g = 2\sqrt{zv_-}, \quad d \le 1-g.$$

For a currently proved $\theta \le \Theta$, lower-bound the binary charge by

$$b_I = \frac{(6+2\sigma_l)(1-\Theta)^{3/2}}{\sqrt{v_+}} \operatorname{dist}(I, [(4-\sigma_u)/2, (4-\sigma_l)/2])^2.$$

All square roots are directed outward. If $r = D_I - b_I < 0$, the entire closed panel is impossible. Otherwise set

$$\bar{A}_l = \sqrt{(1-\Theta)l/(1-l)}, \quad \bar{B}_l = \sqrt{(1-\Theta)(1-u)/u},$$

$$K = \min\{(\sigma_l - 1)(B_l + 2\bar{B}_l) + 6B_l,\ (\sigma_l+1)(A_l + 2\bar{A}_l) + 6A_l\} > 0.$$

Equation (44) gives $\theta \le r/K$. Intersect this bound with Θ, then recompute b_I, r. Put $m_l = \sqrt{(1-\Theta)v_-}$ and $\ell_l = \sqrt{(1-\Theta)/v_+}$. Since $E_0 \ge 6m\ell^2 gd$, we may update

$$d \le \min\{d_{\text{old}}, r/(6m_l\ell_l^2 g)\}, \quad g \ge 1-d,$$

$$A_l \leftarrow \max\{A_l, \ell_l(l-d)_+\}, \quad B_l \leftarrow \max\{B_l, \ell_l(1-u-d)_+\}.$$

Only previously proved bounds enter each update. Zero residual forces zero noise and displacement; no division by that residual is needed. The implementation performs eight rounds, rounding upper bounds upward and lower bounds downward to 10^{-10}.

5.3 Signed moments and a weighted noise bound

For every orientation, including $p > 1/2$,

$$1 - 2u - 2(1-u)\Theta \le \beta \le 1 - 2l + 2l\Theta. \tag{45}$$

This follows from $\beta = (1-\theta)(1-2p) + \mu_+(\mathbb{R}) - \mu_-(\mathbb{R})$. The formerly convenient upper replacement $1 - 2l + \Theta$ requires $l \le 1/2$ and must not be continued past that point.

For a coherent third-moment cap, put

$$L = R + 2\max(\bar{A}_u, \bar{B}_u), \quad k = \min\{\sigma_l - 1 + 6B_l/(R + 2\bar{B}_u),\ \sigma_l + 1 + 6A_l/(R + 2\bar{A}_u)\}.$$

The exact identity

$$\rho = \rho_B(p)(1-\theta)^{3/2} + \int (x + 2\bar{B})d\mu_+ + \int (x + 2\bar{A})d\mu_-.$$

gives a noise charge at most $\min(L\theta, r/k)$. Let $r_B = (1 - 2v_-)/\sqrt{v_-}$ and $s \le (3/2)\sqrt{1-\Theta}$. Integrating the derivative proves $(1-\theta)^{3/2} \le 1 - s\theta$ for $0 \le \theta \le \Theta$. The maximum of this affine, piecewise expression is therefore bounded by

$$\rho \le r_B + (L - sr_B)_+ \min\{\Theta, r/(kL)\}.$$

Intersect with the prior rho cap.

Finally (44) implies the joint inequality

$$D \ge b_I + 6m(A_l + B_l)\ell d + K\theta.$$

Multiplying by the actual n, using $nD \le ND_I$, $m\ell = 1 - \theta$, and $n\theta = (1-\theta)\lambda/(pq)$, proves, when $r > 0$,

$$\frac{d}{r/[6(1-\Theta)(A_l + B_l)]} + \frac{\lambda}{Nv_+ r/[K(1-\Theta)]} \le 1.$$

This retains the weighted resource; it is not obtained by multiplying an arbitrary unweighted noise cap by N.

Theorem 5.2 (Conditional all-panel contraction)**.** *Under Condition 5.1, the exact eight-round calculation on the selected inputs excludes panels 24 and 53. On all 28 remaining panels,*

$$\frac{1}{4} \le p \le \frac{53}{100}, \qquad \theta \le 0.0594425514, \qquad d \le 0.0719539227,$$
$$\rho \le 1.4547288545, \qquad \lambda \le 0.6314459997,$$
$$1.957063175 \le \ell \le 2.3094010768.$$

All endpoints and both signs are included.

Proof. The previous inequalities justify each recorded rational update. The input is the full accepted closed cover, and only a strictly negative residual excludes a panel. All 28 retained outputs and every update are in the exact log; the stated constants are their outward maxima or minima. No sampled parameter or favorable moment optimizer is used. □

5.4 Acquiring the incoming strip

The retained fixed-target contact theorem, with sharp predecessor and nonnegative excess, bounds a support contact by $H_n(H, \pm t) + cQ(t)$, where $Q(t) = -t^3 + 3rt^2/2 + 3bt + (8/875)r$. Here r is a rho upper bound, and $b = \beta_u$ on the positive ray, $b = -\beta_l$ on the negative ray. For $t > \sqrt{3}$, set

$$A_t = (t^3-3t)/6, \quad H_4 = t^4-6t^2+3, \quad S = 81/80, \quad E_t = t^5/4100+3t/820+17(3t^2-2)/78720.$$

The heat expansion used in the preceding entry proof remains valid here: its fifth-derivative error has coefficient $3t^2 - 2$, since $t^2 - 1 > 0$. Using $\|\phi'''\|_\infty < 2\phi(0)$, $\phi(0) < 2/5$ and $\sqrt{n-1} > 6$, without any sign assumption on H_4, yields

$$H_n(H, \pm t) \le \tfrac{2}{5} S^3 (A_t + |H_4|/72) + S^3 E_t.$$

Thus a negative boundary value of this expression plus $(819/2000)Q(t)$ excludes that boundary, provided $Q(t) < 0$. The exact derivative estimate is $|\partial_t H_n| < \frac{2}{5}(t^2/2 + 1/2 + 8/875)$. Check $Q, Q', Q'' < 0$ and

$$L = \tfrac{2}{5}(t^2/2 + 1/2 + 8/875) + (819/2000)Q', \quad L', L'' < 0.$$

Their higher derivatives are negative constants, so both rays are excluded beyond the boundary, for every raw threshold H. The complete Gaussian expansion and its domain are proved in Lemma 6.2; the finite boundary and continuation inequalities remain explicit certificate conditions.

Choose the two boundaries $t_- = \ell_l(l + 3/5)$ and $t_+ = \ell_l(1 - u + 3/5)$. All satisfy $t_\pm > \sqrt{3}$. Both pass on panels 40–52, proving $Y = X/\ell + p \in (-3/5, 8/5)$ there. On panels 25–39 the negative ray passes but the positive boundary upper estimate is nonnegative. This is failure of that sufficient estimate, not a counterexample to the strip theorem. No strip claim is made on those fifteen panels by this test. On panels 40–52 the inward bound already gives $U_0 \le d < 0.072$ and $U_1 \ge -d > -0.072$; together with the outer strip this proves $|U| < 3/5$. This acquisition precedes the half-amplitude argument in Section 6.

For later use, the panel $[0.41, 0.42]$ has the following convenient strict outward bounds. They are weaker than its eight-round output and its retained entry phase bound:

$$\begin{gathered} \theta < 0.054, \quad d < 0.069, \quad \rho < 1.22, \quad 0.09736 \le \beta \le 0.234, \\ 1.9706 < \ell < 2.034, \qquad \lambda < 0.6, \qquad |H| < 1.814, \qquad |U| < 0.6. \end{gathered} \tag{46}$$

For example, its recorded noise and displacement bounds are $\theta \le 0.053026061$, $d \le 0.0672060498$, $\lambda \le 0.5728987925$ and $\rho \le 1.2049635459$. These numerical consequences still require the complete selected entry and the finite contraction checks of Condition 5.1.

5.5 Signed Fourier terms from one actual moment cell

Write $M_j = \mathbb{E}|X|^j$ when an absolute moment is intended, and define

$$K_1 = \mathbb{E}[X|X|^3] - 3\beta - \tfrac{3}{2}\rho\gamma, \qquad \Omega = J_1 + (M_4 - 3)/6, \qquad J_1 = \sum_{(a,b)} \tau_{a,b} \int_a^b (t-a)t\,dt.$$

The moment coefficient Ω is unrelated to the physical noise W. The complete Fourier estimate in the ordered entry supplies, on each retained cell,

$$\mathcal{R}_c \le K_F + cA_F + (cK_1 + \varphi\Omega)H + s_F|H| - (c\rho + \varphi\gamma/2)H^2, \qquad \varphi = \phi(H/\sqrt{n}). \tag{47}$$

Here $K_F, s_F \geq 0$ are complete Fourier error bounds and A_F bounds the constant moment/gap term. This inequality retains all frequency windows and their gates from Section 4.7; the following algebra does not replace those analytic or finite inputs. The cell contains ranges for $\rho, \gamma, K_1, \Omega, \varphi$ belonging to this same law. Write their lower moment endpoints as r_l, g_l.

There are two ordered raw ceilings. The preceding pass supplies the positive density floor φ_l used in (47). After bounding that polynomial one obtains a new ceiling $\mathcal{R}_c \leq T < Nc_- r_l$, where $c_- \leq c \leq c_+$. Only now set

$$\eta = \frac{c_- r_l - T/N}{2} > 0, \qquad \mathcal{U} = (n + H^2)a + n\phi(0)D/6 + n^{3/2}\mathcal{L}.$$

All three costs in $\mathcal{U}$ are nonnegative. The phase-price formula of Corollary 2.4 gives

$$\mathcal{U} \leq \mathcal{R}_c - \eta H^2. \tag{48}$$

To see the ordering directly, put $w = H^2/(2n)$ and $M = c\rho - \phi(0)D/6$. Conservation gives $e^{-w}M \geq c\rho - T/n > 0$, and therefore

$$\mathcal{R}_c - \mathcal{U} = n(1 - e^{-w})M = n(e^w - 1)e^{-w}M \geq \tfrac{1}{2}H^2(c\rho - T/n) \geq \eta H^2.$$

The density floor is not inferred from this new bound and then fed back into the polynomial that produced it.

For each $t \in \{c_-, c_+\}$ define

$$B_t = K_F + tA_F, \qquad Q_t = tr_l + \min_{v \in [\varphi_l, 2/5]} vg_l/2 + \eta, \tag{49}$$
$$F_{t,+} = \max(tK_1 + v\Omega) + s_F, \qquad F_{t,-} = -\min(tK_1 + v\Omega) + s_F.$$

The extrema in the second line range over the recorded rectangular intervals for K_1, v, Ω. The expression is affine in each variable separately, so its extrema occur at the eight corners. The separate gate $Q_t > 0$ must hold for both target endpoints.

Proposition 5.3 (One-cell signed resource)**.** *For $h = |H|$ and the actual sign ε of H,*

$$0 \leq \mathcal{U} \leq \min\left\{T, \max_{t \in \{c_-, c_+\}} (B_t + F_{t,\varepsilon}h - Q_t h^2)\right\}. \tag{50}$$

At $H = 0$ both signed branches give the same value.

Proof. For fixed $h \geq 0$, the maximum or negative minimum defining the signed linear coefficient is a maximum of affine functions of t and hence convex. The remaining terms in the polynomial are affine in t. Its maximum on $[c_-, c_+]$ is bounded by the larger endpoint value. Apply (47) and (48); independently, $\mathcal{U} \leq \mathcal{R}_c \leq T$. This proves the minimum with T. In particular a negative signed slope remains negative. Replacing it by zero is unnecessary and discards information. □

5.6 Closed-region maxima and necessary noise filters

On $p \in [l, u] \subset (0, 1)$ and $0 \leq \lambda \leq \Lambda$, the exact variance identities give

$$\theta = \frac{\lambda}{nv + \lambda}, \qquad m^2 = \frac{v^2}{v + \lambda/n}, \qquad v = p(1 - p).$$

Thus with the actual minimum and maximum v_-, v_+ over the interval,

$$\theta \leq \Theta_\Lambda := \frac{\Lambda}{Nv_- + \Lambda}, \qquad \sqrt{\frac{v_-^2}{v_- + \Lambda/N}} \leq m \leq \sqrt{v_+}, \tag{51}$$
$$1 - 2u - 2(1 - l)\Theta_\Lambda \leq \beta \leq 1 - 2l + 2u\Theta_\Lambda.$$

Indeed $v^2/(v+s)$ increases with $v > 0$ and decreases with $s \geq 0$. The beta bounds follow by enlarging each coefficient in the exact signed second-moment inequalities outward. This deliberately conservative filter is valid on either side of $1/2$. It also includes the limit $n \to \infty$; no finite upper bound on n is needed.

Consider a closed rectangle $h \in [h_0, h_1]$, $\lambda \in [\lambda_0, \lambda_1]$ on the probability panel. Retain every moment cell whose p, m, β intervals meet the necessary ranges in (51), with $\Lambda = \lambda_1$, and whose previously acquired phase ceiling is at least h_0. Only disjoint ranges can be discarded. In each retained cell clip h_1 by that phase ceiling. A concave quadratic $B + Fh - Qh^2$, $Q > 0$, has its maximum at the vertex clipped to this closed interval:

$$h_* = \max\{h_0, \min(h_1, F/(2Q))\}.$$

Take the maximum of the two target-endpoint maxima, intersect with that cell's T, and finally take the maximum over every retained cell. Denote the result by S_ε. Then

$$\mathcal{U} \leq S_\varepsilon \tag{52}$$

on the whole closed rectangle. Shared endpoints remain in every intersecting cell; both phase signs are retained at zero. An empty list excludes a rectangle only relative to the complete ancestor cover. A negative S_ε excludes it because $\mathcal{U} \geq 0$.

Certificate condition 5.4 (Complete signed resource export)**.** The selected fourth pass covers all 2,048 moment boxes and retains every one of its 682 active cells. Both endpoint quadratics are reconstructed on each active cell, giving 1,364 quadratics with their preceding density floors and current positive phase prices. Rejected cells retain their valid exclusion reasons; no missing or unclassified box is interpreted as impossible. Every resource rectangle uses all incident active cells and the complete signed maxima described above.

This finite condition is checked by the exact replay and separate coefficient reconstruction accompanying the selected entry. Its role is to certify the actual-law cover and the rational bounds used here. It does not prove contact, conservation or a Fourier remainder theorem. Retaining the signed quadratics requires no new frequency integration; it preserves information already contained in that last signed pass.

6 Conditional confinement in Bernoulli coordinates

Two different statements will be proved. Low noise gives half-amplitude confinement on the acquired probability range $[0.40, 0.53]$ by one continuous polynomial argument. A separate signed-deletion calculation covers the full acquired noise range on $[0.41, 0.42]$. Neither statement asserts the subsequent strict count comparison. Both preserve the orientation selected for the fixed-target maximizing pair.

6.1 Conditional statement and acquired application

Retain the fixed-target hypotheses of Section 5: $n \geq 42$, nonnegative excess a, $c = C_{\mathrm{E}}$, closed-support contact, and the true predecessor $C_{n-1} \leq c$. The actual raw threshold H can have either sign. Assume an actual central gap $(-A, B)$ with $A, B > 0$, and write

$$p = \mathbb{P}(X > 0), \quad q = 1 - p, \quad m = \mathbb{E}X_+ = \mathbb{E}X_-, \quad \bar{A} = m/q, \quad \bar{B} = m/p, \quad \ell = \bar{A} + \bar{B},$$

$$X = \ell(Y - p), \qquad Y = B_0 + U, \qquad B_0 = \mathbf{1}_{\{X>0\}}, \qquad \mathbb{E}(U \mid B_0) = 0.$$

Define U_i as the conditional value of U on $\{B_0 = i\}$. Assume the already acquired bounds

$$p \in [2/5, 3/5], \qquad |U| < 3/5, \qquad d = 1 - (A+B)/\ell < 9/125, \qquad \lambda = n\mathbb{E}U^2 \leq 3/40. \tag{53}$$

The gap implies $A \leq \bar{A}$, $B \leq \bar{B}$, and hence $d \geq 0$.

Theorem 6.1. *Under these hypotheses, the entire conditional supports satisfy* $|U| < 1/2$. *The conclusion is uniform in all raw* H *and all* $n \geq 42$. *It includes the closed noise endpoint* $\lambda = 3/40$ *and zero noise.*

The probability interval in this abstract theorem is $[0.4, 0.6]$. The currently acquired entry supplies the incoming strip and inward bounds only on the union of panels 40–52, namely $[0.40, 0.53]$. The application below is restricted to that smaller interval. In particular this section does not acquire the strip on $(0.53, 0.6]$ or treat the larger-noise region.

6.2 One common moment enclosure

Put $v = pq$, $s = \mathbb{E}U^2 = \lambda/n$, and $\theta = \mathbb{E}(X - \mathbb{E}[X \mid B_0])^2$. Conditional centering and variance give the exact identities

$$\ell^2 = (v+s)^{-1}, \qquad \theta = \frac{s}{v+s} = \frac{\lambda}{nv+\lambda}, \qquad \ell^2 = \frac{1-\theta}{v}. \tag{54}$$

Throughout (53), $6/25 \leq v \leq 1/4$ and $\max(p, q) \leq 3/5$. Use the rational constants

$$\Theta = \frac{37}{5000}, \qquad E = \frac{249}{125}, \qquad r = \frac{221}{200}.$$

Then

$$\theta \leq \frac{5}{677} < \Theta, \qquad \ell^2 \geq 4(1-\Theta) > E^2,$$

with exact differences $\Theta - 5/677 = 49/3385000$ and $4(1-\Theta) - E^2 = 73/31250$.

Because B_0 is the actual sign of X, the absolute cubic expansion has no sign approximation:

$$\rho := \mathbb{E}|X|^3 = \ell^3 \left\{ v(1-2v) + 3\mathbb{E}[|B_0 - p|U^2] + \mathbb{E}[\operatorname{sgn}(B_0 - p)U^3] \right\}. \tag{55}$$

The linear term vanishes by conditional centering. The incoming strip, used before the conclusion, gives $|U|^3 \leq (3/5)U^2$ and $|B_0 - p| \leq 3/5$. Also $\ell^3 \leq v^{-3/2}$. Therefore

$$\rho \leq \rho_B(v) + \frac{12/5}{v^{3/2}} \frac{\lambda}{n}, \qquad \rho_B(v) = \frac{1-2v}{\sqrt{v}}.$$

The function ρ_B decreases on $(0, 1/4]$, since its derivative is $-\frac{1}{2}v^{-3/2} - v^{-1/2} < 0$. The exact squared comparison

$$24^2(6/25)^3 - (12/5)^2 = \frac{34416}{15625} > 0$$

gives $\rho \leq \rho_B(6/25) + 24\lambda/n < r$. For the last step,

$$r - \frac{3}{70} = \frac{1487}{1400} > 0, \qquad \left(r - \frac{3}{70}\right)^2 \frac{6}{25} - \left(\frac{13}{25}\right)^2 = \frac{8707}{24500000} > 0.$$

If $\theta_\pm$ are the two nonnegative conditional variance contributions in physical coordinates, their sum is θ. Expanding the signed second moment gives

$$\beta := \mathbb{E}[X|X|] = (1-\theta)(1-2p) + \theta_+ - \theta_-.$$

Consequently, for the actual p, including $p > 1/2$,

$$1 - 2p - 2(1-p)\Theta \leq \beta \leq 1 - 2p + 2p\Theta. \tag{56}$$

All these bounds concern one common marginal. No assertion about independent conditional laws or a moment relaxation is used.

6.3 All-threshold contact and exterior rays

Lemma 6.2 (Uniform heat bound and ray continuation)**.** *Under fixed-target contact, nonnegative excess and the sharp predecessor, let $r \geq \rho$ and let b bound the directional signed moment: β on the positive ray, $-\beta$ on the negative ray. At a radius $u > \sqrt{3}$, the polynomials* (58)–(59) *below satisfy the following implication. If $Q_{r,b}(u) < 0$ and $P_{r,b}(u) < 0$, that boundary is absent from the closed support. If also $Q', Q'', L, L', L'' < 0$ there, with the moment cap held fixed, the entire closed exterior ray is absent. This assertion requires no probability panel or incoming noise strip and is uniform in all raw H and $n \geq 42$.*

Proof. For $w = n-1$, $s_n = \sqrt{n/w}$, $F_w(t) = \Phi(t/\sqrt{w})$ and $e_n = n(s_n - 1) - 1/2$, fixed-target contact and the predecessor bound give, at every closed-support point t,

$$0 \leq \mathcal{H}_n(H,t) + c\{-|t|^3 + \tfrac{3}{2}\rho t^2 + 3\beta t + \rho e_n\} - na, \tag{57}$$

where

$$\frac{\mathcal{H}_n(H,t)}{n^{3/2}} = F_w(H-t) - F_n(H) + tF_n'(H) - \frac{t^2-1}{2}F_n''(H).$$

The term $-na$ is nonpositive and may be dropped. This is a fixed-target statement; no equality between c and the actual score is assumed.

The spatial and variance expansions were derived in (27), together with each of their four remainder bounds. They apply for every $u \geq 1$, before any conditional strip is used. For $t = \pm u$, write $w = n-1$, $z = H/\sqrt{w}$, $A_u = (u^3 - 3u)/6$ and $H_4(u) = u^4 - 6u^2 + 3$. Then

$$\frac{\mathcal{H}_n(H, \pm u)}{s_n^3} = \pm A_u(1-z^2)\phi(z) + \frac{H_4(u)}{24\sqrt{w}}\phi'''(z) + R,$$

$$|R| \leq \frac{u^5/100 + 3u/20}{w} + \frac{51(3u^2-2)}{960w^{3/2}}.$$

For $u > \sqrt{3}$, $A_u > 0$. The bounds $\|(1-z^2)\phi(z)\|_\infty = \phi(0)$ and $\|\phi'''\|_\infty \leq 2/\pi < 2\phi(0)$ control both signs. Unlike the initial large-radius calculation, we retain $|H_4(u)|$; its sign is not assumed positive.

Set $S = 81/80$, $c_0 = 819/2000 < c$ and $e = 8/875$. For $n \geq 42$, $s_n < S$, $\sqrt{w} > 6$, and $e_n < e$. For completeness, $ne_n = s_n^2(2s_n+1)/(2(s_n+1)^2)$ is increasing for $s_n \geq 1$; its value at S is $793881/2073680 < 48/125$. Define

$$E_u = u^5/4100 + 3u/820 + 17(3u^2-2)/78720,$$

$$Q_{r,b}(u) = -u^3 + \tfrac{3}{2}ru^2 + 3bu + er, \tag{58}$$

$$P_{r,b}(u) = \tfrac{2}{5}S^3\{A_u + |H_4(u)|/72\} + S^3E_u + c_0Q_{r,b}(u). \tag{59}$$

If the directional signed moment is at most b, $Q < 0$ licenses replacing the positive multiplier c by c_0. Then $P < 0$ contradicts (57) at that boundary for every raw H.

To continue along a physical ray, hold r, b fixed. The derivative estimate established in the proof of Theorem 4.1 is

$$|\partial_t \mathcal{H}_n| < \tfrac{2}{5}(t^2/2 + 1/2 + e).$$

Put $L_{r,b}(u) = \frac{2}{5}(u^2/2 + 1/2 + e) + c_0Q_{r,b}'(u)$. If Q, Q', Q'', L, L', L'' are negative at a boundary, the signs persist along the whole ray: $Q''' = -6$ and $L'' = 2/5 - 6c_0 < 0$. Thus the contact upper bound strictly decreases there. Boundary atoms and the entire closed exterior ray are excluded. □

6.4 A continuous one-parameter certificate

For the negative ray use $z = p$ and the signed moment $-\beta$; for the positive ray use $z = 1 - p$ and β. In both cases $z \in [2/5, 3/5]$, and (56) gives the directional cap

$$b_z = 2z - 1 + 2(1 - z)\Theta.$$

This folds the actual probability, not the endpoints of a probability panel. It does not reflect the chosen discrepancy or change its law. Take the boundary $u = E(z + 1/2)$. Thus

$$u \in [u_0, u_1] = [2241/1250, 2739/1250], \qquad b_z = \alpha u + \delta, \quad \alpha = \frac{4963}{4980}, \quad \delta = -\frac{9889}{5000}.$$

Here $u_0^2 > 3$, so $H_4'(u) = 4u(u^2 - 3) > 0$; $H_4(u_1) < -2.75$ proves $H_4 < 0$ on this whole interval. Let $\mathcal{P}(u) = P_{r,\alpha u+\delta}(u)$. Replacing $|H_4|$ by $-H_4$ in (59) gives an explicit quintic $\mathcal{P}(u) = \sum_{j=0}^{5} a_j u^j$, with

$$\begin{aligned}
a_5 &= S^3/4100, & a_4 &= -\tfrac{2}{5}S^3/72,\\
a_3 &= \tfrac{2}{5}S^3/6 - c_0, & a_2 &= \tfrac{2}{5}S^3/12 + 51S^3/78720 + c_0(3r/2 + 3\alpha),\\
a_1 &= -\tfrac{2}{5}S^3/2 + 3S^3/820 + 3c_0\delta, & a_0 &= -\tfrac{2}{5}S^3/24 - 34S^3/78720 + c_0 er.
\end{aligned}$$

The exact rational endpoint inequalities are

$$\mathcal{P}'''(u_0) < -2.24, \qquad \mathcal{P}'''(u_1) < -2.27, \qquad \mathcal{P}'(u_0) > 0.915, \quad \mathcal{P}'(u_1) > 0.745.$$

Indeed $\mathcal{P}'''$ is a convex quadratic, with second derivative $120a_5 > 0$. Its negative endpoint values make it negative throughout. Therefore $\mathcal{P}'$ is concave and exceeds the chord between its positive endpoint values. It follows that $\mathcal{P}$ increases and

$$\mathcal{P}(u) \le \mathcal{P}(u_1) = -\frac{4560620039489147528734839}{25625000000000000000000000} < -0.1779. \tag{60}$$

One stronger-looking shortcut is false: $\mathcal{P}''(u_0) > 0.0228$. It is not used.

The multiplier and ray signs require a separate calculation. The boundary family

$$\mathcal{Q}(u) = Q_{r,\alpha u+\delta}(u) = -u^3 + (3r/2 + 3\alpha)u^2 + 3\delta u + er$$

has total derivatives $\mathcal{Q}''(u_0) < -1.46$ and $\mathcal{Q}'(u_1) = 179223/6250000 > 0.0286$. Since $\mathcal{Q}''$ decreases, $\mathcal{Q}'$ decreases and stays positive. Hence $\mathcal{Q}$ increases, and $\mathcal{Q}(u) \le \mathcal{Q}(u_1) < -1.198$.

For each physical ray the directional cap is fixed. Its derivatives, evaluated across the boundary family, are instead

$$\begin{aligned}
q_1(u) &= -3u^2 + (3r + 3\alpha)u + 3\delta, & q_2(u) &= -6u + 3r,\\
l(u) &= \tfrac{2}{5}(u^2/2 + 1/2 + e) + c_0 q_1(u), & l_1(u) &= \tfrac{2}{5}u + c_0 q_2(u), \qquad l_2 = \tfrac{2}{5} - 6c_0.
\end{aligned}$$

They are not the total derivatives of $\mathcal{Q}$. The following rational upper bounds certify their signs on the interval:

family	value at u_0	total derivative at u_0
q_1	< -4.272	< -4.452
q_2	< -7.441	$= -6$
l	< -0.903	< -1.105
l_1	< -2.330	$= -2.057$

The two quadratic families have negative constant second derivatives $q_1'' = -6$ and $l'' = -2.057$; the others are affine decreasing. Thus all four stay negative, as does $l_2 = -2.057$. Together with $\mathcal{Q} < 0$, these are the six required signs at every boundary. Holding its actual directional cap fixed, the preceding ray argument excludes the whole exterior ray beyond each boundary.

6.5 Completion, application and remaining premises

For each actual p the two excluded rays start at

$$u_- = E(p + 1/2), \qquad u_+ = E(3/2 - p).$$

Because $\ell > E$, the remaining support satisfies $-1/2 < Y = X/\ell + p < 3/2$ strictly. The already proved actual gap supplies

$$U_0 \le (\bar{A} - A)/\ell \le d < 9/125, \qquad U_1 \ge -(\bar{B} - B)/\ell \ge -d > -9/125.$$

For $B_0 = 0$ the new lower bound is $U_0 > -1/2$; for $B_0 = 1$ the new upper bound is $U_1 < 1/2$. This proves Theorem 6.1. If $\lambda = 0$, then $\mathbb{E}U^2 = 0$ and $U = 0$ directly. There is no division by noise or limiting argument.

Corollary 6.3. *Retain the accepted fixed-target entry and the ordered elementary moment/gap contraction on its 30 probability panels. On its acquired subrange $p \in [0.40, 0.53]$, all actual $n \ge 42$ with $\lambda \in [0, 3/40]$ satisfy $|U| < 1/2$ for every raw H. The previously acquired inward bound $d < 0.072 < 1/4$ is retained.*

Proof. That antecedent provides the strip $|U| < 0.6$ on the 13 closed panels 40–52 and $d \le 0.0719539227 < 0.072$. Their closed union is $[0.40, 0.53]$. Apply the theorem at the actual p. □

6.6 The central panel at larger noise

We now restrict to the oriented panel $p \in [l, h] = [0.41, 0.42]$ and use (46). Set

$$N = 42, \quad \Theta = 0.054, \quad d_0 = 0.069, \quad E = 1.9706, \quad F_\ell = 2.034, \quad \sigma_0 = 0.6,$$

$$r_0 = 1/2, \quad \tau = 1/5, \quad \zeta = 10^{-5}, \quad \kappa = 997/15000 < \phi(0)/6,$$

$$c_l = 0.4095 < c = C_{\mathrm{E}} < c_u = 0.40974.$$

Here E is a lower scale bound and σ_0 is the incoming strip width, not $\sqrt{10}$. These constants replace the older floor-47, width-0.55 confinement parameters; every gate below uses floor 42 and the already acquired width 0.6.

Use distinct count-coordinate inner endpoints $a_0 = p - A/\ell$, $b_0 = p + B/\ell$. They are support points with $0 \le a_0 \le d$, $1 - d \le b_0 \le 1$ and $b_0 - a_0 = 1 - d$. The inner inequalities $U_0 \le a_0$, $U_1 \ge b_0 - 1$ are weak; strictness comes from $d < d_0$. Let

$$x = np + H/\ell, \quad \nu = \mathrm{Law}\left(x - \sum_{i<n} Y_i\right), \qquad J_{\mathrm{gap}} = \int_{[a_0, b_0)} (y - a_0)\nu(dy).$$

The central gap contributes $m\ell J_{\mathrm{gap}} = (1 - \theta)J_{\mathrm{gap}}$ to the scalar leakage, so

$$\mathcal{L} \ge (1 - \theta)J_{\mathrm{gap}}. \tag{61}$$

Write $g(y) = \nu([y, \infty))$ and let Q_c be the count-coordinate contact majorant, with $g \le Q_c$ everywhere and equality on the closed support. Differentiating fixed-target contact gives

$$\sqrt{n}(-Q_c'(y)) = \ell\phi(H/\sqrt{n}) + \frac{\ell}{n}\{3c\beta + (H\phi(H/\sqrt{n}) + 3c\rho)t - 3ct|t|\}, \qquad t = \ell(y - p). \tag{62}$$

The excess changes only an additive constant. The multiplier here is c, not an upper bound on the actual score.

6.6.1 One resource for displacement, noise and leakage

Define the nonnegative endpoint distances

$$M_+ = \mathbb{E}[X - B; X > 0] = m - pB, \qquad M_- = \mathbb{E}[-X - A; X < 0] = m - qA.$$

With $\sigma = \sqrt{10}$, the endpoint expansion of the actual defect gives

$$\begin{aligned} D \geq m\{\sqrt{\sigma+1}A - \sqrt{\sigma-1}B\}^2 + 2(\sigma+2)B^2M_+ + 2(\sigma+4)A^2M_- \\ + 3(\sigma+1)B\theta_+ + 3(\sigma+3)A\theta_-. \end{aligned}$$

Indeed put $Z = X - B \geq 0$ on the positive class. Then

$$T_+ = 2B^2M_+ + 3B\mathbb{E}[Z^2; X > 0] + \mathbb{E}[Z^3; X > 0], \quad R_+ = BM_+ + \mathbb{E}[Z^2; X > 0].$$

The combination $(\sigma-1)T_+ + 6BR_+$ gives the stated positive terms because $\mathbb{E}[Z^2; X > 0] \geq \theta_+$. The negative expansion is the same magnitude calculation with $\sigma+1$. The remaining constant term is the square since $2\sqrt{(\sigma+1)(\sigma-1)} = 6$. Discard this nonnegative square and use $A \geq E(l - d_0)$, $B \geq E(1 - h - d_0)$. The positive coefficients

$$\begin{aligned} c_+ &= 2(5.16)E^2(1-h-d_0)^2, & c_- &= 2(7.16)E^2(l-d_0)^2, \\ b_* &= \min\{3(4.16)E(1-h-d_0), 3(6.16)E(l-d_0)\} \end{aligned}$$

satisfy

$$D \geq c_+M_+ + c_-M_- + b_*\theta, \qquad d = \frac{M_+/p + M_-/q}{\ell}, \qquad \epsilon := \mathbb{E}|U| \leq \frac{2(M_+ + M_-)}{\ell}. \quad (63)$$

The last bound follows signwise from $\mathbb{E}|Z - \mathbb{E}Z| \leq 2\mathbb{E}Z$ for $Z \geq 0$.

On a closed noise band $[\lambda_0, \lambda_1]$, let S_ε be the whole-rectangle resource from (52). Because $n\theta = \ell^2\lambda \geq E^2\lambda_0$, the single common budget implies

$$nc_+M_+ + nc_-M_- + n^{3/2}\mathcal{L}/\kappa \leq R_*, \qquad R_* = S_\varepsilon/\kappa - b_*E^2\lambda_0. \quad (64)$$

A strictly negative R_* excludes a law; equality is retained. With $\alpha = \min\{c_+lE, c_-(1-h)E\} > 0$ and $G = R_*/\alpha$, the same simplex gives $nd \leq G$. These costs are simultaneous, so the defect budget cannot be spent independently on each one.

6.6.2 Concentration at both deleted indices

Independently couple the two conditional noises and sample an independent Bernoulli label. This realizes the actual marginal as $Y = U_0 + B_0(1 + U_1 - U_0)$, with increment at least $1 - d > 0$. Conditional on all pairs, an interval of diameter strictly less than $1 - d$ selects an antichain of label sets. For an antichain $\mathcal{A}$ in j labels, a uniformly random permutation gives a chain meeting it at most once. Taking expectations yields

$$\sum_{A\in\mathcal{A}} \binom{j}{|A|}^{-1} \leq 1, \qquad \mathbb{P}(\mathcal{A}) = \sum_{A\in\mathcal{A}} \binom{j}{|A|}^{-1} \binom{j}{|A|} p^{|A|}q^{j-|A|} \leq \max_k \binom{j}{k} p^k q^{j-k}.$$

Conditioning back proves the bound for the actual iid sum, also after an independent shift. Strict diameter is essential: a closed interval equal to a Bernoulli increment can contain both atoms.

Both actual deleted indices satisfy $j \geq n - 2 \geq 40$. Put

$$r_L = l - (1-l)/40, \quad r_R = h + h/40, \qquad v_* = \min\{r_L(1-r_L), r_R(1-r_R)\}.$$

The gate $0 < r_L < r_R < 1$ holds. Adjacent binomial-mass ratios give $(j+1)p-1 \le k \le (j+1)p$ for a mode; hence $r = k/j \in [r_L, r_R]$ and the mode is interior. The classical finite Stirling inequalities $\sqrt{2\pi j}(j/e)^j < j! < \sqrt{2\pi j}(j/e)^j e^{1/(12j)}$ give

$$\sqrt{j}\binom{j}{k}p^k q^{j-k} < \frac{\phi(0)e^{1/(12j)-jD(r\|p)}}{\sqrt{r(1-r)}} < \frac{0.4}{\sqrt{v_*}(1-1/480)} < M.$$

Here entropy is nonnegative by Jensen's inequality, and $e^z \le (1-z)^{-1}$ for $0 \le z < 1$ follows termwise from its power series. Choose M by taking the downward 10^{-10} grid square root of $0.16/[v_*(1-1/480)^2]$ and adding one grid unit; thus M is strictly larger than the exact root. Thus every interval of strict width below $1-d$ has probability less than $M/\sqrt{j}$. No bound on C_{n-2} is assumed. Finally

$$F_d = 1.037, \qquad F_d^2 > \max\{(42/41)^3, 42/40\}. \tag{65}$$

These ratios decrease with n and control the distance factor from $n-1$ summands and the noise/pair-event factor from $n-2$ summands.

6.6.3 Signed deletion, including endpoint atoms

Let $m_n = n-1$, $Z = x - \sum_{i=1}^{m_n} Y_i$, $\mu = x - m_n p = p + H/\ell$, and let K count the labels. For a removed label define $T_i(I) = \mathbb{P}(x - \sum_{k<m_n} Y_k - i - U_i \in I)$. Exchangeability and $m_n p - K = Z - \mu + \sum U_k$ give exactly

$$\nu(I) - T_1(I) = \frac{\mathbb{E}[(Z - \mu + \sum U_k)\mathbf{1}_I(Z)]}{m_n p},$$
$$T_0(I) - \nu(I) = \frac{\mathbb{E}[(Z - \mu + \sum U_k)\mathbf{1}_I(Z)]}{m_n q}.$$

For a removed full pair, the remaining conditional interval probability is $f(Y_k) \in [0, M/\sqrt{n-2}]$. Since $\mathbb{E}U_k = 0$,

$$\mathbb{E}[U_k f(Y_k)] \le \frac{M}{\sqrt{n-2}}\mathbb{E}(U_k)_+ = \frac{\epsilon M}{2\sqrt{n-2}}.$$

The signed term gives the factor $\epsilon/2$ and also covers deterministic conditional noise.

For independently coupled noises let $\Delta = U_1 - U_0$. Then $\Delta \ge -d$, $\mathbb{E}\Delta = 0$, and $\operatorname{Var}\Delta \le s/p$, since $p < q$ and $s = \lambda/n$. On $\Delta \le \tau$, the point for T_1 in I maps to the point for T_0 in $I^+ = I + 1 + [-d, \tau]$. Conditional concentration, Cantelli and Markov applied to $\Delta + d \ge 0$ give

$$T_1(I) \le T_0(I^+) + \frac{M}{\sqrt{n-2}}\mathbb{P}(\Delta > \tau),$$
$$\mathbb{P}(\Delta > \tau) \le \min\left\{\frac{s}{s+p\tau^2}, \frac{d}{d+\tau}\right\} \le B_\tau := \min\left\{\frac{\lambda_1}{\lambda_1 + Nl\tau^2}, \frac{G}{N\tau + G}\right\}. \tag{66}$$

The exceptional event is strict; $\Delta = \tau$ is in the good event. All intervals below have strict widths below $1-d$. If D_I, D_J bound the positive parts of their upper endpoints minus μ, the two deletion identities imply

$$\sqrt{n}\,\nu(I) \le \sqrt{n}\,\nu(I^+) + A_*/n + a_e\epsilon + B_*^{\mathrm{err}}\mathbb{P}(\Delta > \tau), \tag{67}$$

where

$$a_e = \frac{F_d M}{2}\left(\frac{1}{l} + \frac{1}{1-h}\right), \quad A_* = F_d M\left(\frac{D_I}{l} + \frac{D_J}{1-h}\right), \quad B_*^{\mathrm{err}} = F_d M.$$

Indeed $\sqrt{n}/(n-1)^{3/2} \le F_d/n$ and $\sqrt{n/(n-2)} \le F_d$. Half-open intervals may be enlarged to closed ones for upper concentration bounds; no CDF continuity is assumed.

6.6.4 Whole-band slope bounds

Fix $|H| \in [h_0, h_1]$ with its original sign and a noise band $[\lambda_0, \lambda_1]$. Let r_B be the outward upper bound for $(1-2l(1-l))/\sqrt{l(1-l)}$ obtained by dividing its numerator by the downward 10^{-10} grid square root of $l(1-l)$, and set

$$\rho_* = \min\{1.22, r_B + 24\lambda_1/N\}, \quad \beta_- = 0.09736, \quad \beta_+ = 0.234,$$

$$\varphi_- = 0.39886(1 - h_1^2/(2N)) > 0, \qquad v_H^+ = \begin{cases} 0.4h_1, & H \geq 0, \\ -\varphi_- h_0, & H \leq 0. \end{cases}$$

The coefficient 24 is licensed by the incoming 0.6 strip: $24^2[l(1-l)]^3 > [3(1-l)+0.6]^2$. It follows from the same-law cubic expansion before the smaller strip is acquired. The density floor follows from $e^{-z} \geq 1 - z$; v_H^+ bounds $H\phi(H/\sqrt{n})$ in the correct direction for both signs.

Take $B^* = 1.2 > \sqrt{(1-l)/l}$ and $f = 0.4F_\ell$. Define

$$\begin{aligned} C_{\mathrm{sl},-} &= F_\ell((v_H^+ + 3c_u\rho_*)^2/(12c_l) - 3c_l\beta_-)_+, \\ C_{\mathrm{sl},+} &= F_\ell(0.4h_1B^* + 3c_uB^*(B^*-1) - 3c_l\beta_-)_+, \\ L_g &= E\varphi_- - \max(C_{\mathrm{sl},-}, C_{\mathrm{sl},+})/N, \\ U_+ &= f + \frac{F_\ell}{N}\left\{3c_u\beta_+ + \frac{(0.4h_1 + 3c_u\rho_*)^2}{12c_l}\right\}. \end{aligned}$$

For $t \leq 0$, the bracket in (62) is bounded below by $3c_l\beta_- + (v_H^+ + 3c_u\rho_*)t + 3c_lt^2$. The gate $v_H^+ + 3c_u\rho_* > 0$ permits this square completion. On the central positive interval $0 \leq t \leq B^*$ use $\rho \geq 1$ and $t(t-1) \leq B^*(B^*-1)$ for $t \geq 1$; for $t \leq 1$ the latter term has favorable sign. Thus L_g is a lower slope throughout the central gap. On the full positive half-line, completing the downward quadratic gives the upper slope U_+. Negative corrections in lower bounds and positive corrections in upper bounds use $\ell/n \leq F_\ell/N$.

The lower exterior interval has a sharper slope. Put

$$t_* = -E(l + \tau + \zeta), \quad B(t) = \beta_- + \rho_* t + t^2, \quad B_{\min} = B(\min\{t_*, -\rho_*/2\}).$$

For $t \leq t_* < 0$,

$$3c(\beta + \rho t + t^2) + H\phi t \geq 3c_lB(t) + v_H^+ t + 3(c_u - c_l)\min(0, B_{\min}).$$

First use $\beta \geq \beta_-$, $\rho \leq \rho_*$ and $t < 0$; when varying c, only the possible negative part of B incurs the final correction. Minimize the remaining upward quadratic at $\min\{t_*, -(v_H^+ + 3c_l\rho_*)/(6c_l)\}$ and denote the total minimum by q_*. Then

$$L = E\varphi_- - F_\ell(-q_*)_+/N$$

is a lower slope on the entire lower interval. The finite gates require $L > 0$, $L_g > 0$ and $U_+ > 0$ on every box.

6.6.5 Two strict interval contradictions

For a putative support point $a_0 - r$, $r \geq r_0$, take

$$I = [a_0 - r, a_0 - d - \tau - \zeta], \qquad I^+ = [a_0 - r + 1 - d, b_0 - \zeta].$$

The incoming strip gives $r < \sigma_0 + d_0$. The exact gates $g_0 = 1 - \sigma_0 - 2d_0 > 0$ and $\sigma_0 + d_0 < 1 - d_0$ place $I^+ \subset [a_0 + g_0, b_0)$ and make both widths strictly less than $1 - d$. The intervals are disjoint. Contact at the left endpoint and the global majorant at the right endpoint give

$$\sqrt{n}\,\nu(I) \geq L(r - d - \tau - \zeta), \qquad \nu(I^+) \leq J_{\mathrm{gap}}/g_0.$$

Indeed $\nu([a_0 - r, a_0 - d - \tau - \zeta)) \geq Q_c(a_0 - r) - Q_c(a_0 - d - \tau - \zeta)$; a closed interval only increases its mass. No contact at the right endpoint is needed. For the directed distances use

$$v_H = \begin{cases} -h_0/F_\ell, & H \geq 0, \\ h_1/E, & H \leq 0, \end{cases} \qquad D_I = (v_H - l - \tau - \zeta)_+, \quad D_J = (v_H + 1 - l - \zeta)_+.$$

These are independent of r. Substitution in the deletion bound shows that a lower support point would require

$$L(r - \tau - \zeta) \leq Ld + a_e\epsilon + \sqrt{n}J_{\mathrm{gap}}/g_0 + A_*/n + B_*^{\mathrm{err}}\mathbb{P}(\Delta > \tau). \tag{68}$$

As $L > 0$, a strict contradiction at $r = r_0$ excludes the entire lower part of the acquired strip.

For a putative support point $y \geq 1 + r_0$, take

$$I = [a_0, a_0 + \tau], \qquad I^+ = [b_0, a_0 + 1 + 2\tau].$$

The strict gates $d_0 + 2\tau < r_0$, $1 - d_0 - \tau > 0$ and $1 - 2d_0 - 2\tau > 0$ give disjointness, widths below $1 - d$, and $I^+ \subset [b_0, y)$. Since $\nu([a_0, b_0)) = Q_c(a_0) - Q_c(b_0)$ and $\nu([a_0 + \tau, b_0)) \leq J_{\mathrm{gap}}/\tau$, contact implies

$$\sqrt{n}\,\nu(I) \geq L_g(1 - d) - \sqrt{n}J_{\mathrm{gap}}/\tau, \qquad \sqrt{n}\,\nu(I^+) \leq U_+(\sigma_0 + d).$$

The last bound uses contact at b_0, y, includes an atom at b_0 and excludes the contacting atom at y. The strict containment is essential. For this side use

$$D_I = (d_0 + \tau + h_1/E - l)_+, \qquad D_J = (1 + d_0 + 2\tau + h_1/E - l)_+.$$

An upper support point would therefore require

$$L_g - \sigma_0 U_+ \leq (L_g + U_+)d + a_e\epsilon + \sqrt{n}J_{\mathrm{gap}}/\tau + A_*/n + B_*^{\mathrm{err}}\mathbb{P}(\Delta > \tau). \tag{69}$$

6.6.6 Two error bounds on the same resource simplex

For either side its right-hand error is $a_d d + a_e\epsilon + a_J\sqrt{n}J_{\mathrm{gap}} + A_*/n + B_*^{\mathrm{err}}\mathbb{P}(\Delta > \tau)$, with

$$(a_d, a_J) = \begin{cases} (L, 1/g_0), & \text{lower side}, \\ (L_g + U_+, 1/\tau), & \text{upper side}. \end{cases}$$

Equations (63) and (61) give the three linear costs per unit of (64):

$$z_+ = \frac{a_d/l + 2a_e}{Ec_+}, \qquad z_- = \frac{a_d/(1 - h) + 2a_e}{Ec_-}, \qquad z_J = \frac{\kappa a_J}{1 - \Theta}.$$

For example the positive contribution is at most $(a_d/l + 2a_e)M_+/E = z_+(nc_+M_+)/n$; the leakage contribution is at most $z_J(n^{3/2}\mathcal{L}/\kappa)/n$. Let $z = \max(z_+, z_-, z_J)$. The gate $zR_* + A_* > 0$ licenses replacing n by N, giving

$$\mathcal{E}_0 = (zR_* + A_*)/N + B_*^{\mathrm{err}}B_\tau. \tag{70}$$

This includes retained $R_* = 0$ boxes; resource equality is not excluded.

A second bound pays for the pair event inside the same simplex. At the three nonzero vertices, upper linear-cost/displacement pairs are

$$(v_+, w_+) = \left(z_+R_*/N, \frac{R_*}{Ec_+l}\right), \qquad (v_-, w_-) = \left(z_-R_*/N, \frac{R_*}{Ec_-(1 - h)}\right),$$
$$(v_J, w_J) = (z_JR_*/N, 0).$$

They bound the linear error and nd. The nonnegative origin contributes zero and can be included without changing their maximum. For $t, x \geq 0$,

$$\frac{x}{N\tau + x} \leq \frac{N\tau}{(N\tau + t)^2} x + \frac{t^2}{(N\tau + t)^2},$$

because the difference is $N\tau(x-t)^2/[(N\tau+x)(N\tau+t)^2] \geq 0$. Use $d/(d+\tau) = nd/(n\tau+nd) \leq nd/(N\tau + nd)$. Every tangent gives

$$\mathcal{E}_t = \frac{A_*}{N} + \max_{i \in \{+,-,J\}} \left\{ v_i + B_*^{\mathrm{err}} \frac{N\tau}{(N\tau + t)^2} w_i \right\} + B_*^{\mathrm{err}} \frac{t^2}{(N\tau + t)^2}. \tag{71}$$

Take the minimum of (70) and the eleven valid bounds (71) at $t = jG/10$, $j = 0, \ldots, 10$. Each bounds one actual resource simplex; no incompatible maximizing allocations are combined.

Certificate condition 6.4 (Complete central-panel confinement gates)**.** Use both signs of H, eight closed phase bands $[907j/4000, 907(j+1)/4000]$, $j = 0, \ldots, 7$, and eight closed noise bands $[3k/40, 3(k+1)/40]$, $k = 0, \ldots, 7$. Retain all 128 boxes in the coverage list. On each use the complete incident-cell resource of Condition 5.4. Exclude only an empty incident list, strictly negative S_ε or strictly negative R_*. Otherwise check all domains, directed roots, positive slopes and inverse-index gates above, then require both strict inequalities, with the error computed for the respective side,

$$L(r_0 - \tau - \zeta) > \mathcal{E}, \qquad L_g - \sigma_0 U_+ > \mathcal{E}. \tag{72}$$

For the lowest noise band the all-threshold low-noise theorem may replace these two ray gates.

Theorem 6.5 (Complete conditional confinement on the central panel)**.** *Under the fixed-target entry and contact hypotheses, the panel contraction and signed-resource conditions, and Condition 6.4, every actual $n \geq 42$ with $p \in [0.41, 0.42]$ satisfies $|U| < 1/2$ on the entire conditional supports and $d < 0.069 < 1/4$.*

Proof. The acquired $|H| < 1.814$ and $\lambda < 0.6$ lie inside the closed cover. The 16 first-band boxes are covered by Corollary 6.3, uniformly in H. On the 112-box complement the exact gate evaluation has this census; the finite inequalities themselves, not just the count, are required:

necessary exclusion or strict conclusion	boxes
no incident actual moment cell	14
negative whole-rectangle resource	5
noise lower-endpoint cost exceeds the resource	65
both strict contact-ray inequalities	28

All 56 inequalities in the last row have strict margin greater than 0.00892011917. The lower contradiction extends to every $r \in [r_0, \sigma_0 + d_0)$ because its error is independent of r and $L > 0$. The upper contradiction already applies to every support point $y \geq 1 + r_0$ in the acquired strip. No support point therefore lies at or below $a_0 - r_0$ or at or above $1 + r_0$. Since $a_0 \geq 0$, $b_0 \geq 1 - d$ and $d < d_0 < r_0$, both conditional noises have strict amplitude below r_0. Boundary atoms are included in the contradictions. The seam $3/40$ belongs to both adjacent closed bands; both phase signs are retained at $H = 0$. □

The finite gates in the final condition were checked by exact replay and a separate reconstruction of the interval, slope and common-resource formulas. This computational statement is separate from the analytic derivation. The unassisted confinement diagnostics remain failed sufficient bounds in their recorded low-noise boxes; the composition uses the proved low-noise theorem there without changing those failures.

Thus low-noise confinement is proved conditionally on $[0.40, 0.53]$, and complete acquired-noise confinement conditionally on $[0.41, 0.42]$. Higher noise on the other panels and strict count comparison remain separate obligations. In particular, count formulas requiring $p \le 1/2$ are not transferred to above-half panels by reflecting the already chosen discrepancy.

7 Conditional counts and discrete curvature

The sign decomposition turns an iid sum into a binomial mixture of conditional noise sums. This identity supplies a comparison with the Bernoulli law having the same sign probability. The main issue is to control the error proportionally to the noise variance, including when that variance is arbitrarily small. Throughout this section the target is $c = C_{\mathrm{E}}$, and a is the fixed-target excess. The identities below hold for any actual marginal with the stated moments; extremality is used only when we apply them to a witness with $a \ge 0$.

7.1 The exact count and score identities

Let $\pi_j = \binom{n}{j} p^j q^{n-j}$ for $0 \le j \le n$, zero otherwise, and F_B be the CDF of $\mathrm{Bin}(n, p)$. For independent copies of the marginal, conditioning on the entire sign vector gives independent conditional residuals. All sign vectors with j positive entries yield the same sum law

$$V_j = U_{1,1} + \cdots + U_{1,j} + U_{0,j+1} + \cdots + U_{0,n}.$$

It follows from (16) that at $x = np + H/\ell = k + u$,

$$\mathbb{P}(S_n \le H) = \sum_{j=0}^{n} \pi_j \mathbb{P}(V_j \le k + u - j). \tag{73}$$

For $0 \le k \le n$, subtracting $F_B(k)$ gives the exact signed identity

$$\begin{aligned} Q &:= \mathbb{P}(S_n \le H) - F_B(k) \\ &= -\sum_{i=0}^{k} \pi_{k-i} \mathbb{P}(V_{k-i} > u + i) + \sum_{i=1}^{n-k} \pi_{k+i} \mathbb{P}(V_{k+i} \le u - i). \end{aligned} \tag{74}$$

The outgoing event is strict and the incoming events are closed. Thus an atom exactly on a count plane has its correct weight.

Choosing a nearest integer gives $|u| \le 1/2$, with either integer allowed at a tie. Restricting to legal counts $0 \le k \le n$ requires either $-1/2 \le np + H/\ell \le n + 1/2$ or a separate treatment of the remaining thresholds. Interior-count normalizations further require $1 \le k \le n - 1$.

The own-count variance is

$$t = \mathbb{E}V_k^2 = (n-k)w_0 + kw_1 = \lambda + \delta(w_1 - w_0), \qquad \delta = k - np = H/\ell - u.$$

Since $|w_1 - w_0| \le s/\min(p, q)$, any established bound $|\delta| \le \Delta$ and $c_v \ge 1/\min(p, q)$ gives, for $n \ge N$,

$$(1 - c_v\Delta/N)\lambda \le t \le (1 + c_v\Delta/N)\lambda. \tag{75}$$

When the lower bound is used in a denominator, $1 - c_v\Delta/N$ must be positive.

For the comparison with the same Bernoulli count, set

$$\rho_B = (1-2v)/\sqrt{v}, \qquad C = \frac{\sqrt{n}}{\rho}\{\mathbb{P}(S_n \le H) - \Phi(H/\sqrt{n})\},$$

$$C_B = \frac{\sqrt{n}}{\rho_B}\{F_B(k) - \Phi(\delta/\sqrt{nv})\}, \qquad f(u) = \frac{\Phi((\delta+u)/\sqrt{n(v+s)}) - \Phi(\delta/\sqrt{n(v+s)})}{\pi_k}.$$

For a legal count $\pi_k > 0$. Define the exact repair

$$\mathcal{R} = \sqrt{n}\{\Phi(\delta/\sqrt{nv}) - \Phi(\delta/\sqrt{n(v+s)})\} + c(\rho_B - \rho).$$

Direct subtraction, using $H/\sqrt{n} = (\delta+u)/\sqrt{n(v+s)}$, yields

$$a + \rho_B(c - C_B) = \sqrt{n}\pi_k\{Q/\pi_k - f(u)\} + \mathcal{R}. \tag{76}$$

Both terms on the right must be estimated for the same law, threshold and phase.

Proposition 7.1 (Two conditional local witnesses)**.** *Let $\lambda > 0$, $n \ge N$, and suppose $\sqrt{n}\pi_k \ge m_0 > 0$. The following are sufficient for $a + \rho_B(c - C_B) < -\varepsilon\lambda$.*

1. *Bounds O, T, R_0, κ satisfy*

$$\mathbb{P}(V_k > u) + f(u) \ge O, \quad \sum_{i\ge 1}\mathbb{P}(V_{k+i} \le u - i) \le T, \quad \pi_{k+i}/\pi_k \le R_0, \quad \mathcal{R} \le \kappa\lambda/n,$$

   where the sum and ratio involve legal counts. If $D_0 = O - R_0T > 0$ and $m_0D_0 - \kappa\lambda/n > \varepsilon\lambda$, the conclusion holds.

2. *A proved whole-count estimate gives*

$$a + \rho_B(c - C_B) \le \sqrt{n}\pi_k\lambda B(\lambda) + \lambda I(\lambda)/n.$$

   If $B(\lambda) < 0$ and $m_0B(\lambda) + \max(0, I(\lambda))/N < -\varepsilon$, the conclusion holds.

Proof. For the first witness, discard only the outgoing terms with $i > 0$ in (74); then $Q/\pi_k - f(u) \le -D_0$. Use $D_0 > 0$ before replacing the count mass by its lower bound in (76). For the second, $B < 0$ similarly licenses the mass replacement, and $I/n \le \max(0, I)/N$ holds for either sign of I. Multiplication by $\lambda > 0$ gives the asserted strict inequality. □

For a witness with $a \ge 0$, Schulz's theorem $C_B \le c$ makes $a + \rho_B(c - C_B)$ nonnegative. Either strict negative bound therefore excludes that witness [19]. This is a comparison of actual iid sums; it is not an extremal principle for a moment relaxation.

At zero noise the threshold may be taken at a lattice point: at an attained positive maximum for a Bernoulli law, the threshold is a binomial atom. Between consecutive atoms, lowering the threshold to the preceding atom preserves the CDF and strictly improves the normal subtraction. Above the final atom the same argument applies, while below the first atom the discrepancy is nonpositive. Thus $u = 0$ and $C = C_B$ at such a maximum. The sharp Bernoulli upper bound gives $a \le 0$. To exclude equality as well, one needs its strict finite-index form; we keep that requirement explicit in the conditional global statement.

7.2 Whole-count coupling and its curvature term

Part of the whole-count estimate follows from a coupling. Suppose $U_0 \le a_0$, $U_1 \ge -a_1$, where $d_Y = a_0 + a_1 < 1/2$, and put $\zeta = 1 - d_Y$. Independently realize all $2n$ variables in the definition of V_j above, and extend V_j constantly for $j < 0$ and $j > n$. The increments of $j + V_j$ are at least $\zeta > 1/2$ inside the legal range and equal one outside. Consequently

$$K = \max\{j \in \mathbb{Z} : j + V_j \le k + u\}, \qquad R = K - k$$

are well defined. An independent binomial selector has the correct law for each count, so

$$\mathbb{P}\Big(\sum_{i=1}^{n} Y_i \le k + u\Big) = \mathbb{E}F_B(k + R). \tag{77}$$

Indeed, conditional on the coupled residuals, the closed event for the selector occurs exactly at counts not exceeding K.

Lemma 7.2 (Discrete curvature and crossing moments)**.** *Let $\chi = (n+1)p - k < 2$ and $m_B = \max_j \pi_j$. For every integer r,*

$$F_B(k + r) - F_B(k) - r\pi_k \le \frac{m_B}{(n+1)v}\left\{(\chi - 1)_+ r_+ + \chi_+ \binom{r_-}{2} + \binom{r_-}{3}\right\}, \tag{78}$$

where $r_+ = \max(r, 0)$ and $r_- = \max(-r, 0)$. If $|u| \le 1/2$ and $t = \mathbb{E}V_k^2$, then

$$\mathbb{E}R_+ \le \frac{t}{(\zeta - u)^2}, \qquad \mathbb{E}\binom{R_-}{2} \le \frac{t}{[\zeta + \min(u, \zeta/3)]^2}.$$

If additionally $|U_0|, |U_1| \le A_U$, then

$$\mathbb{E}\binom{R_-}{3} \le \frac{t\sqrt{A_U^2 + 3t}}{(2\zeta - 1/2)^3}.$$

Proof. For $0 \le j \le n$, the binomial difference identity is

$$\pi_j - \pi_{j-1} = \pi_j \frac{(n+1)p - j}{p(n - j + 1)}.$$

When its numerator is positive, $p(n - j + 1) \ge (n + 1)v$. Thus its positive part is bounded by $m_B((n+1)p - j)_+/((n+1)v)$; zero extension gives the same upper bound outside the legal range. For $r > 0$, double summation gives weights $r - i + 1$ on these differences at $j = k + i$. Since $\chi < 2$, only $i = 1$ can contribute positively, giving $(\chi - 1)_+ r$. For $r = -h < 0$, the corresponding weights are $h - 1 - i$ on the differences at $j = k - i$, $0 \le i \le h - 2$. Use $(\chi + i)_+ \le \chi_+ + i$ and sum the weights to obtain $\chi_+\binom{h}{2} + \binom{h}{3}$.

On $R = r \ge 1$, crossing monotonicity gives $-V_k \ge \zeta r - u$. Since $(\zeta r - u)^2 \ge r(\zeta - u)^2$, this proves the first moment bound. On $R = -h \le -2$, the count immediately above K is outside the closed event, giving the strict inequality $V_k > \zeta(h - 1) + u$. Put $j = h - 1$. For $u \le \zeta/3$, differentiation shows that $j(j+1)/(2(\zeta j + u)^2)$ decreases for $j \ge 1$: the derivative has the sign of $j(2u - \zeta) + u \le 0$. Its maximum is $1/(\zeta + u)^2$ at $j = 1$. For $u > \zeta/3$, replace the crossing threshold by the smaller one with $u = \zeta/3$. This proves the second bound.

For the third, use $u \ge -1/2$ and $j \ge 2$. The derivative of $(j^3 - j)/(6(\zeta j - 1/2)^3)$ has the sign of $2\zeta j - (3j^2 - 1)/2 < 0$, because $\zeta \le 1$. Its maximum at $j = 2$ is $(2\zeta - 1/2)^{-3}$. Hence $\binom{R_-}{3} \le (V_k)_+^3/(2\zeta - 1/2)^3$. Independence and centering give $\mathbb{E}V_k^4 \le A_U^2 t + 3t^2$. Finally, $\mathbb{E}(V_k)_+^3 \le \sqrt{\mathbb{E}V_k^2\,\mathbb{E}V_k^4} \le t\sqrt{A_U^2 + 3t}$ by Cauchy–Schwarz. □

Here $\chi = p - H/\ell + u$. Thus an upper bound $b = p_+ - \min(H/\ell)$ yields precisely the phase function

$$\frac{(b+u-1)_+}{(\zeta-u)^2} + \frac{(b+u)_+}{[\zeta+\min(u,\zeta/3)]^2}.$$

We maximize this function in Section 7.3. The conditions $\chi < 2$ and $1 - c_v\Delta/N > 0$, and any normalization such as $\sqrt{n}\, m_B \le 1$, must be checked separately. The marked-crossing estimates needed to bound the remaining linear term $\pi_k \mathbb{E}R$ in (77) are given in Section B; they form a separate part of the local comparison.

7.3 An exact curvature calculation

The local comparison contains the supremum of the phase function of Section 7 over an interval. Despite the positive parts and the truncated denominator, it is attained at an endpoint or at one explicit interior point.

For real b, $1/2 < \zeta \le 1$ and $[l,h] \subseteq [-1/2, 1/2]$, define

$$f(u) = \frac{(b+u-1)_+}{(\zeta-u)^2} + \frac{(b+u)_+}{[\zeta+\min(u,\zeta/3)]^2}, \qquad u_* = \zeta - 2b.$$

All denominators are strictly positive. In the conditional-count comparison $\zeta = 1 - d_L$, where d_L is an upper bound for d_Y on the current noise band, and $b = p_+ - \min(H/\ell)$, where the minimum uses all four signed endpoint quotients.

Theorem 7.3 (Exact three-candidate supremum)**.** *The supremum of f over $[l,h]$ is the maximum of $f(l)$, $f(h)$, and the value $1/[4(\zeta-b)]$ if*

$$\zeta > b, \qquad l \le u_* \le h, \qquad u_* \le \zeta/3, \qquad u_* \le 1 - b.$$

An ineligible third value is omitted. The formula includes singleton intervals, coincident kinks and $\zeta = 1$.

Proof. Write $f = r + g$ for the two summands. On $b + u > 1$,

$$r'' = \frac{2[3(b-1)+u+2\zeta]}{(\zeta-u)^4} > 0,$$

since its bracket is $3(b+u-1) + 2(\zeta-u) > 0$. Where $u < \zeta/3$ and $b + u > 0$,

$$g' = \frac{\zeta - 2b - u}{(\zeta+u)^3}, \qquad g'' = \frac{2(3b+u-2\zeta)}{(\zeta+u)^4}.$$

If also $r > 0$, then $3b + u - 2\zeta > 3 - 2u - 2\zeta \ge 0$, so f is convex there. For $u > \zeta/3$, g is convex piecewise linear. At each of $-b$, $1-b$ and $\zeta/3$, the derivative has a nonnegative jump; simultaneous jumps add. Thus none is a strict interior maximum.

Where $r = 0$, either $g = 0$, or below $\zeta/3$ its derivative changes from positive to negative only at $u_* = \zeta - 2b$. Its positive numerator condition is $\zeta > b$, and $r = 0$ there is equivalent to $u_* \le 1 - b$. Its value is $1/[4(\zeta-b)]$. Above $\zeta/3$, g is nondecreasing. These cases locate a global maximum among the stated candidates. Continuity handles boundaries and collisions; constant segments contribute a value already attained at an endpoint or a listed maximizer. □

For any subinterval $[a,d] \subseteq [l,h]$ and $u \in [a,d]$, monotonicity of the numerators and positivity of the denominators give

$$f(u) \le \frac{(b+d-1)_+}{(\zeta-d)^2} + \frac{(b+d)_+}{[\zeta+\min(a,\zeta/3)]^2}.$$

The exact maximum is therefore no larger than the largest of these bounds over a subdivision of the interval. For fixed b and u, f is nonincreasing in ζ, so a bound computed with the smallest admissible ζ remains valid for larger ones.

7.3.1 The inverse-index correction

With the exact curvature bound the inverse-index correction may be negative. Within a fixed positive-noise cell, the bracket and the correction have the forms

$$B(\lambda) = c_{\mathrm{mark}} + \frac{q}{a\lambda + G^2} + s\lambda - t, \qquad I_{\mathrm{new}}(\lambda) = A_0 + \frac{A_1}{\lambda}, \quad q \geq 0,\ a > 0,\ A_1 \geq 0.$$

Only the curvature contribution to A_0 decreases. Put $I_+(\lambda) = \max(0, I_{\mathrm{new}}(\lambda))$. For $n \geq N$,

$$m_0 B(\lambda) + \frac{I_{\mathrm{new}}(\lambda)}{n} \leq m_0 B(\lambda) + \frac{I_+(\lambda)}{N}.$$

Both B and the right-hand expression are convex for $\lambda > 0$, including a possible interior zero-clamp kink, where $m_0 > 0$ is a lower bound for $\sqrt{n}\,\pi_k$. Here c_{mark} is constant within the cell. Endpoint checks thus cover the closed interval. If an earlier correction $I_{\mathrm{old}} \geq I_{\mathrm{new}}$ with $I_{\mathrm{old}} > 0$ satisfies $m_0 B + I_{\mathrm{old}}/N < -1/1000$ at an endpoint, then $0 \leq I_+ \leq I_{\mathrm{old}}$ and the same strict inequality holds with I_+. The condition $B < 0$ is still needed to use the lower bound for the binomial mass.

7.4 The harmonic step only needs span below one

For a real number x, write $\{x\} = x - \lfloor x \rfloor$.

Lemma 7.4 (Extended conditional-span domain)**.** *Let $Z_1, \ldots, Z_j$ be independent real variables, each supported in an interval of length at most b, where $0 < b < 1$. Put $W = \sum_i Z_i$ and $t = \sum_i \mathrm{Var}(Z_i)$. Suppose $K > 0$ satisfies*

$$1 - \cos(2\pi x) \leq Kx \quad (0 \leq x \leq 1).$$

For every $w \in \mathbb{R}$ and every $0 < r \leq 2\sin^2(\pi b)/b^2$,

$$\mathbb{E}\{w - W\} \geq K^{-1}(1 - e^{-rt}). \tag{79}$$

In particular $K = 8$ is valid without a finite scalar certificate. The sharper value $K = 23/5$ may be used with the scalar majorization in Appendix B.

Proof. Let Z_i' be an independent copy. Since $|Z_i - Z_i'| \leq b$ and sine is concave on $[0, \pi]$, the chord from 0 to πb gives

$$\sin^2(\pi(Z_i - Z_i')) \geq \frac{\sin^2(\pi b)}{b^2}(Z_i - Z_i')^2.$$

Consequently, writing $v_i = \mathrm{Var}(Z_i)$,

$$|\mathbb{E}e^{2\pi i Z_i}|^2 = 1 - 2\mathbb{E}\sin^2(\pi(Z_i - Z_i')) \leq 1 - \frac{4\sin^2(\pi b)}{b^2} v_i \leq e^{-2rv_i}.$$

The intermediate right side is nonnegative because it bounds the nonnegative left side. Taking square roots and multiplying, using independence, gives $|\mathbb{E}e^{2\pi i W}| \leq e^{-rt}$. Periodicity then gives

$$K\mathbb{E}\{w - W\} \geq 1 - \Re\mathbb{E}e^{2\pi i(w-W)} \geq 1 - |\mathbb{E}e^{2\pi i W}| \geq 1 - e^{-rt}.$$

No centering or continuous distribution assumption was used. To verify $K = 8$, the function $y - 1 + \cos y$ is nondecreasing for $y \geq 0$, since its derivative is $1 - \sin y \geq 0$. Thus $1 - \cos(2\pi x) \leq 2\pi x < 8x$ for $x > 0$; $\pi < 4$ also follows directly from $\pi = 4\int_0^1 (1 + y^2)^{-1}\, dy < 4$. Equality at $x = 0$ is harmless. □

The sine argument applies throughout $b < 1$; the stronger restriction $b < 1/2$ is unnecessary for this step. For example, $b = 3/4$ permits the exact rate $r = 16/9$. This extends only the harmonic component. It does not extend other denominator, binomial, support or noise gates. In the whole-count notation a sufficient acquired span is $b = A + d_0$; it must be a bound on each actual conditional residual, not on its variance.

For example, on a phase interval $[u_l, u_h]$, the individual first-mark denominators are positive when $1 - 2d_0 - u_h > 0$ and $1 - A + u_l > 0$, with $\zeta = 1 - d_0$. On the full phase interval $[-1/2, 1/2]$, these become $d_0 < 1/4$ and $A < 1/2$. They imply $A + d_0 < 3/4 < 1$, so the extended harmonic lemma adds no further span restriction there. The optional common-mark bound has additional gates. None of these inequalities alone establishes the final score, and a larger span gives a smaller harmonic rate that must be used in that score.

Proposition 7.5 (The boundary is a real obstruction)**.** *No strictly positive uniform rate in* (79) *is possible for all independent positive-variance summands of support span at most one. Even a uniformly positive rate as $b \uparrow 1$ is impossible.*

Proof. For $0 < \varepsilon < 1$, let Z equal $\varepsilon - 1$ with probability ε, and ε otherwise. It is centered, has positive variance $\varepsilon(1 - \varepsilon)$ and span one. For j independent copies, $W = j\varepsilon - M$ with integer-valued M, so $\{j\varepsilon - W\} = 0$ identically. The right side of (79) is positive for every $r > 0$ and finite $K > 0$.

For the boundary limit, the centered variable taking values $\pm b/2$ with equal probabilities has variance $b^2/4$ and characteristic modulus $|\cos(\pi b)|$. As $b \uparrow 1$, this modulus tends to one while the variance tends to $1/4$. A positive uniform exponential decay rate would contradict these limits. Directly for (79), choose $w = -b/2$: then $\mathbb{E}\{w - Z\} = (1 - b)/2 \to 0$, also contradicting a positive uniform rate with finite fixed K. In fact $1 - |\cos(\pi b)|$ vanishes quadratically in $1 - b$, consistently with the rate in Lemma 7.4. □

7.5 Why small defect alone cannot replace confinement

The obstruction can be realized by the actual conditional laws of a centered, variance-one iid marginal with a large central gap. It is not merely an optimizer of a moment relaxation.

For rational $\varepsilon, t > 0$, set

$$q = \frac{1 + \varepsilon - \varepsilon^2}{1 + t^2}, \quad p = 1 - q, \quad r = tq, \qquad 0 < \varepsilon < p < 1. \tag{80}$$

Let Y take the values $\varepsilon - 1, \varepsilon, 1$ with respective probabilities $q\varepsilon, q(1 - \varepsilon), p$, and put $X = (Y - p)/r$. Direct calculation gives

$$\mathbb{E}Y = p, \quad \mathrm{Var}(Y) = q\{p + \varepsilon(1 - \varepsilon)\} = r^2.$$

Therefore X is exactly centered and standardized. The first two atoms are negative, and the third is positive. In the actual sign split, $\ell = 1/r$, $U_1 = 0$, and

$$U_0 = \begin{cases} \varepsilon - 1, & \text{probability } \varepsilon, \\ \varepsilon, & \text{probability } 1 - \varepsilon. \end{cases}$$

Thus the actual conditional amplitude is $1 - \varepsilon$, its span is one, the normalized central-gap increment is $c_{\mathrm{gap}} = 1 - \varepsilon$, and $\lambda = nq\varepsilon(1 - \varepsilon)$. In particular the central gap is wide even when the harmonic estimate fails completely.

For a fixed interior count k, the actual conditional sum has the law

$$V_k = (n - k)\varepsilon - \mathrm{Bin}(n - k, \varepsilon).$$

Thus $\mathbb{E}\{w - V_k\} = 0$ when $w = (n-k)\varepsilon$, despite its positive variance. At $n = 42$, $k = 17$, sufficiently small ε makes $u = \eta = w/2$ a legal marked phase, with $|u| \le 1/2$ and $0 < \eta \le 1$. Here $w = u + \eta$ is precisely the phase used by the marked-count argument.

Corollary 7.6 (No moment-only amplitude implication)**.** *For every $\delta > 0$ there is an actual centered variance-one finite law supported in $[-96/25, 96/25]$, with positive conditional noise, $D < \delta$ and $\lambda < \delta$ at $n = 42$, whose negative conditional residual span is exactly one and whose fractional-part gain vanishes at a legal marked phase. The normalized central-gap increment can be arbitrarily close to one.*

Proof. At $\varepsilon = 0$ the positive-weight part of (80) is the standardized two-point law at $-t, 1/t$, with positive probability $p = t^2/(1+t^2)$. Its defect is

$$D_B = \frac{(6 + 2\sqrt{10})(p - p_*)^2}{\sqrt{p(1-p)}}, \qquad p_* = (4 - \sqrt{10})/2.$$

This follows by substituting ρ_B, γ_B and $J_B = t + 1/t$. Choose positive rational t arbitrarily near $\sqrt{p_*/(1-p_*)}$. Then D_B is arbitrarily small. For this fixed t, let positive rational ε tend to zero. All moments are continuous. The extra outer-gap weight tends to zero, so the definition of J in(19) gives $D \to D_B$ as well; $\lambda \to 0$ while the conditional span remains one. At the limiting value of t, the outer atom tends to $-(1 + 2t^2)/t$. Here $0.84 < t < 0.85$, so its magnitude is less than $2(0.85) + 1/0.84 < 3$. The support is therefore within $[-96/25, 96/25]$ in a neighborhood. The sign, phase and positive-noise conditions also hold for all sufficiently small ε. □

These laws are not asserted to satisfy a maximizing contact inequality, nonnegative fixed-target excess, or all the signed-acquisition outputs. They do not refute the sharp conjecture or the conditional local comparison. They refute the proposed implication from small moment resources and an actual central gap alone to a positive fractional-part gain or a small conditional amplitude. Contact-based confinement or a quantitatively proved replacement for harmonic nonresonance remains necessary for that proposed shortcut.

8 Certificates and the remaining global implication

The analytical results above have two kinds of conclusions. Some are identities or inequalities for every law in a stated class. Others acquire numerical bounds after finitely many rational gates have been verified. Neither kind supplies an omitted hypothesis of the other. We first state the remaining global obligation, then specify the computational boundary.

8.1 The fixed-target global implication

Assumption 8.1 (Independent finite base)**.** For every integer $1 \le j \le 41$, one has $C_j \le C_{\mathrm{E}}$, by a proof independent of the later-index exclusion.

Assumption 8.2 (Fixed-target local exclusion)**.** For every integer $n \ge 42$ satisfying $C_{n-1}, C_{n-2} \le C_{\mathrm{E}}$, there is no pair maximizing (4) with $c = C_{\mathrm{E}}$ and nonnegative excess.

The second assumption is the missing conclusion of the local program, not a result of this paper. Its predecessor hypotheses are consequences of a least-failure choice. The new contact entry and the central-panel confinement use $C_{n-1} \le C_{\mathrm{E}}$; the latter obtains its two-deletion concentration from antichains, without a bound on C_{n-2}. We leave both predecessor bounds in the final interface so that any later comparison using both must declare that use explicitly.

Theorem 8.3 (Conditional all-index bound)**.** *Under Assumptions 8.1 and 8.2,*

$$C_n < C_{\mathrm{E}} \quad (n \geq 42), \qquad C^* = C_{\mathrm{E}}.$$

Proof. If the strict tail bound fails, choose the least $n \geq 42$ with $C_n \geq C_{\mathrm{E}}$. Both predecessors are at most C_{E}; at $n = 42$ they are the base indices 41 and 40. The fixed-target attainment and equality statements of Section 2 give a maximizing pair with nonnegative excess. This contradicts Assumption 8.2. Together with the base, the strict tail gives $C^* \leq C_{\mathrm{E}}$. The single Bernoulli law in (2) gives the reverse inequality. □

For the sharp global constant alone it is enough to exclude positive excess under the same predecessor hypotheses. Indeed, choose the least $n \geq 42$ with $C_n > C_{\mathrm{E}}$ and repeat the argument. This yields $C_n \leq C_{\mathrm{E}}$, hence $C^* = C_{\mathrm{E}}$, without a claim of strictness at each finite index. Equality-inclusive exclusion additionally needs the zero-excess case, including the strict finite-index Bernoulli endpoint. Neither version requires an upper bound for an unknown maximizing ratio, a global positive ceiling, or an astronomically large terminal index.

8.2 What a finite certificate must establish

A certificate has three ingredients. Its domain description contains every actual parameter tuple allowed by the analytic hypotheses. Its local proof establishes the required inequality throughout each closed cell. Its coverage proof shows that the cells exhaust that domain.

The following elementary lemma is the basic tool for tensor certificates.

Lemma 8.4 (Common-marginal expectation and score)**.** *Let $K \subset \mathbb{R}$ contain the support of $P \in \mathcal{P}$. Let $f_0, \ldots, f_d$ be integrable functions under P, with $f_0 = 1$, and let $m_j = \mathbb{E} f_j(X)$. For a finite tensor sum*

$$T(x_1, \ldots, x_n) = \sum_{\alpha} t_\alpha \prod_{i=1}^{n} f_{\alpha_i}(x_i),$$

independence gives

$$\mathbb{E}T(X_1, \ldots, X_n) = \sum_{\alpha} t_\alpha \prod_{i=1}^{n} m_{\alpha_i}. \tag{81}$$

If $T \geq \mathbf{1}_{\{x_1 + \cdots + x_n \leq H\}}$ on all of K^n and

$$\sum_{\alpha} t_\alpha \prod_{i} m_{\alpha_i} - \Phi(H/\sqrt{n}) \leq a\rho/\sqrt{n},$$

then $\mathbb{P}(S_n \leq H) - \Phi(H/\sqrt{n}) \leq a\rho/\sqrt{n}$.

Proof. Each product is absolutely integrable, because independence factors its absolute expectation into the finite quantities $\mathbb{E}|f_j(X)|$. Fubini and linearity prove (81). Integrate the pointwise majorant and use $\rho > 0$ to obtain the score inequality.

□

For example, a tensor product containing a centered factor has zero expectation when all other factors concern different independent variables. Such terms may improve a pointwise majorant without altering its expectation. Polynomial factors of degree above three require the corresponding moments. For the extremal laws used here, these are finite because support is compact.

An exact polynomial positivity certificate often uses Bernstein coefficients. On $[0,1]$, set $B_{j,d}(t) = \binom{d}{j} t^j (1-t)^{d-j}$. They are nonnegative and sum to one. The identity

$$t^k = \sum_{j=k}^{d} \frac{\binom{j}{k}}{\binom{d}{k}} B_{j,d}(t)$$

follows from $\binom{d}{j}\binom{j}{k} = \binom{d}{k}\binom{d-k}{j-k}$ and the binomial theorem. Applying it in each coordinate proves that a polynomial $\sum_\alpha a_\alpha t^\alpha$ of coordinate degrees at most d_i has tensor Bernstein coefficients

$$b_\nu = \sum_{\alpha \le \nu} a_\alpha \prod_i \frac{\binom{\nu_i}{\alpha_i}}{\binom{d_i}{\alpha_i}}.$$

Thus nonnegative coefficients prove nonnegativity on the whole closed box. For $P(x) = \sum_{i=0}^{d} a_i x^i$ on $[l,h]$, the change $x = l + (h-l)t$ gives power coefficients $\widetilde{a}_j = (h-l)^j \sum_{i=j}^{d} a_i \binom{i}{j} l^{i-j}$. Applying the preceding identity gives the Bernstein coefficients on that interval. Rational affine changes and subdivision preserve exactness. A certificate must also verify the polynomial identity and the coverage of the domain.

The event plane needs particular care. On the closed side $\sum x_i \le H$ the required inequality is $T - 1 \ge 0$; on the other side it is $T \ge 0$. A certificate which verifies only $T \ge 0$ at a boundary point has not proved the closed-event majorant there. Sign orthants, event-plane intersections and endpoint atoms must be included in the pointwise proof.

Proposition 8.5 (Closed-cover soundness)**.** *Suppose a family of actual laws and thresholds has an analytically proved map into a covered parameter domain. A finite certificate has finitely many roots covering that domain and a finite subdivision tree at each root. Every split retains both closed children. At each leaf, either a proved necessary condition excludes every actual mapped point in the leaf, or a valid whole-leaf inequality bounds its positive score by a. Then all actual points in the root domain have positive score at most a. If every leaf instead excludes a putative attained score $C \ge C_{\mathrm{E}}$, no such attained witness belongs to the root domain.*

Proof. An actual point at a parent belongs to at least one child, including if it lies on the common boundary. Induction up each finite tree transfers the leaf conclusion to its root. The root union transfers it to the complete mapped domain. In the second version no actual point can lie in any leaf, a contradiction. □

A root may describe a relaxation containing parameters of no actual law; this does not affect soundness, whereas a missing root or an unfinished subdivision does. A minimum of convex bounds need not be convex, so an endpoint maximization must be justified for each bound before the minimum is taken.

In interval computations every transcendental function is enclosed outwards and every denominator is proved positive on the whole cell; a lower bound for a binomial mass multiplies a bracket only after the bracket is known to be negative.

A cover must include both signs of the phase, all admissible nearest counts, the endpoints of probability panels and the boundaries between noise bands. The case $\lambda = n\mathbb{E}U^2 = 0$ is separate: then $U = 0$ almost surely, the law is a two-point law, and Schulz's theorem applies [19].

8.3 What has been checked

The selected coarse entry is the separately named fixed-target component `INITIAL42_FIXED_TARGET_COARSE_V7`. Its accepted proof object reconstructs the two quadratic passes,

verifies every support-radius gate, and evaluates four signed passes with 11264 nominal boxes in total. It keeps all 100 closed probability panels, including all incidences of a law lying at a shared endpoint; 30 panels are nonempty after entry. The subsequent elementary moment calculation excludes two whole panels and retains 28. These are coverage statements about necessary conditions, not assertions that each remaining box is realized by an iid law.

The selected signed resource export contains all 682 active cells of the last pass and their 1364 endpoint quadratics. The low-noise lemma uses 28 rational endpoint and domain gates for one continuous polynomial argument. Its application is bound to the previously acquired .6 strip on $p \in [.40, .53]$. The complementary calculation on $[.41, .42]$ covers all 128 closed sign, threshold and noise boxes: 16 use the low-noise lemma, 84 are impossible by necessary conditions, and 28 pass both ray tests. The 56 retained strict margins exceed 0.00892011917. The seam $\lambda = .075$ belongs to both arguments; $\lambda = 0$ is handled directly.

The recorded independent checks reconstruct the signed coefficients and the continuous quintic without importing their constructors. They also check the actual-law moment contraction and the central-panel resource and ray inequalities. Negative controls include malformed, missing, duplicated, reordered and corrupted proof data, a wrong endpoint-atom convention, and omission of the H^2a conservation term. These controls test specified failure modes; they do not turn arithmetic into a proof of compactness, inversion, or an analytic reduction.

Result	Mathematical status	Execution status
Fixed-target attainment, contact and conservation	Proved at the stated scope, including reflection and support residuals	Finite-law identity and negative controls recorded
Selected Fourier and coarse entry	Analytic implication proved; numerical conclusion uses the stated exact cover	Isolated replay passed; 11264 signed boxes
Elementary moment contraction and signed phase	Same-law implications proved from the acquired entry	All panels and active cells checked
Continuous low-noise confinement	Proved under its incoming strip and predecessor hypotheses	28 exact gates; independent reconstruction
Full confinement on $[.41, .42]$	Conditional panel theorem proved	Complete 128-box join checked
Four-summand original-law bridge	Conditional on genuine whole-cell rows and complete coverage	Algebraic and atom controls; no complete four-summand acceptance
Finite base and complete local exclusion	Open	No complete acceptance

These arithmetic results were recorded during the research checks of 27–28 October 2026. The manuscript build and source-consistency checks are separate document checks. An execution marked successful is not an independent theorem about all real laws unless the mapping from such laws to the checked domain has also been justified.

8.4 Execution and the trusted boundary

The acceptance programs use exact rational arithmetic and directed rational enclosures. They compare the complete ordered proof data, reject duplicate keys and distinguish numerical types. Source hashes identify inputs; they do not establish the mathematical correctness of those inputs.

An execution audit found that disabling Python bytecode writes does not prevent reading an existing valid cache. A hostile cache could therefore alter a phase export while every pinned source hash remained unchanged; module search order supplied another route to the

wrong implementation. The selected execution path now compiles the explicitly named source bytes and runs in an isolated child with a fresh cache location. The unchanged coarse-entry ancestor is copied from its 13 pinned source and data files into a fresh private tree and checked before and after execution. The repaired path reproduces the genuine output and rejects the hostile fixtures. The Python interpreter, standard library, operating system, rational arithmetic semantics and the analytic-to-code correspondence remain part of the trusted boundary. This is not a Lean formalization.

The repository includes the proof objects, exact programs, input hashes, commands, return codes and separate review reports. Bounded reproduction uses `mathematics/fixed-target-v8/2026-10-28/run_checks.py`; the isolated coarse-entry replay is `mathematics/fixed-target-v8/2026-10-28/isolated_entry_replay.py`. The evidence directory is `runs/fixed-target-v8-20261028/`. No native large-scale certificate-generation campaign is part of these manuscript checks.

8.5 Which computations have actually been removed

For the selected coarse entry, the fixed coefficient and signed moment feedback replace the older unknown-score initialization. Its dependency chain no longer includes the degree-six phase ancestor, the six intervening coarse high-frequency updates, or the older twelve-stage signed schedule and its export evaluation. The comparison is 11264 signed boxes in the new component against 239616 prescribed in the old component, plus their respective separate initial calculations. It is not a measured speedup or a percentage reduction of the whole proof. The two quadratic passes, complete Fourier estimates, actual-law event bounds, and ordered support acquisition remain.

After that entry, the moment contraction is elementary and needs no new Fourier integration. The continuous low-noise proof replaces the additional thirteen-panel boundary table by one quintic and its derivative signs; it also removes the phase table from the lowest-noise part of the completed central panel. The incoming .6 strip is still essential. Larger noise, other probability panels, the strict count comparison and the independent finite base are not removed by these results. Existing certificate files and their original targets are preserved as historical alternatives.

8.6 What remains to complete the conjecture

A full local proof must still acquire the required confinement on the other surviving probability panels at larger noise, then prove the strict count comparison on every remaining count, phase and noise cell. In particular, several normalization and cubic-repair formulas in the appendices assume $p < 1/2$, while the acquired range reaches .53. Those assumptions cannot be omitted. Amplitude bounds below $1/2$ must also be quantitatively strong enough for each selected count denominator. The equality case and zero-noise comparison must have their stated scope.

Independently, the finite base through 41 remains open in this work. Appendix F keeps the original marginal in the four-summand response and proves the quadrature and threshold-interpolation reserves. It does not provide all the necessary numerical rows or a complete four-summand cover. A five-summand result which assumes the four-summand bound cannot supply that missing parent. No historical success marker or partial-checkout file list is used to infer otherwise.

9 Discussion

Fixing the target changes the order of the proof. The predecessor bound and the coefficient in the variational problem now agree from the start, and the conservation identity retains the error excess throughout the calculation. Signed moment feedback makes a shorter coarse acquisition possible. At small noise, a continuous contact argument replaces a further probability grid. These are proved reductions at their stated scopes.

The remaining difficulty is the convolution comparison. A small moment defect alone does not prevent conditional lattice resonance, as the explicit family in Section 7.4 shows. Contact-based confinement therefore has a role that cannot be replaced by a favorable moment box. Even after confinement, the error estimates must stay relative to λ as λ tends to zero. An absolute remainder of order $n^{-3/2}$ is not sufficient for that purpose.

The conditional global argument has no large terminal index: a least failure supplies its own sharp predecessors. To complete it, one still needs both an independent finite base and a local exclusion at every later index. The statements in this paper keep those two obligations separate from the intermediate results already proved.

Use of artificial intelligence

AI systems were used extensively in the research and in writing this paper. ChatGPT and Codex assisted with mathematical exploration, exact checking programs, source and bibliography comparisons, and drafting and revising the text. Other AI systems assisted with mathematical and editorial review. Some proposed arguments failed under checking; they were not used as premises. The authors are responsible for the content and presentation. Agreement between AI systems is not treated as mathematical verification.

The work builds in part on an earlier AI-assisted draft on Hexagon [18]. The present version gives the fixed-target argument and the analytical details of its stated reductions, with the remaining obligations listed in Section 8.

Code and data

The certificates, checking programs, review reports and manuscript sources are available in the authors' fork at `https://github.com/kolodziejek/berry-esseen-certificates`. The preserved reference baseline is commit `6346fc1b4432bc5982822aa696eb566ed9e1e1ee`. The new results have separate names and evidence; they do not overwrite an older certificate's target or claim its acceptance.

A A uniform Fourier enclosure with explicit input hypotheses

We prove the compensated Fourier formula and the complete low- and high-frequency estimates used in Section 4. Every assertion concerns an actual marginal with the displayed moment and event bounds. Attainment and acquisition of those bounds are not premises of the Fourier proof itself; their use is ordered in that section.

For each integer $n \geq N \geq 42$, let $X = X_n$ be a real random variable with $\mathbb{E}X = 0$, $\mathbb{E}X^2 = 1$. Neither the law nor the real raw threshold $H = H_n$ must be fixed as n varies. Write

$$f(t) = \mathbb{E}e^{itX}, \quad \gamma = \mathbb{E}X^3, \quad e_n(t) = e^{-nt^2/2}(1 - in\gamma t^3/6), \quad V_n = f^n - e_n.$$

The two complete integrals are

$$I_0(H) = \int_0^{1/2} \frac{|\Im(e^{-itH}V_n(t))|}{t^3}\,dt, \qquad I_\infty = \int_{1/2}^{\infty} \frac{|V_n(t)|}{t^3}\,dt.$$

A.1 The inversion edge, including atoms and moving thresholds

Lemma A.1. *If a finite signed measure μ has mass and first moment zero and $\int x^2\, d|\mu|(x) < \infty$, then, for every $H \in \mathbb{R}$,*

$$\int (x-H)|x-H|\, d\mu(x) = -\frac{4}{\pi}\int_0^\infty \frac{\Im(e^{-itH}\widehat{\mu}(t))}{t^3}\, dt.$$

The integral on the right is absolutely convergent.

Proof. For real y, substitution and integration by parts give

$$\int_0^\infty \frac{ty - \sin(ty)}{t^3}\, dt = \frac{\pi}{4} y|y|.$$

The boundary terms vanish: near zero the numerator is $O(t^3)$, and at infinity it is $O(t)$. The integrand has the sign of y, since $v - \sin v \geq 0$ for $v \geq 0$. To calculate the constant, nonnegative Tonelli and elementary Laplace integration give

$$\int_0^\infty \frac{1-\cos t}{t^2}\, dt = \int_0^\infty s\left(\frac{1}{s} - \frac{s}{s^2+1}\right) ds = \frac{\pi}{2}.$$

The absolute double integral of the compensated expression against $d|\mu|(x)$ equals

$$\frac{\pi}{4}\int (x-H)^2 d|\mu|(x) < \infty.$$

Absolute Fubini is therefore valid. The linear compensation disappears because $\int (x-H)d\mu = 0$, proving both assertions. No continuity of a CDF at H is involved. □

Apply this to the sum law minus the scale-$\sqrt{n}$ pushforward of the signed density

$$\phi(x)\{1 + \gamma(x^3 - 3x)/(6\sqrt{n})\}.$$

This signed density has mass one, first moment zero and finite absolute second moment. Its transform after scaling is e_n. Gaussian integration by parts, using $q_z(x) = (x-z)|x-z|$ and $q_z''' = 4\delta_z$, yields

$$\varepsilon_n := \mathbb{E}q_{H/\sqrt{n}}(S_n/\sqrt{n}) - \mathbb{E}q_{H/\sqrt{n}}(Z) - \frac{2\gamma\phi(H/\sqrt{n})}{3\sqrt{n}} = -\frac{4}{\pi n}\int_0^\infty \frac{\Im(e^{-itH}V_n(t))}{t^3}\, dt.$$

Consequently the error $E_n = -n^{3/2}\varepsilon_n/2$ satisfies

$$|E_n| \leq \frac{2\sqrt{n}}{\pi}\{I_0(H) + I_\infty\}. \tag{82}$$

This is an identity and bound for each n, H; it exchanges no limit in n with an integral or a varying law.

A.2 Sixth-order LOW with its actual coefficient bounds

Assume common bounds

$$\mu_4 := \mathbb{E}X^4 \leq M \leq 111/10, \quad |\gamma| \leq g \leq 37/20, \quad \mu_6 \leq M_6,$$

and put

$$a = (3-\mu_4)/24, \quad z = (10\gamma - \mathbb{E}X^5)/120, \quad |a| \leq A, \quad |z| \leq Z.$$

Here A, Z, M_6 are nonnegative common ceilings, not quantities inferred from bounded third moments. Define

$$E = M_6/720 + g^2/72 + (1/2 + g^2/144)^3/5, \qquad B = A + Z/2 + E/4,$$

$$a_0 = \frac{N-1}{2N}, \quad b_0 = \frac{45-M}{96}(1-2/N) > 0, \quad \lambda_N = \sqrt{\pi/2}\left(\frac{N}{N-1}\right)^{3/2}.$$

Proposition A.2. *For all the laws and thresholds above,*

$$I_0(H) \le \frac{L_0 + L_H|H|}{\sqrt{n}} + \frac{L_2}{n},$$

where, for any $c_0 \ge \lambda_N$ and $c_1 \ge \lambda_N/2$,

$$\begin{aligned} L_0 &= c_0 Z + c_1 A g + 251 g^3/17280, \\ L_H &= c_0 A + 251 g^2/4800, \\ L_2 &= E/(2a_0^2) + B^2/(2b_0^3). \end{aligned}$$

Proof. Let $h(t) = \exp(-t^2/2 - i\gamma t^3/6)$. Taylor's integral remainder for e^{itX}, followed by expectation, gives its expansion through degree five with error at most $\mu_6 t^6/720$. For $w = -t^2/2 - i\gamma t^3/6$ and $0 \le t \le 1/2$,

$$|w| \le t^2(1/2 + g^2/144) < 1/6, \quad |e^w - 1 - w - w^2/2| \le |w|^3 e^{|w|}/6 \le |w|^3/5.$$

The first bound follows from $\sqrt{1/4 + y} \le 1/2 + y$; for the last use $e^{1/6} \le (1 - 1/6)^{-1} = 6/5$. Comparing the two polynomials gives

$$f - h = -at^4 - izt^5 + r_6(t), \qquad |r_6(t)| \le Et^6, \qquad |f - h| \le Bt^4. \tag{83}$$

The degree-six term in $w^2/2$ costs $g^2 t^6/72$; it has not been discarded.

Four integrations of $1 - \cos u \ge 0$ show $\cos u \le 1 - u^2/2 + u^4/24$ for every real u. Independence gives $\mathbb{E}(X - X')^4 = 2\mu_4 + 6$, and hence

$$|f(t)|^2 \le 1 - t^2 + (M+3)t^4/12 \le 1 - (45-M)t^2/48.$$

Thus $|f|, |h| \le e^{-(45-M)t^2/96}$ throughout the low interval. Every point on their complex joining segment obeys the same bound. Taylor's integral formula for the polynomial $v \mapsto v^n$ proves

$$f^n - h^n = n(f-h)h^{n-1} + R_2, \quad |R_2| \le \tfrac{1}{2}n(n-1)B^2 t^8 e^{-(n-2)(45-M)t^2/96}.$$

The leading remainder in (83) and this power remainder cost at most L_2/n after division by t^3 and integration: use $n - 1 \ge 2a_0 n$, $(n-2)(45-M)/96 \ge b_0 n$, $n(n-1) \le n^2$, and $\int_0^\infty t^3 e^{-At^2} dt = 1/(2A^2)$, $\int_0^\infty t^5 e^{-At^2} dt = A^{-3}$.

For the remaining leading terms let $b = n - 1$ and $v = b\gamma t^3/6$. Their imaginary part after rotation by Ht is bounded by

$$ne^{-bt^2/2}\{(Z + A|H|)t^5 + bAgt^7/6\}.$$

The Gaussian moments of orders two and four, after dividing by t^3, give $n\sqrt{\pi/2}b^{-3/2}(Z + A|H| + Ag/2)$. This is at most $\lambda_N(Z + A|H| + Ag/2)/\sqrt{n}$. Finally put $u = n\gamma t^3/6$. Direct sine and cosine remainders give

$$|\Im[e^{-itH}(h^n - e_n)]| \le e^{-nt^2/2}(|u|^3/6 + |H|tu^2/2).$$

The Gaussian moments of orders six and four respectively give $\sqrt{\pi/2}g^3/(432\sqrt{n})$ times 5, and $\sqrt{\pi/2}g^2|H|/(24\sqrt{n})$. Using $\sqrt{\pi/2} < 251/200$ gives precisely the two displayed rational coefficients. Every extended integrand was nonnegative. The only index comparisons were the displayed rational inequalities; in particular the law and H can vary arbitrarily with n. □

In the selected enclosure one has actual support $|X| \le R$ with $1 \le R \le 6$ and $\mathbb{E}|X|^5 \le E_5$. Then $M_6 = RE_5$ and $A = \max(2, M-3)/24$ are valid. Its sharper Z is also licensed: for $Q = X^2 - \gamma X - 1$ direct multiplication gives

$$\mathbb{E}Q^2 = \mu_4 - 1 - \gamma^2, \quad \mathbb{E}X^5 = \gamma^3 + 2\gamma + \mathbb{E}[Q^2(X + 2\gamma)].$$

Reflect this auxiliary calculation if necessary so that $t = |\gamma| \ge 0$. For any $r \ge \max(R, 2g)$ the two signs of $\mathbb{E}X^5 - 10\gamma$ are at most

$$P_r(t) = r(M-1) + (10-2M)t - rt^2 + t^3, \quad Q_r(t) = r(M-1) + (2M-10)t - rt^2 - t^3.$$

The coefficients of μ_4 before replacing it by M are $r - 2t, r + 2t \ge 0$. If $M \le 5$, $P_r \ge Q_r$ and $P_r \le r(M-1)+(10-2M)t-(r-g)t^2$, whose maximum on $[0, g]$ is at the clipped vertex used in the code. If $M > 5$, $P_r \le r(M-1)$ and $Q_r \le r(M-1)+(2M-10)t-rt^2$, again with the stated clipped vertex. For $r = 6$, $P_6'' = -12+6t < 0$ on $[0, g]$. If $M \le 5$ and $P_6'(g) \ge 0$, its maximum is exactly $P_6(g)$; otherwise the preceding quadratic gives $6(M-1) + (10-2M)^2/[4(6-g)]$. For $M > 5$ the unrestrained vertex bound is $6(M-1) + (2M-10)^2/24$. Taking the minimum of these proved ceilings and $E_5 + 10g$, then dividing by 120, gives a valid coefficient ceiling Z. The radius is an actual support input, not a free tuning parameter.

A.3 All six HIGH pieces and their index gates

For independent copies of the same marginal suppose an actual event $\mathcal{E}$ satisfies

$$\mathbb{P}(\mathcal{E}) \ge \alpha > 0, \quad |X - X'| \ge d > 0 \text{ on } \mathcal{E}, \quad \mathbb{E}[(X - X')^2 \mid \mathcal{E}] \le M_E.$$

Fix $b = 9/10$, $0 < r \le 1/4$, and

$$h = 333/106 - (22/7)^3 r^2/6 > 0, \quad \kappa = \alpha(23/48)^2 d^2, \quad L(x) = \frac{x+2}{x^2+4x+2},$$

$$S_3 = \sum_{j=1}^{20} j^{-3} + 1/800, \quad S_5 = \sum_{j=1}^{20} (j-r)^{-5} + \frac{1}{4(20-r)^4}.$$

Let $c(u) = u/2 - (M+3)u^2/24$ and set

$$c_* = \min\{c(1/4), c(81/100)\}, \quad c_q = 1/8 - (M+3)/384, \quad s = [1 - (M+3)53/600]/2.$$

Proposition A.3. *Suppose $c_* > 0$. For every $n \ge N$, $\sqrt{n} I_\infty$ is at most the sum of the following six terms:*

$$T_1(n) = 2\sqrt{n} e^{-nc_*},$$

$$T_2(n) = \frac{M_E}{4\pi^{3/2} h \sqrt{\alpha}} \{S_3 + 3S_5/(n\alpha h^2)\},$$

$$T_3(n) = \frac{M_E \sqrt{n}}{2\pi^2} e^{-n\alpha h^2 r^2},$$

$$T_4(n) = \frac{e^{-n\kappa b^2}}{2b^4 \kappa \sqrt{n}},$$

$$T_5(n) = \frac{\sqrt{n}}{2b^2} e^{-n\alpha(23/48)^2},$$

$$T_6(n) = \sqrt{n} e^{-n/8} \{2L(n/8) + \sqrt{M-1}/3\}.$$

If $s > 0$, T_1 may be replaced by its minimum with $2\sqrt{n}e^{-nc_q}L(ns/4)$. The fourth term may be replaced by its minimum with $\sqrt{n}e^{-n\kappa b^2}L(n\kappa b^2)/(2b^2)$. The common-floor sum at N is valid if

$$2Nc' > 1 \quad \text{for } c' \in \{c_*, \alpha h^2 r^2, \alpha(23/48)^2, 1/8\}.$$

For the optional first term also require $2Nc_q > 1$; for the optional fourth term require $N\kappa b^2 \geq 1/4$. If an optional gate fails, the simpler term remains available.

Proof. The triangle inequality separates $|f^n|$ and the entire comparator. On $[1/2, b]$, symmetrization gives $|f|^n \leq e^{-nc(t^2)}$. The function c is concave, so $c(t^2) \geq c_*$ and the integral of t^{-3} is less than two. This gives T_1. Also

$$c(u) - c_q - s(u - 1/4) = (M + 3)(u - 1/4)(81/100 - u)/24 \geq 0.$$

When $s > 0$ this chord may be integrated to infinity.

For completeness, the tail calculation used here is, for $x = nab^2 > 0$,

$$\sqrt{n}\int_b^\infty e^{-nat^2}t^{-3}dt \leq \frac{\sqrt{n}e^{-x}}{2b^2}L(x). \tag{84}$$

Substitution and positive Fubini reduce the integral, without its prefactor, to $\int_0^\infty u(x + u)^{-1}e^{-u}du$. Put $P(u) = u^2 - 4u + 2$, $D = x^2 + 4x + 2 = P(-x)$. Exact polynomial division gives

$$\frac{1}{x+u} - \frac{-u^3 + (x+8)u^2 - (x^2 + 8x + 20)u + x^3 + 8x^2 + 20x + 16}{D^2} = \frac{P(u)^2}{(x+u)D^2} \geq 0.$$

Integrate using the factorial moments of e^{-u}; the numerator's integral is $(x+3)D$. Subtracting the resulting lower bound times x from one gives $L(x)$. Applying (84) at endpoint $1/2$ and rate s proves the optional first term.

For $t \geq b$, symmetrization, the actual-event probability bound, and convex Jensen give

$$|f(t)|^n \leq e^{-n\mathbb{E}\sin^2(t(X-X')/2)} \leq \mathbb{E}[e^{-n\alpha\sin^2(tD/2)} \mid \mathcal{E}], \quad D = |X - X'|.$$

Positive Tonelli allows a separate partition for every $D > 0$. With $\ell = 2\pi/D$, use the zero cell $[0, \ell/2]$ and nonzero cells $\ell[j - 1/2, j + 1/2]$, $j \geq 1$. In a nonzero cell write $t = \ell(j + v)$. On its core $|v| \leq r$, $|\sin \pi v| \geq h|v|$; on its wings $r \leq |v| \leq 1/2$ it is at least hr. These follow from $\sin x \geq x - x^3/6$ and monotonicity of sin on $[0, \pi/2]$, with directed bounds on π. For $|v| \leq r$ Taylor's integral remainder gives

$$\{(j - v)^{-3} + (j + v)^{-3}\}/2 \leq j^{-3} + 6(j - r)^{-5}v^2.$$

Extend the nonnegative Gaussian majorant over the whole real line. Its zeroth and second moments, the Jacobian $t^{-3}dt = D^2(4\pi^2)^{-1}(j + v)^{-3}dv$, and $\mathbb{E}[D^2 \mid \mathcal{E}] \leq M_E$ give T_2. The two series tails are bounded by their decreasing integrals after 20. The total wing weight is at most $\int_{\ell/2}^\infty t^{-3}dt = D^2/(2\pi^2)$, giving T_3. Extending these sets below b only increases their nonnegative bounds.

In the zero cell put $u = tD/2 \leq \pi/2$. Where $u \leq 1/2$, $\sin u \geq 23u/24$, and $D \geq d$ yields rate κ. Extending the set to $[b, \infty)$ gives T_4 from $t^{-3} \leq t/b^4$; (84) gives its optional improvement. Where $u \geq 1/2$, monotonicity gives $\sin u \geq 23/48$ and the total weight is at most $1/(2b^2)$, proving T_5. Endpoints have zero Lebesgue measure even if the conditional distance distribution has atoms.

Finally $|e_n(t)| \leq e^{-nt^2/2}(1 + n|\gamma|t^3/6)$. The first part on the entire $[1/2, \infty)$ uses (84); the second uses $\int_{1/2}^\infty e^{-nt^2/2}dt \leq 2e^{-n/8}/n$. Pearson's identity $\mu_4 - 1 - \gamma^2 = \mathbb{E}(X^2 - \gamma X - 1)^2 \geq 0$ gives T_6. No comparator tail is dropped.

For floor evaluation, $\sqrt{n}e^{-nc'}$ decreases for $2nc' > 1$, whereas n^{-1} and $n^{-1/2}e^{-nc'}$ decrease for every $c' > 0$. Also $L'(x) = -(x^2 + 4x + 6)/(x^2 + 4x + 2)^2 < 0$. Thus the optional first term decreases under its own gate, and the comparator decreases for $n \geq 5$. The sign of the derivative of $\sqrt{x}e^{-x}L(x)$ is the sign of

$$4 - 10x - 22x^2 - 13x^3 - 2x^4,$$

which is negative at $1/4$ and strictly decreasing on $(0, \infty)$. This proves the optional zero-cell gate. Every comparison is of a common scalar majorant, not of characteristic functions for different laws. □

A.4 The precise closed Fourier conclusion

Under the preceding moment, actual-event and scalar gates, let K_H be any complete HIGH sum evaluated at N, including its permitted optional improvements. Then

$$|E_n| \leq \frac{2}{\pi}\{L_0 + K_H + L_2/\sqrt{N} + L_H|H|\} \leq \frac{16}{25}\{L_0 + K_H + L_2/\sqrt{N} + L_H|H|\}. \tag{85}$$

The second inequality uses $\pi > 25/8$ and nonnegative coefficients. This is the constant Fourier budget and raw-threshold slope used in Section 4. Complete HIGH bounds for several valid windows may be minimized. The LOW coefficients must remain one complete valid triple.

For rational acceptance, positive denominator roots are rounded downward, numerator roots upward, and $\pi > 333/106$, $\sqrt{\pi} > 4431/2500$, $\pi < 22/7$ are directed correctly. These constants require no numerical transcendental oracle: polynomial division gives

$$22/7 - \pi = \int_0^1 \frac{x^4(1-x)^4}{1+x^2}\,dx > 0,$$

and the tangent addition formula with its first-quadrant branch gives

$$\pi = 16\arctan(1/5) - 4\arctan(1/239).$$

Four alternating terms for the first arctangent and the upper bound $\arctan(1/239) < 1/239$ give $\pi > 333/106 > 25/8$ by rational comparison. Squaring then verifies the two Gaussian constant enclosures used above. Its exponential helper replaces $x \geq 0$ by a downward rational $y \leq \min(x, 64)$ and returns an upward rounding of $1/\sum_{j=0}^{64} y^j/j!$. This is at least e^{-x}. The floor argument applies to the underlying analytic bounds, not to a claim that this rounded helper itself preserves strict monotonicity.

The estimates are uniform over all laws and thresholds satisfying the stated inequalities at the actual index. They require no monotonicity of extremal Berry–Esseen constants, no fixed-law asymptotic limit, and no unproved disappearance of a lattice tail.

B Conditional-count estimates

This appendix supplies the analytic estimates needed after confinement. Its statements are conditional on their displayed support, probability, count and sign hypotheses. In particular, the formulas requiring $p < 1/2$ do not cover the entire acquired range $[.25, .53]$. Neither reflection nor the half-amplitude result alone fills that gap. No complete numerical comparison cover is asserted here.

B.1 The comparison and its domain

A law concentrated near two points has a natural integer count and a small centered error. We compare it with its own Bernoulli count. The same exact identity handles every noise regime. There are two useful estimates of its right side: discard all but the nearest outgoing count, or retain all counts through a random cutoff. The estimates share their normalization, moments, and tail bounds; only the final scalar witness changes.

The identity and sufficient inequalities

Let $0 < p \le 1/2$, $q = 1 - p$, $v = pq$, and let

$$Y = B + U, \qquad \mathbb{P}(B = 1) = p, \qquad \mathbb{E}(U \mid B) = 0.$$

The two actual conditional laws U_0, U_1 obey

$$|U_j| \le A, \quad U_0 \le a_0, \quad U_1 \ge -a_1, \quad a_0, a_1 \ge 0, \quad d = a_0 + a_1 \le d_0 < \min(p, q), \quad d_0 \le A. \tag{86}$$

Write $s_j = \mathbb{E}U_j^2$, $s = qs_0 + ps_1$, $\lambda = ns$, $\ell = (v + s)^{-1/2}$, $\theta = s/(v + s)$ and $\rho = \mathbb{E}|\ell(Y - p)|^3$. At the actual threshold choose

$$x = np + H/\ell = k + u, \qquad k \in \{1, \dots, n - 1\}, \quad |u| \le \tfrac{1}{2}, \qquad \delta = k - np = H/\ell - u.$$

Either nearest integer is allowed at a tie. Put

$$\pi_j = \binom{n}{j} p^j q^{n-j}, \quad F_B(j) = \mathbb{P}(\mathrm{Bin}(n, p) \le j), \qquad \rho_B = (1 - 2v)/\sqrt{v},$$

with $\pi_j = 0$ outside $[0, n]$. Let W_j be an independent sum of j copies of U_1 and $n - j$ copies of U_0. The signed scores are

$$C = \frac{\sqrt{n}}{\rho}\{\mathbb{P}(\textstyle\sum Y_i \le x) - \Phi(H/\sqrt{n})\}, \qquad C_B = \tfrac{\sqrt{n}}{\rho_B}\{F_B(k) - \Phi(\delta/\sqrt{nv})\}.$$

The fixed target is $c = C_{\mathrm{E}}$, with $c \ge c_* > 0$; one may take $c_* = 0.4$. The actual ratio C is used only to name the excess $a = \rho(C - c)$. It is not the coefficient in the moment correction. Define

$$\begin{aligned} f(u) &= \frac{\Phi((\delta + u)/\sqrt{n(v + s)}) - \Phi(\delta/\sqrt{n(v + s)})}{\pi_k}, \\ Q &= \mathbb{P}(\textstyle\sum Y_i \le k + u) - F_B(k), \\ \mathcal{R} &= \sqrt{n}\{\Phi(\delta/\sqrt{nv}) - \Phi(\delta/\sqrt{n(v + s)})\} + c(\rho_B - \rho). \end{aligned} \tag{87}$$

Equation (76), with $V_j = W_j$ and the same interior count k, gives the exact identity

$$a + \rho_B(c - C_B) = \sqrt{n}\pi_k\{Q/\pi_k - f(u)\} + \mathcal{R}. \tag{88}$$

In particular, the phase u, the sign of H, and both conditional laws refer to the same marginal throughout.

Theorem B.1 (Parameterized local comparison)**.** *Fix $n \ge N$, $\lambda > 0$ and a closed parameter cell on which* (86) *holds. Suppose $\sqrt{n}\pi_k \ge m > 0$. Either of the following witnesses proves $a + \rho_B(c - C_B) < -\varepsilon\lambda$.*

1. Nearest-count witness. *Valid bounds O, T, R_0, κ satisfy*

$$\mathbb{P}(W_k > u) + f(u) \geq O, \quad \sum_{i\geq 1} \mathbb{P}(W_{k+i} \leq u - i) \leq T, \quad \pi_{k+i}/\pi_k \leq R_0, \quad \mathcal{R} \leq \kappa\lambda/n.$$

Only legal counts occur in the sum; adding positive upper bounds for other counts is allowed. If $D = O - R_0 T > 0$ and

$$mD - \kappa\lambda/n > \varepsilon\lambda, \tag{89}$$

the conclusion follows.

2. Whole-count witness. *Bounds $B(\lambda) < 0$ and $I(\lambda) \geq 0$ satisfy*

$$a + \rho_B(c - C_B) \leq \sqrt{n}\pi_k \lambda B(\lambda) + \lambda I(\lambda)/n.$$

It is enough that

$$mB(\lambda) + I(\lambda)/N < -\varepsilon. \tag{90}$$

The witnesses may differ from cell to cell. A finite closed cover and a separate treatment of $\lambda = 0$ therefore suffice for a complete local comparison.

Proof. The signed count identity (74), with $V_j = W_j$, is

$$Q = -\sum_{i=0}^{k} \pi_{k-i} \mathbb{P}(W_{k-i} > u + i) + \sum_{i=1}^{n-k} \pi_{k+i} \mathbb{P}(W_{k+i} \leq u - i). \tag{91}$$

Apply Proposition 7.1 with $m_0 = m$. The legal counts, closed incoming events and strict outgoing events are identical. In the second alternative the present hypothesis $I \geq 0$ makes its $\max(0, I)/N$ exactly I/N, yielding (90). □

Binomial normalization and the variance correction

Here is an explicit finite recipe for the constants required above. Let $p \in [p_-, p_+] \subset (0, 1/2)$, $n \geq N$, $\lambda \leq \Lambda$, $\theta \leq \Theta$, and $|\delta| \leq \Delta$, where $\Delta \geq 1$ also includes every binomial mode. Choose $0 < r_- < p_- - \Delta/N$ and $p_+ + \Delta/N < r_+ < 1/2$. Let $v_* > 0$ be a lower bound for $p(1-p)$. Choose positive E, L, R bounding, on $[r_-, r_+]$,

$$|D'''(r\|p)| = \frac{1-2r}{r^2(1-r)^2}, \qquad |(\log(r(1-r)))'| = \frac{1-2r}{r(1-r)}, \qquad \frac{1}{12r(1-r)},$$

respectively, where $D(r\|p) = r\log(r/p) + (1-r)\log((1-r)/q)$. Then, with $Z(t) = Lt/2 + R + Et^3/(6N)$,

$$\sqrt{n}\pi_k = \frac{\phi(0)}{\sqrt{v}} e^{-\delta^2/(2nv)+\zeta}, \qquad n|\zeta| < Z(|\delta|). \tag{92}$$

For completeness, the finite Stirling formula used here is

$$\log(j!) = (j + \tfrac{1}{2})\log j - j + \tfrac{1}{2}\log(2\pi) + r_j, \qquad 0 < r_j < 1/(12j).$$

If $a_j = \log(j!) - (j + 1/2)\log j + j$ and $z = (2j+1)^{-1}$, the integrated geometric series gives $0 < a_j - a_{j+1} = \sum_{i\geq 1} z^{2i}/(2i+1) < 1/[12j(j+1)]$. Summing gives the remainder relative to $\lim a_j$. The integration-by-parts recursion for $\int_0^{\pi/2} \sin^j t\, dt$ and the inequalities between three successive integrals give Wallis's product; its finite product equals $e^{4a_j - 2a_{2j}} j/[2(2j+1)]$ and tends to $\pi/2$. Thus $\lim a_j = \frac{1}{2}\log(2\pi)$. Applying this formula to $n!, k!, (n-k)!$ and

Taylor-expanding D and the logarithmic prefactor proves (92); the factorial error is at most $1/[12nr(1-r)]$.

Consequently a valid mass lower bound is any positive

$$m < \frac{0.39886}{\sqrt{p_+(1-p_+)}}\left[1 - \frac{\Delta^2/(2v_*) + Z(\Delta)}{N}\right]. \tag{93}$$

The same formula at a mode proves $\sqrt{n}\max_j \pi_j < 1$ whenever $v_*[1 - Z(\Delta)/N]^2 > 0.4^2$ and $Z(\Delta) < N$. The rational bounds $0.39886 < \phi(0) < 0.4$ follow, for example, by the alternating arctangent series in Machin's identity.

Differentiating (87) gives

$$\log f' = -\tfrac{1}{2}\log(1 + s/v) + \delta^2/(2nv) - (\delta + u)^2/(2n(v + s)) - \zeta. \tag{94}$$

Thus $n|\log f'| \le B_0\lambda + P(D, a)$ on $|\delta| \le D \le \Delta$, $|u| \le a \le 1/2$, where

$$B_0 > \frac{1}{2v_*} + \frac{\Delta^2}{2Nv_*^2}, \qquad P(D, a) = \frac{2Da + a^2}{2v_*} + Z(D).$$

Choose $0 < b < 1$ above $[B_0\Lambda + P(\Delta, 1/2)]/N$ and put $F = (1 - b)^{-1}$. The elementary estimates $e^z \le (1 - z)^{-1}$ and $1 - e^{-z} \le z$ imply

$$|f' - 1| \le (B_1\lambda + P_1)/n, \qquad B_1 = FB_0, \quad P_1 = FP(D, a). \tag{95}$$

This supplies both a derivative interval $l_f \le f' \le h_f$ and the finer phase-error bound $|u - f(u)| \le a(B_1\lambda + P_1)/n$.

The exact count variance is

$$t_j = \mathbb{E}W_j^2 = \lambda + (j - np)(s_1 - s_0), \qquad |s_1 - s_0| \le s/p \le c_v s, \quad c_v \ge 1/p_-. \tag{96}$$

It gives $\omega_-\lambda \le t_k \le \omega_+\lambda$ with $\omega_\pm = 1 \pm c_v D/N$. All uses below require the checked condition $\omega_- > 0$. Also $t_{k+i} \le iV_*\lambda$ for $V_* = 1 + c_v(\Delta + 1)/N$. The binomial recursion gives, for every legal $i \ge 1$,

$$\frac{\pi_{k+i}}{\pi_k} \le R_0, \qquad R_0 > 1 + \frac{p_+ + \Delta - 1}{Nv_*[1 - (\Delta - 1)/(Np_-)]}, \tag{97}$$

provided the denominator is positive and $p_+ + \Delta < 2$. Indeed the first ratio has this bound and all subsequent rising factors are at most one. No count outside $[0, n]$ is used in this recursion.

Finally (86) fixes the signs inside the absolute cubic. Expansion and centering give

$$\mathbb{E}|Y - p|^3 = v(1 - 2v) + 3pq(s_0 + s_1) + p\mathbb{E}U_1^3 - q\mathbb{E}U_0^3 \ge v(1 - 2v) + (3p - d)s.$$

The inequality $(1 + z)^{3/2} \le 1 + 3z/2 + 3z^2/8$ for $z \ge 0$ yields

$$(v + s)^{3/2}(\rho - \rho_B) \ge s\left(6p - 3p^2 - \tfrac{3}{2} - d - \frac{3(1 - 2v)s}{8v}\right). \tag{98}$$

Let $Q_* > 0$ be a lower bound for the bracket and $m_c \le c_*/[p_+(1 - p_+) + \Lambda/N]^{3/2}$. The normal variance correction is nonpositive for $\delta \le 0$; for $\delta > 0$ its derivative in s is at most $0.4\delta/(2v^{3/2})$. Therefore

$$\mathcal{R} \le (P_*D_+ - m_cQ_*)\lambda/n, \qquad P_* \ge \frac{0.4}{2v_*^{3/2}}, \quad D_+ \ge \delta_+. \tag{99}$$

One may take $D_+ = D$, but retaining the positive part is often decisive. All bounds concern $v + s$, which can exceed $1/4$.

Two elementary tools for probabilities

Lemma B.2 (A shifted quartic)**.** *Suppose W is a sum of independent centered variables bounded in absolute value by A, and $t = \mathbb{E}W^2$. Let $K_0 \geq (9+6\sqrt{3})/4$ and $a > 0$. For $r \geq 0$ and $\sigma \in \{-1, 1\}$,*

$$\mathbb{P}(W + \sigma r > 0) \geq \frac{(3a^2 - A^2)t - 3t^2 - 4Art - 6r^2t - r^4 + 3a^2r^2 + 2\sigma a^3 r}{K_0 a^4}. \tag{100}$$

Proof. The polynomial $h_a(z) = (-z^4+3a^2z^2+2a^3z)/(K_0a^4)$ is below $\mathbf{1}_{\{z>0\}}$: it is nonpositive for $z \leq 0$, and its maximum for $z > 0$ is at $z/a = (1+\sqrt{3})/2$ and is at most one. Independence gives $|\mathbb{E}W^3| \leq At$ and $\mathbb{E}W^4 \leq A^2t+3t^2$. Expand $\mathbb{E}h_a(W+\sigma r)$ and apply these two inequalities. The value at the threshold is zero, so atoms require no limiting argument. □

Lemma B.3 (One common tail bound)**.** *If independent centered $X_i \leq A$ have total variance at most V, then, for $b > 0$, their sum has upper tail bounded by the minimum of one and*

$$\frac{V}{V+b^2}, \qquad \mathcal{B}(A,V,b) = e^{b/A}\left(\frac{V}{V+Ab}\right)^{b/A+V/A^2}, \qquad \frac{\mathbb{E}(P_\mu - j)_+^3}{(b/A+\mu-j)^3}, \tag{101}$$
$$\mu = V/A^2, \qquad 0 \leq j < b/A + \mu.$$

where P_μ is Poisson. The same holds for a lower tail when $|X_i| \leq A$. All bounds apply to closed events, hence also to strict events.

Proof. Cantelli's bound follows by applying Markov to $(\sum X_i + V/b)^2$. Taylor's integral remainder gives $\mathbb{E}e^{zX_i} \leq \exp\{\mathbb{E}X_i^2(e^{zA} - 1 - zA)/A^2\}$ for $z \geq 0$. Multiply and use $z = A^{-1}\log(1 + Ab/V)$ to get $\mathcal{B}$. The zero-variance case is immediate.

For the third bound, smooth $f(x) = (x-t)_+^3$ by a nonnegative smooth compactly supported density. Then $f'', f''', f'''' \geq 0$. For $\mathcal{A}f(x) = f(x+A) - f(x) - Af'(x)$, the Taylor quotient

$$\frac{f(x+y) - f(x) - yf'(x)}{y^2} = \int_0^1 (1-r)f''(x+ry)\,dr$$

is nondecreasing in y. A centered $X \leq A$ of variance v consequently satisfies $\mathbb{E}f(x+X) \leq f(x) + (v/A^2)\mathcal{A}f(x)$. For $\mathcal{P}_z f(x) = \mathbb{E}f(x + A(P_z - z))$, $\partial_z \mathcal{P}_z f = \mathcal{P}_z \mathcal{A} f$ and

$$\partial_z^2 \mathcal{P}_z f = \mathcal{P}_z \mathcal{A}^2 f \geq 0, \qquad \mathcal{A}^2 f(x) = \int_0^A \int_0^A (A-r)(A-w) f''''(x+r+w)\,dr\,dw.$$

Thus $\mathcal{P}_{v/A^2} f \geq f + (v/A^2)\mathcal{A}f$. Replace the independent variables successively; translations and expectations preserve the derivative signs and Poisson rates add. Increasing the total variance to V is permitted by Jensen after adding independent centered Poisson noise. Removing the smoothing is justified by bounded actual upper supports and finite Poisson third moments. Markov at $t = A(j - \mu)$ proves the displayed bound. □

These bounds have finite rational evaluations. For example,

$$\mathbb{E}(P_\mu - j)_+^3 = \mu^3 + 3(1-j)\mu^2 + (1 - 3j + 3j^2)\mu - j^3 \tag{102}$$
$$+ e^{-\mu} \sum_{h=0}^{\lfloor j \rfloor} \frac{\mu^h (j-h)^3}{h!}.$$

Replacing $e^{-\mu}$ by $(\sum_{h=0}^{24} \mu^h/h!)^{-1}$ enlarges the nonnegative correction. Alternatively sum the positive Poisson series through 32; the remaining tail is at most its first term divided by $1 - \mu(34-j)^3/[34(33-j)^3]$, provided $j < 33$ and that ratio is below one. For Bennett,

put $x = Ab/V$ and $L = [(b/A + \mu)\log(1 + x) - b/A]$. The positive series $\log(1 + x) = 2\sum_{h\geq 0} z^{2h+1}/(2h+1)$, $z = x/(2+x)$, gives lower bounds for L; any positive lower exponential series gives an upper bound for e^{-L}. Rounding a positive lower L down before exponentiation is also valid. These are choices of rational witnesses, not optimization claims.

The nearest-count witness, including vanishing noise

For $u = -r \leq 0$, add $-h_f r$ to (100) with $\sigma = 1$; for $u = r \geq 0$, add $l_f r$ with $\sigma = -1$. Both bound $\mathbb{P}(W_k > u) + f(u)$ from below. For a cell $t \in [t_-, t_+]$, $r \in [r_-, r_+]$, this expression is concave in t (second derivative $-6/(K_0 a^4)$), so only t_-, t_+ are needed. At each endpoint, bound a monomial with positive coefficient at r_- and one with negative coefficient at r_+. Any rational $a > 0$ is a valid witness. Other valid choices are

$$\frac{r_-^2}{t_+ + r_-^2} - h_f r_+ \quad (u \leq 0), \qquad l_f r_- \quad (u \geq 0). \tag{103}$$

They follow from Cantelli and nonnegativity, respectively. Their maximum or one selected witness is valid without an optimality proof.

For incoming counts use $t_{k+i} \leq iV_*\lambda$. On $u \in [-r_+, -r_-]$ the first distance is $1 + r_-$ and every later distance is at least i. On $u \in [r_-, r_+]$ they are $1 - r_+$ and $i(1 - r_+/2)$ for $i \geq 2$. Let b, b_o be these first and later distances. On $\lambda \in [L, U]$, let T_1 be any bound from Lemma B.3 at (A, V_*U, b). Since $\mathcal{B}(A, iV, ib) = \mathcal{B}(A, V, b)^i$, all later counts total at most $B^2/(1 - B)$ for $B = \mathcal{B}(A, V_*U, b_o) < 1$. Thus

$$D = O - R_0\{T_1 + B^2/(1 - B)\} > 0, \qquad mD - \kappa U/N > \varepsilon U \tag{104}$$

proves the entire cell when $\kappa \geq 0$. This sign condition is required on each cell. The argument applies to both central and left probability panels.

A cell touching $\lambda = 0$ requires estimates proportional to λ; a positive lower endpoint may not be substituted. The following rescaling provides them. Put $r = zt$, $t_* = \omega_+ U$ and $d_a = K_0 a^4$. The shifted quartic plus br, divided by t, becomes

$$F_{\sigma,b}(z, t) = \frac{3a^2 - A^2 - 3t}{d_a} + \left(\frac{2\sigma}{K_0 a} + b - \frac{4At}{d_a}\right) z + \frac{3a^2 t - 6t^2}{d_a} z^2 - \frac{t^3 z^4}{d_a}. \tag{105}$$

It extends to $t = 0$ and is concave in t. On $0 \leq z \leq z_0$, replacing z^4 by $z_0^2 z^2$ gives a lower quadratic at $t = t_*$; at $t = 0$ it is linear. Endpoints and a clipped quadratic vertex therefore give an exact lower bound $J(\sigma, b, z_0)$ for the whole rectangle.

For negative phase choose $z_-, c_- > 0$ and verify

$$J(1, -h_f, z_-) > c_-, \quad \frac{z_-^2}{1 + z_-^2 t_*} - h_f z_- > c_-, \quad \frac{2z_-}{(1 + z_-^2 t_*)^2} > h_f, \tag{106}$$
$$\frac{1}{1 + 4t_*} - \frac{h_f}{2} > c_- t_*, \qquad z_- t_* < 1/2.$$

The quartic handles $r \leq z_- t$. Beyond that point use $r^2/(t + r^2) - h_f r$. Its derivative at $r = z_- t$ is positive by the third gate; its second derivative has just one sign change, so any later critical point is a maximum. Its minimum is at $z_- t$ or $1/2$, covered by the other gates. Hence the own count exceeds $c_- t$. For positive phase choose $z_+, c_+ > 0$ with

$$J(-1, l_f, z_+) > c_+, \qquad l_f z_+ > c_+, \qquad z_+ t_* < 1/2. \tag{107}$$

The quartic handles $r \leq z_+ t$ and the normal term handles the rest. The own count exceeds $c_+ t$ and also exceeds $l_f r$.

For incoming tails proportional to λ, fix a Poisson anchor j. A positive series term of index h divided by $\lambda(b/A+\mu-j)^3$ has logarithmic derivative, multiplied by λ,

$$h-1-\mu-\frac{3\mu}{b/A+\mu-j},\qquad \mu=V_*\lambda/A^2.$$

If $b/A-j>0$ and this is nonnegative at $\lambda=U$ for the first positive index, it is nonnegative for every term and all $0<\lambda\le U$. Therefore the Poisson bound divided by λ is largest at U. Likewise $B^2/[\lambda(1-B)]$ is increasing whenever $\lambda B'/B\ge 1/2$. Differentiating the Bennett formula gives

$$\lambda B'/B = b/A-\mu\log(1+b/(A\mu));$$

The function $g(\mu)=b/A-\mu\log(1+b/(A\mu))$ decreases because $\log(1+x)>x/(1+x)$ for $x>0$. Moreover the logarithmic derivative of $B^2/[\lambda(1-B)]$, multiplied by λ, is $2g-1+gB/(1-B)\ge 0$ when $g\ge 1/2$. Thus checking the gate at U proves the whole interval. A rational upper bound for the logarithm verifies that gate. On cells using the fixed positive bound $l_f r_-$ instead, unscaled endpoint tails suffice: after a positive bracket is checked, divide it by U to obtain a lower bound per λ. These arguments, followed by (89), prove the nearest-count comparison uniformly down to arbitrary positive noise, including a deterministic conditional component.

The whole-count witness

Assume $\zeta=1-d_0>1/2$. Couple all $2n$ conditional variables independently and set $W_j=\sum_{i\le j}U_{1,i}+\sum_{i>j}U_{0,i}$, extending it constantly outside $[0,n]$. Every increment of $j+W_j$ is at least ζ. Define

$$K=\max\{j\in\mathbb{Z}: j+W_j\le k+u\},\qquad R=K-k.$$

The coupling of Section 7, with $V_j=W_j$, gives

$$\mathbb{P}(\textstyle\sum Y_i\le k+u)=\mathbb{E}F_B(k+R).\tag{108}$$

Let $\chi=(n+1)p-k<2$ and $m_B=\max_j\pi_j$. For every integer r,

$$F_B(k+r)-F_B(k)-r\pi_k\le\frac{m_B}{(n+1)v}\left[(\chi-1)_+r_+ +\chi_+\binom{r_-}{2}+\binom{r_-}{3}\right].\tag{109}$$

This is Lemma 7.2, with the same zero extension, $\chi<2$ and interior count. Its proof requires only an increment lower bound $\zeta>1/2$. Here $d\le d_0$ gives increments at least $1-d_0=\zeta$, and $\zeta\le 1$ follows from $d_0\ge 0$.

Write $t=\mathbb{E}W_k^2$. Crossing monotonicity and Cauchy–Schwarz give

$$\mathbb{E}R_+\le\frac{t}{(\zeta-u)^2},\quad \mathbb{E}\binom{R_-}{2}\le\frac{t}{[\zeta+\min(u,\zeta/3)]^2},\quad \mathbb{E}\binom{R_-}{3}\le\frac{t\sqrt{A^2+3t}}{(2\zeta-1/2)^3}.\tag{110}$$

These are the crossing-moment bounds of Lemma 7.2 with $A_U=A$, $V_k=W_k$, $|u|\le 1/2$ and the increment lower bound $\zeta=1-d_0$. The lemma includes the strict outgoing crossing and the fourth-moment estimate from independence and conditional centering.

The marked crossing.

Let $D_i=U_{1,i}-U_{0,i}$, $\vartheta=\mathbb{E}(D_i)_+=\mathbb{E}(D_i)_-$, $0<\eta\le 1$, and $R_f=\mathbb{E}\{u+\eta-W_k\}$. The two deleted laws are

$$A_0\overset{d}{=}kU_1+(n-k-1)U_0,\qquad A_1\overset{d}{=}(k-1)U_1+(n-k)U_0.$$

Then

$$\mathbb{E}R \le u + \eta - R_f + J, \tag{111}$$
$$J = \frac{\vartheta}{\eta} \sum_{i \ge 1} \{\mathbb{P}(A_0 \le u + d_0 - \zeta i) + \mathbb{P}(A_1 > u + \eta + i - A)\}.$$

Here are the pathwise details. Put $z = u - W_k$ and mark

$$Z_f = \sum_{i=1}^{n-k} (D_{k+i})_- \mathbf{1}_{\{W_k \le u - \zeta i\}}, \quad Z_b = \sum_{i=1}^{k} (D_{k-i+1})_+ \mathbf{1}_{\{W_k > u + \eta + i\}}.$$

If $R > z + \eta$, the crossing activates the marks through R: for $R > 0$ use $z \ge \zeta R$, while for $R < 0$ use $z < R - \eta$. In either case $R \le z + Z_f + Z_b$. Hence if $h = R - \lfloor z + \eta \rfloor > 0$, $Z_f + Z_b > h - 1 + \eta \ge \eta h$. Thus $R \le \lfloor z + \eta \rfloor + (Z_f + Z_b)/\eta$ on every path. On an active forward mark $U_0 > U_1 \ge -d_0$; deleting its pair implies $A_0 \le u + d_0 - \zeta i$. On a backward mark delete $U_1 \le A$ to obtain the strict event for A_1. The remaining sum is independent of the entire pair, so the expectations factor. Extending to all $i \ge 1$ adds only nonnegative bounds. Finally $|R| \le 1 + (|W_k| + |u|)/\zeta$ proves integrability and permits expectations. No coupling has replaced either deleted marginal.

A finite bound for every mark.

Fix $\lambda \in [L, U]$, $L > 0$, and $t \le \omega_+ \lambda$; put $T_L = \omega_+ L$, $T_U = \omega_+ U$. On a phase subcell $u \in [v_l, v_h]$, for $i = 1, 2$ let $b_1 = \zeta i - d_0 - v_h > 0$ and $b_2 = i - A + v_l > 0$. Lemma B.3 gives a bound $\omega_+ \lambda K_i^{\mathrm{ind}}$, where

$$K_i^{\mathrm{ind}} = \sum_{j=1}^{2} \min \left\{ \frac{1}{b_j^2 + T_L}, \frac{\overline{P}(A, T_U, b_j)}{T_L} \right\}.$$

The denominator T_L is a lower bound for the chosen upper envelope $\omega_+ \lambda$, not for any actual variance. Alternatively couple $A_0 = Z + U_1$, $A_1 = Z + U_0$ with $Z \stackrel{d}{=} (k-1)U_1 + (n-k-1)U_0$. The two events project into $Z \le -\alpha$ and $Z > \beta$, where $\alpha = \zeta i - 2d_0 - v_h > 0$, $\beta = i - A - d_0 + v_l > 0$. For any $z \in (-\alpha, \beta)$ their sum is at most $(\omega_+ \lambda + z^2)/\min(z + \alpha, \beta - z)^2$. Consequently a second kernel is

$$K_i^{\mathrm{com}} = \frac{1 + z^2/T_L}{\min(z + \alpha, \beta - z)^2}, \qquad z = \max\{-T_L/\beta, \min((\beta - \alpha)/2, T_L/\alpha)\}.$$

This center is inside the interval; differentiating the two quadratic ratios on either side of the midpoint proves its optimality, although any interior rational center suffices. Take the smaller kernel on each phase subcell and the maximum over their closed cover. Denote it K_i. For $i \ge 3$ use fourth moments and the convex midpoint integral for $(cx - b)^{-4}$. On $u \in [u_l, u_h]$ this gives

$$K = K_1 + K_2 + \frac{A^2 + 3T_U}{3\zeta} \left\{ \frac{1}{((5/2)\zeta - A - u_h)^3} + \frac{1}{((5/2)\zeta - A + u_l)^3} \right\}, \qquad J \le \frac{\vartheta}{\eta} \omega_+ \lambda K. \tag{112}$$

All denominators must be positive. Indeed, the tail fourth moment is at most $A^2 t + 3t^2$, and integration from $5/2$ to infinity covers every remaining integer, including the harmless added illegal counts.

If the same-law resource supplies $nd < G$ and $G^2 < c_v N L$, then

$$\vartheta \le \frac{c_v G \lambda}{c_v n \lambda + G^2}. \tag{113}$$

To prove this, for centered $X \ge -b$, $\mathbb{E}X^2 \le V$ and $b^2 \le V$, the quadratic $b(X - V/b)^2/(b + V/b)^2$ majorizes X_- on $[-b, \infty)$. Its expectation is at most $bV/(V + b^2)$. Apply this to $D_i \ge -d$, using $\mathrm{Var}(D_i) = s_0 + s_1 \le c_v\lambda/n$, and monotonicity in $b \le \sqrt{V}$. The case $d = 0$ is immediate. With $\eta = z\lambda$, $z > 0$, $zU \le 1$, (112) becomes

$$\frac{J}{\lambda} \le \frac{\omega_+ K c_v G}{z(c_v N\lambda + G^2)}. \tag{114}$$

The fractional-part gain.

The following implication assumes the scalar majorization and its rational constant enclosures stated below. Their finite arithmetic acceptance is a separate condition. If each centered summand has support span at most $0 < b < 1$ and the sum has variance t, then, for any real w and $0 < r_* \le 2\sin^2(\pi b)/b^2$,

$$\mathbb{E}\{w - W\} \ge \frac{5}{23}(1 - e^{-r_* t}). \tag{115}$$

Indeed $1 - \cos(2\pi x) \le (23/5)x$ for $0 \le x \le 1$. A finite proof of this sole scalar inequality is recorded below. For an independent copy Z' of one summand, concavity of sine on $[0, \pi b]$ gives

$$|\mathbb{E}e^{2\pi i Z}|^2 = 1 - 2\mathbb{E}\sin^2 \pi(Z - Z') \le e^{-4\sin^2(\pi b)\,\mathrm{Var}(Z)/b^2}.$$

Multiply square roots over the independent summands and use periodicity. This argument includes zero variances and integer collisions, and is valid up to $b < 1$ by Lemma 7.4. The other count denominator gates are unchanged. Here the conditional spans are at most $A + d_0$, so this is the b to use.

For a reproducible scalar proof, $x \ge 10/23$ follows from $1 - \cos \le 2$. On $[0, 10/23]$ use the degree-ten alternating upper polynomial for $1 - \cos(2\pi x)$, with directed coefficients:

$$Q(x) = \sum_{j=1}^{5} \frac{(-1)^{j+1}(2\pi_j x)^{2j}}{(2j)!}, \qquad \pi_j = \begin{cases} 3.141592653589794, & j \text{ odd}, \\ 3.141592653589793, & j \text{ even}. \end{cases}$$

Put $P(x) = 23/5 - Q(x)/x$. On the eight intervals with left endpoint $a_j = 5j/92$ and width $w = 5/92$, its degree-nine Bernstein coefficients are

$$b_{jk} = \sum_{i=0}^{k} \frac{\binom{k}{i}}{\binom{9}{i}} \sum_{h=i}^{9} [x^h]P(x) \binom{h}{i} a_j^{h-i} w^i, \qquad 0 \le j < 8, \quad 0 \le k \le 9.$$

All 80 rational coefficients exceed $4492/100000$. This finite comparison is reproduced by the independent rational check accompanying the manuscript; together with the displayed formula it proves the scalar majorization. Expansion in powers of the unit-interval coordinate proves the formula; Bernstein weights are nonnegative and sum to one. The alternating remainder and the stated rational bounds on π follow from the same elementary series above. This completely specifies the scalar comparison. To obtain a rational rate throughout $0 < b < 1$, put $y = \min(b, 1 - b)$ and $x = 3.14159y$. Since $0 < y \le 1/2$, $\sin(\pi b) = \sin(\pi y) \ge \sin x$, and the alternating bound

$$0 < x - x^3/6 + x^5/120 - x^7/5040 \le \sin x$$

is valid. Positivity follows from $x^2 < 3$ for the first two terms and $x^2 < 42$ for the last pair. Squaring this positive rational lower bound therefore gives a valid rate below $2\sin^2(\pi b)/b^2$. Folding at $1/2$ is essential: decreasing the sine argument would not be monotone on the whole unfurled interval.

Set $r = r_*\omega_-$, require $rU < 3$, and define

$$g = \frac{5}{23}\left(\frac{r^3L^2}{6} - \frac{r^4L^3}{24} + \frac{r^5L^4}{120} - \frac{r^6L^5}{720}\right). \tag{116}$$

The sixth-order Taylor lower bound for $1 - e^{-r\lambda}$ gives $R_f/\lambda \geq 5r/23 - 5r^2\lambda/46 + g$. The discarded higher-order part is increasing as a function of λ for $r\lambda < 3$: its derivative has the sign of $48 - 18x + (24/5)x^2 - x^3 > 0$ on $[0, 3]$. For instance this polynomial has positive degree-three Bernstein coefficients $48, 30, 132/5, 51/5$ there. Thus freezing it at L is valid.

One final expression.

On a signed phase cell find an interval $[x_l, x_h]$ containing H/ℓ by its four endpoint quotients, using positive bounds for ℓ. Let

$$D = \max(|x_l - u_h|, |x_h - u_l|), \quad D_+ = \max(0, x_h - u_l), \quad a = \max(|u_l|, |u_h|), \quad b_\chi = p_+ - x_l.$$

The same formula allows coarser upper bounds D, D_+, b_χ, as used by the central certificate. Check $D \leq \Delta$ and $b_\chi + u_h < 2$. With $\omega_\pm = 1 \pm c_v D/N$, the curvature coefficient is

$$V_0 = \frac{\omega_+}{v_*}\left[\sup_{u_l \leq u \leq u_h}\left\{\frac{(b_\chi + u - 1)_+}{(\zeta - u)^2} + \frac{(b_\chi + u)_+}{[\zeta + \min(u, \zeta/3)]^2}\right\} + \frac{\sqrt{A^2 + 3T_U}}{(2\zeta - 1/2)^3}\right]. \tag{117}$$

On a rational phase subcell, both numerators are bounded at the right endpoint; the first denominator is smallest there, and the second is smallest at the left endpoint. This proves the supremum bound, including both kinks. Equations (108)–(110), $\sqrt{n}m_B \leq 1$, and (88) now give the whole-count witness with

$$B(\lambda) = z + \frac{\omega_+ K c_v G}{z(c_v N\lambda + G^2)} - \frac{5r}{23} + \frac{5r^2\lambda}{46} - g, \tag{118}$$

$$I(\lambda) = V_0 + aB_1 + \frac{aP_1}{\lambda} + P_*D_+ - m_cQ_*.$$

We used $|u - f(u)| \leq a(B_1\lambda + P_1)/n$ and $\sqrt{n}\pi_k \leq 1$. Every constant is fixed before λ varies. Both B and $mB + I/N$ are convex on $\lambda > 0$; I is decreasing. Thus their two endpoint gates $B < 0$, $I > 0$, $mB + I/N < -0.001$ prove the entire closed noise cell. Thus this argument supplies the analytic implication for each of the three whole-count estimates, conditional on its own complete cover and endpoint gates.

The zero-noise endpoint and the remaining composition

If $\lambda = 0$, then $p, q > 0$ and conditional centering imply $U_0 = U_1 = 0$ almost surely. The actual law is its Bernoulli comparison law. The zero-noise argument in Section 7 places an attained positive closed-CDF maximum at a binomial atom, so $u = 0$ and $C = C_B$. The external all-probability binomial theorem of [19, Theorem 1], when applied with its stated hypotheses, then supplies the sharp Bernoulli upper bound. An equality-inclusive exclusion requires a separately justified strict finite-index version. For positive noise, the local conclusion requires every parameter cell to satisfy one of the displayed strict tests. No selection of historical table rows, completeness of a finite cover, or fresh acceptance of those tests is asserted here.

The mathematical composition therefore still requires: the actual contact witness; already acquired common-law support, moments and signed resources; any additional concentration hypotheses used in acquiring confinement; valid normalization and positivity of every divisor; a complete cover of all signs, phases and noise values; and acceptance of the stated finite gates on that cover. No step replaces an $n - 2$ convolution by an $n - 1$ convolution or infers a uniform statement from a fixed-law asymptotic limit.

C Binomial normalization and elementary constants

C.1 A Taylor segment crossing one half

A phase cell can place the Taylor segment of a binomial mass on both sides of $1/2$. The required derivative estimates remain valid provided the segment lies in an interval $[a, b]$ with $a + b < 1$. We give the argument because imposing a one-sided segment would unnecessarily exclude such cells.

Lemma C.1 (Symmetric derivative envelopes)**.** *If $0 < a < 1/2$, $a < b < 1$, $a + b < 1$, then for $r \in [a, b]$,*

$$r(1-r) \geq a(1-a), \qquad \frac{|1-2r|}{[r(1-r)]^j} \leq \frac{1-2a}{[a(1-a)]^j} \quad (j = 1, 2).$$

Proof. $[a, b] \subset [a, 1-a]$. Set $s = \min(r, 1-r) \in [a, 1/2]$. The expressions are symmetric; $s(1-s)$ increases and $1-2s$ decreases nonnegatively. Their positive quotients therefore have their maxima at a. □

Let $n \geq N \geq 1$, $p \in [p_-, p_+] \subset (0, 1/2)$, $q = 1-p$, $1 \leq k \leq n-1$, $\delta = k - np$, $r = k/n$, $v = pq$. Suppose the whole segment between p, r lies in $[a, b]$ above. If

$$E \geq \frac{1-2a}{a^2(1-a)^2}, \quad L \geq \frac{1-2a}{a(1-a)}, \quad R \geq \frac{1}{12a(1-a)},$$

then the binomial mass π_k satisfies

$$\sqrt{n}\,\pi_k = \frac{\phi(0)}{\sqrt{v}} e^{-\delta^2/(2nv)+\zeta}, \qquad n|\zeta| < L|\delta|/2 + R + E|\delta|^3/(6N).$$

For proof, relative entropy has $D(p\|p) = D'(p\|p) = 0$, $D''(p\|p) = 1/v$ and $D^{(3)}(r\|p) = (2r-1)/[r^2(1-r)^2]$. Taylor bounds $|nD(r\|p) - \delta^2/(2nv)|$ by $E|\delta|^3/(6n^2)$; the logarithmic variance prefactor changes by at most $L|\delta|/(2n)$. Finite Stirling, $0 < s_j < 1/(12j)$, gives exactly

$$\sqrt{n}\,\pi_k = \frac{\phi(0)}{\sqrt{r(1-r)}} e^{-nD(r\|p)+s_n-s_k-s_{n-k}}.$$

Writing $e = s_n - s_k - s_{n-k}$, use the compatible one-sided estimates $e < 1/(12n)$ and $e > -1/(12k) - 1/(12(n-k)) = -1/[12nr(1-r)]$. Thus $|e| < R/n$, and collecting all three errors proves the claim.

For finite interval applications choose rational endpoints satisfying $a < p_- - \Delta/N$, $p_+ + \Delta/N < b$, $a + b < 1$. For $|\delta| \leq \Delta$ they contain the whole segment. The inequalities $Np_- - \Delta - 2 > 0$ and $N(1-p_+) - \Delta - 2 > 0$ ensure interior/adjacent factorial domains. Positive $v_f < a(1-a)$ and upward-rounded $(1-2a)/v_f^2$, $(1-2a)/v_f$, $1/(12v_f)$ supply E, L, R. Mass bounds use $e^{-x} \geq 1-x$, $e^x \leq 1/(1-x)$ only on their stated positive domains. The original p band remains below $1/2$ when bounding pq; monotonicity is not assumed on the enlarged Taylor segment.

C.2 Elementary Gaussian constants

Lemma C.2 (Elementary Gaussian constants)**.** *For the standard normal density,*

$$\|\phi''\|_\infty = \phi(0) < 2/5, \qquad \|\phi'''\|_\infty \leq 2/\pi, \qquad 1/(4\pi) < 2/25.$$

Moreover $819/2000 < C_{\mathrm{E}} < 20487/50000 < 41/100$ *and* $\phi(0) > 0.39886 > 0.3988 = 997/2500$.

We give the exact comparisons used in Lemma C.2.

Proof. Since $\phi''(x) = (x^2-1)\phi(x)$, the norm on $x^2 \le 1$ is at most $\phi(0)$. On $x^2 \ge 1$, the maximum of $(x^2-1)e^{-x^2/2}$ is $2e^{-3/2} < 1$, attained when $x^2 = 3$. Thus the norm is exactly $\phi(0)$.

For completeness, the Gaussian transform can be obtained without an inversion theorem for the distribution being studied. Differentiating $J(t) = \int_{\mathbb{R}} e^{-x^2/2}e^{itx}\,dx$ under the integral and integrating by parts give $J'(t) = -tJ(t)$. The Gaussian integral, evaluated by squaring it and using polar coordinates, gives $J(0) = \sqrt{2\pi}$. Hence $J(t) = \sqrt{2\pi}e^{-t^2/2}$; applying the same formula with the variables interchanged gives

$$\phi(x) = \frac{1}{2\pi}\int_{\mathbb{R}} e^{-t^2/2}e^{-itx}\,dt.$$

Three differentiations are justified by the integrability of $|t|^3e^{-t^2/2}$. Consequently

$$\|\phi'''\|_\infty \le \frac{1}{2\pi}\int_{\mathbb{R}} |t|^3 e^{-t^2/2}\,dt = \frac{2}{\pi}.$$

The last integral equals four after the substitution $w = t^2/2$ on the positive half-line.

Here are elementary rational comparisons for all numerical constants used in this section. Put $a = \arctan(1/5)$ and $b = \arctan(1/239)$. Tangent addition gives $\tan(2a) = 5/12$, $\tan(4a) = 120/119$ and $\tan(4a-b) = 1$. Also

$$0 < 4(1/5 - 1/(3\cdot 5^3)) - 1/239 < 4a - b < 4/5 < 1 < \pi/2.$$

The arctangent inequalities follow by integrating $1 - t^2 < 1/(1+t^2) < 1$, while $\pi/2 = 2\arctan(1) > 1$ follows by integrating $1/(1+t^2) > 1/2$ on $[0,1)$. It follows that $\pi = 16a - 4b$ and

$$\pi > 16(1/5 - 1/(3\cdot 5^3)) - 4/239 = 281476/89625 > 25/8.$$

This proves $\phi(0) < 2/5$ and $1/(4\pi) < 2/25$.

To bound C_{E} more closely, define

$$s_k(x) = \sum_{j=0}^{k-1} \frac{(-1)^j x^{2j+1}}{2j+1}, \quad \pi_- = 16s_6(1/5) - 4s_1(1/239), \quad \pi_+ = 16s_5(1/5) - 4s_2(1/239).$$

The remainder in the finite geometric expansion of $1/(1+t^2)$ has its stated alternating sign, so integration gives $\pi_- < \pi < \pi_+$. Let $r_- = 3162277660/10^9$ and $r_+ = 3162277661/10^9$. Squaring shows $r_-^2 < 10 < r_+^2$. The following comparisons are exact rational ones:

$$\left(\frac{819}{2000}\right)^2 < \frac{19+6r_-}{72\pi_+} < C_{\mathrm{E}}^2 = \frac{19+6\sqrt{10}}{72\pi} < \frac{19+6r_+}{72\pi_-} < \left(\frac{20487}{50000}\right)^2.$$

For example, each outer comparison follows by multiplying by its positive rational denominator; the displayed finite sums specify those denominators completely. All quantities are positive, so taking square roots proves the claimed bounds. In particular, $0.4095 < C_{\mathrm{E}} < 0.40974 < 0.41$. The same upper endpoint for π is smaller than $22/7$. Since $(0.39886)^2 < 7/44$, we also obtain $\phi(0) > 0.39886 > 0.3988$. □

D Signed refinements of the local comparison

The refinements in this appendix apply to an already acquired common-law resource panel. They preserve the actual count while integrating a Gaussian increment, and retain the sign of the cubic correction. They are possible ingredients of a comparison with floor 42; each retains its own probability and positivity domain. Every numerical positivity and coverage statement below is an acceptance condition. Whenever a normalized binomial mass or a Stirling expression is used, assume $n \ge N \ge 2$ and an interior integer count $1 \le k \le n-1$, so that $\pi_k > 0$.

Write $\xi = H/\ell$ for the raw threshold in Bernoulli coordinates. The normalization constants are supplied by Appendix C. The underlying count identities, moment and tail comparisons are in Section 3 and Appendix B.

D.1 Notation

Let $Y = B + U$, where $\mathbb{P}(B = 1) = p$, $\mathbb{E}(U \mid B) = 0$, $v = p(1-p)$, $s = \mathbb{E}U^2$, $\lambda = ns$, and $\ell = (v+s)^{-1/2}$. The actual threshold has the form

$$x = np + H/\ell = k + u, \qquad |u| \le \tfrac{1}{2}, \qquad \delta = k - np = H/\ell - u.$$

The integer k and the value δ stay fixed when integrating a function of a new variable t below. Write $\pi_k = \mathbb{P}(\mathrm{Bin}(n,p) = k)$ and

$$f(t) = \frac{\Phi((\delta+t)/\sqrt{n(v+s)}) - \Phi(\delta/\sqrt{n(v+s)})}{\pi_k}.$$

The conditional errors satisfy $|U_j| \le A$, $U_0 \le a_0$, $U_1 \ge -a_1$, with $a_0, a_1 \ge 0$ and $d = a_0 + a_1$. All quantities in an application refer to the same iid marginal.

Lemma D.1 (A joint budget on a noise interval)**.** *Suppose the acquired same-law resource inequality is*

$$\frac{d}{d_*} + \frac{\lambda}{\Lambda} \le 1, \qquad d_* > 0, \quad \Lambda > 0.$$

On a closed noise interval $L \le \lambda \le U \le \Lambda$ one may use

$$d \le d_L := d_*(1 - L/\Lambda).$$

Every conditional support, cubic-moment, tail-distance and characteristic function bound whose hypothesis is $d \le d_0$ may therefore be applied with $d_0 = d_L$ on that interval.

Proof. Rearrange the displayed resource inequality and use $\lambda \ge L$. □

For example, the whole-count kernel uses $\zeta = 1 - d_0$. Its moment coefficients are recomputed as

$$\alpha = 6m_0\ell_-^2(1-d_0), \qquad \beta = \min\{3(4.16)\ell_-(1-p_+-d_0), 3(6.16)\ell_-(p_- - d_0)\},$$

where m_0, ℓ_- are the previously justified moment and scale lower bounds. The ordinary cubic correction is recomputed with the same d_0. Likewise, a noise amplitude bound A gives a conditional support span at most $A + d_0$; the characteristic-function rate is recomputed from that span. These substitutions are licensed only when the original coefficient inequalities and positivity gates hold for the acquired moment and scale bounds.

D.2 A binomial mass bound on the actual phase cell

Suppose the finite Stirling argument has supplied positive constants E, L_0, R on a fixed segment containing every k/n under consideration, with

$$\sqrt{n}\,\pi_k = \frac{\phi(0)}{\sqrt{v}}\exp\{-\delta^2/(2nv)+\zeta\}, \qquad n|\zeta| \le Z_N(|\delta|), \qquad Z_N(t) = \frac{L_0 t}{2}+R+\frac{Et^3}{6N}, \quad n \ge N.$$

The constants may have been computed on a much wider global phase range. Let $0 < v_- \le v \le v_+$ and let $q_+ \ge \sqrt{v_+}$.

Lemma D.2 (Mass localization)**.** *On any smaller closed cell where $|\delta| \le D$,*

$$\sqrt{n}\,\pi_k \ge \frac{0.39886}{q_+}\left[1 - \frac{D^2/(2v_-) + Z_N(D)}{N}\right].$$

The maximum of this expression and any previously proved mass lower bound is also a valid mass lower bound. There is no requirement $D \ge 1$.

Proof. Lemma C.2 gives $\phi(0) > 0.39886$. Set $A_* = D^2/(2v_-) + Z_N(D) \ge 0$. The logarithmic formula gives $\sqrt{n}\pi_k \ge (0.39886/q_+)e^{-A_*/n}$. Now $e^{-z} \ge 1 - z$ and $n \ge N$. If the bracket is negative the displayed inequality still holds, and the previously positive bound can be retained. The global Stirling segment, rather than the smaller value D, continues to justify E, L_0, R. □

D.3 A sharper elementary harmonic constant

Lemma D.3. *For every $x \ge 0$,*

$$1 - \cos(2\pi x) \le \frac{4553}{1000}x.$$

Proof. Put $K = 4553/1000$. For $x \ge 2/K$ the assertion follows from $1 - \cos \le 2$. On $[0, 2/K]$ take the degree-fourteen alternating upper Taylor polynomial $Q(x)$ for $1 - \cos(2\pi x)$. Its remainder is nonpositive: the first omitted term is negative, and the remaining term magnitudes decrease, since $(4\pi/K)^2 < 16 \cdot 15$.

Use the directed endpoints

$$\pi_- = 3.141592653589793 < \pi < 3.141592653589794 = \pi_+.$$

In each positive coefficient of Q replace π by π_+; in each negative coefficient replace it by π_-. This only increases Q. Expand $P(x) = K - Q(x)/x$, a polynomial of degree thirteen, on each of the 64 intervals $[2j/(64K), 2(j+1)/(64K)]$. If $P(x) = \sum_{h=0}^{13} p_h x^h$, its Bernstein coefficients on $[a, a+w]$ are

$$b_k = \sum_{i=0}^{k} \frac{\binom{k}{i}}{\binom{13}{i}} \sum_{h=i}^{13} p_h \binom{h}{i} a^{h-i} w^i, \qquad k = 0, \dots, 13.$$

All 896 rational coefficients exceed $123/10^6$, as checked directly from the displayed rational formula. Thus $P > 0$ on the full interval. The value at zero follows by continuity (or directly).

For completeness the stated π endpoints are checked using alternating series for $\arctan(1/5)$ (32 terms) and $\arctan(1/239)$ (12 terms). Machin's identity was established in Lemma C.2; the term counts here give the finer directed endpoints required by this polynomial. The finite recipe uses only rational operations and the Bernstein identity proved in Section 8. □

If the original characteristic-function argument gives $|\mathbb{E}e^{2\pi iW}| \le e^{-rt}$, Lemma D.3 upgrades its fractional part estimate from $(5/23)(1-e^{-rt})$ to

$$\frac{1000}{4553}(1-e^{-rt}).$$

Consequently every occurrence of $5/23$ and $5/46$ in that harmonic gain is replaced by $1000/4553$ and $500/4553$, respectively. The existing degree-six Taylor gain and its monotonicity argument remain unchanged.

D.4 The Gaussian derivative knows the sign of the threshold

Lemma D.4 (Positive phase, with a fixed actual count)**.** *Assume $u \in [a,b] \subset [0,1/2]$, $H/\ell \le \xi_*$, $|\delta| \le D$, $v \ge v_- > 0$, $\lambda \le U$, and the Stirling remainder bound above. Set*

$$G = \max\left(0, \max_{z\in[a,b]}(2\xi_* z - z^2)\right), \qquad \eta = \frac{U+G}{2v_-} + Z_N(D).$$

Then $f(u) \ge u(1-\eta/N)$ for every $n \ge N$. The inner maximum is attained at the projection of ξ_* onto $[a,b]$. If $H \le 0$, one may take $\xi_* = 0$, hence $G = 0$.

Proof. For a fixed actual law and threshold, differentiate while keeping $\delta = k - np$ fixed:

$$\log f'(t) = -\tfrac{1}{2}\log(1+s/v) + \frac{\delta^2}{2nv} - \frac{(\delta+t)^2}{2n(v+s)} - \zeta.$$

For $0 \le t \le u$, use $\log(1+s/v) \le s/v$ and $v+s \ge v$ to obtain

$$\log f'(t) \ge -\frac{\lambda + 2\delta t + t^2}{2nv} - \zeta.$$

The convex function $t \mapsto 2\delta t + t^2$ attains its maximum on $[0,u]$ at an endpoint. Since the actual relation is $\delta = H/\ell - u$,

$$2\delta t + t^2 \le \max(0, 2(H/\ell)u - u^2) \le G.$$

Thus $\log f'(t) \ge -\eta/n$, and $f'(t) \ge e^{-\eta/n} \ge 1 - \eta/N$. Integrate from zero to the actual u. □

A signed H interval supplies the bound D. If $H \ge 0$, one can use $\xi_* = H_+/\ell_-$ and $D = \max(b, \xi_* - a)$; if $H \le 0$, use $D = H_-/\ell_- + b$. The new lower derivative constant is maximized with the older valid constant. Every resulting bound is checked positive.

The normal variance and cubic-moment repair is localized separately. On a phase cell, use its actual upper bound for δ_+, rather than $|\delta|$. If the resulting coefficient is κ, replace it by $\max(0,\kappa)$ before using $\kappa/n \le \kappa/N$. That sign condition is essential for an all-index conclusion.

D.5 Reusing a curvature bound

For fixed probability, threshold and phase intervals, the curvature function in Section 7.3 is nonincreasing in ζ. Lemma D.1 makes $\zeta = 1 - d_L$ increase as the lower noise endpoint L increases. A curvature upper bound for the first band therefore remains valid on later bands. The other ζ-dependent moment and support bounds are evaluated using the value for each band.

D.6 Signed Gaussian corrections on both phase halves

Let $n \geq N$, $v = p(1-p) > 0$, $s \geq 0$, $\lambda = ns$, and let the actual nearest count and phase satisfy

$$\delta = k - np = \xi - u, \qquad f(t) = \frac{\Phi((\delta+t)/\sqrt{n(v+s)}) - \Phi(\delta/\sqrt{n(v+s)})}{\pi_k}.$$

The actual k and δ are held fixed when t varies. Assume the proved finite Stirling formula on this count range is

$$\sqrt{n}\,\pi_k = \frac{\phi(0)}{\sqrt{v}} \exp\{-\delta^2/(2nv) + \zeta\}, \qquad n|\zeta| \leq Z_N(D), \quad |\delta| \leq D.$$

Write $v \geq v_- > 0$ and $Z = Z_N(D) \geq 0$.

Lemma D.5 (Negative phase)**.** *Suppose $u = -r$, $0 \leq a \leq r \leq b$, and $\xi \leq \xi_*$. Define*

$$t_* = \max(0, \min(b, \xi_* + b)), \qquad G_- = 2(\xi_* + b)t_* - t_*^2, \qquad \eta_- = G_-/(2v_-) + Z.$$

If $D^2 \leq Nv_-$ and $\eta_- < N$, then

$$f'(-t) \leq \frac{1}{1 - \eta_-/N} \quad (0 \leq t \leq r), \qquad u - f(u) \leq \frac{b\eta_-}{n(1-\eta_-/N)}.$$

Proof. Direct differentiation and the Stirling formula give

$$\log f'(-t) = -\tfrac{1}{2}\log(1 + s/v) + \frac{\delta^2 s}{2nv(v+s)} + \frac{2\delta t - t^2}{2n(v+s)} - \zeta.$$

Since $\log(1+x) \geq x/(1+x)$ for $x \geq 0$, the first two terms are at most

$$\frac{s}{2(v+s)}\left(\frac{\delta^2}{nv} - 1\right) \leq 0.$$

This cancellation removes the variance term from the upper derivative bound. Because $\delta = \xi + r$, $r \leq b$ and $0 \leq t \leq r \leq b$,

$$2\delta t - t^2 \leq 2(\xi_* + b)t - t^2 \leq G_-.$$

The last maximum is the concave quadratic maximum on $[0, b]$, achieved at the displayed projection t_*. It is nonnegative because $t = 0$ is available. Therefore $\log f'(-t) \leq \eta_-/n$. Using $e^x \leq (1-x)^{-1}$ for $0 \leq x < 1$ gives

$$f'(-t) - 1 \leq \frac{\eta_-/n}{1 - \eta_-/n} \leq \frac{\eta_-}{n(1-\eta_-/N)}.$$

Integrate $u - f(u) = \int_0^r (f'(-t) - 1)\,dt$. □

Equivalently, $G_- = b^2 + 2\xi_* b$ for $\xi_* \geq 0$, and $G_- = \max(0, \xi_* + b)^2$ for $\xi_* < 0$. This sharper vertex is used in the whole-count comparison on the left probability panels at floor 42.

Lemma D.6 (Positive phase)**.** *Suppose $u \in [a, b] \subset [0, 1/2]$, $\xi \leq \xi_*$, and define*

$$G_+ = \max\left(0, \max_{z \in [a,b]} (2\xi_* z - z^2)\right).$$

Then

$$u - f(u) \leq \frac{b}{n}\left(\frac{\lambda}{2v_-} + \frac{G_+}{2v_-} + Z\right).$$

The inner maximum is at the projection of ξ_* onto $[a, b]$.

Proof. The exact derivative identity and $\log(1+s/v) \le s/v$ give

$$\log f'(t) \ge -\frac{\lambda + 2\delta t + t^2}{2nv} - \zeta.$$

For fixed actual u, the function $2\delta t + t^2$ is convex on $[0, u]$. Its maximum is at an endpoint, and $\delta = \xi - u$ gives

$$2\delta t + t^2 \le \max(0, 2\xi u - u^2) \le G_+.$$

Use $e^{-x} \ge 1 - x$ and integrate from zero to u. □

D.6.1 Whole-count insertion and interval coverage

The whole-count identity is split as $Q/\pi_k - f(u) = (Q/\pi_k - u) + (u - f(u))$. Thus one needs an upper bound on the signed correction $u - f(u)$; an absolute-value estimate is unnecessary. After division by λ, the new coefficients replacing the old Gaussian correction are

$$\begin{aligned} u \ge 0: &\quad \frac{b}{2v_-} + \frac{b\{G_+/(2v_-) + Z\}}{\lambda}, \\ u \le 0: &\quad \frac{b\eta_-}{(1 - \eta_-/N)\lambda}. \end{aligned}$$

On each fixed phase cell these have the form $A + B/\lambda$ with $A, B \ge 0$. They therefore preserve the original convex endpoint check on each closed noise interval. Every phase cell is confined to one half-line; zero is a shared closed endpoint. The count displacement bound D comes from the same actual $\xi - u$ cell, and the Stirling constants continue to come from the already proved global count segment.

The other all-index steps are unchanged: the event bracket is checked negative before substituting a binomial mass lower bound, and the entire inverse-n correction is checked positive before replacing n by N. The inverse correction is $I(\lambda) = K + A + B/\lambda$, hence decreases with λ; positivity at both endpoints (in particular the upper endpoint) licenses the all-index step throughout the interval. The final score is convex in λ: its event fraction is a positive constant divided by an affine positive function, its B/λ coefficient is nonnegative, and its remaining terms are affine or constant. These are independent gates in the exact kernel. The formulas apply to both signs of ξ without reflecting the law or changing the count during integration. They also supply an improved upper derivative for the negative-phase nearest-count inequalities.

D.6.2 A complete phase cover below the whole-count threshold

For $\lambda \in [0.05, 0.08]$, the shifted-quartic nearest-count comparisons must be checked on every closed phase interval of length 0.001 in $[0, 1/2]$, for both phase signs. The shifted-quartic polynomial and nearest-count identity are defined at phase zero, so the two closed covers include $u = 0$ directly. This permits the whole-count comparison to start at $\lambda = 0.08$. The four nearest-count bands use phase starts $0, 0, 0.19, 0.27$ at noise bands $[0.05, 0.06]$, $[0.06, 0.08]$, $[0.08, 0.10]$, and $[0.10, 0.12]$, respectively.

D.7 Conditional centering and the exact cubic bound

Let B be Bernoulli with parameter $0 < p < 1/2$, put $q = 1 - p$ and $v = pq$, and write $Y = B + U$, with $\mathbb{E}(U \mid B) = 0$. Let U_0, U_1 denote the conditional laws of U, and assume

$$U_0 \le a_0 < p, \qquad U_1 \ge -a_1 > -(1-p), \qquad a_0, a_1 \ge 0, \qquad d = a_0 + a_1.$$

Put $s_j = \mathbb{E}U_j^2$, $s = qs_0 + ps_1$, $\lambda = ns$, and

$$X = \frac{Y - p}{\sqrt{v + s}}, \quad \rho = \mathbb{E}|X|^3, \quad \rho_B = \frac{1 - 2v}{\sqrt{v}}.$$

Finite third moments are assumed. The support conditions fix the sign of $Y - p$ in each class, which is necessary for the following expansion.

Lemma D.7 (No positivity assumption on the cubic bracket)**.** *With the preceding notation,*

$$(v + s)^{3/2}(\rho - \rho_B) \geq sQ, \qquad Q = 6p - 3p^2 - \frac{3}{2} - d - \frac{3(1 - 2v)s}{8v}. \tag{119}$$

Proof. Expanding within each sign class and using conditional centering gives

$$\mathbb{E}|Y - p|^3 = v(1 - 2v) + 3pq(s_0 + s_1) + p\mathbb{E}U_1^3 - q\mathbb{E}U_0^3.$$

The one-sided support bounds imply $\mathbb{E}U_1^3 \geq -a_1 s_1$ and $\mathbb{E}U_0^3 \leq a_0 s_0$. Moreover $3pq(s_0 + s_1) \geq 3p(qs_0 + ps_1) = 3ps$, because $q \geq p$. Consequently

$$\mathbb{E}|Y - p|^3 \geq v(1 - 2v) + (3p - d)s.$$

For $z \geq 0$, Taylor's formula gives $(1 + z)^{3/2} \leq 1 + 3z/2 + 3z^2/8$: the third derivative is negative. Subtract

$$\rho_B(v + s)^{3/2} = v(1 - 2v)(1 + s/v)^{3/2} \leq v(1 - 2v) + \tfrac{3}{2}(1 - 2v)s + \frac{3(1 - 2v)s^2}{8v}$$

from the preceding moment lower bound and use $3p - \frac{3}{2}(1 - 2v) = 6p - 3p^2 - \frac{3}{2}$. □

D.8 The coefficient must be bounded in the correct direction

Suppose $n \geq N$, $p \in [p_-, p_+] \subset (0, 1/2)$, $0 \leq d \leq d_0$, and $0 \leq \lambda \leq \Lambda$. Choose $0 < v_* \leq p_-(1 - p_-)$ and put

$$V^* = p_+(1 - p_+) + \Lambda/N, \qquad Q_* = 6p_- - 3p_-^2 - \frac{3}{2} - d_0 - \frac{3(1 - 2v_*)\Lambda}{8v_* N}.$$

Then $Q \geq Q_*$: the first term increases with p, and $(1 - 2v)/v$ decreases with v. In particular, Q_* may be negative. Already with $d = s = 0$ its sign changes at $p = 1 - 1/\sqrt{2}$.

Let $c = C_{\mathrm{E}}$ be the fixed coefficient multiplying the moment correction, and choose $0 < c_* \leq c \leq c^*$. Choose positive multipliers satisfying

$$0 < m_- \leq \frac{c_*}{(V^*)^{3/2}}, \qquad m_+ \geq \frac{c^*}{v_*^{3/2}},$$

and define the signed gain

$$G_* := \min\{m_- Q_*, m_+ Q_*\}. \tag{120}$$

Proposition D.8 (Signed gain and nonnegative repair)**.** *Let Φ denote the standard normal distribution function. For an actual count displacement δ, set*

$$\mathcal{R} = \sqrt{n}\left\{\Phi\left(\frac{\delta}{\sqrt{nv}}\right) - \Phi\left(\frac{\delta}{\sqrt{n(v + s)}}\right)\right\} + c(\rho_B - \rho).$$

If $D_+ \geq \max(\delta, 0)$ and $P_* \geq 0.4/(2v_*^{3/2})$, then

$$\mathcal{R} \leq (P_* D_+ - G_*)\frac{\lambda}{n} \leq \kappa\frac{\lambda}{N}, \qquad \kappa = \max\{0, P_* D_+ - G_*\}. \tag{121}$$

Proof. Write $A = c/(v+s)^{3/2}$. The hypotheses give $0 < m_- \le A \le m_+$. By Lemma D.7, $c(\rho - \rho_B) \ge sAQ \ge sAQ_*$. If $Q_* \ge 0$, $AQ_* \ge m_- Q_*$; if $Q_* < 0$, $AQ_* \ge m_+ Q_*$. This proves $c(\rho - \rho_B) \ge sG_*$ in both cases. One must use the *upper* multiplier in the negative case.

The normal part is nonpositive when $\delta \le 0$. For $\delta > 0$, differentiation with respect to an intermediate variance $t \in [0, s]$ gives

$$\frac{d}{dt}\left[-\sqrt{n}\,\Phi\left(\frac{\delta}{\sqrt{n(v+t)}}\right)\right] = \frac{\delta}{2(v+t)^{3/2}}\phi\left(\frac{\delta}{\sqrt{n(v+t)}}\right) \le \frac{0.4\delta}{2v_*^{3/2}} \le P_* D_+.$$

Integrate over $[0, s]$, and combine with the moment bound. Finally $s = \lambda/n$ and $\kappa \ge 0$, so $(P_* D_+ - G_*)\lambda/n \le \kappa\lambda/n \le \kappa\lambda/N$. The clamp at zero is needed before replacing n by its lower bound. □

D.9 What must be checked on a new cover

For the fixed target one can take

$$N = 42, \qquad c_* = 819/2000, \qquad c^* = 20487/50000, \qquad c_* < C_{\mathrm{E}} < c^*.$$

On each actual probability and noise cell, verify positivity of the multipliers as well as

$$m_-^2 (V^*)^3 \le c_*^2, \qquad m_+^2 v_*^3 \ge (c^*)^2, \qquad 4P_*^2 v_*^3 \ge 0.4^2.$$

The upper multiplier is indispensable when $Q_* < 0$. All constants must be computed from the same acquired support, moment and signed resource bounds. The new entry range is $p \in [.25, .53]$, whereas the formulas above assume a probability interval below $1/2$. A complete comparison still needs every surviving probability, phase and positive-noise interval, including the part above $1/2$, and a separate zero-noise argument.

An earlier proposed schedule used narrower probability rows and noise bands with stronger amplitude caps. Those rows are not outputs of the entry proved in this paper. We therefore do not import that schedule as a certificate of the new comparison. Every new amplitude cap must first be acquired by contact, and every seam between noise and phase intervals must be covered as a closed boundary.

E Closed-half-line smoothing

The finite-index reduction in Appendix F needs a majorant for the closed event, including equality at an atom. We give its construction and normalization here. This appendix is independent of the Fourier remainder used in the fixed-target coarse entry.

E.1 Exact half-line smoothing

We construct a majorant of the closed half-line and justify its expectation under an integrable random variable. Its value at the boundary is part of the proof, because the finite-index laws can have atoms.

Write ϕ and Φ for the standard normal density and distribution function. All terminating decimals are exact rational numbers. The imaginary unit is i. Symbols introduced within a subsection, including D, R, L, s and K, are local to that subsection unless explicitly carried forward. In particular the moment defect D, an auxiliary distance bound, and the Bernoulli inward displacement d are different quantities.

The one-sided kernel is

$$K(t) = \frac{1-t}{2} + \frac{i}{2}\left((1-t)\cot(\pi t) + \frac{1}{\pi}\right), \qquad 0 < t < 1.$$

For $\kappa \in \{K, \overline{K}\}$, $0 < b \le 1$ and $g(t) = g(0) + O(t)$ with real $g(0)$, the convention throughout is

$$\mathfrak{I}_{\kappa,b}(g) = 2 \lim_{\epsilon \downarrow 0} \mathrm{Re} \int_{\epsilon}^{b} \kappa(t) g(t)\, dt. \tag{122}$$

Only the real-part improper limit is intended; the unmodified complex integral need not converge. The proof below establishes directly that

$$M(x) = \tfrac{1}{2} + \mathfrak{I}_{K,1}(e^{2\pi i t x}) \ge \mathbf{1}_{(-\infty,0]}(x), \qquad 1 - M(-x) \le \mathbf{1}_{(-\infty,0)}(x),$$

including atoms at the boundary, and justifies the expectation identity for an integrable random variable.

E.1.1 A real-variable Beurling construction

Set

$$W(x) = \left(\frac{\sin \pi x}{\pi x}\right)^2 \quad (x \ne 0), \qquad W(0) = 1,$$

and define the continuous odd function

$$H_B(x) = \sum_{m=1}^{\infty} \{W(x-m) - W(x+m)\} + 2xW(x). \tag{123}$$

The series converges absolutely and locally uniformly on the real line: for $|x| \le R$ and $m > 2R + 1$, each shifted term is bounded by a constant times m^{-2}. Thus the function and all the integer values in what follows are unambiguous. In particular,

$$H_B(0) = 0, \qquad H_B(j) = \mathrm{sgn}(j) \quad (j \in \mathbb{Z} \setminus \{0\}). \tag{124}$$

Here $\mathrm{sgn}(0) = 0$. The corresponding classical Beurling majorant is $B = H_B + W$. For noninteger x, its series is

$$B(x) = \left(\frac{\sin \pi x}{\pi}\right)^2 \left\{ \frac{2}{x} + \sum_{m=0}^{\infty} \frac{1}{(x-m)^2} - \sum_{m=1}^{\infty} \frac{1}{(x+m)^2} \right\}. \tag{125}$$

We prove the needed inequality directly, without importing an extremality or Fourier-transform theorem about B.

The elementary partial-fraction identity

$$\sum_{m \in \mathbb{Z}} \frac{1}{(y-m)^2} = \frac{\pi^2}{\sin^2 \pi y}, \qquad y \notin \mathbb{Z}, \tag{126}$$

has the following short proof. Integrate $\pi \cot(\pi z)/(z-y)^2$ counterclockwise over the square with vertices $\pm(N + \frac{1}{2}) \pm i(N + \frac{1}{2})$. The cotangent is uniformly bounded on these edges: on the vertical edges it is a multiple of a hyperbolic tangent, and on the horizontal edges its modulus is bounded by $\coth(\pi(N + \frac{1}{2}))$. The denominator is of order N^2 and the perimeter of order N, so the integral tends to zero. Its residues are $(m-y)^{-2}$ at $m = -N, \dots, N$, and $-\pi^2 \csc^2(\pi y)$ at y. The residue theorem and absolute convergence give (126).

For $y > 0$, $y \notin \mathbb{Z}$, put

$$A(y) = \left(\frac{\sin \pi y}{\pi}\right)^2, \qquad S(y) = \sum_{m=1}^{\infty} (y+m)^{-2}.$$

Subtracting (126) from (123) gives exactly

$$H_B(y) - 1 = A(y)\left\{\frac{2}{y} - \frac{1}{y^2} - 2S(y)\right\}. \tag{127}$$

The decreasing positive function $s \mapsto (y+s)^{-2}$ satisfies

$$S(y) \le \int_0^\infty (y+s)^{-2}\,ds = \frac{1}{y} \le \frac{1}{y^2} + S(y). \tag{128}$$

Consequently the expression in braces in (127) lies between $-y^{-2}$ and y^{-2}. Hence

$$|H_B(y) - 1| \le W(y) \quad (y > 0).$$

Positive integer points follow directly from (124). Oddness of H_B and evenness of W, together with their values at zero, now prove

$$|H_B(x) - \operatorname{sgn}(x)| \le W(x) \qquad (x \in \mathbb{R}). \tag{129}$$

This is the required sign-function majorization, with its normalization proved rather than assumed.

E.1.2 Recovering the exact Fourier kernel

Elementary integration gives, for every real x,

$$W(x) = 2\int_0^1 (1-t)\cos(2\pi t x)\,dt. \tag{130}$$

For $0 < r < 1$, insert (130) into the absolutely convergent Abel-weighted series. Termwise integration is legitimate because $\sum_{m\ge1} r^m < \infty$. The cosine subtraction formula gives

$$\begin{aligned}\sum_{m\ge1} r^m\{W(x-m) - W(x+m)\} &= 4\int_0^1 (1-t)\sin(2\pi t x)\sum_{m\ge1} r^m \sin(2\pi m t)\,dt \\ &= 4\int_0^1 (1-t)\sin(2\pi t x)\frac{r\sin(2\pi t)}{1 - 2r\cos(2\pi t) + r^2}\,dt.\end{aligned} \tag{131}$$

For $0 < t < 1$,

$$\left|\frac{r\sin(2\pi t)}{1 - 2r\cos(2\pi t) + r^2}\right| \le \frac{1}{2}|\cot(\pi t)|, \tag{132}$$

since the denominator is $(1-r)^2 + 4r\sin^2(\pi t)$. The right side of (131), including its remaining factors, is therefore dominated by

$$2(1-t)|\sin(2\pi t x)|\,|\cot(\pi t)|.$$

This is integrable at both endpoints: the sine is $O_x(t)$ at zero, and $(1-t)\cot(\pi t)$ is bounded at one. Dominated convergence on the integral side and absolute convergence on the series side allow $r \uparrow 1$. The fraction in (131) tends to $\frac{1}{2}\cot(\pi t)$. Also, direct integration gives

$$2xW(x) = \frac{2}{\pi}\int_0^1 \sin(2\pi t x)\,dt, \tag{133}$$

including $x = 0$. Combining (123) and (131)–(133),

$$H_B(x) = 2\int_0^1 R(t)\sin(2\pi t x)\,dt, \qquad R(t) = (1-t)\cot(\pi t) + \frac{1}{\pi}. \tag{134}$$

Every integral in (134) is a convergent real integral. No principal-value Fourier manipulation has been used.

Now take exactly the manuscript's kernel

$$K(t) = \frac{1-t}{2} + \frac{i}{2}R(t), \qquad 0 < t < 1. \tag{135}$$

Equations (130) and (134) establish the exact identity

$$\boxed{M(x) := \frac{1}{2} + 2\lim_{\epsilon\downarrow 0} \operatorname{Re} \int_\epsilon^1 K(t) e^{2\pi i t x}\, dt = \frac{1 - H_B(x) + W(x)}{2} = \frac{1 + B(-x)}{2}.} \tag{136}$$

In particular, the exponent has the *positive* sign and the imaginary part of K has the *positive* sign. The minus sign on H_B comes from taking the real part of $iR(t)e^{2\pi i t x}$.

E.1.3 Closed boundary, cardinal values and the minorant

For $x < 0$, (129) gives $M(x) \geq 1$; for $x > 0$, it gives $M(x) \geq 0$. At the boundary, (124) and $W(0) = 1$ give

$$M(0) = 1. \tag{137}$$

Thus this majorizes the *closed* half-line, including a point mass at zero, rather than merely the half-height convention for a jump:

$$M(x) \geq 1_{(-\infty,0]}(x), \qquad 1 - M(-x) \leq 1_{(-\infty,0)}(x). \tag{138}$$

The second inequality follows by subtracting the first inequality at $-x$ from one; its endpoint is open. Direct cardinal values are

$$M(j) = 1_{\{j\leq 0\}}, \qquad 1 - M(-j) = 1_{\{j<0\}} \quad (j \in \mathbb{Z}). \tag{139}$$

The proof also gives the useful global error bound

$$0 \leq M(x) - 1_{(-\infty,0]}(x) \leq W(x) \quad (x \in \mathbb{R}). \tag{140}$$

At zero the left-hand error is exactly zero; it is not obtained by assigning half of an atom to each side. In particular $0 \leq M \leq 2$.

E.1.4 Expectations and the compensated identity

Suppose U is real and $\mathbb{E}|U| < \infty$, and use the convention $\psi_U(t) = \mathbb{E}e^{2\pi i t U}$. Since $2\int_0^1 \operatorname{Re} K(t)\, dt = 1/2$, (136) can be written

$$M(u) = 1 + 2\operatorname{Re} \int_0^1 K(t)\{e^{2\pi i t u} - 1\}\, dt. \tag{141}$$

The complex integral in (141) is absolutely convergent. Indeed, $K(t) = i/(2\pi t) + 1/2 + O(t)$ at zero, K is bounded near one, and

$$\mathbb{E}|e^{2\pi i t U} - 1| \leq 2\pi t\, \mathbb{E}|U|.$$

Consequently $\int_0^1 |K(t)|\mathbb{E}|e^{2\pi i t U} - 1|\, dt < \infty$, so ordinary absolute Fubini applies to (141). Restoring the constant real part gives

$$\mathbb{E}M(U) = \frac{1}{2} + 2\lim_{\epsilon\downarrow 0} \operatorname{Re} \int_{\epsilon}^{1} K(t)\psi_U(t)\,dt. \tag{142}$$

Since $\psi_{-U} = \overline{\psi_U}$, conjugation similarly yields

$$\mathbb{E}\{1 - M(-U)\} = \frac{1}{2} - 2\lim_{\epsilon\downarrow 0} \operatorname{Re} \int_{\epsilon}^{1} \overline{K(t)}\psi_U(t)\,dt. \tag{143}$$

These are the two expectation identities under the real-part improper convention (122). All expectations are finite; (140) even gives boundedness of the majorant without a moment assumption. The first absolute moment is a sufficient hypothesis for the stated compensated Fourier/Fubini argument. The uncompensated complex integral itself need not converge.

E.1.5 Four-summand smoothing normalization

Let Y be integrable, let $h \in \mathbb{R}$, and let $T > 0$. For $\chi_Y(s) = \mathbb{E}e^{isY}$, apply (138) and (142) to $U = T(Y - h)/(2\pi)$. The resulting exact normalization is

$$\mathbb{P}(Y \le h) \le \frac{1}{2} + 2\lim_{\epsilon\downarrow 0} \operatorname{Re} \int_{\epsilon}^{1} K(t)e^{-iTth}\chi_Y(Tt)\,dt. \tag{144}$$

For the four-summand application, $Z = (X_1 + \cdots + X_4)/2$, $\varphi_P(s) = \mathbb{E}e^{isX_1}$, and iid summands, choose $T = 6$. Independence gives $\chi_Z(6t) = \varphi_P(3t)^4$. Thus (144) is exactly

$$F_Z(x) \le \frac{1}{2} + \int_0^1 \operatorname{Re}\{\widehat{K}(t)e^{-6itx}\varphi_P(3t)^4\}\,dt, \qquad \widehat{K} = 2K, \tag{145}$$

the physical four-summand smoothing normalization. Writing $D = x - Z$ and $S(t) = tR(t)$, its real integrand is exactly

$$(1 - t)\mathbb{E}\cos(6Dt) + S(t)\mathbb{E}[6D\operatorname{sinc}(6Dt)], \qquad \operatorname{sinc}(v) = \sin(v)/v, \tag{146}$$

with the continuous value at zero. This verifies both the factor six and the sign of the sine term. No non-atomicity, finite-support, density, or continuity-at-the-threshold assumption is needed. For (145), iid independence and integrability are sufficient; the manuscript's bounded support assumptions imply integrability.

F An original-law reduction for four summands

This appendix proves a conditional analytic implication for the finite base. It does not supply a complete four-summand certificate. The smoothing inequality used below is proved in Appendix E. The remaining steps are proved for the original iid marginal, including laws with infinite support inside a bounded interval and atoms at zero or at a threshold. No optimizer or contact assumption is used.

Let X have mean zero, variance one, $|X| \le R = 17/4$, and $\rho = \mathbb{E}|X|^3 < 46/25$. Set

$$u = \mathbb{E}X_+ = \mathbb{E}X_- > 0, \quad v_\pm = \mathbb{E}X_\pm^2, \quad a = v_-/u, \quad b = v_+/u,$$

$$w = u/a + u/b, \qquad \vartheta = v_+ - v_-.$$

Both $v_\pm$ and a, b are positive; also $a, b \le R$. Signwise Cauchy gives $u/a \le \mathbb{P}(X < 0)$ and $u/b \le \mathbb{P}(X > 0)$, so $0 < w \le 1$. The canonical probability

$$Q = (u/a)\delta_{-a} + (1 - w)\delta_0 + (u/b)\delta_b$$

has the same first and second moments on each strict half-line as P. Put $r = \rho(Q)$ and $\eta = \rho - r$.

F.1 One actual measure, including all contact atoms

For $y = |X|$, let $c = b$ on the positive tag and $c = a$ on the negative tag, and put $W = y(y-c)^2$ (zero at $X = 0$). Direct expansion gives

$$\mathbb{E}W = \eta \geq 0, \tag{147}$$

$$\mathbb{E}X^3 = b - a + \mathbb{E}[\mathrm{sgn}(X)W],$$

$$\mathbb{E}X^4 = a^2 - ab + b^2 + \mathbb{E}[(y + 2c)W]. \tag{148}$$

Indeed $y^3 = W + 2cy^2 - c^2y$ and $y^4 = (y + 2c)W + 3c^2y^2 - 2c^3y$; integrate on each tag using $\mathbb{E}[y; \mathrm{tag}] = u$ and $\mathbb{E}[y^2; \mathrm{tag}] = uc$. If $\eta = 0$, then P is supported on $\{-a, 0, b\}$ and its first two moments force $P = Q$. This case requires no defect probability.

For $\eta > 0$, define the single tagged probability $\nu = W\,dP/\eta$. It charges neither zero nor either contact. On its domain set

$$k_0 = (c^2y)^{-1}, \qquad k_2 = (y - c)^{-2}, \qquad k_1 = (y - c)^{-1}.$$

Then, writing $p_0 = 1 - w$,

$$\eta \int k_0\,d\nu = p_0 - P\{0\},$$

$$\eta \int_\sigma k_2\,d\nu = u - c_\sigma P\{\sigma c_\sigma\}, \qquad \int_\sigma k_1\,d\nu = 0. \tag{149}$$

To check the first formula, $Wk_0 = (y - c)^2/c^2$ off zero and contacts; its integral on a tag is $P(\mathrm{tag}) - u/c$. For k_2, cancellation gives y only away from the contact, so the omitted contact mass must be subtracted. For k_1 it gives $y(y - c)$, whose integral is zero even after omitting the contact. All integrals exist; for k_1, Cauchy uses the finite k_2 integral.

Let $\psi(q) = \mathbb{E}e^{\mathrm{i}qX}$ and $g(q) = \int e^{\mathrm{i}qx}\,dQ(x)$. Hermite interpolation of $f(y) = e^{\mathrm{i}\sigma qy}$ at $0, c, c$ gives

$$f(y) - L_f(y) = y(y - c)^2 r_q(\sigma, y), \qquad r_q = f[0, c, c, y].$$

The interpolant is quadratic with constant $f(0) = 1$. Its linear and quadratic integrals match on each tag; its constants match only after both tags and the zero atoms are combined. Consequently

$$\psi(q) - g(q) = \eta \int r_q\,d\nu. \tag{150}$$

This identity uses the same measure at all frequencies. The integral form of a third divided difference gives $|r_q| \leq |q|^3/6$. For completeness another bound is $|r_q| \leq q^2/y$: write

$$L_f(y) = f(c) + f'(c)(y - c) + \frac{f(0) - f(c) + cf'(c)}{c^2}(y - c)^2.$$

The quadratic coefficient and the first-order Taylor remainder at c are bounded by $q^2/2$ and $q^2(y - c)^2/2$, respectively. Division gives the claim. Thus the following three radii are valid:

$$|\psi(q) - g(q)| \leq \min\{\eta|q|^3/6,\ \max(a, b)^2 p_0 q^2,\ 1 + |g(q)|\}. \tag{151}$$

The mass radius follows from $1/y = c^2k_0$ and (149).

For arbitrary finite coefficients κ_j, define $A = \Re \sum_j 4\kappa_j g(q_j)^3 r_{q_j}$. If fixed prices $\lambda_0, \lambda_+, \lambda_- \geq 0$ and signed $\theta_\pm$ satisfy

$$A - \lambda_0 k_0 - \lambda_\sigma k_2 - \theta_\sigma k_1 \leq S$$

on the entire tagged domain, then

$$\Re \sum_j 4\kappa_j g(q_j)^3(\psi(q_j) - g(q_j)) \le p_0\lambda_0 + u(\lambda_+ + \lambda_-) + S\eta =: \Gamma(\eta). \tag{152}$$

This holds for each actual P, not for a purported sufficiency characterization of all measures satisfying the budgets. Enlarging that class in an upper-bound problem is safe. Price choices must precede the supremum; signed equality prices and a negative S are retained. At $\eta = 0$ the response is zero and the displayed right side is nonnegative, so the same bound applies directly.

F.2 The physical smoothing and its two reserves

Let $Z = (X_1 + \cdots + X_4)/2$ with independent copies of this same law. The input Prawitz inequality is, for every real x,

$$\mathbb{P}(Z \le x) \le J_P(x) := \frac{1}{2} + \int_0^1 \Re\{\widehat{K}(t)e^{-6itx}\psi(3t)^4\}\,dt, \quad \widehat{K}(t) = 1 - t + \mathrm{i}[(1-t)\cot(\pi t) + 1/\pi]. \tag{153}$$

Only the real integrand is integrated. Put $S(t) = t\Im\widehat{K}(t)$. Its endpoint-continuous extension satisfies

$$0 \le S \le 1/3, \qquad |S'| < 2, \qquad |S''| < 5. \tag{154}$$

Here are explicit sufficient bounds. The cotangent partial fractions give

$$S = \pi^{-1} - 2\pi^{-1}(A + MG), \quad A = t^2/(1+t),\ M = t^2(1-t),\ G = \sum_{j\ge 2}(j^2 - t^2)^{-1}.$$

On $[0,1]$, $|A'| \le 3/4$, $|A''| \le 2$, $|M| \le 1/6$, $|M'| \le 1$, $|M''| \le 4$, and $G \le 3/4$, $|G'| \le 10/27$, $|G''| \le 106/135$. The first sum telescopes; the latter two follow from $j^2 - t^2 \ge 3j^2/4$ and $\sum_{j\ge 2} j^{-k} \le 2^{-k} + \int_2^\infty s^{-k}\,ds$. The derivative series converge uniformly. With $\pi > 3$, these imply $|S'| < 19/18$, $|S''| < 223/54$. Nonnegativity follows directly from $(1-t)\cot(\pi(1-t)) \le 1/\pi$ when $t > 1/2$, and from the nonnegative cotangent when $t \le 1/2$. The positive partial-fraction terms give $S \le 1/\pi < 1/3$.

Set $D = x - Z$, $A_x(t) = \mathbb{E}\cos(6Dt)$ and $L_x(t) = \mathbb{E}[6D\,\mathrm{sinc}(6Dt)]$. The regular real integrand is $(1-t)A_x(t) + S(t)L_x(t)$. Since $\mathrm{sinc}(v) = \int_0^1 \cos(sv)\,ds$,

$$|A_x'| \le 6m_1,\ |A_x''| \le 36m_2, \quad |L_x| \le 6m_1,\ |L_x'| \le 18m_2,\ |L_x''| \le 72m_3, \quad m_j = \mathbb{E}|D|^j.$$

Differentiation under these bounded-moment integrals and the product rule therefore give

$$|F_x''(t)| \le 42m_1 + 108m_2 + 24m_3. \tag{155}$$

All four copies are used to obtain

$$\mathbb{E}D^2 = x^2 + 1 =: D_2, \qquad \mathbb{E}D^4 = x^4 + 6x^2 - 2x\mathbb{E}X^3 + (\mathbb{E}X^4 + 9)/4.$$

Thus for valid whole-cell moment bounds $\kappa_* \ge |\mathbb{E}X^3|$ and $M_* \ge \mathbb{E}X^4$, put $D_4 = x^4 + 6x^2 + 2|x|\kappa_* + (M_* + 9)/4$. Then $m_1 \le \sqrt{D_2}$ and $m_3 \le \sqrt{D_2 D_4}$. The bounds (148) supply these without an additional fourth-moment premise. On $|x| \le 2$, $m_2 \le 5$, $\mathbb{E}D^4 < 54$, $m_1 < 9/4$, $m_3 < 33/2$, so (155) is less than $2061/2$.

Choose $N \ge 1$, $t_j = (j + 1/2)/N$, $q_j = 3t_j$ and $\kappa_j(H) = \widehat{K}(t_j)e^{-\mathrm{i}q_j H}/N$. Composite midpoint quadrature has error at most $\sup|F''|/(24N^2)$. Consequently the raw numerator $2(J_P(H/2) - \Phi(H/2))$ has the single reserve

$$Q_H = \min\left\{\frac{687}{8N^2}, \frac{42\sqrt{D_2} + 108D_2 + 24\sqrt{D_2 D_4}}{12N^2}\right\}. \tag{156}$$

This quadrature is for the original law once, before reference expansion. Neither the reference nor its response/remainder needs another quadrature.

To control threshold joins, suppose $a_j \geq |g(q_j)|$ and e_j bounds $|\psi(q_j) - g(q_j)|$ uniformly on the entire excess cell. The actual characteristic function is 1-Lipschitz, since $\mathbb{E}|X| \leq 1$. Hence

$$P_j = \min(1, a_j + e_j + 3/(2N))$$

bounds $|\psi(3t)|$ throughout its frequency cell. Define d_j as the smaller of the two integrals over that cell of

$$t^2(1-t) + t/3, \qquad t^2(1-t) + t(1-t)/2.$$

Both dominate $t^2|\widehat{K}(t)|$: use $|\widehat{K}| \leq 1 - t + \Im\widehat{K}$ and, for the second, $S \leq (1-t)(1+2t)/(\pi(1+t)) \leq (1-t)/2$. Their full integrals are $1/4$ and $1/6$. Thus

$$|J_P''(x)| \leq B := 36 \sum_j d_j P_j^4.$$

For the smooth physical function $G_P(H) = 2(J_P(H/2) - \Phi(H/2))$, $|G_P''| \leq (B + 1/4)/2$, since $\sup|\Phi''| < 1/4$. On any raw-threshold interval of length h, its chord lies above G_P up to the single reserve

$$E_H = h^2(B + 1/4)/16. \tag{157}$$

There is no differentiation or interpolation of the CDF. The moment reserve (156) may be computed separately at each node; the curvature reserve is common to the whole geometry/excess cell.

F.3 A precise conditional certificate implication

Theorem F.1 (Low-excess original-law bridge)**.** *Fix a geometry cell, an excess interval $[L, U] \subset [0, \infty)$, and raw nodes $-4 \leq H_0 < \cdots < H_m \leq 25/8$ of spacing at most h. Assume all the following bounds are valid for every actual standardized P in this cell, with $\rho = r + \eta < 46/25$ and $\eta \in [L, U]$. At each node let $M_H \geq C_H(Q) - Tr$, where*

$$C_H(Q) = 1 + 2\Re \sum_j \kappa_j(H) g(q_j)^4 - 2\Phi(H/2).$$

Let $\Gamma_H(\eta)$ be a whole-domain price bound (152)*, and let $R_{2,H}(\eta)$ bound*

$$\sum_j |\kappa_j(H)||\psi(q_j)^4 - g(q_j)^4 - 4g(q_j)^3(\psi(q_j) - g(q_j))|.$$

Suppose the complete row

$$P_H(\eta) = M_H + Q_H + E_H + 2\Gamma_H(\eta) + 2R_{2,H}(\eta) - T\eta$$

is a convex quadratic, nonpositive at L and U, at every node. Then $2\{\mathbb{P}(X_1 + \cdots + X_4 \leq H) - \Phi(H/2)\} \leq T\rho$ for every $H \in [H_0, H_m]$ and every such law, including $\eta = 0$.

Proof. Expand the actual fourth power at each frequency. The linear part is controlled by the one common measure, and the remaining complex part by its full nonnegative modulus bound. Midpoint quadrature yields $G_P(H) - T\rho \leq P_H(\eta) - E_H$ at each node. Convexity gives $P_H(\eta) \leq 0$ on $[L, U]$. The chord inequality (157) now implies $G_P(H) - T\rho \leq 0$ between nodes. Finally use (153). All these inequalities hold for one arbitrary law throughout; no law is reconstructed from a relaxation. □

An elementary permitted remainder is obtained from

$$\psi^4 - g^4 - 4g^3(\psi - g) = (\psi - g)^2(\psi^2 + 2\psi g + 3g^2).$$

If $|g| \le a \le 1$ and $|\psi - g| \le A + B\eta$ nonnegatively on the whole cell, freeze $p = \min(1, a + \max(A + BL, A + BU))$. Then $(A + B\eta)^2(p^2 + 2pa + 3a^2)$ is a convex quadratic remainder. Other complete bounds, including a proven disk/lens bound, may replace it. Their correctness remains a premise if not proved separately. Selecting a whole valid row independently at each node is safe; interpolating a pointwise minimum of candidate curves is not needed.

For $H \le -4$, Cantelli gives $F_Z(H/2) \le 1/5$. For $H \ge 25/8$, the elementary normal bound $\Phi(25/16) > 4/5$ gives the same raw numerator bound $2/5$. Since $\rho \ge 1$, both positive-score tails are at most $2/5 < T = 4097/10000$. To verify that normal bound directly, $\pi < 22/7$ gives $\phi(0) > 997/2500$ by squaring. With $x = 25/16$, the exponential Taylor lower bound gives

$$\Phi(x) \ge \frac{1}{2} + \frac{997}{2500}\left(x - \frac{x^3}{6} + \frac{x^5}{40} - \frac{x^7}{336}\right) > \frac{4}{5}.$$

F.4 The independent count route and reflection

For this separate count route assign zero to the nonnegative class: let $p = P(X \ge 0)$, $q = 1 - p$. This differs from the strict-positive convention of the main argument when there is a zero atom. Both are positive. Conditioning on the number j of nonnegative observations gives mass $\binom{4}{j}p^j q^{4-j}$. Conditional on any fixed sign pattern, independence retains independent conditional marginals. Thus Z has conditional mean and variance

$$\mu_j = \frac{u(j - 4p)}{2pq}, \qquad V_j = \frac{1}{4}\left[j\left(\frac{v_+}{p} - \frac{u^2}{p^2}\right) + (4 - j)\left(\frac{v_-}{q} - \frac{u^2}{q^2}\right)\right].$$

The same moments hold after mixing patterns with that count. Closed-event Cantelli yields the source bound $A_j(x) = 1$ for $x \ge \mu_j$ and $A_j(x) = V_j/(V_j + (\mu_j - x)^2)$ otherwise. This includes $V_j = 0$. Cauchy gives $w(1 - \vartheta)/2 \le p \le 1 - w(1 + \vartheta)/2$. Taking upper bounds on all five A_j on a cell and then a valid upper bound of their single binomial mixture preserves the original-law bound. No arbitrary dependent sign allocation can replace the iid weights.

Any complete count-based above-cut bound, or independent all-law large-ρ bound, can join the low-excess theorem when both closed ranges cover their common cutoff. A parent geometry may be replaced by all four closed children, each with its own complete low/high cover. No unexecuted terminal inequality is supplied by this observation.

For the reflected law $\widetilde{P} = \mathcal{L}(-X)$, a, b are swapped, ϑ changes sign, η and ρ are unchanged, and

$$\widetilde{p} = P(X \le 0) = 1 - p + P\{0\}, \qquad \widetilde{\psi}(q) = \overline{\psi(q)}.$$

The raw phase e^{-iqH} is retained when evaluating its own threshold H. Applying both positive closed bounds covers the negative CDF discrepancy because

$$\Phi(x) - F_Z(x) \le \Phi(x) - F_Z(x-) = F_{-Z}(-x) - \Phi(-x).$$

Equality at an atom is not assumed. These facts supply the analytic coverage implication only. Complete numerical enclosures, their selected implementation, and the unrestricted-owner localization remain separate.

References

[1] BerryEsseenLean: Berry–Esseen eventual sharpness in Lean. Lean 4 formalization accompanying the manuscript of Hengzhi He and Guang Cheng, 2026. URL https://github.com/UCLA-Trustworthy-AI-Lab/BerryEsseenLean/tree/2b97fa7400dd85fc98b829224c89b0f986789395. Commit 2b97fa7400dd85fc98b829224c89b0f986789395; designated endpoints retain explicit external mathematical inputs; Lean v4.28.0.

[2] Logan Bell. Extremizers for the independent Berry–Esseen inequality. Manuscript dated 2026-10-06; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/extremizers/extremizers.pdf.

[3] Logan Bell. Berry–Esseen bounds for two summands and Bernoulli arrays. Manuscript dated 2026-10-06; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/finite-arrays/finite-arrays.pdf.

[4] Logan Bell. An upper bound on the Berry–Esseen constant for independent summands. Manuscript dated 2026-10-06; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/independent-berry-esseen.pdf.

[5] Logan Bell. The sharp Berry–Esseen bound for independent arrays with small Lyapunov ratio. Manuscript dated 2026-10-06; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/small-lyapunov/small-lyapunov.pdf.

[6] Logan Bell. On the structural conjecture for extremizers of the independent Berry–Esseen inequality. Manuscript dated 2026-10-18; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/structural-conjecture/structural-conjecture.pdf.

[7] Logan Bell. Zero-bias transport and extremal Bernoulli arrays. Manuscript dated 2026-10-06; collection version 0.7.0; source snapshot f61bc638774bfa1165065462f98fe31efade30bf, 2026. URL https://github.com/epicycloids/independent-berry-esseen-bounds/blob/f61bc638774bfa1165065462f98fe31efade30bf/paper/transport/transport.pdf.

[8] Andrew C. Berry. The accuracy of the Gaussian approximation to the sum of independent variates. *Transactions of the American Mathematical Society*, 49(1):122–136, 1941. doi: 10.1090/S0002-9947-1941-0003498-3.

[9] Carl-Gustav Esseen. On the Liapunoff limit of error in the theory of probability. *Arkiv för Matematik, Astronomi och Fysik*, 28A(9):1–19, 1942.

[10] Carl-Gustav Esseen. Fourier analysis of distribution functions. A mathematical study of the Laplace–Gaussian law. *Acta Mathematica*, 77:1–125, 1945. doi: 10.1007/BF02392223.

[11] Carl-Gustav Esseen. A moment inequality with an application to the central limit theorem. *Scandinavian Actuarial Journal*, 1956(2):160–170, 1956. doi: 10.1080/03461238.1956.10414946.

[12] Larry Goldstein and Gesine Reinert. Stein's method and the zero bias transformation with application to simple random sampling. *The Annals of Applied Probability*, 7(4):935–952, 1997. doi: 10.1214/aoap/1043862419.

[13] Hengzhi He and Guang Cheng. The Berry–Esseen constant conjecture is eventually true, 2026. URL https://arxiv.org/abs/2609.06358v3. Version 3, revised 30 September 2026.

[14] Christian Hipp and Lutz Mattner. On the normal approximation to symmetric binomial distributions. *Theory of Probability and its Applications*, 52(3):516–523, 2008. doi: 10.1137/S0040585X97983213. Original edition: Teoriya Veroyatnostei i ee Primeneniya 52 (2007), 610–617.

[15] Victor Yu. Korolev and Irina G. Shevtsova. On the upper bound for the absolute constant in the Berry–Esseen inequality. *Theory of Probability and its Applications*, 54(4):638–658, 2010. doi: 10.1137/S0040585X97984449. English translation of the Russian article published in 2009.

[16] Yongxi Lin. Berry–Esseen bounds in Lean. GitHub repository, 2026. URL https://github.com/CoolRmal/BerryEsseen. Accompanying proof document: The i.i.d. Berry–Esseen constant is at most 0.423: a machine-checked proof, 13 September 2026, paper/upper-0423.pdf. Commit bfb22608ad18b4fa9fda8dd9a1ed2c2b6fbc29e7; accessed 27 October 2026.

[17] Håkan Prawitz. Limits for a distribution, if the characteristic function is given in a finite domain. *Scandinavian Actuarial Journal*, 1972(2):138–154, 1972. doi: 10.1080/03461238.1972.10404645.

[18] Kacper Rodziewicz and Bartosz Kołodziejek. Moment defects and sharp iid Berry–Esseen bounds. Hexagon preprint hexagon:2610.00147v3, 2026. URL https://hexagonmath.org/2610.00147v3. Working draft, version 3, 17 October 2026; accessed 27 October 2026.

[19] Jona Schulz. *The optimal Berry–Esseen constant in the binomial case*. PhD thesis, Universität Trier, 2016. URL https://ubt.opus.hbz-nrw.de/frontdoor/index/index/docId/732.

[20] Irina Shevtsova. On the absolute constants in the Berry–Esseen type inequalities for identically distributed summands, 2011. URL https://arxiv.org/abs/1111.6554v1.

[21] Irina Shevtsova. On the accuracy of the approximation of the complex exponent by the first terms of its Taylor expansion with applications. *Journal of Mathematical Analysis and Applications*, 418(1):185–210, 2014. doi: 10.1016/j.jmaa.2014.03.075. URL https://arxiv.org/abs/1301.2783v3.

[22] Irina G. Shevtsova. Sharpening of the upper bound of the absolute constant in the Berry–Esseen inequality. *Theory of Probability and its Applications*, 51(3):549–553, 2007. doi: 10.1137/S0040585X97982591. English translation of the Russian article published in 2006.

[23] Irina G. Shevtsova. On the absolute constants in the Berry–Esseen inequality and its structural and nonuniform improvements. *Informatics and its Applications*, 7(1):124–125, 2013. URL https://www.mathnet.ru/eng/ia252. In Russian.

[24] I. S. Shiganov. Refinement of the upper bound of the constant in the central limit theorem. *Journal of Soviet Mathematics*, 35:2545–2550, 1986. doi: 10.1007/BF01121471. English translation of the Russian original, 1982, pp. 109–115.

[25] Terence Tao and contributors. The Berry–Esseen constant. Optimization Constants in Mathematics, Problem 19a, 2026. URL https://github.com/teorth/optimizationproblems/blob/e95af44bc6ae938f545a0fc62cd1b6c1f6088a56/constants/19a.md. Repository snapshot e95af44bc6ae938f545a0fc62cd1b6c1f6088a56; accessed 27 October 2026.

[26] I. S. Tyurin. Refinement of the upper bounds of the constants in Lyapunov's theorem. *Russian Mathematical Surveys*, 65(3):586–588, 2010. doi: 10.1070/RM2010v065n03ABEH004688.

[27] Ilya Tyurin. New estimates of the convergence rate in the Lyapunov theorem, 2009. URL https://arxiv.org/abs/0912.0726v1.

[28] Paul van Beek. An application of Fourier methods to the problem of sharpening the Berry–Esseen inequality. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 23: 187–196, 1972. doi: 10.1007/BF00536558.

[29] Haonan Xiao and Chunlin Li. A 0.4395 upper bound for the i.i.d. Berry–Esseen constant. GitHub repository, 2026. URL https://github.com/haonan-xiao/iid-berry-esseen. Accompanying paper: paper/berry-esseen-04395.pdf. Commit f00d781c8c42908bf4aef2739129442a0919fa38; accessed 27 October 2026.

[30] Asher Yan. ProbabilityApproximation: Berry–Esseen bounds for independent sums. Lean 4 software, release v0.9.6, 2026. URL https://github.com/Polarnova/ProbabilityApproximation/releases/tag/v0.9.6. Commit c89fba908a33ba741836407f43c67609e2d05279; Lean and Mathlib v4.32.0. Cite the original mathematical sources for theorems used.

[31] Vladimir M. Zolotarev. An absolute estimate of the remainder term in the central limit theorem. *Theory of Probability and its Applications*, 11(1):95–105, 1966. doi: 10.1137/1111005.

[32] Vladimir M. Zolotarev. A sharpening of the inequality of Berry–Esseen. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 8:332–342, 1967. doi: 10.1007/BF00531598.