# One acquired strip and two noise estimates

29 October 2026

The new comparison has one probability interval and one support strip. Below noise 1/100, the loss at a single count dominates the entire incoming tail by fourth moments. Above that point, five neighboring counts retain enough crossing loss, and four rational endpoint checks finish the argument. The acquisition reaches this common strip directly from the coarse defect budget and two contact rays. This note gives the exact joins and the remaining conditional boundary. No claim about arbitrary laws follows from a strip theorem without the acquisition step.

For a centered variance-one marginal $P$ and its independent copies, write

$$\mathcal{E}(P,n,H)=\sqrt{n}\{\mathbb{P}(X_1+\cdots+X_n\le H)-\Phi(H/\sqrt{n})\}-C_{\mathrm{E}}\mathbb{E}|X|^3,\qquad C_{\mathrm{E}}=\frac{3+\sqrt{10}}{6\sqrt{2\pi}}.$$

Here $H$ is a raw sum threshold. In particular its normal argument is $H/\sqrt{n}$, not $H$.

**Theorem 1** (Complete local comparison)**.** *Let $n\ge 700$ be an integer and $Y=B+U$, where $B$ is Bernoulli of parameter $p\in[3/8,93/200]$ and $\mathbb{E}(U\mid B)=0$. Suppose the conditional noises obey*

$$-3/5\le U_0\le a_0,\qquad -a_1\le U_1\le 3/5,\qquad a_0,a_1\ge 0,\quad a_0+a_1\le 1/250.$$

*Put $v=p(1-p)$, $s=\mathbb{E}U^2$, $\lambda=ns$, and $X=(Y-p)/\sqrt{v+s}$. If $0<\lambda\le 7/20$ and $|H|\le 46/25$, then $\mathcal{E}(P,n,H)<-\lambda/100$. At $\lambda=0$ one has $\mathcal{E}(P,n,H)<0$.*

*Proof.* For $0<\lambda\le 1/100$, apply Theorem 1 of `small-common.tex`. Choose an integer $k$ so that $np+H\sqrt{v+s}=k+u$, with $u\in[-3/10,7/10]$. The interval has length one, and the proof checks that the anchor is interior. At $C=C_{\mathrm{E}}\ge 0$, that theorem bounds the difference between $\mathcal{E}$ and the signed binomial excess at $k$ by $-\lambda/100$. Schulz's strict finite-binomial theorem (Theorem 1, 2016 thesis) bounds the latter excess by zero for every $0<p<1$ and finite $n$.

For $1/100\le\lambda\le 7/20$, use the Compact row of Theorem 1 in `unified-comparison.tex`. All probability, support, inward and threshold hypotheses agree literally. Its floor anchor need not equal the small-noise anchor: both conclusions concern the same $\mathcal{E}(P,n,H)$. The shared noise endpoint is included in both statements; there is no phase or noise gap. At zero noise, $U=0$ almost surely, so the same strict binomial theorem applies directly. A limit of positive-noise margins is not used. □

**Theorem 2** (Conditional acquired-witness exclusion)**.** *Suppose $n\ge 700$ and $(P,H)$ is a compact, centered variance-one attained fixed-target witness at $C_{\mathrm{E}}$, with $\mathcal{E}(P,n,H)\ge 0$ and true sharp predecessor $C_{n-1}\le C_{\mathrm{E}}$. Assume the retained closed-support contact and all-threshold heat bounds hold for this witness. Suppose the complete coarse entry theorem supplies, for this same oriented law,*

$$nD\le 32.20665378,\qquad |H|\le 1.83605680,\quad D=(3+\sqrt{10})\rho-\gamma-3J,$$

*where $\rho=\mathbb{E}|X|^3$, $\gamma=\mathbb{E}X^3$ and $J$ is its actual weighted support-gap sum. Then such a witness cannot exist.*

*Proof.* The scalar theorem in `acquisition-direct.tex` first establishes an actual gap across zero, with $p = \mathbb{P}(X > 0) \in (3/8, 93/200)$. Write $m_j = \mathbb{E}(X \mid B = j)$ and $\ell = m_1 - m_0 > 0$, where $B = \mathbf{1}_{X>0}$. Centering gives $m_0 = -p\ell$ and $m_1 = (1-p)\ell$. With $U = (X - m_B)/\ell$ and $\theta = \mathbb{E}(X - m_B)^2$, total variance gives

$$1 = v\ell^2 + \theta, \qquad s = \mathbb{E}U^2 = \theta/\ell^2, \qquad v + s = \ell^{-2}, \qquad X = \frac{B + U - p}{\sqrt{v+s}}.$$

Thus the normalized marginal in Theorem 1 is exactly the original law, with $\lambda = n\theta/\ell^2 < 7/20$ and inward sum $d < 1/250$. It is not an optimizer of a relaxation or a separately chosen product law. The two contact rays in the second acquisition theorem give the common amplitude $3/5$. The raw threshold obeys $|H| < 46/25$. These are every hypothesis of Theorem 1, which yields $\mathcal{E} < 0$ and the contradiction. Neither the old left/right probability panels nor an additional noise-localization table enters this implication. □

## The global boundary is unchanged

For any sum $S$,

$$\mathbb{P}(-S \le -H) - \Phi(-H/\sqrt{n}) = \Phi(H/\sqrt{n}) - \mathbb{P}(S < H) \ge \Phi(H/\sqrt{n}) - \mathbb{P}(S \le H).$$

The gap between the last two expressions is the atom at $H$. This selects the positive closed-event orientation; it does not transport the signed $D$ budget to a reflected law. Attainment and entry must apply afresh to the chosen oriented witness.

If a genuine base establishes $C_j \le C_{\mathrm{E}}$ through $j = 41$, and the retained attainment/contact/entry implications hold at every least failure, then any first failure of the sharp bound is in $42, \dots, 699$. Minimality supplies the predecessor needed in Theorem 2. This does not prove an unconditional cutoff at 700: an unresolved earlier failure invalidates that predecessor argument at later rows. A genuine base through 699, or an independent argument closing the remaining range, is still needed.

The coarse Fourier certificate, its two coarse and four signed entry passes, and their internal cover obligations are retained. The new proof removes only downstream probability-panel localization, old numerical noise tables, the separate left/right local union, the Poisson-tail comparisons and the 291-cell negative-inner certificate from this acquired $n \ge 700$ route. The old results at smaller indices and all their proof objects are preserved. Exact scalar checks and actual-law controls accompany these proofs; they do not independently establish the retained analytic inputs or the finite base.