

AN EXPONENTIAL IDENTITY IN TERMS OF PARTIAL DERIVATIVES

PETRO KOLOSOV

ABSTRACT. Your abstract here.

CONTENTS

1. Introduction and Main Results	1
2. Conclusions	3
References	3

1. INTRODUCTION AND MAIN RESULTS

This manuscript provides an exponential identity in terms of partial derivatives, extending the main idea explained in [Kol22] that gives polynomial identity in a form as follows

$$n^{2m+1} = \sum_{k=1}^n \sum_{r=0}^m \mathbf{A}_{m,r} k^r (n-k)^r, \quad (m, n) \in \mathbb{N}, \quad (1)$$

where $\mathbf{A}_{m,r}$ are real coefficients defined recursively, see [Kol16]. Define the function f such that based on the identity (1) with the only difference that values of n, m in the right part of (1) appear to be parameters of the function f . In contrast to the equation (1), upper bound n of the sum $\sum_{k=1}^n$ turned into function's parameter as well, so that f defined as follows

Definition 1.1. (*Polynomial function f .*)

$$f(x, y, z) = \sum_{k=1}^z \sum_{r=0}^y \mathbf{A}_{y,r} k^r (x-k)^r, \quad (2)$$

where definition space of (x, y, z) in (2) is $x \in \mathbb{R}$, $y \in \mathbb{N}$, $z \in \mathbb{R}$ so that the triple $(x, y, z) \in \mathbb{R} \times \mathbb{N} \times \mathbb{R}$. At first glance, equation (2) might look complex and not immediately understood, so in order to understand the function f and polynomials it produces clearly

Date: October 25, 2022.

2010 Mathematics Subject Classification. 32W50, 33B10.

Key words and phrases. Partial differential equations, PDE, Exponential function .

let's show few examples. Substituting the values of $y = 1, 2, 3$ to the function f we get the following polynomials

$$f(x, 1, z) = 3xz - 3z^2 + 3xz^2 - 2z^3$$

$$f(x, 2, z) = 5x^2z - 15xz^2 + 15x^2z^2 + 10z^3 - 30xz^3 + 10x^2z^3 + 15z^4 - 15xz^4 + 6z^5$$

$$f(x, 3, z) = -7xz + 14x^2z + 7z^2 - 42xz^2 + 35x^3z^2 + 28z^3 - 140x^2z^3 + 70x^3z^3 + 175xz^4 \\ - 210x^2z^4 + 35x^3z^4 - 70z^5 + 210xz^5 - 84x^2z^5 - 70z^6 + 70xz^6 - 20z^7$$

According to the main topic of the current manuscript, it provides an odd-exponential identity in terms of partial derivatives. Therefore, define the exponential function we work in context of. Exponential function g is a function of two variables defined as follows

Definition 1.2. (*Exponential function g .*)

$$g(x, y) = x^{2y+1}, \quad (x, y) \in \mathbb{R} \times \mathbb{N} \quad (3)$$

One more important thing is to conclude on partial derivative notation, more precisely the following notation for the partial derivative is used across the manuscript and remains unchanged

Notation 1.3. (*Partial derivative.*) Let be a function $f(x_1, x_2, \dots, x_n)$ defined over the real space \mathbb{R}^n . We denote partial derivative of the function f with respect to x_i , $1 \leq i \leq n$ as follows

$$f'_{x_i} = \lim_{\Delta x_i \rightarrow 0} \frac{f(x_1, x_2, \dots, x_i + \Delta x_i, \dots, x_n) - f(x_1, x_2, \dots, x_n)}{\Delta x_i}$$

Derivative of the function f with respect to x_i , $1 \leq i \leq n$ evaluated in point $(y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ is denoted as follows

$$f'_{x_i}(y_1, y_2, \dots, y_n)$$

Therefore, the following identity in terms of partial derivatives shows the relation between exponential function g and polynomial function f

Theorem 1.4. *Derivative of the odd-exponential function $g(x, y) = x^{2y+1}$ with respect to x evaluated at point $(u, v) \in \mathbb{R} \times \mathbb{N}$ equals to derivative of the function f with respect to x evaluated at point $(u, v, u) \in \mathbb{R} \times \mathbb{N} \times \mathbb{R}$ plus derivative of the function f with respect to z evaluated at point $(u, v, u) \in \mathbb{R} \times \mathbb{N} \times \mathbb{R}$*

$$g'_x(u, v) = f'_x(u, v, u) + f'_z(u, v, u) \quad (4)$$

Particularly, it follows that for every triple $(u, v, u) \in \mathbb{R} \times \mathbb{N} \times \mathbb{R}$ an identity holds

$$(2v + 1)u^{2v} = f'_x(u, v, u) + f'_z(u, v, u) \quad (5)$$

that is also an ordinary derivative of odd-power function $f(u) = u^{2v+1}$. To summarize and clarify all above, we provide a few examples that show an identity (4) in action.

2. CONCLUSIONS

Conclusions of your manuscript.

REFERENCES

- [Kol16] Petro Kolosov. On the link between Binomial Theorem and Discrete Convolution of Polynomials. *arXiv preprint arXiv:1603.02468*, 2016. <https://arxiv.org/abs/1603.02468>.
- [Kol22] Petro Kolosov. 106.37 An unusual identity for odd-powers. *The Mathematical Gazette*, 106(567):509–513, 2022. <https://doi.org/10.1017/mag.2022.129>.

Email address: kolosovp94@gmail.com

URL: <https://razumovsky.me/>