

# IDENTITIES IN ITERATED RASCAL TRIANGLES

PETRO KOLOSOV

ABSTRACT. In this manuscript we show new binomial identities in iterated rascal triangles. In particular, iterated rascal numbers are closely related to  $(1, q)$ -binomial coefficients. Finally, we state an open conjecture about the relation between iterated rascal numbers and  $(p, q)$ -binomial coefficients.

## CONTENTS

1. Introduction	2
2. Binomial identities in Iterated Rascal Triangles	3
3. Q-Binomial identities in Iterated Rascal Triangles	6
4. Conclusions	8
References	8

---

*Date:* July 1, 2024.

2010 *Mathematics Subject Classification.* 11B25, 11B99.

*Key words and phrases.* Pascal's triangle, Rascal triangle, Binomial coefficients, Binomial identities, Binomial theorem, Generalized Rascal triangles, Iterated rascal triangles, Iterated rascal numbers, Vandermonde identity, Vandermonde convolution .

Sources: <https://github.com/kolosovpetro/IdentitiesInRascalTriangle>

## 1. INTRODUCTION

In 2010, three middle school students, Alif Anggoro, Eddy Liu, and Angus Tulloch [1], were challenged to provide the next row for the number triangle shown below:

$$\begin{array}{ccccccc}
 & & & & 1 & & & & \\
 & & & & & & 1 & & \\
 & & & 1 & & & & 1 & \\
 & & 1 & & 2 & & 1 & & \\
 & 1 & & 3 & & 3 & & 1 & 
 \end{array}$$

The expected answer that matches Pascal's triangle [2] was “1 4 6 4 1”. However, Anggoro, Liu, and Tulloch suggested “1 4 5 4 1” instead. They devised this new row via so-called diamond formula:

$$\text{South} = \frac{\text{East} \cdot \text{West} + 1}{\text{North}}$$

So that upcoming rows of the triangle are

$n/k$	0	1	2	3	4	5	6	7
0	1							
1	1	1						
2	1	2	1					
3	1	3	3	1				
4	1	4	5	4	1			
5	1	5	7	7	5	1		
6	1	6	9	10	9	6	1	
7	1	7	11	13	13	11	7	1

**Table 1.** Rascal triangle. See the OEIS sequence [A077028](#) [3].

Since then, a lot of work has been done over the topic of rascal triangles. Numerous identities and relations have been revealed. For instance, few combinatorial interpretations of rascal numbers provided at [4], in particular, these interpretations establish a relation between rascal numbers and combinatorics of binary words. Few generalization approaches were proposed, namely generalized and iterated rascal triangles [5, 6]. In particular, the

concept of iterated rascal numbers establishes a close connection between rascal numbers and binomial coefficients.

## 2. BINOMIAL IDENTITIES IN ITERATED RASCAL TRIANGLES

Prior we begin our discussion it is worth to introduce a few preliminary facts and statements. Define the iterated rascal number

**Definition 2.1.** *Iterated rascal number* [6]

$$\binom{n}{k}_i = \sum_{m=0}^i \binom{n-k}{m} \binom{k}{m} \quad (2.1)$$

First important thing is to notice that iterated rascal number is a partial case of Vandermonde convolution [7]. Consider Vandermonde convolution

$$\binom{a+b}{r} = \sum_{m=0}^r \binom{a}{m} \binom{b}{r-m}$$

Thus,

$$\binom{n}{k}_i = \sum_{m=0}^i \binom{n-k}{m} \binom{k}{m} = \sum_{m=0}^i \binom{n-k}{m} \binom{k}{k-m} \quad (2.2)$$

Therefore, iterated rascal number is partial case of Vandermonde convolution with upper summation bound equals to  $i$ . Without further hesitation consider our findings.

**Proposition 2.2.** *Iterated rascal triangle equals to Pascal's triangle up to  $i$ -th column.*

$$\binom{n}{k}_i = \binom{n}{k}, \quad 0 \leq k \leq i \quad (2.3)$$

*Proof.* Proof is given by [6]. □

Then binomial identity follows

$$\binom{n}{i-k}_i = \binom{n}{i-k}$$

Applying binomial coefficients symmetry principle we obtain

$$\binom{n}{n-i+k}_i = \binom{n}{n-i+k}$$

**Proposition 2.3.** *Iterated rascal triangle equals to Pascal's triangle up to  $2i + 1$ -th row*

$$\binom{n}{k}_i = \binom{n}{k}, \quad 0 \leq n \leq 2i + 1$$

Therefore, for every fixed  $i \geq 0$

$$\binom{2i + 1 - n}{k}_i = \binom{2i + 1 - n}{k} \quad (2.4)$$

Equation (2.4) is of interest because in contrast to rascal column identity (2.3) it gives relation over  $k$  for each  $i$ , so that it is true for all cases in  $i, k$ :  $i < k$ ,  $i = k$  and  $k > i$ .

Taking  $t \geq 2i + 1$  for every fixed  $i \geq 0$

$$\binom{t - n}{k}_{t-i-1} = \binom{t - n}{k}$$

*Proof.* Proof of proposition (2.3). We have three possible relations between  $i, k$ :  $k < i$ ,  $k = i$ ,  $k > i$ . So we have to prove that for every  $i, k$

$$\sum_{m=0}^k \binom{2i + 1 - n - k}{m} \binom{k}{m} - \sum_{m=0}^i \binom{2i + 1 - n - k}{m} \binom{k}{m} = 0$$

For the case  $k < i$  proof is given in Jenna Gregory et al. [6]. For the case  $k = i$  proof is trivial. Thus, the remaining case is  $k > i$  yields that

$$\sum_{m=i+1}^k \binom{2i + 1 - n - k}{m} \binom{k}{m} = 0$$

Considering the constraints,

$$\begin{cases} n \geq 0 \\ k \geq i + 1 \\ 2i + 1 - n - k \leq i - n \\ m \geq i + 1 \end{cases}$$


Thus,

$$\sum_{m=i+1}^k \binom{2i + 1 - n - k}{m} \binom{k}{m}$$

Moreover, equation (2.4) gives Vandermonde-like identity

$$\binom{2i+1-n}{k} = \sum_{m=0}^i \binom{2i+1-n-k}{m} \binom{k}{m}$$
$$\binom{2i+1}{k} = \sum_{m=0}^i \binom{2i+1-k}{m} \binom{k}{m}$$

```
In[67]:= Grid[Table[Binoimial[n, k] - RascalNumber[n, k, 3], {n, 0, 20}, {k, 0, n}], Frame -> All]
```



We can spot that having  $i = 3$  the  $k = 4$ -th column gives binomial coefficient  $\binom{n}{4}$ . Indeed, this rule is true for every  $i$ .

**Proposition 2.5.** (*Row-column difference.*) For every fixed  $i \geq 0$

$$\binom{n+2i}{i} - \binom{n+2i}{i}_{i-1} = \binom{n+i}{i}$$

*Proof.* We have previously stated that iterated rascal numbers are closely related to Vandermonde convolution (2.2). Thus, proposition (2.5) can be rewritten as

$$\sum_{m=0}^i \binom{n+i}{m} \binom{i}{i-m} - \sum_{m=0}^{i-1} \binom{n+i}{m} \binom{i}{m}$$

Therefore,  $\binom{n+2i}{i} - \binom{n+2i}{i}_{i-1} = \binom{n+i}{i}$  is indeed true.  $\square$

Proposition (2.5) yields to few more identities. Applying binomial coefficients symmetry

$$\binom{n+2i}{n+i} - \binom{n+2i}{n+i}_{i-1} = \binom{n+i}{n}$$

Taking  $j = n + i$  gives

$$\begin{aligned} \binom{j+i}{j} - \binom{j+i}{j}_{i-1} &= \binom{j}{j-i} \\ \binom{j+i}{i} - \binom{j+i}{i}_{i-1} &= \binom{j}{i} \end{aligned}$$

Proposition (2.5) can be generalized even further, for every fixed  $i < k$ .

**Proposition 2.6.** (*Binomial coefficient difference iterated rascal number.*) For every fixed  $i < k$

$$\binom{n}{k} - \binom{n}{k}_i = \sum_{m=i+1}^k \binom{n-k}{m} \binom{k}{k-m}$$

*Proof.* It is true by means of Vandermonde convolution.  $\square$

### 3. Q-BINOMIAL IDENTITIES IN ITERATED RASCAL TRIANGLES

Consider the table of differences of binomial coefficients and iterated rascal numbers one more time as there is another pattern we can spot.

```
In[67]:= Grid[Table[Binomial[n, k] - RascalNumber[n, k, 3], {n, 0, 20}, {k, 0, n}], Frame -> All]
```

0																				
0	0																			
0	0	0																		
0	0	0	0																	
0	0	0	0	0																
0	0	0	0	0	0															
0	0	0	0	0	0	0														
0	0	0	0	0	0	0	0													
0	0	0	0	0	1	0	0	0	0											
0	0	0	0	0	5	5	0	0	0	0										
0	0	0	0	0	15	26	15	0	0	0	0									
0	0	0	0	0	35	81	81	35	0	0	0	0								
0	0	0	0	0	70	196	262	196	70	0	0	0	0							
0	0	0	0	0	126	406	658	658	406	126	0	0	0	0						
0	0	0	0	0	210	756	1414	1716	1414	756	210	0	0	0	0					
0	0	0	0	0	330	1302	2730	3830	3830	2730	1302	330	0	0	0	0				
0	0	0	0	0	495	2112	4872	7680	8885	7680	4872	2112	495	0	0	0	0			
0	0	0	0	0	715	3267	8184	14232	18525	18525	14232	8184	3267	715	0	0	0	0		
0	0	0	0	0	1001	4862	13101	24816	35697	40186	35697	24816	13101	4862	1001	0	0	0	0	
0	0	0	0	0	1365	7007	20163	41217	64713	80587	80587	64713	41217	20163	7007	1365	0	0	0	0
0	0	0	0	0	1820	9828	30030	65780	111705	152020	168230	152020	111705	65780	30030	9828	1820	0	0	0

**Figure 2.** Difference  $\binom{n}{k} - \binom{n}{k}_3$ . Highlighted column is  $(1, 5)$ -binomial coefficient  $\begin{bmatrix} n \\ k \end{bmatrix}^5$ . Sequence **A096943** in the OEIS [9].

The  $(1, q)$ -binomial coefficients  $\begin{bmatrix} n \\ k \end{bmatrix}^q$  are special kind of binomial coefficients defined by

**Definition 3.1.**  $(1, q)$ -Binomial coefficient [10]

$$\begin{bmatrix} n \\ k \end{bmatrix}^q = \begin{cases} q & \text{if } k = 0, n = 0 \\ 1 & \text{if } k = 0 \\ 0 & \text{if } k > n \\ \begin{bmatrix} n-1 \\ k \end{bmatrix}^q + \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}^q & \end{cases} \quad (3.1)$$

Indeed, the relation shown in Figure (2) is true for every  $i$ , so that it establishes a relation between  $(1, q)$ -binomial coefficients and iterated rascal numbers.

**Proposition 3.2.** (Relation between iterated rascal numbers and  $(1, q)$ -binomial coefficients.)

For every fixed  $i \geq 0$

$$\binom{2i+3+j}{i+2} - \binom{2i+3+j}{i+2}_i = \begin{bmatrix} i+2+j \\ i+2 \end{bmatrix}^{i+2}$$

Taking  $t = i + 2$  in (3.2) yields

$$\binom{2t-1+j}{t} - \binom{2t-1+j}{t}_{t-2} = \left[ \begin{matrix} t+j \\ t \end{matrix} \right]^t$$

In particular, having  $i = 1$  proposition (3.2) gives the OEIS sequence A006503 [11] such that third column of (1,3)-Pascal triangle A095660 [12].

#### 4. CONCLUSIONS

Conclusions of your manuscript.

#### REFERENCES

- [1] Anggoro, Alif and Liu, Eddy and Tulloch, Angus. The Rascal Triangle. *The College Mathematics Journal*, 41(5):393–395, 2010. <https://doi.org/10.4169/074683410X521991>.
- [2] Sloane, N. J. A. Pascal’s triangle read by rows. Entry A007318 in The On-Line Encyclopedia of Integer Sequences, 1994. <https://oeis.org/A007318>.
- [3] Sloane, N. J. A. The Rascal triangle read by rows. Entry A077028 in The On-Line Encyclopedia of Integer Sequences, 2002. <https://oeis.org/A077028>.
- [4] Amelia Gibbs and Brian K. Miceli. Two Combinatorial Interpretations of Rascal Numbers. *arXiv preprint arXiv:2405.11045*, 2024. <https://arxiv.org/abs/2405.11045>.
- [5] Hotchkiss, Philip K. Generalized Rascal Triangles. *arXiv preprint arXiv:1907.11159*, 2019. <https://arxiv.org/abs/1907.11159>.
- [6] Gregory, Jena and Kronholm, Brandt and White, Jacob. Iterated ascal triangles. *Aequationes mathematicae*, pages 1–18, 2023. <https://doi.org/10.1007/s00010-023-00987-6>.
- [7] Andrews, George E. and Askey, Richard and Roy, Ranjan and Roy, Ranjan and Askey, Richard. *Special functions*, volume 71. Cambridge university press Cambridge, 2000.
- [8] Sloane, N. J. A. Binomial coefficient binomial(n,4). Entry A000332 in The On-Line Encyclopedia of Integer Sequences, 2009. <https://oeis.org/A000332>.
- [9] Sloane, N. J. A. Sixth column of (1,5)-Pascal triangle. Entry A096943 in The On-Line Encyclopedia of Integer Sequences, 2004. <https://oeis.org/A096943>.
- [10] Sloane, N. J. A. Pascal (1,5) triangle. Entry A096940 in The On-Line Encyclopedia of Integer Sequences, 2004. <https://oeis.org/A096940>.



- [11] Sloane, N. J. A. Entry A006503 in The On-Line Encyclopedia of Integer Sequences, 1995. <https://oeis.org/A006503>.
- [12] Sloane, N. J. A. Pascal (1,3) triangle. Entry A095660 in The On-Line Encyclopedia of Integer Sequences, 2004. <https://oeis.org/A095660>.

**Version:** Local-0.1.0

SOFTWARE DEVELOPER, DEVOPS ENGINEER

*Email address:* kolosovp94@gmail.com

*URL:* <https://kolosovpetro.github.io>