

# COUNTING PRIMES AND TWIN-PRIMES USING MINIMAL GOLDBACH PAIRS

PETRO KOLOSOV

ABSTRACT. Assuming Goldbach's Conjecture holds, every even integer  $2N \geq 4$  can be written as  $2N = p_i + p_j$  where  $(p_i, p_j)$  is called Goldbach pair. The minimal Goldbach pair is a pair  $(p_i, p_j)$  having the minimal  $p_i$  such that  $p_j = 2N - p_i$  is also a prime. We define a function  $F_{2N}(P)$  that counts occurrences of  $p_j = P$  within the range  $6 \leq 2k \leq 2N$ . In particular, the function  $F_{2N}(P)$  provides the following identities in terms of prime counting  $\pi(2N)$  and twin-prime counting  $\pi_2(2N)$

$$\pi(2N) = F_{2N}(3) + 1, \quad \pi_2(2N) = F_{2N}(3) - F_{2N}(5)$$

## CONTENTS

1. Introduction	1
2. Conclusions	5
References	5

## 1. INTRODUCTION

This manuscript provides a comprehensive review of the work [1] done by Michel Yamagishi. The Goldbach conjecture asserts that every even integer  $2N \geq 4$  is a sum of two

---

*Date:* April 19, 2025.

*2010 Mathematics Subject Classification.* 11N05, 11P32.

*Key words and phrases.* Goldbach conjecture, Goldbach pairs, Minimal Goldbach pairs, Primes, Twin primes .

Sources: <https://github.com/kolosovpetro/MinimalGoldbachPairsInPrimesCounting>

primes

$$2N = p_i + p_j$$

where  $(p_i, p_j)$  is called a Goldbach pair.

A Goldbach pair is not unique for even integers greater than 6, which means that there can be multiple Goldbach pairs for even integer  $2N \geq 8$ . For example:  $10 = 3 + 7$  and  $10 = 5 + 5$  and  $10 = 7 + 3$ , where the Goldbach pairs are  $(3, 7)$ ,  $(5, 5)$ ,  $(7, 3)$ .

The minimal Goldbach pair is the pair with the smallest  $p_i$  among all goldbach pairs for even integer  $2N$ . For the even integer 10, we have three pairs:  $(3, 7)$ ,  $(5, 5)$ ,  $(7, 3)$  and the minimal one is  $(3, 7)$  because 3 is the smallest among all  $p_i$  values: 3, 5, 7.

Consider the following minimal goldbach pairs for even integer  $2k$  within the range  $6 \leq 2k \leq 50$

$6 = 3 + 3,$	$8 = 3 + 5,$
$10 = 3 + 7,$	$12 = 5 + 7,$
$14 = 3 + 11,$	$16 = 3 + 13,$
$18 = 5 + 13,$	$20 = 3 + 17,$
$22 = 3 + 19,$	$24 = 5 + 19,$
$26 = 3 + 23,$	$28 = 5 + 23,$
$30 = 7 + 23,$	$32 = 3 + 29,$
$34 = 3 + 31,$	$36 = 5 + 31,$
$38 = 7 + 31,$	$40 = 3 + 37,$
$42 = 5 + 37,$	$44 = 3 + 41,$
$46 = 3 + 43,$	$48 = 5 + 43,$
$50 = 3 + 47$	

We can notice that minimal Goldbach pairs having  $p_i = 3$  produce a sequence of odd prime numbers  $p_j = 3, 5, 7, 11, 13, 17 \dots$  which is quite remarkable:

$$\begin{array}{ll}
 6 = 3 + 3, & 8 = 3 + 5, \\
 10 = 3 + 7, & 14 = 3 + 11, \\
 16 = 3 + 13, & 20 = 3 + 17, \\
 22 = 3 + 19, & 26 = 3 + 23, \\
 32 = 3 + 29, & 34 = 3 + 31, \\
 40 = 3 + 37, & 44 = 3 + 41, \\
 46 = 3 + 43, & 50 = 3 + 47
 \end{array}$$

Another interesting observation is that by selecting the pairs with minimal  $p_i = 5$  yields the sequence of primes  $p_j$  such that  $p_j + 2$  is not prime

$$\begin{array}{ll}
 12 = 5 + 7, & 18 = 5 + 13, \\
 24 = 5 + 19, & 28 = 5 + 23, \\
 36 = 5 + 31, & 42 = 5 + 37, \\
 48 = 5 + 43.
 \end{array}$$

To formalize and clarify our discussion, we define a few functions. Let  $G_{\min}(2N)$  be a function that returns a set of minimal Goldbach pairs  $(p_i, p_j)$  having  $\min p_i$  over the range  $6 \leq 2k \leq 2N$

$$G_{\min}(2N) = \{(p_i, p_j) \mid p_i + p_j = 2k \mid 6 \leq 2k \leq 2N \mid \min p_i\}.$$

For example,

$$G_{\min}(20) = \{(3, 3), (3, 5), (3, 7), (5, 7), (3, 11), (3, 13), (5, 13), (3, 17)\}$$

Let  $W_{2N}(P)$  be a function that returns the set of elements  $p_j$  from  $G_{\min}(2N)$  having  $p_i = P$

$$W_{2N}(P) = \{p_j \mid (p_i, p_j) \in G_{\min}(2N) \text{ and } p_i = P\}$$

Then, the sequence of odd prime numbers [2] is given by  $W_{2N}(3)$

$$\{3, 5, 7, 11, \dots, p \leq 2N - 3\} = W_{2N}(3)$$

Now we can easily count the number of primes within the interval  $6 \leq 2k \leq 2N$  because  $\pi(2N)$  is equal to the total number of elements inside the set  $W_{2N}(3)$

$$\pi(2N) = F_{2N}(3) + 1$$

where  $F_{2N}(3)$  is the function that counts the number of elements inside the set  $W_{2N}(3)$ . In general  $F_{2N}(P) = |W_{2N}(3)|$ .

Taking  $P = 5$  in  $W_{2N}(P)$  yields a sequence of primes  $p_j$  such that  $p_j + 2$  is not a prime [3]

$$W_{2N}(5) = \{7, 13, 19, 23, 31, 37, 43, 47, 53, \dots, p \leq 2N - 5\}$$

This implies that the number of twin primes in range  $6 \leq 2k \leq 2N$  can be expressed in terms of  $F_{2N}(P)$

$$\pi_2(2N) = F_{2N}(3) - F_{2N}(5)$$

where  $2N = 10^k + 2$ ,  $k = 1, 2, 3, 4, \dots$ . For example,

$$\pi_2(12) = F_{12}(3) - F_{12}(5) = 2$$

$$\pi_2(102) = F_{102}(3) - F_{102}(5) = 8$$

$$\pi_2(1002) = F_{1002}(3) - F_{1002}(5) = 35$$

$$\pi_2(10002) = F_{10002}(3) - F_{10002}(5) = 205$$

$$\pi_2(100002) = F_{100002}(3) - F_{100002}(5) = 1224$$

$$\pi_2(1000002) = F_{1000002}(3) - F_{1000002}(5) = 8169$$

These results match the sequence [4].

Having  $P = 7$  function  $W_{2N}(P)$  yields the sequence of primes such that  $p_j - p_i \geq 6$ , where  $p_j$  is the next prime after  $p_i$ , see [5]

$$W_{2N}(7) = \{23, 31, 47, 53, 61, 73, 83, 89, 113, \dots, p \leq 2N - 7\}$$

This allows us to count the primes for which the next-prime gap at least 6:  $\delta_6(2N) = F_{2N}(7)$ .

## 2. CONCLUSIONS

Assuming Goldbach's Conjecture holds, we introduced a framework based on minimal Goldbach pairs to derive expressions for key prime-related functions. Specifically, we defined the function  $F_{2N}(P)$  that counts occurrences of primes  $p_j$  in minimal Goldbach pairs  $(p_i, p_j)$  where  $p_i = P$ . Using this framework, we obtained

- The prime-counting function:  $\pi(2N) = F_{2N}(3) + 1$
- The twin-prime counting function:  $\pi_2(2N) = F_{2N}(3) - F_{2N}(5)$
- The count of primes with next-prime gap at least 6:  $\delta_6(2N) = F_{2N}(7)$

These identities establish a novel connection between Goldbach partitions and classical prime number theory. Computational examples confirm alignment with known integer sequences, reinforcing the potential of this approach for analytical and numerical exploration of prime distributions. All the results validated up to  $N = 10^8$  with source code available on GitHub [6].

## REFERENCES

- [1] Michel Yamagishi. Goldbach's conjecture and how to calculate  $\pi(n)$  and  $\pi_2(n)$ , 2025. <http://dx.doi.org/10.13140/RG.2.2.22119.76963>.
- [2] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A065091: Odd primes. <https://oeis.org/A065091>, 2025. Accessed: 2025-04-18.
- [3] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A049591: Odd primes  $p$  such that  $p+2$  is composite. <https://oeis.org/A049591>, 2025. Accessed: 2025-04-18.
- [4] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A007508: Number of twin prime pairs below  $10$  to power  $N$ . <https://oeis.org/A007508>, 2025. Accessed: 2025-04-18.

- [5] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A124582: Primes  $p$  such that  $q-p$  greater of equal 6, where  $q$  is the next prime after  $p$ . <https://oeis.org/A124582>, 2025. Accessed: 2025-04-18.
- [6] Petro Kolosov. Minimal Goldbach Pairs (GitHub). <https://github.com/kolosovpetro/MinimalGoldbachPairs>, 2025. Accessed: 2025-04-18.

**Version:** Local-0.1.0

SOFTWARE DEVELOPER, DEVOPS ENGINEER

*Email address:* kolosovp94@gmail.com

*URL:* <https://kolosovpetro.github.io>