# MINIMAL GOLDBACH PAIRS IN PRIME AND TWIN-PRIME COUNTING

#### PETRO KOLOSOV

ABSTRACT. Goldbach's Conjecture assumes that every even integer  $2N \geq 4$  can be written as the sum of two primes  $2N = p_i + p_j$ , where  $(p_i, p_j)$  is called a Goldbach pair. The minimal Goldbach pair is a pair  $(p_i, p_j)$  such that  $p_i$  is minimal and  $p_j = 2N - p_i$  is also a prime. We define a function  $F_{2N}(P)$  that counts the occurrences of  $p_i = P$  in a set of minimal Goldbach pairs up to 2N, where P is a fixed prime number. In particular, the function  $F_{2N}(P)$  provides the following identities in terms of prime counting  $\pi(2N)$ , twin-prime counting  $\pi_2(2N)$  and cousin prime counting  $\pi_4(2N)$ 

$$\pi(2N) = F_{2N+3}(3) + 1$$

$$\pi_2(2N) = F_{2N+3}(3) - F_{2N+5}(5)$$

$$\pi_4(2N) = F_{2N}(5) - F_{2N}(7)$$

## 1. Introduction

This manuscript provides a comprehensive review of the work [1] done by Michel Yamagishi, extending it with additional results. The Goldbach conjecture asserts that every even integer  $2N \ge 4$  is a sum of two primes

$$2N = p_i + p_i$$

where  $(p_i, p_j)$  is called a Goldbach pair.

Date: April 22, 2025.

2010 Mathematics Subject Classification. 11N05, 11P32.

Key words and phrases. Goldbach conjecture, Goldbach pairs, Minimal Goldbach pairs, Primes, Twin primes .

Sources: https://github.com/kolosovpetro/MinimalGoldbachPairsInPrimesCounting

A Goldbach pair is not unique for even integers greater than 6, which means that there can be multiple Goldbach pairs for even integer  $2N \ge 8$ . For example: 10 = 3 + 7 and 10 = 5 + 5 and 10 = 7 + 3, where the Goldbach pairs are (3,7), (5,5), (7,3).

The minimal Goldbach pair is the pair with the smallest  $p_i$  among all goldbach pairs for even integer 2N. For the even integer 10, we have three pairs: (3,7), (5,5), (7,3) and the minimal one is (3,7) because 3 is the smallest among all  $p_i$  values: 3,5,7.

Consider the following minimal Goldbach pairs for even integer 2k within the range  $6 \le 2k \le 50$ 

6 = 3 + 3,	30 = 7 + 23,
8 = 3 + 5,	32 = 3 + 29,
10 = 3 + 7,	34 = 3 + 31,
12 = 5 + 7,	36 = 5 + 31,
14 = 3 + 11,	38 = 7 + 31,
16 = 3 + 13,	40 = 3 + 37,
18 = 5 + 13,	42 = 5 + 37,
20 = 3 + 17,	44 = 3 + 41,
22 = 3 + 19,	46 = 3 + 43,
24 = 5 + 19,	48 = 5 + 43,
26 = 3 + 23,	50 = 3 + 47,
28 = 5 + 23,	

We can notice that minimal Goldbach pairs having  $p_i = 3$  produce a sequence of odd prime numbers  $p_j = 3, 5, 7, 11, 13, 17...$  which is quite remarkable:

6 = 3 + 3,	32 = 3 + 29,
8 = 3 + 5,	34 = 3 + 31,
10 = 3 + 7,	40 = 3 + 37,
14 = 3 + 11,	44 = 3 + 41,
16 = 3 + 13,	46 = 3 + 43,
20 = 3 + 17,	50 = 3 + 47,
22 = 3 + 19,	
26 = 3 + 23,	

Another interesting observation is that by selecting the pairs with minimal  $p_i = 5$  yields the sequence of primes  $p_j$  such that  $p_j + 2$  is not prime

$$12 = 5 + 7,$$
  $36 = 5 + 31,$   $18 = 5 + 13,$   $42 = 5 + 37,$   $24 = 5 + 19,$   $48 = 5 + 43,$   $28 = 5 + 23,$ 

To formalize and clarify our discussion, we define a few functions. Let  $G_{\min}(2N)$  be a function that returns a set of minimal Goldbach pairs  $(p_i, p_j)$  having  $\min p_i$  over the range  $6 \le 2k \le 2N$ 

$$G_{\min}(2N) = \{(p_i, p_j) \mid p_i + p_j = 2k \mid 6 \le 2k \le 2N \mid \min p_i\}.$$

For example,

$$G_{\min}(20) = \{(3,3), (3,5), (3,7), (5,7), (3,11), (3,13), (5,13), (3,17)\}$$

Let  $W_{2N}(P)$  be a function that returns the set of elements  $p_j$  from  $G_{\min}(2N)$  having  $p_i = P$ 

$$W_{2N}(P) = \{ p_j \mid (p_i, p_j) \in G_{\min}(2N) \text{ and } p_i = P \}$$

Then, the sequence of odd prime numbers [2] is given by  $W_{2N}(3)$ 

$$\{3, 5, 7, 11, \ldots, p \le 2N - 3\} = W_{2N}(3)$$

Now we can easily count the number of primes within the interval  $6 \le 2k \le 2N$  because  $\pi(2N)$  is equal to the total number of elements inside the set  $W_{2N}(3)$ , which corresponds to the sequence [3]

$$\pi(2N) = F_{2N+3}(3) + 1$$

where  $F_{2N}(3)$  is the function that counts the number of elements inside the set  $W_{2N}(3)$ . In general  $F_{2N}(P) = |W_{2N}(P)|$ .

Taking P = 5 in  $W_{2N}(P)$  yields a sequence of primes  $p_j$  such that  $p_j + 2$  is not a prime [4]

$$W_{2N}(5) = \{7, 13, 19, 23, 31, 37, 43, 47, 53, \dots, p \le 2N - 5\}$$

Hence, by excluding the elements of the set  $W_{2N}(5)$  from the set  $W_{2N}(3)$  yields the sequence of lesser twin primes [5]

$$W_{2N}(3) \setminus W_{2N}(5) = \{3, 5, 11, 17, 29, 41, 59, 71, 101, 107, 137, \dots, p \le 2N - 3\}$$

This implies that the number of twin primes in range  $6 \le 2k \le 2N$  can be expressed in terms of  $F_{2N}(P)$ 

$$\pi_2(2N) = F_{2N+3}(3) - F_{2N+5}(5)$$

where  $2N = 10^k$ , k = 1, 2, 3, 4, ... For example,

$$\pi_2(10) = F_{10+3}(3) - F_{10+5}(5) = 2$$

$$\pi_2(100) = F_{100+3}(3) - F_{100+5}(5) = 8$$

$$\pi_2(1000) = F_{1000+3}(3) - F_{1000+5}(5) = 35$$

$$\pi_2(10000) = F_{10000+3}(3) - F_{10000+5}(5) = 205$$

$$\pi_2(100000) = F_{100000+3}(3) - F_{100000+5}(5) = 1224$$

$$\pi_2(1000000) = F_{1000000+3}(3) - F_{1000000+5}(5) = 8169$$

These results match the sequence [6].

In addition, the functions W, F provide a way to count twin-primes p with the difference of 4. We can observe this by excluding the elements of the set  $W_{2N}(7)$  from the set  $W_{2N}(5)$  which gives the sequence [7] of twin primes  $p_4$ 

$$W_{2N}(5) \setminus W_{2N}(7) = \{7, 13, 19, 37, 43, 67, 79, 97, 103, 109, 127, 163, \dots, p \le 2N - 5\}$$

Hence, the number of twin-primes  $p_4$  up to 2N is

$$\pi_4(2N) = F_{2N}(5) - F_{2N}(7)$$

For example,

$$\pi_4(10) = F_{10}(5) - F_{10}(7) = 0$$

$$\pi_4(100) = F_{100}(5) - F_{100}(7) = 7$$

$$\pi_4(1000) = F_{1000}(5) - F_{1000}(7) = 40$$

$$\pi_4(10000) = F_{10000}(5) - F_{10000}(7) = 202$$

$$\pi_4(100000) = F_{100000}(5) - F_{100000}(7) = 1215$$

$$\pi_4(1000000) = F_{1000000}(5) - F_{1000000}(7) = 8143$$

These results match the sequence [8].

Having P = 7 function  $W_{2N}(P)$  yields the sequence of primes such that  $p_j - p_i \ge 6$ , where  $p_j$  is the next prime after  $p_i$ , see [9]

$$W_{2N}(7) = \{23, 31, 47, 53, 61, 73, 83, 89, 113, \dots, p \le 2N - 7\}$$

This allows us to count the primes for which the next-prime gap at least 6:  $\delta_6(2N) = F_{2N+7}(7)$ .

### 2. Conclusions

Assuming Goldbach's Conjecture holds, we introduced a framework based on minimal Goldbach pairs to derive expressions for key prime-related functions. Specifically, we defined the function  $F_{2N}(P)$  that counts occurrences of primes  $p_j$  in minimal Goldbach pairs  $(p_i, p_j)$  where  $p_i = P$ . Using this framework, we obtained

- The prime-counting function:  $\pi(2N) = F_{2N+3}(3) + 1$
- The twin-prime  $p_2$  counting function:  $\pi_2(2N) = F_{2N+3}(3) F_{2N+5}(5)$
- The twin-prime  $p_4$  counting function:  $\pi_4(2N) = F_{2N}(5) F_{2N}(7)$
- The count of primes with next-prime gap at least 6:  $\delta_6(2N) = F_{2N+7}(7)$

These identities establish a novel connection between Goldbach partitions and classical prime number theory. Computational examples confirm alignment with known integer sequences, reinforcing the potential of this approach for analytical and numerical exploration of prime distributions. All the results validated up to  $N = 10^8$  with source code available on GitHub [10].

## REFERENCES

- [1] Michel Yamagishi. Goldbach's conjecture and how to calculate  $\pi(n)$  and  $\pi_2(n)$ , 2025. http://dx.doi.org/10.13140/RG.2.2.22119.76963.
- [2] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A065091: Odd primes. https://oeis.org/A065091, 2025. Accessed: 2025-04-18.
- [3] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A000720: Number of primes less than or equal to N. https://oeis.org/A000720, 2025. Accessed: 2025-04-18.

MINIMAL GOLDBACH PAIRS IN PRIME AND TWIN-PRIME COUNTING

7

[4] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A049591: Odd primes p

such that p+2 is composite. https://oeis.org/A049591, 2025. Accessed: 2025-04-18.

[5] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A001359: Lesser of twin

primes. https://oeis.org/A001359, 2025. Accessed: 2025-04-18.

[6] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A007508: Number of twin

prime pairs below 10 to power N. https://oeis.org/A007508, 2025. Accessed: 2025-04-18.

[7] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A023200: Primes p such

that p + 4 is also prime. https://oeis.org/A023200, 2025. Accessed: 2025-04-18.

[8] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A093737: Number of cousin

primes (prime pairs differing by 4) less than or equal to n. https://oeis.org/A093737, 2025. Accessed:

2025-04-18.

[9] The OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, A124582: Primes p such

that q-p greater of equal 6, where q is the next prime after p. https://oeis.org/A124582, 2025.

Accessed: 2025-04-18.

[10] Petro Kolosov. Minimal Goldbach Pairs (GitHub). https://github.com/kolosovpetro/

MinimalGoldbachPairs, 2025. Accessed: 2025-04-18.

Version: Local-0.1.0

SOFTWARE DEVELOPER, DEVOPS ENGINEER

 $Email\ address: {\tt kolosovp94@gmail.com}$ 

URL: https://kolosovpetro.github.io