

ODD-POWER IDENTITY VIA MULTIPLICATION OF CERTAIN MATRICES

PETRO KOLOSOV

ABSTRACT. In this manuscript, we show an odd-power identity in terms of certain matrix multiplication. More precisely, the matrix of size 1×1 such that $a_{1,1} = N^{2M+1}$ is a result of multiplication of three matrices $\mathbf{J}_N \times \mathbf{K}_{N,M} \times \mathbf{T}_M$, where \mathbf{J}_N is unit row vector of size $1 \times N$, $\mathbf{K}_{N,M}$ is a matrix of size $N \times M$, and \mathbf{T}_M is a column vector of size $M \times 1$, so that

$$\begin{bmatrix} N^{2M+1} \end{bmatrix} = \mathbf{J}_N \times \mathbf{K}_{N,M} \times \mathbf{T}_M$$

CONTENTS

Definitions	1
1. Introduction	2
2. Conclusions	4
References	4

DEFINITIONS

- \mathbf{J}_N – unit row vector of all 1's having the size $1 \times N$. For example,

$$\mathbf{J}_5 = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

- $\mathbf{K}_{N,M}$ – matrix of size $N \times M$ defined by

$$\mathbf{K}_{N,M} = (k^r (N - k)^r)_{0 \leq k \leq N, 0 \leq r \leq M}$$

Date: March 9, 2024.

2010 Mathematics Subject Classification. 26E70, 05A30.

Key words and phrases. Keyword1, Keyword2 .

- \mathbf{T}_M – column vector of size $M \times 1$ defined by

$$\mathbf{T}_M = (\mathbf{A}_{M,r})_{M=\text{const}, 0 \leq r \leq M}$$

where $\mathbf{A}_{M,r}$ is rational value (literature).

- Row vector
- Column vector
- $\mathbf{A} = (a_{ij})$

1. INTRODUCTION

Your introduction here. Include some references [1, 2, 3, 4, 5, 6]. Lorem Ipsum is simply dummy text of the printing and typesetting industry. Lorem Ipsum has been the industry's standard dummy text ever since the 1500s, when an unknown printer took a galley of type and scrambled it to make a type specimen book. It has survived not only five centuries, but also the leap into electronic typesetting, remaining essentially unchanged. It was popularised in the 1960s with the release of Letraset sheets containing Lorem Ipsum passages, and more recently with desktop publishing software like Aldus PageMaker including versions of Lorem Ipsum.

Figure example




Figure 1. Figure example.

$$\prod_m^a b$$

And for any natural m we have polynomial identity

$$x^m = \sum_{k=1}^m T(m, k) x^{[k]} \tag{1.1}$$

where $x^{[k]}$ denotes central factorial defined by

$$x^{[n]} = x \left(x + \frac{n}{2} - 1 \right)^{n-1}$$

where $(n)^k = n(n-1)(n-2)\cdots(n-k+1)$ denotes falling factorial in Knuth's notation. In particular,

$$x^{[n]} = x \left(x + \frac{n}{2} - 1\right) \left(x + \frac{n}{2} - 1\right) \cdots \left(x + \frac{n}{2} - n + 1\right) = x \prod_{k=1}^{n-1} \left(x + \frac{n}{2} - k\right)$$

2. CONCLUSIONS

Conclusions of your manuscript.

REFERENCES

- [1] Benaoumeur Bayour, Ahmed Hammoudi, and Delfim FM Torres. A truly conformable calculus on time scales. *arXiv preprint arXiv:1705.08928*, 2017. <https://arxiv.org/abs/1705.08928>.
- [2] Nadia Benkhettou, Salima Hassani, and Delfim FM Torres. A conformable fractional calculus on arbitrary time scales. *Journal of King Saud University-Science*, 28(1):93–98, 2016.
- [3] M Cristina Caputo. Time scales: from nabla calculus to delta calculus and vice versa via duality. *arXiv preprint arXiv:0910.0085*, 2009.
- [4] Nat á lia Martins and Delfim FM Torres. Calculus of variations on time scales with nabla derivatives. *Nonlinear Analysis: Theory, Methods & Applications*, 71(12):e763–e773, 2009.
- [5] Petro Kolosov. "Github Template" Source files. Available electronically at <https://github.com/kolosovpetro/github-latex-template>, 2022.
- [6] N. J. A. Sloane. The on-line encyclopedia of integer sequences. published electronically at <https://oeis.org>, 1964.

Version: Local-0.1.0

SOFTWARE DEVELOPER, DEVOPS ENGINEER

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>