

# DEFINITIONS

PETRO KOLOSOV

ABSTRACT. Definitions

## CONTENTS

### 1. Definitions

1

#### 1. DEFINITIONS

- $T(n, k)$  recursively defines central factorial numbers of the second kind (in the context of Knuth and Riordan (see references)). It is defined in mathematica package as

`CentralFactorialNumber1`

$$\left\{ \begin{array}{l} T(n, 1) = 1 \\ T(n, n) = 1 \\ T(n, k) = T(n-1, k-1) + k^2 T(n-1, k) \end{array} \right.$$

- $T(n, k)$  is central factorial number defined as `CentralFactorialNumber2` in mathematica package

$$T(n, k) = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^j \left( \frac{1}{2}k - j \right)^n$$

---

*Date:* July 22, 2023.

*2010 Mathematics Subject Classification.* 26E70, 05A30.

*Key words and phrases.* Polynomials, Polynomial identities, Faulhaber's formula, Cental Factorial Numbers .

- Identity in central factorial numbers defined as `CFNIdentity1` in mathematica package

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^j \binom{2k}{j} (k-j)^{2n}$$

- Identity in central factorial numbers defined as `CFNIdentity2` in mathematica package

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^{k-j} \binom{2k}{k-j} j^{2n}$$

- Identity in central factorial numbers defined as `CFNIdentity3` in mathematica package

$$(2k-1)!T(2n, 2k) = \frac{1}{2k} \sum_{j=0}^{2k} (-1)^j \binom{2k}{j} (k-j)^{2n}$$