

# POLYNOMIAL IDENTITIES AUXILIARY

PETRO KOLOSOV

ABSTRACT. Polynomial identities auxiliary

## CONTENTS

1. Polynomial identities auxiliary	1
1.1. Central factorial numbers	1
1.2. Knuth's formula - approach 1 (to be verified all)	2

## 1. POLYNOMIAL IDENTITIES AUXILIARY

### 1.1. Central factorial numbers.

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^j \binom{2k}{j} (k-j)^{2n} \quad (CFNIdentity1)$$

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^{k-j} \binom{2k}{k-j} j^{2n} \quad (CFNIdentity2)$$

$$(2k-1)!T(2n, 2k) = \frac{1}{2k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^j (k-j)^{2n} \quad (CFNIdentity3)$$

$$T(n, k) = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^j \left(\frac{1}{2}k - j\right)^n \quad (CentralFactorialNumber2)$$

---

*Date:* July 22, 2023.

2010 *Mathematics Subject Classification.* 26E70, 05A30.

*Key words and phrases.* Polynomials, Polynomial identities, Faulhaber's formula, Cental Factorial Numbers .

### 1.2. Knuth's formula - approach 1 (to be verified all).

$$n^{2m-1} = \sum_{k=1}^m (2k-1)! T(2m, 2k) \binom{n+k-1}{2k-1} \quad (1.1)$$

Equation (1.1) is validated via Mathematica functions: *OddPowerIdentity1*, *OddPowerIdentity2*, *OddPowerIdentity3*. Substituting  $(2k-1)! T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^{k-j} \binom{2k}{k-j} j^{2n}$  we get

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{k} \binom{n+k-1}{2k-1} \binom{2k}{k-j} j^{2m} \quad (\text{OddPowerIdentity21})$$

By means of binomial identity  $\frac{k}{n} \binom{n}{k} = \binom{n-1}{k-1}$

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{k} \frac{2k}{n+k} \binom{n+k}{2k} \binom{2k}{k-j} j^{2m} \quad (\text{OddPowerIdentity22})$$

Collapsing common terms we get

$$n^{2m-1} = 2 \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{n+k} \binom{n+k}{2k} \binom{2k}{k-j} j^{2m} \quad (\text{OddPowerIdentity23})$$

By means of binomial identity  $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}$  we get

$$n^{2m-1} = 2 \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{n+k} \binom{n+k}{k-j} \binom{n+j}{k+j} j^{2m} \quad (\text{OddPowerIdentity24})$$

Because the symmetry of binomial coefficients  $\binom{n+k}{k-j} = \binom{n+k}{n+k-(k-j)}$  holds, we get

$$n^{2m-1} = 2 \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{n+k} \binom{n+k}{n+j} \binom{n+j}{k+j} j^{2m} \quad (\text{OddPowerIdentity25})$$