POLYNOMIAL IDENTITIES INVOLVING CENTRAL FACTORIAL NUMBERS

PETRO KOLOSOV

Abstract.

Contents

1. Formulae 1

1. Formulae

From OEIS, note that this is not Central factorial number itself, this formula is in the mathematica package as OEISFormula

$$T_{\text{OEIS}}(n,k) = \frac{1}{m} \sum_{j=0}^{m} (-1)^j \binom{2m}{j} (m-j)^{2n}$$

where m = n - k + 1. So that

$$T_{\text{OEIS}} = \frac{1}{n-k+1} \sum_{j=0}^{n-k+1} (-1)^j \binom{2n-2k+2}{j} (n-k+1-j)^{2n}$$

Also, OEIS sequence is defined by

$$T_{\text{OEIS}} = (2(n-k)+1)!T(2n,2n-2k)$$

Date: July 19, 2023.

2010 Mathematics Subject Classification. 26E70, 05A30.

 $Key\ words\ and\ phrases.$ Polynomials, Polynomial identities, Faulhaber's formula, Cental Factorial Numbers .

where T(2n, 2n - 2k) are central factorial numbers. From stackoverflow, these are pure central factorial numbers already

$$k!T(n,k) = \sum_{j=0}^{n} {k \choose j} (-1)^j \left(\frac{1}{2}k - j\right)^n$$

So that central factorial number is, this is the function Central1(n,k) in mathematica package and it is true and holds in mathematica program

$$T(n,k) = \frac{1}{k!} \sum_{j=0}^{n} {k \choose j} (-1)^j \left(\frac{1}{2}k - j\right)^n$$

Let be (k-1)!T(n,k)

$$(k-1)!T(n,k) = \frac{1}{k} \sum_{j=0}^{n} {k \choose j} (-1)^j \left(\frac{1}{2}k - j\right)^n$$

Let be (2k-1)!T(2n,2k) in is true and checked in mathematica as KnuthCoefficient2

$$(2k-1)!T(2n,2k) = \frac{1}{2k} \sum_{j=0}^{\infty} {2k \choose j} (-1)^j (k-j)^{2n}$$

 $Email\ address: {\tt kolosovp94@gmail.com}$

URL: https://kolosovpetro.github.io