

POLYNOMIAL IDENTITIES INVOLVING CENTRAL FACTORIAL NUMBERS

PETRO KOLOSOV

ABSTRACT. Central factorial numbers appear in mathematical literature quite frequently. Few of prominent works are Riordan's "Combinatorial identities" and Knuth's "Johann Faulhaber and sums of powers" along as many others. In this manuscript, we start our discussion from the definition of central factorial numbers (both, recursive and iterative) continuing with a set of identities shown further in this manuscript. Afterward, based on odd-power identities given by Knuth, we show new variations of odd-power identities applying certain relations in terms of central factorial numbers. Finally, we provide a comprehensive way to validate the results of the manuscript via supplementary Mathematica programs.

CONTENTS

1. Introduction and main results	2
2. Conclusions	6
References	6

Date: November 8, 2024.

2010 *Mathematics Subject Classification.* 26E70, 05A30.

Key words and phrases. Polynomials, Polynomial identities, Central factorial numbers, Central factorials, Binomial identities, Riordan Combinatorial identities, Falling factorials, Power sums, Faulhaber's formula .

Sources: <https://github.com/kolosovpetro/PolynomialIdentitiesInvolvingCentralFactorialNumbers>

1. INTRODUCTION AND MAIN RESULTS

Central factorial numbers appear in mathematical literature quite frequently. In Riordan's Combinatorial identities [1, p. 217] they defined as

$$k!T(n, k) = \sum_{j=0}^k \binom{k}{j} (-1)^j \left(\frac{1}{2}k - j\right)^n \quad (1.1)$$

where $T(n, k)$ is central factorial number. Also, in the book *The Divided Central Differences of Zero* [2, eq. (10a)] central factorial numbers are referenced as

$$K_{rs} = \frac{1}{(2s)!} \sum_{t=0}^{2s} (-1)^t \binom{2s}{t} (s - t)^{2r+2}$$

J.F. Steffenson mentions central factorial numbers in his book *Interpolation* [3] in context of central difference of polynomial x^r at zero

$$k!T(n, k) = \delta^m 0^r = \sum_{v=0}^m (-1)^v \binom{m}{v} \left(\frac{m}{2} - v\right)^r$$

D. E. Knuth gives the following recurrence for the central factorial numbers

$$T(2m + 2, 2k) = k^2 T(2m, 2k) + T(2m, 2k - 2)$$

See *Johann Faulhaber and sums of powers* [4, p. 284].

Central factorial numbers appear in The On-Line Encyclopedia of Integer Sequences [5] as the following recurrence relation. Let be $U(n, k) = T(2n, 2k)$ then

$$U(n, k) = \begin{cases} 1, & \text{if } k = 1 \\ 1, & \text{if } k = n \\ U(n - 1, k - 1) + k^2 U(n - 1, k), & \text{otherwise} \end{cases}$$

Moreover, the OEIS sequence [6] gives polynomial representation of central factorial numbers

$$T(2n, 2k) = 2 \sum_{j=1}^k (-1)^{k-j} \frac{j^{2n}}{(k-j)!(k+j)!}$$

It is important to note that central factorial numbers are closely related to the central difference operator δ , Newton interpolation formula and central factorials $x^{[n]}$. A process of

derivation of central factorial numbers by applying central difference operator δ , Newton interpolation formula and central factorials $x^{[n]}$ is provided by the MathStackExchange discussion [7]. So there are a few identities in central factorial numbers at our disposal. The first is

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^j \binom{2k}{j} (k-j)^{2n} \quad (1.2)$$

Such that it is derived from the formula given by the OEIS sequence A303675 [8], that is

$$A303675(n, k) = \frac{1}{n-k+1} \sum_{j=0}^{n-k+1} (-1)^j \binom{2(n-k+1)}{j} ((n-k+1)-j)^{2n}$$

Note that the term $n-k+1$ is given only to serve proper sequence offset and can be omitted by iterator k which yields the identity (1.2). The second identity is actively used in current manuscript is

$$(2k-1)!T(2n, 2k) = \frac{1}{k} \sum_{j=0}^k (-1)^{k-j} \binom{2k}{k-j} j^{2n} \quad (1.3)$$

which is derived by inversion of the sum (1.2). The third one and the final identity we base our results on involves a partial case of the equation (1.1)

$$(2k-1)!T(2n, 2k) = \frac{1}{2k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^j (k-j)^{2n} \quad (1.4)$$

So that now we review the results of Knuth's work *Johann Faulhaber and sums of powers* [4, p. 10], in particular odd-power polynomial identities. For odd powers we have

$$n^{2m-1} = \sum_{k=1}^m (2k-1)!T(2m, 2k) \binom{n+k-1}{2k-1} \quad (1.5)$$

And for any integer power m there is an identity in terms of central factorial numbers and central factorials

$$x^m = \sum_{k=1}^m T(m, k) x^{[k]} \quad (1.6)$$

where $x^{[k]}$ denotes central factorial defined by

$$x^{[n]} = x \left(x + \frac{n}{2} - 1 \right)^{\frac{n-1}{2}}$$

where $(n)^k = n(n-1)(n-2)\cdots(n-k+1)$ denotes falling factorial in Knuth's notation. In particular,

$$x^{[n]} = x \left(x + \frac{n}{2} - 1\right) \left(x + \frac{n}{2} - 1\right) \cdots \left(x + \frac{n}{2} - n + 1\right) = x \prod_{k=1}^{n-1} \left(x + \frac{n}{2} - k\right)$$

So that having the whole context of the topic, we can derive and discuss few other polynomial identities smoothly.

For example, given odd power identity (1.5) and identity in central factorial numbers (1.2) we easily obtain

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^j}{k} \binom{2k}{j} \binom{n+k-1}{2k-1} (k-j)^{2m}$$

Furthermore, by means of binomial identity $\frac{k}{n} \binom{n}{k} = \binom{n-1}{k-1}$ we get ¹

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^j}{k} \frac{2k}{n+k} \binom{n+k}{2k} \binom{2k}{j} (k-j)^{2m}$$

Collapsing common terms k and applying binomial identity $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}$ another polynomial identity follows

$$n^{2m-1} = 2 \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^j}{n+k} \binom{n+k}{j} \binom{n+k-j}{2k-j} (k-j)^{2m}$$

It is possible to derive more polynomial identities based on binomial coefficients symmetry $\binom{n}{k} = \binom{n}{n-k}$ and an identity in binomial coefficients $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}$.

So for now, we smoothly switch our discussion to another example. Given an odd power identity (1.5) we can observe that odd powered polynomial can be expressed in terms of central factorial numbers and falling factorials as follows

$$n^{2m-1} = \sum_{k=1}^m T(2m, 2k) (n+k-1)^{\underline{2k-1}}$$

Because $\binom{n}{k} = \frac{1}{k!} (n)^k$ where $(n)^k$ is falling factorial.

¹The majority of binomial identities used in this paper are given by Gross J. L. in his book *Combinatorial methods with computer applications* [9], chapter 4.

Yet another odd power identity follows by means of (1.5) and (1.3), that is

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{k} \binom{n+k-1}{2k-1} \binom{2k}{k-j} j^{2m}$$

By means of binomial identity $\frac{k}{n} \binom{n}{k} = \binom{n-1}{k-1}$ we get

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{k} \frac{2k}{n+k} \binom{n+k}{2k} \binom{2k}{k-j} j^{2m}$$

Collapsing common terms k and applying the binomial identity $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}$ yields

$$n^{2m-1} = 2 \sum_{k=1}^m \sum_{j=0}^k \frac{(-1)^{k-j}}{n+k} \binom{n+k}{k-j} \binom{n+j}{k+j} j^{2m}$$

As the final part of our discussion on odd power identities, consider the polynomial identities based on odd power identity (1.5) and identity in central factorial numbers (1.4).

Replacing the term $(2k-1)!T(2n, 2k)$ in (1.5) we get

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^{2k} \frac{(-1)^j}{2k} \binom{n+k-1}{2k-1} \binom{2k}{j} (k-j)^{2m}$$

Continuing similarly with the binomial identity $\frac{k}{n} \binom{n}{k} = \binom{n-1}{k-1}$ yields

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^{2k} \frac{(-1)^j}{n+k} \binom{n+k}{2k} \binom{2k}{j} (k-j)^{2m}$$

So that now we are free applying the binomial identity $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}$ to obtain

$$n^{2m-1} = \sum_{k=1}^m \sum_{j=0}^{2k} \frac{(-1)^j}{n+k} \binom{n+k}{j} \binom{n+k-j}{2k-j} (k-j)^{2m}$$

The remaining opportunity to show a few more power identities lays in replacing the corresponding coefficient $T(n, k)$ in (1.6) by identities in central factorial numbers (1.2), (1.3), (1.4).

2. CONCLUSIONS

In this manuscript, we have provided and discussed polynomial identities in terms of central factorial number. We started from literature overview including various forms and definitions of central factorial numbers continuing with polynomial identities based on those definitions. Finally, all the results of this manuscript can be validated using supplementary Mathematica programs at this [GitHub repository](#).

REFERENCES

- [1] J. Riordan. *Combinatorial Identities*. Wiley series in probability and mathematical statistics. Wiley, 1968.
- [2] L. Carlitz and John Riordan. The Divided Central Differences of Zero. *Canadian Journal of Mathematics*, 15:94–100, 1963.
- [3] Johan Frederik Steffensen. *Interpolation*. Williams & Wilkins, 1927.
- [4] Knuth, Donald E. Johann Faulhaber and sums of powers. *Mathematics of Computation*, 61(203):277–294, 1993. <https://arxiv.org/abs/math/9207222>.
- [5] Sloane, N. J. A. Triangle of central factorial numbers, Entry A008957 in The On-Line Encyclopedia of Integer Sequences. Published electronically at <https://oeis.org/A008957>, 2000.
- [6] Sloane, N. J. A. Triangle of central factorial numbers, Entry A036969 in The On-Line Encyclopedia of Integer Sequences. Published electronically at <https://oeis.org/A036969>, 2000.
- [7] Scheuer, Markus. MathStackExchange answer 3665722/463487. Published electronically at <https://math.stackexchange.com/a/3665722/463487>, 2020.
- [8] Kolosov, Petro and Luschny, Peter. Coefficients in the sum of odd powers as expressed by Faulhaber’s theorem, Entry A303675 in The On-Line Encyclopedia of Integer Sequences. Published electronically at <https://oeis.org/A303675>, 2018.
- [9] Jonathan L. Gross. *Combinatorial methods with computer applications*. CRC Press, 2016. <https://kolosovpetro.github.io/pdf/CM4.pdf>.

Version: Local-0.1.0

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>