

POLYNOMIAL IDENTITY INVOLVING BINOMIAL THEOREM AND FAULHABER'S FORMULA

PETRO KOLOSOV

CONTENTS

1. Approach via system of linear equations	1
--	---

1. APPROACH VIA SYSTEM OF LINEAR EQUATIONS

One approach to prove the conjecture was proposed by Albert Tkaczyk in his series of preprints (two references). The essence of the approach lays in construction and solving of the particular system of linear equations. Such system linear equations is constructed using Binomial theorem and Faulhaber's formula that allows us to find closed forms of power sums as part of identity (reference to equation). Consider the polynomial relation

$$n^{2m+1} = \sum_{r=0}^m \mathbf{A}_{m,r} \sum_{k=1}^n k^r (n-k)^r \quad (1.1)$$

Expanding the $(n-k)^r$ part via Binomial theorem we get

$$\begin{aligned} n^{2m+1} &= \sum_{r=0}^m \mathbf{A}_{m,r} \sum_{k=1}^n k^r (n-k)^r \\ &= \sum_{r=0}^m \mathbf{A}_{m,r} \sum_{k=1}^n k^r \left[\sum_{t=0}^r (-1)^t \binom{r}{t} n^{r-t} k^t \right] \\ &= \sum_{r=0}^m \mathbf{A}_{m,r} \left[\sum_{t=0}^r (-1)^t \binom{r}{t} n^{r-t} \sum_{k=1}^n k^{t+r} \right] \end{aligned} \quad (1.2)$$

Date: July 15, 2023.

2010 *Mathematics Subject Classification.* 26E70, 05A30.

Key words and phrases. Binomial theorem, Polynomial identities, Binomial coefficients, Bernoulli numbers, Pascal's triangle, Faulhaber's formula .

For arbitrary m we have

$$\begin{aligned}
n^{2m+1} &= \sum_{r=0}^m \mathbf{A}_{m,r} \left[\sum_{t=0}^r (-1)^t \binom{r}{t} n^{r-t} \sum_{k=1}^n k^{t+r} \right] \\
&= \mathbf{A}_{m,0}n + \mathbf{A}_{m,1} \left[\frac{1}{6}(-n + n^3) \right] + \mathbf{A}_{m,2} \left[\frac{1}{30}(-n + n^5) \right] + \mathbf{A}_{m,3} \left[\frac{1}{420}(-10n + 7n^3 + 3n^7) \right] \\
&+ \mathbf{A}_{m,4} \left[\frac{1}{630}(-21n + 20n^3 + n^9) \right] + \mathbf{A}_{m,5} \left[\frac{1}{2772}(-210n + 231n^3 - 22n^5 + n^{11}) \right] \\
&+ \mathbf{A}_{m,6} \left[\frac{1}{60060}(-15202n + 18200n^3 - 3003n^5 + 5n^{13}) \right] \\
&+ \mathbf{A}_{m,7} \left[\frac{1}{51480}(-60060n + 76010n^3 - 16380n^5 + 429n^7 + n^{15}) \right] \\
&+ \mathbf{A}_{m,8} \left[\frac{1}{218790}(-1551693n + 2042040n^3 - 516868n^5 + 26520n^7 + n^{17}) \right] + \dots
\end{aligned} \tag{1.3}$$

Example 1.1. Let be fixed $m = 1$ so that we have the following relation defined by (eqref)

$$\mathbf{A}_{m,0}n + \mathbf{A}_{m,1} \left[\frac{1}{6}(-n + n^3) \right] - n^3 = 0$$

Multiplying by 6 both, right hand side and left hand side, we get

$$6\mathbf{A}_{1,0}n + \mathbf{A}_{1,1}(-n + n^3) - 6n^3 = 0$$

Opening brackets and rearranging the terms gives

$$6\mathbf{A}_{1,0} - \mathbf{A}_{1,1}n + \mathbf{A}_{1,1}n^3 - 6n^3 = 0$$

Combining the terms yields

$$n(6\mathbf{A}_{1,0} - \mathbf{A}_{1,1}) + n^3(\mathbf{A}_{1,1} - 6) = 0$$

Therefore, the system of linear equations follows

$$\begin{cases} 6\mathbf{A}_{1,0} - \mathbf{A}_{1,1} = 0 \\ \mathbf{A}_{1,1} - 6 = 0 \end{cases}$$

Solving it we get

$$\begin{cases} \mathbf{A}_{1,1} = 6 \\ \mathbf{A}_{1,0} = 1 \end{cases}$$

Example 1.2. Let be fixed $m = 2$ so that we have the following relation defined by (eqref)

$$\mathbf{A}_{m,0}n + \mathbf{A}_{m,1} \left[\frac{1}{6}(-n + n^3) \right] + \mathbf{A}_{m,2} \left[\frac{1}{30}(-n + n^5) \right] - n^5 = 0$$

Multiplying by 30 both, right hand side and left hand side, we get

$$30\mathbf{A}_{2,0}n + 5\mathbf{A}_{2,1}(-n + n^3) + \mathbf{A}_{2,2}(-n + n^5) - 30n^5 = 0$$

Opening brackets and rearranging the terms gives

$$30\mathbf{A}_{2,0} - 5\mathbf{A}_{2,1}n + 5\mathbf{A}_{2,1}n^3 - \mathbf{A}_{2,2}n + \mathbf{A}_{2,2}n^5 - 30n^5 = 0$$

Combining the terms yields

$$n(30\mathbf{A}_{2,0} - 5\mathbf{A}_{2,1} - \mathbf{A}_{2,2}) + 5\mathbf{A}_{2,1}n^3 + n^5(\mathbf{A}_{2,2} - 30) = 0$$

Therefore, the system of linear equations follows

$$\begin{cases} 30\mathbf{A}_{2,0} - 5\mathbf{A}_{2,1} - \mathbf{A}_{2,2} = 0 \\ \mathbf{A}_{2,1} = 0 \\ \mathbf{A}_{2,2} - 30 = 0 \end{cases}$$

Solving it we get

$$\begin{cases} \mathbf{A}_{2,2} = 30 \\ \mathbf{A}_{2,1} = 0 \\ \mathbf{A}_{2,0} = 1 \end{cases}$$

Example 1.3. Let be fixed $m = 3$ so that we have the following relation defined by (eqref)

$$\mathbf{A}_{m,0}n + \mathbf{A}_{m,1} \left[\frac{1}{6}(-n + n^3) \right] + \mathbf{A}_{m,2} \left[\frac{1}{30}(-n + n^5) \right] + \mathbf{A}_{m,3} \left[\frac{1}{420}(-10n + 7n^3 + 3n^7) \right] - n^7 = 0$$

Multiplying by 420 both, right hand side and left hand side, we get

$$420\mathbf{A}_{3,0}n + 70\mathbf{A}_{2,1}(-n + n^3) + 14\mathbf{A}_{2,2}(-n + n^5) + \mathbf{A}_{3,3}(-10n + 7n^3 + 3n^7) - 420n^7 = 0$$

Opening brackets and rearranging the terms gives

$$420\mathbf{A}_{3,0}n - 70\mathbf{A}_{3,1} + 70\mathbf{A}_{3,1}n^3 - 14\mathbf{A}_{3,2}n + 14\mathbf{A}_{3,2}n^5 - 10\mathbf{A}_{3,3}n + 7\mathbf{A}_{3,3}n^3 + 3\mathbf{A}_{3,3}n^7 - 420n^7 = 0$$

Combining the terms yields

$$n(420\mathbf{A}_{3,0} - 70\mathbf{A}_{3,1} - 14\mathbf{A}_{3,2} - 10\mathbf{A}_{3,3}) + n^3(70\mathbf{A}_{3,1} + 7\mathbf{A}_{3,3}) + n^5 14\mathbf{A}_{3,2} + n^7(3\mathbf{A}_{3,3} - 420) = 0$$

Therefore, the system of linear equations follows

$$\begin{cases} 420\mathbf{A}_{3,0} - 70\mathbf{A}_{3,1} - 14\mathbf{A}_{3,2} - 10\mathbf{A}_{3,3} = 0 \\ 70\mathbf{A}_{3,1} + 7\mathbf{A}_{3,3} = 0 \\ \mathbf{A}_{3,2} - 30 = 0 \\ 3\mathbf{A}_{3,3} - 420 = 0 \end{cases}$$

Solving it we get

$$\begin{cases} \mathbf{A}_{3,3} = 140 \\ \mathbf{A}_{3,2} = 0 \\ \mathbf{A}_{3,1} = -\frac{7}{70}\mathbf{A}_{3,3} = -14 \\ \mathbf{A}_{3,0} = \frac{(70\mathbf{A}_{3,1} + 10\mathbf{A}_{3,3})}{420} = 1 \end{cases}$$

Example 1.4. Let be fixed $m = 4$ so that we have the following relation defined by (eqref)

$$\begin{aligned} &\mathbf{A}_{m,0}n + \mathbf{A}_{m,1} \left[\frac{1}{6}(-n + n^3) \right] + \mathbf{A}_{m,2} \left[\frac{1}{30}(-n + n^5) \right] + \mathbf{A}_{m,3} \left[\frac{1}{420}(-10n + 7n^3 + 3n^7) \right] \\ &+ \mathbf{A}_{m,4} \left[\frac{1}{630}(-21n + 20n^3 + n^9) \right] - n^9 = 0 \end{aligned}$$

Multiplying by 630 both, right hand side and left hand side, we get

$$630\mathbf{A}_{4,0}n + 105\mathbf{A}_{4,1}(-n + n^3) + 21\mathbf{A}_{4,2}(-n + n^5) + \frac{3}{2}\mathbf{A}_{4,3}(-10n + 7n^3 + 3n^7) + \mathbf{A}_{4,4}(-21n + 20n^3 + n^9) - 630n^9 = 0$$

Opening brackets and rearranging the terms gives

$$630\mathbf{A}_{4,0}n - 105\mathbf{A}_{4,1}n + 105\mathbf{A}_{4,1}n^3 - 21\mathbf{A}_{4,2}n + 21\mathbf{A}_{4,2}n^5 - \frac{3}{2}\mathbf{A}_{4,3} \cdot 10n + \frac{3}{2}\mathbf{A}_{4,3} \cdot 7n^3 + \frac{3}{2}\mathbf{A}_{4,3} \cdot 3n^7 - 21\mathbf{A}_{4,4}n + 20\mathbf{A}_{4,4}n^3 + \mathbf{A}_{4,4}n^9 - 630n^9 = 0$$

Combining the terms yields

$$n(630\mathbf{A}_{4,0} - 105\mathbf{A}_{4,1} - 21\mathbf{A}_{4,2} - 15\mathbf{A}_{4,3} - 21\mathbf{A}_{4,4}) + n^3 \left(105\mathbf{A}_{4,1} + \frac{21}{2}\mathbf{A}_{4,3} + 20\mathbf{A}_{4,4} \right) + n^5(21\mathbf{A}_{4,2}) + n^7 \left(\frac{9}{2}\mathbf{A}_{4,3} \right) + n^9(\mathbf{A}_{4,4} - 630) = 0$$

Therefore, the system of linear equations follows

$$\begin{cases} 630\mathbf{A}_{4,0} - 105\mathbf{A}_{4,1} - 21\mathbf{A}_{4,2} - 15\mathbf{A}_{4,3} - 21\mathbf{A}_{4,4} = 0 \\ 105\mathbf{A}_{4,1} + \frac{21}{2}\mathbf{A}_{4,3} + 20\mathbf{A}_{4,4} = 0 \\ \mathbf{A}_{4,2} = 0 \\ \mathbf{A}_{4,3} = 0 \\ \mathbf{A}_{4,4} - 630 = 0 \end{cases}$$

Solving it we get

$$\begin{cases} \mathbf{A}_{4,4} = 630 \\ \mathbf{A}_{4,3} = 0 \\ \mathbf{A}_{4,2} = 0 \\ \mathbf{A}_{4,1} = -\frac{20}{105}\mathbf{A}_{4,4} = -120 \\ \mathbf{A}_{4,0} = \frac{105\mathbf{A}_{4,1} + 21\mathbf{A}_{4,4}}{630} = 1 \end{cases}$$

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>