

ROW SUMS CONJECTURE IN ITERATED RASCAL TRIANGLES

PETRO KOLOSOV

ABSTRACT. In [1], Gregory et al. provide the following conjecture for row sums of iterated rascal triangles. For every i

$$\sum_{k=0}^{4i+3} \binom{4i+3}{k}_i = 2^{4i+2}$$

where $\binom{n}{k}_i$ is an iterated rascal number.

CONTENTS

1. Introduction	1
2. Row sums conjecture	3
References	3

1. INTRODUCTION

Rascal triangle is Pascal-like numeric triangle developed in 2010 by three middle school students, Alif Anggoro, Eddy Liu, and Angus Tulloch [?]. During math classes they were

Date: July 6, 2024.

2010 Mathematics Subject Classification. 11B25, 11B99.

Key words and phrases. Pascal's triangle, Rascal triangle, Binomial coefficients, Binomial identities, Binomial theorem, Generalized Rascal triangles, Iterated rascal triangles, Iterated rascal numbers, Number triangle, Arithmetic sequence, Vandermonde identity, Vandermonde convolution .

Sources: <https://github.com/kolosovpetro/RowSumsConjectureInRascalTriangle>

challenged to provide the next row for the following number triangle

$$\begin{array}{ccccccc}
 & & & & 1 & & & \\
 & & & & & & 1 & \\
 & & 1 & & & 2 & & 1 \\
 & 1 & & 3 & & & 3 & 1 \\
 & & & & \dots & & &
 \end{array}$$

Teacher's expected answer was the one that matches Pascal's triangle, e.g. "1 4 6 4 1". However, Anggoro, Liu, and Tulloch suggested "1 4 5 4 1" instead. They devised this new row via what they called diamond formula

$$\mathbf{South} = \frac{\mathbf{East} \cdot \mathbf{West} + 1}{\mathbf{North}}$$

So they obtained the following triangle

n/k	0	1	2	3	4	5	6	7
0	1							
1	1	1						
2	1	2	1					
3	1	3	3	1				
4	1	4	5	4	1			
5	1	5	7	7	5	1		
6	1	6	9	10	9	6	1	
7	1	7	11	13	13	11	7	1

Table 1. Rascal triangle.

Indeed, the forth row is "1 4 5 4 1" because $4 = \frac{1 \cdot 3 + 1}{1}$ and $5 = \frac{3 \cdot 3 + 1}{2}$. Since then, a lot of work has been done over the topic of rascal triangles. In this article we stick our attention to the one of rascal triangles generalizations, namely iterated rascal triangles [1]. Iterated rascal number is defined via a sum of binomial coefficients multiplication

Definition 1.1. *Iterated rascal number*

$$\binom{n}{k}_i = \sum_{m=0}^i \binom{n-k}{m} \binom{k}{m} \quad (1.1)$$

Thus, the rascal triangle (1) is the triangle generated by $\binom{n}{k}_1$.

2. ROW SUMS CONJECTURE

In [1], Gregory et al. provide the following conjecture for row sums of iterated rascal triangles.

Conjecture 2.1. *For every i*

$$\sum_{k=0}^{4i+3} \binom{4i+3}{k}_i = 2^{4i+2}$$

where $\binom{n}{k}_i$ is an iterated rascal number.

REFERENCES

- [1] Gregory, Jena and Kronholm, Brandt and White, Jacob. Iterated rascal triangles. *Aequationes mathematicae*, pages 1–18, 2023. <https://doi.org/10.1007/s00010-023-00987-6>.

Version: Local-0.1.0

SOFTWARE DEVELOPER, DEVOPS ENGINEER

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>