

SUMS OF POWERS OF INTEGERS: A COMPLETE FRAMEWORK FOR CLOSED FORMULAS

PETRO KOLOSOV

ABSTRACT.

CONTENTS

1. Historical context	1
2. Definitions and notations	3
3. Auxiliary lemmas	4
4. Ordinary sums of powers	6
5. Remark on central factorial numbers of the second kind	8
6. Generalized central factorial numbers of the second kind	9
7. Double and triple sums of powers in terms of central differences	12
8. Multifold sums of powers in terms of central differences	13
9. Multifold sums of powers in generalized central factorial numbers	14
10. Multifold sums of powers in binomial form	16
11. Framework for sums of powers	17
12. Conclusions	20
References	20
Mathematica programs	22

1. HISTORICAL CONTEXT

The story begins in ancient Greek times, with discrete approximations to obtain continuous areas by the method of exhaustion. In particular, Archimedes (287-212 BC) evaluated sums

Date: May 30, 2026.

of squares while finding the spiral's area [1]

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

Moving to medieval age (1636), Pierre de Fermat (1601-1665) calls the problem of finding formulas for sums of powers ... *perhaps the most beautiful problem of all arithmetics*, in his correspondence with Blaise Pascal (1623-1662). Fermat's contribution to the topic was quite remarkable indeed, as he introduced the concept of figurate numbers $F_{n,k}$, which is extensively discussed in [2, (2020)].

Meanwhile, Johann Faulhaber (1580-1635) worked on the problem of sums of powers independently, in *Academia Algebrae* [3, (1631)]. Faulhaber's contribution was great indeed, because Johann managed to compute formulas up to $\Sigma^1 n^{25} = 1^{25} + 2^{25} + \dots + n^{25}$, which is quite impressive. Imagine to derive the formula back in 17-th century, without computers. Although, he published formulas up to $\Sigma^1 n^{17}$, keeping $\Sigma^1 n^{25}$ as an exercise for readers. In fact, this exercise was solved in 1993, by Donald Knuth, around three centuries later. Yet again, Faulhaber's contribution was indeed impressive; however, it was not complete, as he did not manage to find more general pattern.

The real breakthrough was done by Jacob Bernoulli (1665-1705) in his work entitled *Ars coniectandi* [4, (1713)], published postmortem. In the work, Jacob introduced Bernoulli numbers (named after himself of course), thus allowing for arbitrary powers to be summed up

$$\sum_{k=1}^n k^p = \frac{1}{p+1} \sum_{r=0}^p \binom{p+1}{r} B_r n^{p+1-r}$$

It assumes that $B_0 = +\frac{1}{2}$. The argument why $B_0 = +\frac{1}{2}$ is quite a huge topic, the reader is encouraged to familiarize with the whole story [5, 6]. Therefore, Bernoulli managed to contribute the missing capstone to Faulhaber's work, making it complete. In fact, the formula above is widely known as Faulhaber's formula, although Bernoulli has done the final great work.

Since then, the topic of power sums was abandoned for nearly three centuries. In 1993, Donald Knuth resurrected the topic of sums of powers, giving it new life (with remarkable results). Based on works of Carl Gustav Jacobi (1804-1851), and Gessel-Viennot [7, (1989)], Knuth reviews Faulhaber's work from completely new point of view, from the point of Modern enumerative combinatorics. Remarkably, Knuth's contribution does not necessary involves Bernoulli numbers, instead he derived formulas for powers sums by utilizing special numbers (Stirling, central factorials), and binomial coefficients. Apart that, Knuth introduced a really handy notation for r -fold sums of powers $\Sigma^r n^m$, which we use in this work.

2. DEFINITIONS AND NOTATIONS

Allow us to define the mathematical objects we utilize throughout the manuscript. For multifold sums of powers, we use the notation provided by Donald Knuth in [8], because it is simple and beautiful.

Definition 2.1 (Multifold sums of powers). *For non-negative integers r, n, m*

$$\begin{aligned}\Sigma^0 n^m &= n^m, \\ \Sigma^1 n^m &= \Sigma^0 1^m + \Sigma^0 2^m + \cdots + \Sigma^0 n^m, \\ \Sigma^{r+1} n^m &= \Sigma^r 1^m + \Sigma^r 2^m + \cdots + \Sigma^r n^m.\end{aligned}$$

We utilize the following notation for falling factorials

Definition 2.2 (Falling factorials). *For integers x, n*

$$(x)_n = \begin{cases} 1, & \text{if } k = 0, \\ x(x-1)(x-2) \cdots (x-n+1) = \prod_{k=0}^{n-1} (x-k), & \text{if } k > 0, \\ 0, & \text{else.} \end{cases}$$

We utilize the following notation for central factorials

Definition 2.3 (Central factorials). *For integers n, k*

$$n^{[k]} = \begin{cases} 0, & \text{if } k < 0, \\ 1, & \text{if } k = 0, \\ n \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = n \left(n + \frac{k}{2} - 1\right)_{k-1}, & \text{if } k > 0, \end{cases}$$

where $\left(n + \frac{k}{2} - 1\right)_{k-1} = \prod_{j=0}^{k-2} \left(n + \frac{k}{2} - 1 - j\right)$ are falling factorials.

We encourage the audience to read J. F. Steffensen's work on the definition of generalized factorials [9].

3. AUXILIARY LEMMAS

Lemma 3.1 (Central Newton's formula). *For a function f defined on integers and arbitrary integers x, t*

$$f(x) = \sum_{k=0}^{\infty} \frac{(x-t)^{[k]}}{k!} \delta^k f(t),$$

where $\delta^k f(t)$ denotes the k -th central finite difference

$$\delta^k f(t) = \sum_{j=0}^k (-1)^j \binom{k}{j} f\left(t + \frac{k}{2} - j\right),$$

and $(x-t)^{[k]}$ are central factorials defined by (2.3).

Proof. See [10, p. 462], and [11, section 2, eq. (19)], and [12, section 6, eq. (21)]. □

Lemma 3.2 (Newton's formula for powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \sum_{k=0}^m \frac{(n-t)^{[k]}}{k!} \delta^k t^m.$$

Lemma 3.3 (Central factorials binomial form). *For integers n and $k \geq 1$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Proof. We have

$$\frac{n^{[k]}}{k!} = \frac{n}{k!} \left(n + \frac{k}{2} - 1 \right)_{k-1} = \frac{n}{k(k-1)!} \left(n + \frac{k}{2} - 1 \right)_{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1},$$

because of the identity in falling factorials $\frac{(x)_n}{n!} = \binom{x}{n}$. $\square$

Lemma 3.4 (Central factorials binomial form decomposition). *For integers $n, k \geq 0$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \binom{n + \frac{k}{2}}{k} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k}.$$

Proof. We have $\binom{a}{k} = \frac{a}{k} \binom{a-1}{k-1}$. Let $a = n + \frac{k}{2}$, then

$$\binom{n + \frac{k}{2}}{k} = \frac{n + \frac{k}{2}}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Thus,

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \binom{n + \frac{k}{2}}{k} - \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

By moving under common denominator, and applying binomial recurrence yields

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \left[2 \binom{n + \frac{k}{2}}{k} - \binom{n + \frac{k}{2} - 1}{k-1} \right] = \frac{1}{2} \left[\binom{n + \frac{k}{2}}{k} + \binom{n + \frac{k}{2} - 1}{k} + \binom{n + \frac{k}{2} - 1}{k-1} - \binom{n + \frac{k}{2} - 1}{k-1} \right].$$

Thus, the statement follows. $\square$

Lemma 3.5 (Newton's formula for powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n - t + \frac{k}{2}}{k} + \binom{n - t + \frac{k}{2} - 1}{k} \right] \delta^k t^m.$$

Lemma 3.6 (Generalized hockey-stick identity). *For integers a, b and j ,*

$$\sum_{k=a}^b \binom{k}{j} = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

Proof. We have, $\sum_{k=a}^b \binom{k}{j} = \binom{a}{j} + \binom{a+1}{j} + \dots + \binom{b}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right)$. By hockey stick identity $\sum_{k=0}^n \binom{k}{j} = \binom{n+1}{j+1}$ yields,

$$\sum_{k=a}^b \binom{k}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right) = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

This completes the proof. $\square$

Lemma 3.7 (Centered hockey-stick identity). *For integers $n \geq 1$, and arbitrary integers t, k, r*

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Proof. By setting $a = 1 - t + \frac{k}{2} + r$, and $b = n - t + \frac{k}{2} + r$ into Lemma (3.6) yields

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

This completes the proof. □

4. ORDINARY SUMS OF POWERS

By decomposition of Newton's formula for powers (3.5), for integers $n, m \geq 0$, and an arbitrary integer t , we get

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\sum_{j=1}^n \binom{j-t+\frac{k}{2}}{k} + \sum_{j=1}^n \binom{j-t+\frac{k}{2}-1}{k} \right] \delta^k t^m. \quad (1)$$

Now we reach the transition to closed form of ordinary sums of powers. By setting $r = 0$, and $r = -1$ into Lemma (3.7), we get closed forms of the sums $\sum_{j=1}^n \binom{j-t+\frac{k}{2}}{k}$, and $\sum_{j=1}^n \binom{j-t+\frac{k}{2}-1}{k}$, respectively. Therefore,

Proposition 4.1 (Ordinary sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} + \binom{n-t+\frac{k}{2}}{k+1} - \binom{-t+\frac{k}{2}}{k+1} \right] \delta^k t^m.$$

For example,

Example 4.2 (Sums of squares in central differences).

$$\begin{aligned}
t = 0 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(1+n) + \frac{\delta^2 t^2}{2} \frac{n(1+3n+2n^2)}{6}, \\
t = 1 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-1) + \frac{\delta^2 t^2}{2} \frac{n(1-3n+2n^2)}{6}, \\
t = 2 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-3) + \frac{\delta^2 t^2}{2} \frac{n(13-9n+2n^2)}{6}, \\
t = 3 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-5) + \frac{\delta^2 t^2}{2} \frac{n(37-15n+2n^2)}{6}, \\
t = 4 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-7) + \frac{\delta^2 t^2}{2} \frac{n(73-21n+2n^2)}{6}.
\end{aligned}$$

Example 4.3 (Sums of cubes in central differences).

$$\begin{aligned}
t = 0 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(1+n) + \frac{\delta^2 t^3}{2} \frac{n(1+3n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-1+n+4n^2+2n^3)}{24}, \\
t = 1 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-1) + \frac{\delta^2 t^3}{2} \frac{n(1-3n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(1+n-4n^2+2n^3)}{24}, \\
t = 2 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-3) + \frac{\delta^2 t^3}{2} \frac{n(13-9n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-21+25n-12n^2+2n^3)}{24}, \\
t = 3 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-5) + \frac{\delta^2 t^3}{2} \frac{n(37-15n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-115+73n-20n^2+2n^3)}{24}, \\
t = 4 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-7) + \frac{\delta^2 t^3}{2} \frac{n(73-21n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-329+145n-28n^2+2n^3)}{24}.
\end{aligned}$$

It is indeed interesting to notice the connection between the decomposition of ordinary sums of powers (4.1) and Riordan's central factorial numbers of the second kind, that is

Proposition 4.4 (Central factorial numbers sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n + \frac{k}{2} + 1}{k+1} - \binom{1 + \frac{k}{2}}{k+1} + \binom{n + \frac{k}{2}}{k+1} - \binom{\frac{k}{2}}{k+1} \right] k! T(m, k),$$

where $T(n, k)$ denotes the central factorial numbers of the second kind (in Riordan's notation). See [12, p. 213], and [13].

These central factorial numbers of the second kind $T(n, k)$ are defined by central Newton's formula for powers (3.2) evaluated in zero, such that

$$n^m = \sum_{k=0}^{\infty} T(m, k) n^{[k]}.$$

Which implies

$$T(n, k) = \frac{\delta^k 0^m}{k!},$$

by Newton's formula (3.2).

5. REMARK ON CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND

In [8] Donald Knuth gives a remarkable formula for sums of odd powers

Proposition 5.1 (Knuth's odd powers sum). *For non-negative integers n, m*

$$\Sigma^1 n^{2m-1} = \sum_{k=1}^m (2k-1)! \binom{n+k}{2k} T(2m, 2k),$$

where $T(2m, 2k)$ are central factorial numbers of the second kind.

It is quite interesting to notice that Proposition (5.1) originates from centered Newton's formula for powers (3.2) evaluated at zero. By Newton's formula, we have

$$n^{2m} = \sum_{k=0}^{2m} \frac{n^{[k]}}{k!} \cdot \delta^k 0^{2m}$$

The iteration for $k = 0$ can be omitted because $\delta^0 0^{2m} = 0$. Next, we observe that $\delta^k 0^{2m} = 0$ for every odd k , thus

$$n^{2m} = \sum_{k=1}^m \frac{n^{[2k]}}{(2k)!} \cdot \delta^{2k} 0^{2m}.$$

By definition of central factorial numbers of the second kind, yields

$$n^{2m} = \sum_{k=1}^m n^{[2k]} T(2m, 2k).$$

By applying identity (3.3), we get

$$\begin{aligned} n^{2m} &= \sum_{k=1}^m \frac{(2k)!}{(2k)!} n^{[2k]} T(2m, 2k) = \sum_{k=1}^m (2k)! \frac{n}{2k} \binom{n+k-1}{2k-1} T(2m, 2k) \\ &= \sum_{k=1}^m (2k-1)! n \binom{n+k-1}{2k-1} T(2m, 2k). \end{aligned}$$

By factoring out n , we obtain

$$n^{2m-1} = \sum_{k=1}^m (2k-1)! \binom{n+k-1}{2k-1} T(2m, 2k).$$

By hockey stick identity $\sum_{k=0}^n \binom{k}{j} = \binom{n+1}{j+1}$ yields

$$\Sigma^1 n^{2m-1} = \sum_{k=1}^m (2k-1)! \binom{n+k}{2k} T(2m, 2k),$$

which is exactly the Proposition (5.1). Clearly, sum of odd powers above can be generalized to multifold case, by applying hockey stick pattern repeatedly

Proposition 5.2 (Knuth's odd powers sum multifold). *For non-negative integers r, n, m*

$$\Sigma^{r+1} n^{2m-1} = \sum_{k=1}^m (2k-1)! \binom{n+k+r}{2k+r} T(2m, 2k),$$

where $T(2m, 2k)$ are central factorial numbers of the second kind.

6. GENERALIZED CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND

Definition 6.1 (Generalized central factorial numbers of the second kind). *For integers $0 \leq k \leq m$, and an arbitrary integer t , the generalized central factorial numbers of the second kind are defined by the relation*

$$n^m = \sum_{k=0}^{\infty} T_t(m, k) \cdot (n-t)^{[k]}.$$

It is also straightforward that

$$T_t(m, k) = \frac{\delta^k t^m}{k!},$$

by Newton's formula (3.2). For example,

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	0	1							
2	0	0	1						
3	0	$\frac{1}{4}$	0	1					
4	0	0	1	0	1				
5	0	$\frac{1}{16}$	0	$\frac{5}{2}$	0	1			
6	0	0	1	0	5	0	1		
7	0	$\frac{1}{64}$	0	$\frac{91}{16}$	0	$\frac{35}{4}$	0	1	
8	0	0	1	0	21	0	14	0	1

Table 1. Values of generalized central factorial numbers $T_0(n, k)$. See also the sequences [A395862](#), and [A370703](#) in the OEIS [\[14\]](#).

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	1	1							
2	1	2	1						
3	1	$\frac{13}{4}$	3	1					
4	1	5	7	4	1				
5	1	$\frac{121}{16}$	15	$\frac{25}{2}$	5	1			
6	1	$\frac{91}{8}$	31	35	20	6	1		
7	1	$\frac{1093}{64}$	63	$\frac{1491}{16}$	70	$\frac{119}{4}$	7	1	
8	1	$\frac{205}{8}$	127	$\frac{483}{2}$	231	126	42	8	1

Table 2. Values of generalized central factorial numbers $T_1(n, k)$. See also the sequences [A395860](#), and [A395861](#) in the OEIS [\[14\]](#).

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	2	1							
2	4	4	1						
3	8	$\frac{49}{4}$	6	1					
4	16	34	25	8	1				
5	32	$\frac{1441}{16}$	90	$\frac{85}{2}$	10	1			
6	64	$\frac{931}{4}$	301	190	65	12	1		
7	128	$\frac{37969}{64}$	966	$\frac{12411}{16}$	350	$\frac{371}{4}$	14	1	
8	256	$\frac{6001}{4}$	3025	3003	1701	588	126	16	1

Table 3. Values of generalized central factorial numbers $T_2(n, k)$. See also the sequences [A394466](#), and [A395314](#) in the OEIS [14].

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	3	1							
2	9	6	1						
3	27	$\frac{109}{4}$	9	1					
4	81	111	55	12	1				
5	243	$\frac{6841}{16}$	285	$\frac{185}{2}$	15	1			
6	729	$\frac{12753}{8}$	1351	585	140	18	1		
7	2187	$\frac{372709}{64}$	6069	$\frac{53011}{16}$	1050	$\frac{791}{4}$	21	1	
8	6561	$\frac{167943}{8}$	26335	$\frac{35049}{2}$	6951	1722	266	24	1

Table 4. Values of generalized central factorial numbers $T_3(n, k)$. See also the sequences [A396559](#), and [A395861](#) in the OEIS [14].

Thus, we have the formula for sums of powers in terms of generalized central factorial numbers of the second kind.

t	$T_t(2m, 2k)$	OEIS ID
0	1, 0, 1, 0, 1, 1, 0, 1, 5, 1, 0, 1, 21, 14, 1, 0, 1, 85, 147, 30, 1, $\dots$	A269945
1	1, 1, 1, 1, 7, 1, 1, 31, 20, 1, 1, 127, 231, 42, 1, 1, 511, $\dots$	A394692
2	1, 4, 1, 16, 25, 1, 64, 301, 65, 1, 256, 3025, 1701, 126, 1, $\dots$	A395456
3	1, 9, 1, 81, 55, 1, 729, 1351, 140, 1, 6561, 26335, 6951, 266, 1, $\dots$	A395457

Table 5. Sequences of $T_t(2m, 2k)$.

Proposition 6.2 (Generalized Central factorial sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} + \binom{n-t+\frac{k}{2}}{k+1} - \binom{-t+\frac{k}{2}}{k+1} \right] k! T_t(m, k).$$

7. DOUBLE AND TRIPLE SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES

Proposition 7.1 (Double sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^2 n^m = \frac{1}{2} \sum_{k=0}^m \left[+ \binom{n-t+\frac{k}{2}+2}{k+2} - \binom{2-t+\frac{k}{2}}{k+2} \Sigma^0 n^0 - \binom{1-t+\frac{k}{2}}{k+1} \Sigma^1 n^0 \right. \\ \left. + \binom{n-t+\frac{k}{2}+1}{k+2} - \binom{1-t+\frac{k}{2}}{k+2} \Sigma^0 n^0 - \binom{0-t+\frac{k}{2}}{k+1} \Sigma^1 n^0 \right] \delta^k t^m \end{aligned}$$

For example,

Example 7.2 (Double sum of squares in central differences).

$$\begin{aligned} t=0: \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(1+n)^2(2+n)}{12}, \\ t=1: \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-1+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n^2(-1+n^2)}{12}, \\ t=2: \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-4-3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(10+5n-4n^2+n^3)}{12}, \\ t=3: \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-7-6n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(32+23n-8n^2+n^3)}{12}, \\ t=4: \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-10-9n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(66+53n-12n^2+n^3)}{12}. \end{aligned}$$

Proposition 7.3 (Triple sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^3 n^m = & \frac{1}{2} \sum_{k=0}^m \left[\right. \\ & + \binom{n-t+\frac{k}{2}+3}{k+3} - \binom{3-t+\frac{k}{2}}{k+3} \Sigma^0 n^0 - \binom{2-t+\frac{k}{2}}{k+2} \Sigma^1 n^0 - \binom{1-t+\frac{k}{2}}{k+1} \Sigma^2 n^0 \\ & \left. + \binom{n-t+\frac{k}{2}+2}{k+3} - \binom{2-t+\frac{k}{2}}{k+3} \Sigma^0 n^0 - \binom{1-t+\frac{k}{2}}{k+2} \Sigma^1 n^0 - \binom{0-t+\frac{k}{2}}{k+1} \Sigma^2 n^0 \right] \delta^k t^m \end{aligned}$$

For example,

Example 7.4 (Triple sum of squares in central differences).

$$\begin{aligned} t = 0 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(6+11n+6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(18+45n+40n^2+15n^3+2n^4)}{120}, \\ t = 1 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-2-n+2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(-2-5n+5n^3+2n^4)}{120}, \\ t = 2 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-10-13n-2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(58+65n-5n^3+2n^4)}{120}, \\ t = 3 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-18-25n-6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(198+255n+40n^2-15n^3+2n^4)}{120}, \\ t = 4 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-26-37n-10n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(418+565n+120n^2-25n^3+2n^4)}{120}. \end{aligned}$$

8. MULTIFOLD SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES

Proposition 8.1 (Multifold sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k t^m. \end{aligned}$$

For example,

Example 8.2 (Examples of multifold sums).

$$\begin{aligned}
t = 0 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n+\frac{k}{2}+r}{k+r} + \binom{n+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s+\frac{k}{2}}{k+r-s} + \binom{r-1-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 0^m, \\
t = 1 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-1+\frac{k}{2}+r}{k+r} + \binom{n-1+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-1+\frac{k}{2}}{k+r-s} + \binom{r-2-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 1^m, \\
t = 2 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-2+\frac{k}{2}+r}{k+r} + \binom{n-2+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-2+\frac{k}{2}}{k+r-s} + \binom{r-3-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 2^m.
\end{aligned}$$

9. MULTIFOLD SUMS OF POWERS IN GENERALIZED CENTRAL FACTORIAL NUMBERS

Proposition 9.1 (Multifold central factorial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned}
\Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} k! T_t(m, k).
\end{aligned}$$

Example 9.2 (Sums of squares in central factorial numbers).

$$\begin{aligned}
t = 0 : \quad \Sigma^1 n^2 &= n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1, \\
t = 1 : \quad \Sigma^1 n^2 &= n \cdot 1 + \frac{1}{2}n(n-1) \cdot 2 + \frac{1}{6}n(1-3n+2n^2) \cdot 1, \\
t = 2 : \quad \Sigma^1 n^2 &= n \cdot 4 + \frac{1}{2}n(n-3) \cdot 4 + \frac{1}{6}n(13-9n+2n^2) \cdot 1, \\
t = 3 : \quad \Sigma^1 n^2 &= n \cdot 9 + \frac{1}{2}n(n-5) \cdot 6 + \frac{1}{6}n(37-15n+2n^2) \cdot 1, \\
t = 4 : \quad \Sigma^1 n^2 &= n \cdot 16 + \frac{1}{2}n(n-7) \cdot 8 + \frac{1}{6}n(73-21n+2n^2) \cdot 1, \\
t = 5 : \quad \Sigma^1 n^2 &= n \cdot 25 + \frac{1}{2}n(n-9) \cdot 10 + \frac{1}{6}n(121-27n+2n^2) \cdot 1.
\end{aligned}$$

From the example above, we may observe that

- For $t = 0$ the coefficients 0, 0, 1 correspond to the second row of the triangle (1).
- For $t = 1$ the coefficients 1, 2, 1 correspond to the second row of the triangle (2).
- For $t = 2$ the coefficients 4, 4, 1 correspond to the second row of the triangle (3).
- For $t = 3$ the coefficients 9, 6, 1 correspond to the second row of the triangle (4).

Example 9.3 (Sums of cubes in central factorial numbers).

$$\begin{aligned}
t = 0 : \quad \Sigma^1 n^4 &= n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1 \\
&\quad + \frac{1}{8}n(-1+n+4n^2+2n^3) \cdot 0 + \frac{1}{10}n(-2-5n+5n^3+2n^4) \cdot 1, \\
t = 1 : \quad \Sigma^1 n^4 &= n \cdot 1 + \frac{1}{2}n(n-1) \cdot 5 + \frac{1}{6}n(1-3n+2n^2) \cdot 7 \\
&\quad + \frac{1}{8}n(1+n-4n^2+2n^3) \cdot 4 + \frac{1}{10}n(-2+5n-5n^3+2n^4) \cdot 1, \\
t = 2 : \quad \Sigma^1 n^4 &= n \cdot 16 + \frac{1}{2}n(n-3) \cdot 34 + \frac{1}{6}n(13-9n+2n^2) \cdot 25 \\
&\quad + \frac{1}{8}n(-21+25n-12n^2+2n^3) \cdot 8 \\
&\quad + \frac{1}{10}n(18-45n+40n^2-15n^3+2n^4) \cdot 1, \\
t = 3 : \quad \Sigma^1 n^4 &= n \cdot 81 + \frac{1}{2}n(n-5) \cdot 111 + \frac{1}{6}n(37-15n+2n^2) \cdot 55 \\
&\quad + \frac{1}{8}n(-115+73n-20n^2+2n^3) \cdot 12 \\
&\quad + \frac{1}{10}n(298-275n+120n^2-25n^3+2n^4) \cdot 1
\end{aligned}$$

From the example above, we may observe that

- For $t = 0$ the coefficients 0, 0, 1, 0, 1 correspond to the fourth row of the triangle (1).
- For $t = 1$ the coefficients 1, 5, 7, 4, 1 correspond to the fourth row of the triangle (2).
- For $t = 2$ the coefficients 16, 34, 25, 8, 1 correspond to the fourth row of the triangle (3).
- For $t = 3$ the coefficients 81, 111, 55, 12, 1 correspond to the fourth row of the triangle (4).

10. MULTIFOLD SUMS OF POWERS IN BINOMIAL FORM

Lemma 10.1 (Multifold sum of zero powers). *For integers $r \geq 0$ and $n \geq 1$*

$$\Sigma^r n^0 = \binom{r+n-1}{r}$$

Proof. (1) Let $r = 0$, then we have $\Sigma^0 n^0 = 1$, by definition (2.1).

(2) Let $r = 1$, then we have $\Sigma^1 n^0 = \sum_{k=1}^n 1 = \binom{n}{1}$.

(3) Let $r = 2$, then we have $\Sigma^2 n^0 = \sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$.

(4) Let $r = 3$, then we have $\Sigma^3 n^0 = \sum_{k=1}^n \binom{k+1}{2} = \binom{n+2}{3}$.

(5) Similarly, for arbitrary integer r , by hockey stick identity $\sum_{k=1}^n \binom{k}{r} = \binom{n+1}{r+1}$, yields

$$\Sigma^r n^0 = \sum_{k=1}^n \Sigma^{r-1} k^0 = \sum_{k=1}^n \binom{k+r-2}{r-1} = \binom{r+n-1}{r}.$$

□

Proposition 10.2 (Multifold binomial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta_{t^m}^k \end{aligned}$$

For example,

Example 10.3 (Examples of multifold binomial sums).

$$\begin{aligned}
t = 0 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n+\frac{k}{2}+r}{k+r} + \binom{n+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s+\frac{k}{2}}{k+r-s} + \binom{r-1-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 0^m, \\
t = 1 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-1+\frac{k}{2}+r}{k+r} + \binom{n-1+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-1+\frac{k}{2}}{k+r-s} + \binom{r-2-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 1^m, \\
t = 2 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-2+\frac{k}{2}+r}{k+r} + \binom{n-2+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-2+\frac{k}{2}}{k+r-s} + \binom{r-3-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 2^m.
\end{aligned}$$

11. FRAMEWORK FOR SUMS OF POWERS

In this manuscript, we derive formulas for sums of powers by combining centered Newton's interpolation formula for powers (3.2) with hockey-stick identities (3.6), and (3.7). This approach can be generalized further. In particular, while using Newton's formulas

$$\begin{cases} f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^{[k]}}{k!} \delta^k f(a) = \sum_{k=0}^{\infty} \frac{x-a}{k} \binom{x-a+\frac{k}{2}-1}{k-1} \delta^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)_k}{k!} \Delta^k f(a) = \sum_{k=0}^{\infty} \binom{x-a}{k} \Delta^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^{(k)}}{k!} \nabla^k f(a) = \sum_{k=0}^{\infty} \binom{x-a+k-1}{k} \nabla^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^k}{k!} \frac{d^k f(a)}{dx^k} \end{cases} \quad (2)$$

one may replace linear difference operators δ, Δ, ∇ with their non-linear dynamic analogs, allowing more generic approach for sums of powers. It is indeed so remarkable that Taylor's series $f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^k}{k!} \frac{d^k f(a)}{dx^k}$ is a continuous analogue of Newton's formula! Back to the topic, such non-linear difference operators arise naturally in the theory of dynamic equations on time scales, developed by Bohner and Peterson in [15]. Thus, the discrete change $x + h$ of a function $f(x + h)$ is replaced by function $g(h)$ which implies that dynamic difference

operator is

$$D_g f = \frac{f(g(x)) - f(x)}{g(x) - x}. \quad (3)$$

A well-known example is Jackson's q -difference [16]

$$D_q f(x) = \frac{f(qx) - f(x)}{qx - x},$$

where $g(x) = qx$. In particular, for Newton's formula in linear differences δ, Δ, ∇ , their respective falling, rising, and central factorials are related to binomial coefficients via

$$\begin{cases} \frac{(x)_n}{n!} &= \frac{1}{n!} x(x-1)(x-2) \cdots (x-n+1) = \binom{x}{n}, \\ \frac{x^{(n)}}{n!} &= \frac{1}{n!} x(x+1)(x+2) \cdots (x+n-1) = \binom{x+n-1}{n}, \\ \frac{x^{[n]}}{n!} &= \frac{1}{n!} \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = \frac{x}{n} \binom{x+\frac{n}{2}-1}{n-1}. \end{cases} \quad (4)$$

Motivated by this viewpoint, we can see that dynamic version of Newton's formula is

$$f(x) = \sum_{k=0}^{\infty} \frac{(x \mid \alpha, \beta)_k}{k!} \cdot D_g f(a), \quad (5)$$

where a is an arbitrary integer, $D_g f(a)$ is dynamic difference operator, and $(x \mid \alpha, \beta)_n$ is generalized factorial which uses the combination of Nörlund's notation [17], and definition given by Steffensen [9]

Definition 11.1 (Generalized factorial). *For non-negative integers x, n, α, β*

$$(x \mid \alpha, \beta)_n = x(x - n\alpha - \beta)(x - n\alpha - 2\beta) \cdots (x - n\alpha - n - I\beta).$$

For instance,

$$\begin{cases} (x)_n &= (x \mid \alpha, \beta)_n, & \alpha = 0, & \beta = 1, & \text{(forward Newton's formula)} \\ x^{(n)} &= (x \mid \alpha, \beta)_n, & \alpha = 0, & \beta = -1, & \text{(backward Newton's formula)} \\ x^{[n]} &= (x \mid \alpha, \beta)_n, & \alpha = \frac{1}{2}, & \beta = 1, & \text{(central Newton's formula)} \\ x^n &= (x \mid \alpha, \beta)_n, & \alpha = 0, & \beta = 0, & \text{(Taylor's formula)} \end{cases}$$

More formally, the algorithm derive closed form formula for sums of powers is:

- Algorithm 11.2.** (1) Define the function $f(x) = x^m$.
- (2) Pick generalized difference operator $D_g f(x)$, see (3).
- (3) Derive generalized Newton's formula (5) in terms of $D_g f(x)$, for example (2).
- (4) Find an identity that connects respective factorials $(x \mid \alpha, \beta)_n$ with binomial coefficients (as per Newton's formula for powers), for example (4).
- (5) Apply hockey stick identities (3.6), and (3.7) to derive closed form for sums of powers $\Sigma^1 n^m$.
- (6) Apply identities (3.6), and (3.7) r -times to reach r -fold closed form for sums of powers $\Sigma^r n^m$.

The algorithm above has already been utilized in practice, in the works

- [18] Newtons interpolation formula and sums of powers (2026)
- [19] Sums of powers via backward finite differences and Newton's formula (2026)
- [20] Sums of powers via central finite differences and Newton's formula (2026)

The work entitled *Sums of powers of integers and generalized Stirling numbers of the second kind* [21] discusses closed formulas for sums of powers in terms of generalized Stirling numbers of the second kind, which originates from forward Newton's formula for powers, around arbitrary integer point t . It is quite remarkable that both [21], and [18] reached the same result independently. In [21] we get

$$\Sigma^1 n^k = \sum_{j=0}^k j! \left[\binom{n+1-r}{j+1} + (-1)^j \binom{r+j-1}{j+1} \right] \left\{ k \right\}_r, \quad (6)$$

$$\Sigma^1 n^k = \sum_{j=0}^k j! \left[\binom{n+1-r}{j+1} + \binom{r+1}{j+1} \right] \left\{ k \right\}_{-r}. \quad (7)$$

Formula (6) is a special case of proposition [18, Prop. (1.9)]

Proposition 11.3. For integers $n \geq 0$, and $m \geq 0$, and an arbitrary integer t

$$\Sigma^1 n^m = \sum_{j=0}^m \Delta^j t^m \left[\binom{n-t+1}{j+1} + (-1)^j \binom{j+t-1}{j+1} \right].$$

in terms of generalized Stirling numbers $\left\{ \begin{smallmatrix} k \\ j \end{smallmatrix} \right\}_r$. This implies that finite differences of powers can be expressed in terms of generalized Stirling numbers of the second kind, $\Delta^j t^m = j! \left\{ \begin{smallmatrix} m \\ j \end{smallmatrix} \right\}_t$. Formula (7) is a special case of proposition [18, Prop. (1.8)]

Proposition 11.4. *For integers $n \geq 0$, and $m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \sum_{j=0}^m \Delta^j t^m \left[\binom{n-t+1}{j+1} - \binom{1-t}{j+1} \right].$$

with setting $t = r$.

12. CONCLUSIONS

REFERENCES

- [1] David J Pengelley. The bridge between the continuous and the discrete via original sources. In *Study the Masters: The Abel-Fauvel Conference, Kristiansand*. Citeseer, 2002. <https://sites.google.com/view/davidpengelley/david-pengelley>.
- [2] František Marko and Semyon Litvinov. Geometry of figurate numbers and sums of powers of consecutive natural numbers. *The American Mathematical Monthly*, 127(1):4–22, 2020.
- [3] Johannes Faulhaber. *Academia Algebrae: darinnen dir miraculosische Gvontiones zu den höchsten Costen weiters continuirt u. profitiert werden*. Johann Remelins.
- [4] Jakob Bernoulli. *Ars coniectandi*. Impensis Thurnisiorum, fratrum, 1713.
- [5] Luschny, Peter HN. The Bernoulli Manifesto. 2021. <http://luschny.de/math/zeta/The-Bernoulli-Manifesto.html>.
- [6] Luschny, Peter HN. An introduction to the Bernoulli function. *arXiv preprint arXiv:2009.06743*, 2020. <https://arxiv.org/abs/2009.06743>.
- [7] Ira M Gessel and Xavier Viennot. Determinants, paths, and plane partitions. *preprint*, 132(197.15), 1989.
- [8] Knuth, Donald E. Johann Faulhaber and sums of powers. *Mathematics of Computation*, 61(203):277–294, 1993. <https://arxiv.org/abs/math/9207222>.
- [9] J. F. Steffensen. On the definition of the central factorial. *Journal of the Institute of Actuaries (1886-1994)*, 64(2):165–168, 1933. <https://www.jstor.org/stable/41137516>.

- [10] Paul Leo Butzer, K. Schmidt, E.L. Stark, and L. Vogt. Central factorial numbers; their main properties and some applications. *Numerical Functional Analysis and Optimization*, 10(5-6):419–488, 1989. <https://doi.org/10.1080/01630568908816313>.
- [11] Steffensen, Johan Frederik. *Interpolation*. Williams & Wilkins, 1927. <https://www.amazon.com/-/de/Interpolation-Second-Dover-Books-Mathematics-ebook/dp/B00GHQVON8>.
- [12] John Riordan. *Combinatorial identities*, volume 217. Wiley New York, 1968. <https://www.amazon.com/-/de/Combinatorial-Identities-Probability-Mathematical-Statistics/dp/0471722758>.
- [13] L. Carlitz and John Riordan. The divided central differences of zero. *Canadian Journal of Mathematics*, 15:94–100, 1963. <https://doi.org/10.4153/CJM-1963-010-8>.
- [14] Sloane, Neil J.A. and others. The on-line encyclopedia of integer sequences. <https://oeis.org/>, 2003.
- [15] Bohner, Martin and Peterson, Allan. *Dynamic equations on time scales: An introduction with applications*. Springer Science & Business Media, 2001. <https://web.mst.edu/~bohner/sample.pdf>.
- [16] Jackson, Frederick H. XI.—on q-functions and a certain difference operator. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 46(2):253–281, 1909. <https://doi.org/10.1017/S0080456800002751>.
- [17] Niels Erik Nörlund. *Vorlesungen über Differenzenrechnung*. Springer, Berlin, 1924. <https://link.springer.com/book/10.1007/978-3-642-50824-0>.
- [18] Petro Kolosov. Newton’s interpolation formula and sums of powers, January 2026. <https://doi.org/10.5281/zenodo.18040979>.
- [19] Petro Kolosov. Sums of powers via backward finite differences and newton’s formula, January 2026. <https://doi.org/10.5281/zenodo.18118011>.
- [20] Petro Kolosov. Sums of powers via central finite differences and newton’s formula, January 2026. <https://doi.org/10.5281/zenodo.18361364>.
- [21] José L. Cereceda. Sums of powers of integers and generalized stirling numbers of the second kind, 2022. <https://arxiv.org/abs/2211.11648>.

MATHEMATICA PROGRAMS

Use this [GitHub](#) repository to validate the results using Mathematica programs.

Mathematica Function	Validates / Prints
<code>MultifoldSumOfPowersRecurrence[r, n, m]</code>	Computes $\sum^r n^m$, (2.1)
<code>CentralFactorial[n, k]</code>	Computes $n^{[k]}$, (2.3)
<code>CentralFactorialsBinomialFormDecomposition[20]</code>	Validates Lem. (3.4)
<code>ValidateNewtonsFormulaForPowers[10]</code>	Validates Lem. (3.5)
<code>ValidateCenteredHockeyStickIdentity[5]</code>	Validates Lem. (3.7)
<code>ValidateOrdinarySumsOfPowersDecomposition[10]</code>	Validates Prop. (4.1)
<code>ValidateDoubleSumsOfPowersDecomposition[5]</code>	Validates Prop. (7.1)
<code>ValidateTripleSumsOfPowersDecomposition[5]</code>	Validates Prop. (7.3)
<code>ValidateMultifoldSumsOfPowers[3]</code>	Validates Prop. (8.1)