

A NEW FORMULA FOR SUMS OF POWERS USING CENTRAL NEWTON'S INTERPOLATION

PETRO KOLOSOV

ABSTRACT.

CONTENTS

Preliminaries	2
1. Historical context	11
2. Central Newton's formula	11
3. Definitions and notations	11
4. Helper lemmas	11
5. Ordinary sums of powers	11
6. Remark about central factorial numbers of the second kind	11
7. Generalized central factorial numbers of the second kind	11
8. Double and triple sums of powers in terms of central differences	11
9. Multifold sums of powers in terms of central differences	11
10. Multifold sums of powers in generalized central factorial numbers	11
11. Multifold sums of powers in binomial form	11
12. Related works	11
13. Framework for sums of powers	11
14. Conclusions	11
15. Mathematica programs	11

PRELIMINARIES

Definition 0.1 (Multifold sums of powers). *For non-negative integers r, n, m*

$$\Sigma^0 n^m = n^m,$$

$$\Sigma^1 n^m = \Sigma^0 1^m + \Sigma^0 2^m + \cdots + \Sigma^0 n^m,$$

$$\Sigma^{r+1} n^m = \Sigma^r 1^m + \Sigma^r 2^m + \cdots + \Sigma^r n^m.$$

Definition 0.2 (Central factorials). *For integers n, k*

$$n^{[k]} = \begin{cases} 0, & \text{if } k < 0, \\ 1, & \text{if } k = 0, \\ n \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = n \left(n + \frac{k}{2} - 1\right)_{k-1}, & \text{if } k > 0, \end{cases}$$

where $\left(n + \frac{k}{2} - 1\right)_{k-1} = \prod_{j=1}^{k-1} \left(n + \frac{k}{2} - j\right)$ are falling factorials.

Proposition 0.3. (Central Newton's formula [?, p. 462]). *For arbitrary integers x , and t*

$$f(x) = \sum_{k=0}^{\infty} \frac{(x-t)^{[k]}}{k!} \delta^k f(t),$$

where $\delta^k f(t)$ denotes central finite difference

$$\delta^k f(t) = \sum_{j=0}^k (-1)^j \binom{k}{j} f\left(t + \frac{k}{2} - j\right).$$

Lemma 0.4 (Binomial form of central factorials). *For integers n and $k \geq 1$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Proof. We have

$$\frac{n^{[k]}}{k!} = \frac{n}{k!} \left(n + \frac{k}{2} - 1\right)_{k-1} = \frac{n}{k(k-1)!} \left(n + \frac{k}{2} - 1\right)_{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1},$$

because of the identity in falling factorials $\frac{(x)_n}{n!} = \binom{x}{n}$. □

Lemma 0.5 (Halved central factorials). *For integers $n, k \geq 0$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \binom{n + \frac{k}{2}}{k} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k}.$$

Proof. We have $\binom{a}{k} = \frac{a}{k} \binom{a-1}{k-1}$. Let $a = n + \frac{k}{2}$, then

$$\binom{n + \frac{k}{2}}{k} = \frac{n + \frac{k}{2}}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Thus,

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \binom{n + \frac{k}{2}}{k} - \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

By moving under common denominator, and applying binomial recurrence yields

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \left[2 \binom{n + \frac{k}{2}}{k} - \binom{n + \frac{k}{2} - 1}{k-1} \right] = \frac{1}{2} \left[\binom{n + \frac{k}{2}}{k} + \binom{n + \frac{k}{2} - 1}{k} + \binom{n + \frac{k}{2} - 1}{k-1} - \binom{n + \frac{k}{2} - 1}{k-1} \right].$$

Thus, the statement follows. $\square$

Proposition 0.6 (Central Newton's formula for powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \sum_{k=0}^m \frac{(n-t)^{[k]}}{k!} \delta^k t^m.$$

Proposition 0.7 (Halved central factorial powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t + \frac{k}{2}}{k} + \binom{n-t + \frac{k}{2} - 1}{k} \right] \delta^k t^m$$

Proposition 0.8 (Generalized hockey-stick identity). *For integers a, b and j ,*

$$\sum_{k=a}^b \binom{k}{j} = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

Proof. We have, $\sum_{k=a}^b \binom{k}{j} = \binom{a}{j} + \binom{a+1}{j} + \cdots + \binom{b}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right)$. By hockey stick identity $\sum_{k=0}^n \binom{k}{j} = \binom{n+1}{j+1}$ yields,

$$\sum_{k=a}^b \binom{k}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right) = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

This completes the proof. $\square$

Lemma 0.9 (Centered hockey-stick identity). *For integers $n \geq 1$, and arbitrary integers t, k, r*

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Proof. By setting $j = 1 - t + \frac{k}{2} + r$, and $b = n - t + \frac{k}{2} + r$ yields

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k}.$$

By Generalized hockey-stick identity (0.8),

$$\sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Thus, the claim follows. □

Proposition 0.10 (Ordinary sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \delta^k t^m \sum_{j=1}^n \left[\binom{j-t+\frac{k}{2}}{k} + \binom{j-t+\frac{k}{2}-1}{k} \right]$$

Proposition 0.11 (Ordinary sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} + \binom{n-t+\frac{k}{2}}{k+1} - \binom{-t+\frac{k}{2}}{k+1} \right] \delta^k t^m$$

Proposition 0.12 (Central factorial numbers sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n+\frac{k}{2}+1}{k+1} - \binom{1+\frac{k}{2}}{k+1} + \binom{n+\frac{k}{2}}{k+1} - \binom{\frac{k}{2}}{k+1} \right] k! T(m, k),$$

where $T(n, k) = \frac{\delta^k 0^n}{k!}$ denote Central factorial numbers of the second kind (in Riordan's notation, see [?, p. 213], and [?].), defined by central Newton's formula for powers, evaluated

in $t = 0$

$$n^m = \sum_{k=0}^{\infty} T(m, k) n^{[k]}.$$

Definition 0.13 (Generalized Central factorial numbers of the second kind). *For integers $0 \leq k \leq m$, and an arbitrary integer t ,*

$$T_t(m, k) = \frac{\delta^k t^m}{k!},$$

defined by central Newton's formula for powers, evaluated at arbitrary integer t

$$n^m = \sum_{k=0}^{\infty} T_t(m, k) (n - t)^{[k]}.$$

Proposition 0.14 (Generalized Central factorial sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n - t + \frac{k}{2} + 1}{k + 1} - \binom{1 - t + \frac{k}{2}}{k + 1} + \binom{n - t + \frac{k}{2}}{k + 1} - \binom{-t + \frac{k}{2}}{k + 1} \right] k! T_t(m, k)$$

Proposition 0.15 (Double sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^2 n^m = \frac{1}{2} \sum_{k=0}^m \left[+ \binom{n - t + \frac{k}{2} + 2}{k + 2} - \binom{2 - t + \frac{k}{2}}{k + 2} \Sigma^0 n^0 - \binom{1 - t + \frac{k}{2}}{k + 1} \Sigma^1 n^0 \right. \\ \left. + \binom{n - t + \frac{k}{2} + 1}{k + 2} - \binom{1 - t + \frac{k}{2}}{k + 2} \Sigma^0 n^0 - \binom{0 - t + \frac{k}{2}}{k + 1} \Sigma^1 n^0 \right] \delta^k t^m \end{aligned}$$

Proposition 0.16 (Triple sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^3 n^m = \frac{1}{2} \sum_{k=0}^m \left[\right. \\ + \binom{n - t + \frac{k}{2} + 3}{k + 3} - \binom{3 - t + \frac{k}{2}}{k + 3} \Sigma^0 n^0 - \binom{2 - t + \frac{k}{2}}{k + 2} \Sigma^1 n^0 - \binom{1 - t + \frac{k}{2}}{k + 1} \Sigma^2 n^0 \\ \left. + \binom{n - t + \frac{k}{2} + 2}{k + 3} - \binom{2 - t + \frac{k}{2}}{k + 3} \Sigma^0 n^0 - \binom{1 - t + \frac{k}{2}}{k + 2} \Sigma^1 n^0 - \binom{0 - t + \frac{k}{2}}{k + 1} \Sigma^2 n^0 \right] \delta^k t^m \end{aligned}$$

Proposition 0.17 (Multifold sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k t^m \end{aligned}$$

Lemma 0.18 (Multifold sum of zero powers). *For integers $r \geq 0$ and $n \geq 1$*

$$\Sigma^r n^0 = \binom{r+n-1}{r}$$

Proof. (1) Let $r = 0$, then we have $\Sigma^0 n^0 = 1$, by definition (0.1).

(2) Let $r = 1$, then we have $\Sigma^1 n^0 = \sum_{k=1}^n 1 = \binom{n}{1}$.

(3) Let $r = 2$, then we have $\Sigma^2 n^0 = \sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$.

(4) Let $r = 3$, then we have $\Sigma^3 n^0 = \sum_{k=1}^n \binom{k+1}{2} = \binom{n+2}{3}$.

(5) Similarly, for arbitrary r , by hockey stick identity $\sum_{k=1}^n \binom{k}{r} = \binom{n+1}{r+1}$, thus, the claim follows $\Sigma^r n^0 = \sum_{k=1}^n \Sigma^{r-1} k^0 = \sum_{k=1}^n \binom{k+r-2}{r-1} = \binom{r+n-1}{r}$.

□

Proposition 0.19 (Multifold binomial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k t^m \end{aligned}$$

Proposition 0.20 (Multifold central factorial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} k! T_t(m, k) \end{aligned}$$

Definition 0.21.

$$R(n, t, k, r) = \binom{n - t + \frac{k}{2} + r}{k + r} + \binom{n - t + \frac{k}{2} + r - 1}{k + r} \\ - \sum_{s=0}^{r-1} \left[\binom{r - s - t + \frac{k}{2}}{k + r - s} + \binom{r - 1 - s - t + \frac{k}{2}}{k + r - s} \right] \Sigma^s n^0$$

Proposition 0.22 (Multifold sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^r n^m = \sum_{k=0}^m \frac{\delta^k t^m}{2} R(n, t, k, r)$$

Example 0.23 (Sums of squares in central differences).

$$\begin{aligned} t = 0 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(1 + n) + \frac{\delta^2 t^2}{2} \frac{n(1 + 3n + 2n^2)}{6}, \\ t = 1 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n - 1) + \frac{\delta^2 t^2}{2} \frac{n(1 - 3n + 2n^2)}{6}, \\ t = 2 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n - 3) + \frac{\delta^2 t^2}{2} \frac{n(13 - 9n + 2n^2)}{6}, \\ t = 3 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n - 5) + \frac{\delta^2 t^2}{2} \frac{n(37 - 15n + 2n^2)}{6}, \\ t = 4 : \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n - 7) + \frac{\delta^2 t^2}{2} \frac{n(73 - 21n + 2n^2)}{6}. \end{aligned}$$

Example 0.24 (Sums of cubes in central differences).

$$\begin{aligned} t = 0 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(1 + n) + \frac{\delta^2 t^3}{2} \frac{n(1 + 3n + 2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-1 + n + 4n^2 + 2n^3)}{24}, \\ t = 1 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n - 1) + \frac{\delta^2 t^3}{2} \frac{n(1 - 3n + 2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(1 + n - 4n^2 + 2n^3)}{24}, \\ t = 2 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n - 3) + \frac{\delta^2 t^3}{2} \frac{n(13 - 9n + 2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-21 + 25n - 12n^2 + 2n^3)}{24}, \\ t = 3 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n - 5) + \frac{\delta^2 t^3}{2} \frac{n(37 - 15n + 2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-115 + 73n - 20n^2 + 2n^3)}{24}, \\ t = 4 : \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n - 7) + \frac{\delta^2 t^3}{2} \frac{n(73 - 21n + 2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-329 + 145n - 28n^2 + 2n^3)}{24}. \end{aligned}$$

Example 0.25 (Double sum of squares in central differences).

$$\begin{aligned}
t = 0 : \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(1+n)^2(2+n)}{12}, \\
t = 1 : \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-1+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n^2(-1+n^2)}{12}, \\
t = 2 : \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-4-3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(10+5n-4n^2+n^3)}{12}, \\
t = 3 : \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-7-6n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(32+23n-8n^2+n^3)}{12}, \\
t = 4 : \quad \Sigma^2 n^2 &= \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-10-9n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(66+53n-12n^2+n^3)}{12}.
\end{aligned}$$

Example 0.26 (Triple sum of squares in central differences).

$$\begin{aligned}
t = 0 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(6+11n+6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(18+45n+40n^2+15n^3+2n^4)}{120}, \\
t = 1 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-2-n+2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(-2-5n+5n^3+2n^4)}{120}, \\
t = 2 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-10-13n-2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(58+65n-5n^3+2n^4)}{120}, \\
t = 3 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-18-25n-6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(198+255n+40n^2-15n^3+2n^4)}{120}, \\
t = 4 : \quad \Sigma^3 n^2 &= \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-26-37n-10n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(418+565n+120n^2-25n^3+2n^4)}{120}.
\end{aligned}$$

Lemma 0.27 (Stirling form of Binomial). *For integers n, k*

$$\binom{n}{k} = \sum_{j=0}^k \frac{(-1)^{k-j}}{k!} \begin{bmatrix} k \\ j \end{bmatrix} n^j$$

Example 0.28 (Examples of multifold binomial sums).

$$\begin{aligned}
t = 0 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n+\frac{k}{2}+r}{k+r} + \binom{n+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s+\frac{k}{2}}{k+r-s} + \binom{r-1-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 0^m, \\
t = 1 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-1+\frac{k}{2}+r}{k+r} + \binom{n-1+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-1+\frac{k}{2}}{k+r-s} + \binom{r-2-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 1^m, \\
t = 2 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-2+\frac{k}{2}+r}{k+r} + \binom{n-2+\frac{k}{2}+r-1}{k+r} \right. \\
&\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-2+\frac{k}{2}}{k+r-s} + \binom{r-3-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 2^m.
\end{aligned}$$

t	$T_t(2m, 2k)$	OEIS
0	1, 0, 1, 0, 1, 1, 0, 1, 5, 1, 0, 1, 21, 14, 1, 0, 1, 85, 147, 30, 1, $\dots$	A269945
1	1, 1, 1, 1, 7, 1, 1, 31, 20, 1, 1, 127, 231, 42, 1, 1, 511, $\dots$	A394692
2	1, 4, 1, 16, 25, 1, 64, 301, 65, 1, 256, 3025, 1701, 126, 1, $\dots$	A395456
3	1, 9, 1, 81, 55, 1, 729, 1351, 140, 1, 6561, 26335, 6951, 266, 1, $\dots$	A395457

Example 0.29 (Sums of squares in central factorial numbers).

$$t = 0 : \quad \Sigma^1 n^2 = n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1,$$

$$t = 1 : \quad \Sigma^1 n^2 = n \cdot 1 + \frac{1}{2}n(n-1) \cdot 2 + \frac{1}{6}n(1-3n+2n^2) \cdot 1,$$

$$t = 2 : \quad \Sigma^1 n^2 = n \cdot 4 + \frac{1}{2}n(n-3) \cdot 4 + \frac{1}{6}n(13-9n+2n^2) \cdot 1,$$

$$t = 3 : \quad \Sigma^1 n^2 = n \cdot 9 + \frac{1}{2}n(n-5) \cdot 6 + \frac{1}{6}n(37-15n+2n^2) \cdot 1,$$

$$t = 4 : \quad \Sigma^1 n^2 = n \cdot 16 + \frac{1}{2}n(n-7) \cdot 8 + \frac{1}{6}n(73-21n+2n^2) \cdot 1,$$

$$t = 5 : \quad \Sigma^1 n^2 = n \cdot 25 + \frac{1}{2}n(n-9) \cdot 10 + \frac{1}{6}n(121-27n+2n^2) \cdot 1.$$

Example 0.30 (Sums of cubes in central factorial numbers).

$$t = 0 : \quad \Sigma^1 n^4 = n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1$$

$$+ \frac{1}{8}n(-1+n+4n^2+2n^3) \cdot 0 + \frac{1}{10}n(-2-5n+5n^3+2n^4) \cdot 1,$$

$$t = 1 : \quad \Sigma^1 n^4 = n \cdot 1 + \frac{1}{2}n(n-1) \cdot 5 + \frac{1}{6}n(1-3n+2n^2) \cdot 7$$

$$+ \frac{1}{8}n(1+n-4n^2+2n^3) \cdot 4 + \frac{1}{10}n(-2+5n-5n^3+2n^4) \cdot 1,$$

$$t = 2 : \quad \Sigma^1 n^4 = n \cdot 16 + \frac{1}{2}n(n-3) \cdot 34 + \frac{1}{6}n(13-9n+2n^2) \cdot 25$$

$$+ \frac{1}{8}n(-21+25n-12n^2+2n^3) \cdot 8 + \frac{1}{10}n(18-45n+40n^2-15n^3+2n^4) \cdot 1,$$

$$t = 3 : \quad \Sigma^1 n^4 = n \cdot 81 + \frac{1}{2}n(n-5) \cdot 111 + \frac{1}{6}n(37-15n+2n^2) \cdot 55$$

$$+ \frac{1}{8}n(-115+73n-20n^2+2n^3) \cdot 12 + \frac{1}{10}n(298-275n+120n^2-25n^3+2n^4) \cdot 1,$$

$$t = 4 : \quad \Sigma^1 n^4 = n \cdot 256 + \frac{1}{2}n(n-7) \cdot 260 + \frac{1}{6}n(73-21n+2n^2) \cdot 97$$

$$+ \frac{1}{8}n(-329+145n-28n^2+2n^3) \cdot 16 + \frac{1}{10}n(1318-805n+240n^2-35n^3+2n^4) \cdot 1,$$

$$t = 5 : \quad \Sigma^1 n^4 = n \cdot 625 + \frac{1}{2}n(n-9) \cdot 505 + \frac{1}{6}n(121-27n+2n^2) \cdot 151$$

$$+ \frac{1}{8}n(-711+241n-36n^2+2n^3) \cdot 20 + \frac{1}{10}n(3798-1755n+400n^2-45n^3+2n^4) \cdot 1.$$

1. HISTORICAL CONTEXT
2. CENTRAL NEWTON'S FORMULA
3. DEFINITIONS AND NOTATIONS
4. HELPER LEMMAS
5. ORDINARY SUMS OF POWERS
6. REMARK ABOUT CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND
7. GENERALIZED CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND
8. DOUBLE AND TRIPLE SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES
9. MULTIFOLD SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES
10. MULTIFOLD SUMS OF POWERS IN GENERALIZED CENTRAL FACTORIAL NUMBERS
11. MULTIFOLD SUMS OF POWERS IN BINOMIAL FORM
12. RELATED WORKS
13. FRAMEWORK FOR SUMS OF POWERS
14. CONCLUSIONS
15. MATHEMATICA PROGRAMS

Use this [GitHub](#) repository to validate the results using Mathematica programs.

Mathematica Function	Validates / Prints
<code>MultifoldSumOfPowersRecurrence[r, n, m]</code>	Computes $\Sigma^r n^m$, (0.1)
<code>CentralFactorial[n, k]</code>	Computes $n^{[k]}$, (0.2)
<code>ValidateHalvedCentralBinomials[20]</code>	Validates Lem. (0.5)
<code>ValidateNewtonsFormulaForPowers[10]</code>	Validates Prop. (0.6)
<code>ValidateHalvedCentralFactorialPowers[5]</code>	Validates Prop. (0.7)
<code>ValidateCenteredHockeyStickIdentity[5]</code>	Validates Lem. (0.9)
<code>ValidateOrdinarySumsOfPowersDecomposition[10]</code>	Validates Prop. (0.11)
<code>ValidateDoubleSumsOfPowersDecomposition[5]</code>	Validates Prop. (0.15)
<code>ValidateTripleSumsOfPowersDecomposition[5]</code>	Validates Prop. (0.16)
<code>ValidateMultifoldSumsOfPowers[3]</code>	Validates Prop. (0.17)