
SUMS OF POWERS OF INTEGERS: A COMPLETE FRAMEWORK FOR CLOSED FORMULAS

PETRO KOLOSOV

ABSTRACT.

CONTENTS

1. Definitions and notations	2
2. Historical context	2
3. Central Newton's formula	2
4. Helper lemmas	3
5. Ordinary sums of powers	4
6. Remark on central factorial numbers of the second kind	6
7. Generalized central factorial numbers of the second kind	6
8. Double and triple sums of powers in terms of central differences	9
9. Multifold sums of powers in terms of central differences	10
10. Multifold sums of powers in generalized central factorial numbers	11
11. Multifold sums of powers in binomial form	13
12. Related works	15
13. Framework for sums of powers	15
14. Conclusions	15
References	15
15. Mathematica programs	16

Date: April 27, 2026.

1. DEFINITIONS AND NOTATIONS

Allow us to define the mathematical objects we utilize throughout the manuscript. For multifold sums of powers, we use the notation provided by Donald Knuth in [1], because it is simple and beautiful.

Definition 1.1 (Multifold sums of powers). *For non-negative integers r, n, m*

$$\Sigma^0 n^m = n^m,$$

$$\Sigma^1 n^m = \Sigma^0 1^m + \Sigma^0 2^m + \cdots + \Sigma^0 n^m,$$

$$\Sigma^{r+1} n^m = \Sigma^r 1^m + \Sigma^r 2^m + \cdots + \Sigma^r n^m.$$

We utilize the following notation for central factorials, see also [2, 3, 4]

Definition 1.2 (Central factorials). *For integers n, k*

$$n^{[k]} = \begin{cases} 0, & \text{if } k < 0, \\ 1, & \text{if } k = 0, \\ n \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = n \left(n + \frac{k}{2} - 1\right)_{k-1}, & \text{if } k > 0, \end{cases}$$

where $\left(n + \frac{k}{2} - 1\right)_{k-1} = \prod_{j=1}^{k-1} \left(n + \frac{k}{2} - j\right)$ are falling factorials.

2. HISTORICAL CONTEXT

3. CENTRAL NEWTON'S FORMULA

Proposition 3.1. (Central Newton's formula [2, p. 462]). *For arbitrary integers x , and t*

$$f(x) = \sum_{k=0}^{\infty} \frac{(x-t)^{[k]}}{k!} \delta^k f(t),$$

where $\delta^k f(t)$ denotes central finite difference

$$\delta^k f(t) = \sum_{j=0}^k (-1)^j \binom{k}{j} f\left(t + \frac{k}{2} - j\right),$$

and $(x-t)^{[k]}$ are central factorials defined by (1.2).

4. HELPER LEMMAS

Lemma 4.1 (Newton's formula for powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \sum_{k=0}^m \frac{(n-t)^{[k]}}{k!} \delta^k t^m.$$

Lemma 4.2 (Central factorials binomial form). *For integers n and $k \geq 1$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Proof. We have

$$\frac{n^{[k]}}{k!} = \frac{n}{k!} \binom{n + \frac{k}{2} - 1}{k-1}_{k-1} = \frac{n}{k(k-1)!} \binom{n + \frac{k}{2} - 1}{k-1}_{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1},$$

because of the identity in falling factorials $\frac{(x)_n}{n!} = \binom{x}{n}$. □

Lemma 4.3 (Central factorials binomial form decomposition). *For integers $n, k \geq 0$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \binom{n + \frac{k}{2}}{k} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k}.$$

Proof. We have $\binom{a}{k} = \frac{a}{k} \binom{a-1}{k-1}$. Let $a = n + \frac{k}{2}$, then

$$\binom{n + \frac{k}{2}}{k} = \frac{n + \frac{k}{2}}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} + \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Thus,

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \binom{n + \frac{k}{2}}{k} - \frac{1}{2} \binom{n + \frac{k}{2} - 1}{k-1}.$$

By moving under common denominator, and applying binomial recurrence yields

$$\frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} = \frac{1}{2} \left[2 \binom{n + \frac{k}{2}}{k} - \binom{n + \frac{k}{2} - 1}{k-1} \right] = \frac{1}{2} \left[\binom{n + \frac{k}{2}}{k} + \binom{n + \frac{k}{2} - 1}{k} + \binom{n + \frac{k}{2} - 1}{k-1} - \binom{n + \frac{k}{2} - 1}{k-1} \right].$$

Thus, the statement follows. □

Lemma 4.4 (Newton's formula for powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t + \frac{k}{2}}{k} + \binom{n-t + \frac{k}{2} - 1}{k} \right] \delta^k t^m.$$

Lemma 4.5 (Generalized hockey-stick identity). *For integers a, b and j ,*

$$\sum_{k=a}^b \binom{k}{j} = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

Proof. We have, $\sum_{k=a}^b \binom{k}{j} = \binom{a}{j} + \binom{a+1}{j} + \cdots + \binom{b}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right)$. By hockey stick identity $\sum_{k=0}^n \binom{k}{j} = \binom{n+1}{j+1}$ yields,

$$\sum_{k=a}^b \binom{k}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right) = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

This completes the proof. □

Lemma 4.6 (Centered hockey-stick identity). *For integers $n \geq 1$, and arbitrary integers t, k, r*

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Proof. By setting $j = 1 - t + \frac{k}{2} + r$, and $b = n - t + \frac{k}{2} + r$ yields

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k}.$$

By Generalized hockey-stick identity (4.5),

$$\sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Thus, the claim follows. □

5. ORDINARY SUMS OF POWERS

Proposition 5.1 (Ordinary sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \delta^k t^m \sum_{j=1}^n \left[\binom{j-t+\frac{k}{2}}{k} + \binom{j-t+\frac{k}{2}-1}{k} \right].$$

Proposition 5.2 (Ordinary sums of powers decomposition). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} + \binom{n-t+\frac{k}{2}}{k+1} - \binom{-t+\frac{k}{2}}{k+1} \right] \delta^k t^m.$$

For example,

Example 5.3 (Sums of squares in central differences).

$$\begin{aligned} t=0: \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(1+n) + \frac{\delta^2 t^2}{2} \frac{n(1+3n+2n^2)}{6}, \\ t=1: \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-1) + \frac{\delta^2 t^2}{2} \frac{n(1-3n+2n^2)}{6}, \\ t=2: \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-3) + \frac{\delta^2 t^2}{2} \frac{n(13-9n+2n^2)}{6}, \\ t=3: \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-5) + \frac{\delta^2 t^2}{2} \frac{n(37-15n+2n^2)}{6}, \\ t=4: \quad \Sigma^1 n^2 &= \frac{t^2}{2}(2n) + \frac{\delta t^2}{2} n(n-7) + \frac{\delta^2 t^2}{2} \frac{n(73-21n+2n^2)}{6}. \end{aligned}$$

Example 5.4 (Sums of cubes in central differences).

$$\begin{aligned} t=0: \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(1+n) + \frac{\delta^2 t^3}{2} \frac{n(1+3n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-1+n+4n^2+2n^3)}{24}, \\ t=1: \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-1) + \frac{\delta^2 t^3}{2} \frac{n(1-3n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(1+n-4n^2+2n^3)}{24}, \\ t=2: \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-3) + \frac{\delta^2 t^3}{2} \frac{n(13-9n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-21+25n-12n^2+2n^3)}{24}, \\ t=3: \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-5) + \frac{\delta^2 t^3}{2} \frac{n(37-15n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-115+73n-20n^2+2n^3)}{24}, \\ t=4: \quad \Sigma^1 n^3 &= \frac{t^3}{2}(2n) + \frac{\delta t^3}{2} n(n-7) + \frac{\delta^2 t^3}{2} \frac{n(73-21n+2n^2)}{6} + \frac{\delta^3 t^3}{2} \frac{n(-329+145n-28n^2+2n^3)}{24}. \end{aligned}$$

It is indeed interesting to notice the connection between the Proposition (5.2), and Rior-dan's central factorial numbers of the second kind, that is

Proposition 5.5 (Central factorial numbers sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n + \frac{k}{2} + 1}{k+1} - \binom{1 + \frac{k}{2}}{k+1} + \binom{n + \frac{k}{2}}{k+1} - \binom{\frac{k}{2}}{k+1} \right] k! T(m, k),$$

where $T(n, k)$ denotes the central factorial numbers of the second kind (in Riordan's notation, see [3, p. 213], and [4]).

These central factorial numbers $T(n, k)$ are defined by centered Newton's formula for powers (4.1) evaluated in zero, such that

$$n^m = \sum_{k=0}^{\infty} T(m, k) n^{[k]}.$$

It is also quite clear that

$$T(n, k) = \frac{\delta^k 0^m}{k!},$$

by Newton's formula (4.1).

6. REMARK ON CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND

7. GENERALIZED CENTRAL FACTORIAL NUMBERS OF THE SECOND KIND

Definition 7.1 (Generalized central factorial numbers of the second kind). *For integers $0 \leq k \leq m$, and an arbitrary integer t , the generalized central factorial numbers of the second kind are defined by the relation*

$$n^m = \sum_{k=0}^{\infty} T_t(m, k) \cdot (n - t)^{[k]}.$$

It is also straightforward that

$$T_t(m, k) = \frac{\delta^k t^m}{k!},$$

by Newton's formula (4.1). For example,

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	0	1							
2	0	0	1						
3	0	$\frac{1}{4}$	0	1					
4	0	0	1	0	1				
5	0	$\frac{1}{16}$	0	$\frac{5}{2}$	0	1			
6	0	0	1	0	5	0	1		
7	0	$\frac{1}{64}$	0	$\frac{91}{16}$	0	$\frac{35}{4}$	0	1	
8	0	0	1	0	21	0	14	0	1

Table 1. Values of $T_0(n, k)$ for $0 \leq k \leq n \leq 8$.

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	1	1							
2	1	2	1						
3	1	$\frac{13}{4}$	3	1					
4	1	5	7	4	1				
5	1	$\frac{121}{16}$	15	$\frac{25}{2}$	5	1			
6	1	$\frac{91}{8}$	31	35	20	6	1		
7	1	$\frac{1093}{64}$	63	$\frac{1491}{16}$	70	$\frac{119}{4}$	7	1	
8	1	$\frac{205}{8}$	127	$\frac{483}{2}$	231	126	42	8	1

Table 2. Values of $T_1(n, k)$ for $0 \leq k \leq n \leq 8$.

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	2	1							
2	4	4	1						
3	8	$\frac{49}{4}$	6	1					
4	16	34	25	8	1				
5	32	$\frac{1441}{16}$	90	$\frac{85}{2}$	10	1			
6	64	$\frac{931}{4}$	301	190	65	12	1		
7	128	$\frac{37969}{64}$	966	$\frac{12411}{16}$	350	$\frac{371}{4}$	14	1	
8	256	$\frac{6001}{4}$	3025	3003	1701	588	126	16	1

Table 3. Values of $T_2(n, k)$ for $0 \leq k \leq n \leq 8$.

n/k	0	1	2	3	4	5	6	7	8
0	1								
1	3	1							
2	9	6	1						
3	27	$\frac{109}{4}$	9	1					
4	81	111	55	12	1				
5	243	$\frac{6841}{16}$	285	$\frac{185}{2}$	15	1			
6	729	$\frac{12753}{8}$	1351	585	140	18	1		
7	2187	$\frac{372709}{64}$	6069	$\frac{53011}{16}$	1050	$\frac{791}{4}$	21	1	
8	6561	$\frac{167943}{8}$	26335	$\frac{35049}{2}$	6951	1722	266	24	1

Table 4. Values of $T_3(n, k)$ for $0 \leq k \leq n \leq 8$.

Thus, we have the formula for sums of powers in terms of generalized central factorial numbers of the second kind.

Proposition 7.2 (Generalized Central factorial sums). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} + \binom{n-t+\frac{k}{2}}{k+1} - \binom{-t+\frac{k}{2}}{k+1} \right] k! T_t(m, k).$$

83 8. DOUBLE AND TRIPLE SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES

84 **Proposition 8.1** (Double sums of powers decomposition). *For integers $n, m \geq 0$, and an*
85 *arbitrary integer t ,*

$$\begin{aligned} \Sigma^2 n^m = \frac{1}{2} \sum_{k=0}^m \left[& + \binom{n-t+\frac{k}{2}+2}{k+2} - \binom{2-t+\frac{k}{2}}{k+2} \Sigma^0 n^0 - \binom{1-t+\frac{k}{2}}{k+1} \Sigma^1 n^0 \right. \\ & \left. + \binom{n-t+\frac{k}{2}+1}{k+2} - \binom{1-t+\frac{k}{2}}{k+2} \Sigma^0 n^0 - \binom{0-t+\frac{k}{2}}{k+1} \Sigma^1 n^0 \right] \delta^k t^m \end{aligned}$$

86 For example,

Example 8.2 (Double sum of squares in central differences).

$$t = 0 : \quad \Sigma^2 n^2 = \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(1+n)^2(2+n)}{12},$$

$$t = 1 : \quad \Sigma^2 n^2 = \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-1+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n^2(-1+n^2)}{12},$$

$$t = 2 : \quad \Sigma^2 n^2 = \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-4-3n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(10+5n-4n^2+n^3)}{12},$$

$$t = 3 : \quad \Sigma^2 n^2 = \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-7-6n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(32+23n-8n^2+n^3)}{12},$$

$$t = 4 : \quad \Sigma^2 n^2 = \frac{t^2}{2} n(1+n) + \frac{\delta t^2}{2} \frac{n(-10-9n+n^2)}{3} + \frac{\delta^2 t^2}{2} \frac{n(66+53n-12n^2+n^3)}{12}.$$

87 **Proposition 8.3** (Triple sums of powers decomposition). *For integers $n, m \geq 0$, and an*
88 *arbitrary integer t ,*

$$\begin{aligned} \Sigma^3 n^m = \frac{1}{2} \sum_{k=0}^m \left[& + \binom{n-t+\frac{k}{2}+3}{k+3} - \binom{3-t+\frac{k}{2}}{k+3} \Sigma^0 n^0 - \binom{2-t+\frac{k}{2}}{k+2} \Sigma^1 n^0 - \binom{1-t+\frac{k}{2}}{k+1} \Sigma^2 n^0 \right. \\ & \left. + \binom{n-t+\frac{k}{2}+2}{k+3} - \binom{2-t+\frac{k}{2}}{k+3} \Sigma^0 n^0 - \binom{1-t+\frac{k}{2}}{k+2} \Sigma^1 n^0 - \binom{0-t+\frac{k}{2}}{k+1} \Sigma^2 n^0 \right] \delta^k t^m \end{aligned}$$

89 For example,

Example 8.4 (Triple sum of squares in central differences).

$$t = 0 : \quad \Sigma^3 n^2 = \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(6+11n+6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(18+45n+40n^2+15n^3+2n^4)}{120},$$

$$t = 1 : \quad \Sigma^3 n^2 = \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-2-n+2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(-2-5n+5n^3+2n^4)}{120},$$

$$t = 2 : \quad \Sigma^3 n^2 = \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-10-13n-2n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(58+65n-5n^3+2n^4)}{120},$$

$$t = 3 : \quad \Sigma^3 n^2 = \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-18-25n-6n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(198+255n+40n^2-15n^3+2n^4)}{120},$$

$$t = 4 : \quad \Sigma^3 n^2 = \frac{t^2}{2} \frac{n(2+3n+n^2)}{3} + \frac{\delta t^2}{2} \frac{n(-26-37n-10n^2+n^3)}{12} + \frac{\delta^2 t^2}{2} \frac{n(418+565n+120n^2-25n^3+2n^4)}{120}.$$

9. MULTIFOLD SUMS OF POWERS IN TERMS OF CENTRAL DIFFERENCES

Proposition 9.1 (Multifold sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ &\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k t^m. \end{aligned}$$

For example,

Example 9.2 (Examples of multifold sums).

$$\begin{aligned}
 t = 0 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n+\frac{k}{2}+r}{k+r} + \binom{n+\frac{k}{2}+r-1}{k+r} \right. \\
 &\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s+\frac{k}{2}}{k+r-s} + \binom{r-1-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 0^m, \\
 t = 1 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-1+\frac{k}{2}+r}{k+r} + \binom{n-1+\frac{k}{2}+r-1}{k+r} \right. \\
 &\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-1+\frac{k}{2}}{k+r-s} + \binom{r-2-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 1^m, \\
 t = 2 : \quad \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-2+\frac{k}{2}+r}{k+r} + \binom{n-2+\frac{k}{2}+r-1}{k+r} \right. \\
 &\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-2+\frac{k}{2}}{k+r-s} + \binom{r-3-s+\frac{k}{2}}{k+r-s} \right] \Sigma^s n^0 \right\} \delta^k 2^m.
 \end{aligned}$$

10. MULTIFOLD SUMS OF POWERS IN GENERALIZED CENTRAL FACTORIAL NUMBERS

Proposition 10.1 (Multifold central factorial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned}
 \Sigma^r n^m &= \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\
 &\quad \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} k! T_t(m, k).
 \end{aligned}$$

t	$T_t(2m, 2k)$	OEIS ID
0	1, 0, 1, 0, 1, 1, 0, 1, 5, 1, 0, 1, 21, 14, 1, 0, 1, 85, 147, 30, 1, $\dots$	A269945
1	1, 1, 1, 1, 7, 1, 1, 31, 20, 1, 1, 127, 231, 42, 1, 1, 511, $\dots$	A394692
2	1, 4, 1, 16, 25, 1, 64, 301, 65, 1, 256, 3025, 1701, 126, 1, $\dots$	A395456
3	1, 9, 1, 81, 55, 1, 729, 1351, 140, 1, 6561, 26335, 6951, 266, 1, $\dots$	A395457

Example 10.2 (Sums of squares in central factorial numbers).

$$t = 0 : \quad \Sigma^1 n^2 = n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1,$$

$$t = 1 : \quad \Sigma^1 n^2 = n \cdot 1 + \frac{1}{2}n(n-1) \cdot 2 + \frac{1}{6}n(1-3n+2n^2) \cdot 1,$$

$$t = 2 : \quad \Sigma^1 n^2 = n \cdot 4 + \frac{1}{2}n(n-3) \cdot 4 + \frac{1}{6}n(13-9n+2n^2) \cdot 1,$$

$$t = 3 : \quad \Sigma^1 n^2 = n \cdot 9 + \frac{1}{2}n(n-5) \cdot 6 + \frac{1}{6}n(37-15n+2n^2) \cdot 1,$$

$$t = 4 : \quad \Sigma^1 n^2 = n \cdot 16 + \frac{1}{2}n(n-7) \cdot 8 + \frac{1}{6}n(73-21n+2n^2) \cdot 1,$$

$$t = 5 : \quad \Sigma^1 n^2 = n \cdot 25 + \frac{1}{2}n(n-9) \cdot 10 + \frac{1}{6}n(121-27n+2n^2) \cdot 1.$$

We may notice that

- For $t = 0$ the coefficients ... correspond to the second row of the triangle [reference].
- For $t = 1$ the coefficients ... correspond to the second row of the triangle [reference].
- For $t = 2$ the coefficients ... correspond to the second row of the triangle [reference].
- For $t = 3$ the coefficients ... correspond to the second row of the triangle [reference].

Example 10.3 (Sums of cubes in central factorial numbers).

$$\begin{aligned}
t = 0 : \quad \Sigma^1 n^4 &= n \cdot 0 + \frac{1}{2}n(1+n) \cdot 0 + \frac{1}{6}n(1+3n+2n^2) \cdot 1 \\
&\quad + \frac{1}{8}n(-1+n+4n^2+2n^3) \cdot 0 + \frac{1}{10}n(-2-5n+5n^3+2n^4) \cdot 1, \\
t = 1 : \quad \Sigma^1 n^4 &= n \cdot 1 + \frac{1}{2}n(n-1) \cdot 5 + \frac{1}{6}n(1-3n+2n^2) \cdot 7 \\
&\quad + \frac{1}{8}n(1+n-4n^2+2n^3) \cdot 4 + \frac{1}{10}n(-2+5n-5n^3+2n^4) \cdot 1, \\
t = 2 : \quad \Sigma^1 n^4 &= n \cdot 16 + \frac{1}{2}n(n-3) \cdot 34 + \frac{1}{6}n(13-9n+2n^2) \cdot 25 \\
&\quad + \frac{1}{8}n(-21+25n-12n^2+2n^3) \cdot 8 \\
&\quad + \frac{1}{10}n(18-45n+40n^2-15n^3+2n^4) \cdot 1, \\
t = 3 : \quad \Sigma^1 n^4 &= n \cdot 81 + \frac{1}{2}n(n-5) \cdot 111 + \frac{1}{6}n(37-15n+2n^2) \cdot 55 \\
&\quad + \frac{1}{8}n(-115+73n-20n^2+2n^3) \cdot 12 \\
&\quad + \frac{1}{10}n(298-275n+120n^2-25n^3+2n^4) \cdot 1
\end{aligned}$$

We may notice that

- For $t = 0$ the coefficients ... correspond to the fourth row of the triangle [reference].
- For $t = 1$ the coefficients ... correspond to the fourth row of the triangle [reference].
- For $t = 2$ the coefficients ... correspond to the fourth row of the triangle [reference].
- For $t = 3$ the coefficients ... correspond to the fourth row of the triangle [reference].

11. MULTIFOLD SUMS OF POWERS IN BINOMIAL FORM

Lemma 11.1 (Multifold sum of zero powers). *For integers $r \geq 0$ and $n \geq 1$*

$$\Sigma^r n^0 = \binom{r+n-1}{r}$$

Proof. (1) Let $r = 0$, then we have $\Sigma^0 n^0 = 1$, by definition (1.1).

(2) Let $r = 1$, then we have $\Sigma^1 n^0 = \sum_{k=1}^n 1 = \binom{n}{1}$.

(3) Let $r = 2$, then we have $\Sigma^2 n^0 = \sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$.

(4) Let $r = 3$, then we have $\Sigma^3 n^0 = \sum_{k=1}^n \binom{k+1}{2} = \binom{n+2}{3}$.

(5) Similarly, for arbitrary r , by hockey stick identity $\sum_{k=1}^n \binom{k}{r} = \binom{n+1}{r+1}$, thus, the claim follows $\Sigma^r n^0 = \sum_{k=1}^n \Sigma^{r-1} k^0 = \sum_{k=1}^n \binom{k+r-2}{r-1} = \binom{r+n-1}{r}$.

□

Proposition 11.2 (Multifold binomial sums of powers). *For integers $n, m \geq 0$, and an arbitrary integer t ,*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-t+\frac{k}{2}+r}{k+r} + \binom{n-t+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-t+\frac{k}{2}}{k+r-s} + \binom{r-1-s-t+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k t^m \end{aligned}$$

For example,

Example 11.3 (Examples of multifold binomial sums).

$$\begin{aligned} t = 0 : \quad \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n+\frac{k}{2}+r}{k+r} + \binom{n+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s+\frac{k}{2}}{k+r-s} + \binom{r-1-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 0^m, \\ t = 1 : \quad \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-1+\frac{k}{2}+r}{k+r} + \binom{n-1+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-1+\frac{k}{2}}{k+r-s} + \binom{r-2-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 1^m, \\ t = 2 : \quad \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left\{ \binom{n-2+\frac{k}{2}+r}{k+r} + \binom{n-2+\frac{k}{2}+r-1}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left[\binom{r-s-2+\frac{k}{2}}{k+r-s} + \binom{r-3-s+\frac{k}{2}}{k+r-s} \right] \binom{s+n-1}{s} \right\} \delta^k 2^m. \end{aligned}$$

12. RELATED WORKS

13. FRAMEWORK FOR SUMS OF POWERS

14. CONCLUSIONS

REFERENCES

- [1] Knuth, Donald E. Johann Faulhaber and sums of powers. *Mathematics of Computation*, 61(203):277–294, 1993. <https://arxiv.org/abs/math/9207222>.
- [2] Paul Leo Butzer, K. Schmidt, E.L. Stark, and L. Vogt. Central factorial numbers; their main properties and some applications. *Numerical Functional Analysis and Optimization*, 10(5-6):419–488, 1989. <https://doi.org/10.1080/01630568908816313>.
- [3] John Riordan. *Combinatorial identities*, volume 217. Wiley New York, 1968. <https://www.amazon.com/-/de/Combinatorial-Identities-Probability-Mathematical-Statistics/dp/0471722758>.
- [4] L. Carlitz and John Riordan. The divided central differences of zero. *Canadian Journal of Mathematics*, 15:94–100, 1963. <https://doi.org/10.4153/CJM-1963-010-8>.

15. MATHEMATICA PROGRAMS

Use this [GitHub](#) repository to validate the results using Mathematica programs.

Mathematica Function	Validates / Prints
<code>MultifoldSumOfPowersRecurrence[r, n, m]</code>	Computes $\Sigma^r n^m$, (1.1)
<code>CentralFactorial[n, k]</code>	Computes $n^{[k]}$, (1.2)
<code>ValidateHalvedCentralBinomials[20]</code>	Validates Lem. (4.3)
<code>ValidateNewtonsFormulaForPowers[10]</code>	Validates Prop. (4.1)
<code>ValidateHalvedCentralFactorialPowers[5]</code>	Validates Prop. (4.4)
<code>ValidateCenteredHockeyStickIdentity[5]</code>	Validates Lem. (4.6)
<code>ValidateOrdinarySumsOfPowersDecomposition[10]</code>	Validates Prop. (5.2)
<code>ValidateDoubleSumsOfPowersDecomposition[5]</code>	Validates Prop. (8.1)
<code>ValidateTripleSumsOfPowersDecomposition[5]</code>	Validates Prop. (8.3)
<code>ValidateMultifoldSumsOfPowers[3]</code>	Validates Prop. (9.1)