

SUMS OF POWERS VIA CENTRAL FINITE DIFFERENCES AND NEWTON'S FORMULA

PETRO KOLOSOV

ABSTRACT.

1. INTRODUCTION

Theorem 1.1 (Newton series via central difference).

$$f(x) = \sum_{k=0}^{\infty} \frac{x^{[k]}}{k!} \delta^k f(0)$$

Theorem 1.2 (Newton series for power via central difference).

$$x^n = \sum_{k=0}^{\infty} \frac{x^{[k]}}{k!} \delta^k 0^n$$

Where

Corollary 1.3 (Central difference of power in zero).

$$\delta^k 0^n = \sum_{j=0}^k (-1)^j \binom{k}{j} \left(\frac{k}{2} - j\right)^n$$

Lemma 1.4 (Central factorial).

$$n^{[k]} = n \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = n \prod_{j=1}^{k-1} \left(n + \frac{k}{2} - j\right)$$

Such that

Date: December 29, 2025.

2010 *Mathematics Subject Classification.* 05A19, 05A10, 41A15, 11B68, 11B73, 11B83.

Key words and phrases. Sums of powers, Newton's interpolation formula, Finite differences, Binomial coefficients, Faulhaber's formula, Bernoulli numbers, Bernoulli polynomials, Interpolation, Discrete convolution, Combinatorics, Polynomial identities, Central factorial numbers, Stirling numbers, Eulerian numbers, Worpitzky identity, Pascal's triangle, OEIS.

Proposition 1.5.

$$n^{[k]} = n \left(n + \frac{k}{2} - 1 \right)_{k-1}$$

Proposition 1.6. For $k \geq 1$

$$\begin{aligned} \frac{n^{[k]}}{k!} &= \frac{1}{k!} n \left(n + \frac{k}{2} - 1 \right)_{k-1} = \frac{n}{k(k-1)!} \left(n + \frac{k}{2} - 1 \right)_{k-1} \\ &= \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1} \end{aligned}$$

Because

Proposition 1.7.

$$\frac{(x)_n}{n!} = \binom{x}{n}$$

Thus

Proposition 1.8.

$$x^n = \sum_{k=1}^n \frac{x}{k} \binom{x + \frac{k}{2} - 1}{k-1} \delta^k 0^n$$

Thus

Proposition 1.9.

$$x^{n-1} = \sum_{k=1}^n \frac{1}{k} \binom{x + \frac{k}{2} - 1}{k-1} \delta^k 0^n$$

Thus

Proposition 1.10.

$$x^n = \sum_{k=0}^n \frac{1}{k+1} \binom{x + \frac{k+1}{2} - 1}{k} \delta^{k+1} 0^{n+1}$$

Thus

Proposition 1.11.

$$\sum^r n^m = \sum_{k=0}^m \frac{1}{k+1} \binom{n + \frac{k+1}{2} - 1 + r}{k+r} \delta^{k+1} 0^{m+1}$$

Proof. By hockey stick identity. \square

Lemma 1.12 (Central factorial numbers).

$$\delta^k 0^n = k! T(n, k)$$

Thus

Proposition 1.13.

$$\sum^r n^m = \sum_{k=0}^m \frac{1}{k+1} \binom{n + \frac{k+1}{2} - 1 + r}{k+r} (k+1)! T(m+1, k+1)$$

Thus

Proposition 1.14.

$$\sum^r n^{m-1} = \sum_{k=0}^m k! \binom{n + \frac{k+1}{2} - 1 + r}{k+r} T(m, k+1)$$

2. CONCLUSIONS

3. ACKNOWLEDGEMENTS

The author is grateful to [Full Name] for his valuable contribution [contribution] about the fact that [interesting claim].

REFERENCES

- [1] Petro Kolosov. Finding the derivative of polynomials via double limit. *GithHub*, 2024. <https://kolosovpetro.github.io/pdf/FindingTheDerivativeOfPolynomialsViaDoubleLimit.pdf>.
- [2] Alekseyev, Max. MathOverflow answer 297916/113033, 2018. <https://mathoverflow.net/a/297916/113033>.
- [3] Petro Kolosov. The coefficients U(m, l, k), m = 3 defined by the polynomial identity, 2018. <https://oeis.org/A316387>.

- [4] Petro Kolosov. Entry A302971 in The On-Line Encyclopedia of Integer Sequences, 2018. <https://oeis.org/A302971>.
- [5] Petro Kolosov. Entry A304042 in The On-Line Encyclopedia of Integer Sequences, 2018. <https://oeis.org/A304042>.

Version: Local-0.1.0

License: This work is licensed under a [CC BY 4.0 License](#).

Sources: github.com/kolosovpetro/SumsOfPowersViaCentralFiniteDifferencesAndNewtonFormula

ORCID: [0000-0002-6544-8880](https://orcid.org/0000-0002-6544-8880)

Email: kolosovp94@gmail.com

Email address: kolosovp94@gmail.com

SOFTWARE DEVELOPER, DEVOPS ENGINEER

URL: <https://kolosovpetro.github.io>