

SUMS OF POWERS VIA CENTRAL FINITE DIFFERENCES AND NEWTON'S FORMULA

PETRO KOLOSOV

ABSTRACT. In this manuscript, we derive closed formulas for multifold sums of powers of integers by combining the central Newton interpolation formula with hockey-stick identities for binomial coefficients. We further obtain representations of multifold sums of powers in terms of Stirling numbers of the second kind and Eulerian numbers. Finally, we provide Wolfram Mathematica programs for the efficient verification of the derived identities.

CONTENTS

1. Auxiliary lemmas	2
2. Introduction	5
3. Main results	6
4. Conclusions	11
References	11
Mathematica programs	13

Date: July 1, 2026.

2010 *Mathematics Subject Classification.* 05A19, 05A10, 11B73, 11B83.

Key words and phrases. Sums of powers, Newton's interpolation formula, Finite differences, Binomial coefficients, Faulhaber's formula, Bernoulli numbers, Bernoulli polynomials, Interpolation, Discrete convolution, Combinatorics, Polynomial identities, Central factorial numbers, Stirling numbers, Eulerian numbers, Worpitzky identity, Pascal's triangle, OEIS.

DOI: <https://doi.org/10.5281/zenodo.18096789>

1. AUXILIARY LEMMAS

In this section, allow us to define a few lemmas, definitions, and conventions that serve as foundation for main results of this manuscript. For multifold sums of powers, we use the notation provided by Donald Knuth in [1], because it is simple and beautiful

Definition 1.1 (Multifold sums of powers). *For non-negative integers r, n, m*

$$\Sigma^0 n^m = n^m,$$

$$\Sigma^1 n^m = \Sigma^0 1^m + \Sigma^0 2^m + \cdots + \Sigma^0 n^m,$$

$$\Sigma^{r+1} n^m = \Sigma^r 1^m + \Sigma^r 2^m + \cdots + \Sigma^r n^m.$$

We utilize the following notation for falling factorials

Definition 1.2 (Falling factorials). *For integers x, n*

$$(n)_k = \begin{cases} 1, & \text{if } k = 0, \\ n(n-1)(n-2) \cdots (n-k+1) = \prod_{j=0}^{k-1} (n-j), & \text{if } k > 0, \\ 0, & \text{else.} \end{cases}$$

We utilize the following notation for central factorials

Definition 1.3 (Central factorials). *For integers n, k*

$$n^{[k]} = \begin{cases} 0, & \text{if } k < 0, \\ 1, & \text{if } k = 0, \\ n \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = n \left(n + \frac{k}{2} - 1\right)_{k-1}, & \text{if } k > 0, \end{cases}$$

where $\left(n + \frac{k}{2} - 1\right)_{k-1} = \prod_{j=0}^{k-2} \left(n + \frac{k}{2} - 1 - j\right)$ are falling factorials.

We encourage the audience to read J. F. Steffensen's work on the definition of generalized factorials [2].

Lemma 1.4 (Central Newton's formula). *For a function f defined on integers and arbitrary integers x, t*

$$f(x) = \sum_{k=0}^{\infty} \frac{(x-t)^{[k]}}{k!} \delta^k f(t),$$

where $\delta^k f(t)$ denotes the k -th central finite difference

$$\delta^k f(t) = \sum_{j=0}^k (-1)^j \binom{k}{j} f\left(t + \frac{k}{2} - j\right),$$

and $(x-t)^{[k]}$ are central factorials defined by (1.3).

Proof. See [3, p. 462], and [4, section 2, eq. (19)], and [5, section 6, eq. (21)]. □

We have

Lemma 1.5 (Newton's formula for powers). *For non-negative integers n, m , and an arbitrary integer t ,*

$$n^m = \sum_{k=0}^m \frac{(n-t)^{[k]}}{k!} \delta^k t^m.$$

Proof. Special case of Lemma (1.4) for $f(n) = n^m$. □

We have

Lemma 1.6 (Central factorials binomial form). *For integers n and $k \neq 0$,*

$$\frac{n^{[k]}}{k!} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1}.$$

Proof. We have

$$\frac{n^{[k]}}{k!} = \frac{n}{k!} \binom{n + \frac{k}{2} - 1}{k-1}_{k-1} = \frac{n}{k(k-1)!} \binom{n + \frac{k}{2} - 1}{k-1}_{k-1} = \frac{n}{k} \binom{n + \frac{k}{2} - 1}{k-1},$$

because of the identity in falling factorials $\frac{(x)_n}{n!} = \binom{x}{n}$. □

We have

Lemma 1.7 (Generalized hockey-stick identity). *For arbitrary integers a, b and j ,*

$$\sum_{k=a}^b \binom{k}{j} = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

Proof. We have, $\sum_{k=a}^b \binom{k}{j} = \binom{a}{j} + \binom{a+1}{j} + \cdots + \binom{b}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right)$. By hockey-stick identity $\sum_{k=0}^n \binom{k}{j} = \binom{n+1}{j+1}$ yields,

$$\sum_{k=a}^b \binom{k}{j} = \left(\sum_{k=0}^b \binom{k}{j} \right) - \left(\sum_{k=0}^{a-1} \binom{k}{j} \right) = \binom{b+1}{j+1} - \binom{a}{j+1}.$$

This completes the proof. $\square$

We have

Lemma 1.8 (Centered hockey-stick identity). *For integer $n \geq 1$, and arbitrary integers t, k, r*

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

Proof. By setting $a = 1 - t + \frac{k}{2} + r$, and $b = n - t + \frac{k}{2} + r$ into Lemma (1.7) yields

$$\sum_{j=1}^n \binom{j-t+\frac{k}{2}+r}{k} = \sum_{j=1-t+\frac{k}{2}+r}^{n-t+\frac{k}{2}+r} \binom{j}{k} = \binom{n-t+\frac{k}{2}+r+1}{k+1} - \binom{1-t+\frac{k}{2}+r}{k+1}.$$

This completes the proof. $\square$

We have

Lemma 1.9 (Multifold sums of zero powers). *For non-negative integers r, n*

$$\Sigma^r n^0 = \binom{r+n-1}{r}.$$

Proof.

- (1) Let $r = 0$, then we have $\Sigma^0 n^0 = 1$, by definition (1.1).
- (2) Let $r = 1$, then we have $\Sigma^1 n^0 = \sum_{k=1}^n 1 = \binom{n}{1}$.
- (3) Let $r = 2$, then we have $\Sigma^2 n^0 = \sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$.
- (4) Let $r = 3$, then we have $\Sigma^3 n^0 = \sum_{k=1}^n \binom{k+1}{2} = \binom{n+2}{3}$.

(5) Similarly, for arbitrary integer r , by hockey-stick identity $\sum_{k=1}^n \binom{k}{r} = \binom{n+1}{r+1}$, yields

$$\Sigma^r n^0 = \sum_{k=1}^n \Sigma^{r-1} k^0 = \sum_{k=1}^n \binom{k+r-2}{r-1} = \binom{r+n-1}{r}.$$

This completes the proof. $\square$

We follow the convention

Convention 1.10. *For all x*

$$x^0 = 1.$$

Donald Knuth give extensive discussion on the convention $x^0 = 1$ for all x , including zero, in Concrete Mathematics [6, p. 162], and in [7].

2. INTRODUCTION

In this manuscript, we derive formulas for sums of powers by combining central Newton's interpolation formula for powers (1.5) with hockey-stick identity (1.8). We implement the algorithm for finding closed forms of sums of powers of integers, proposed in [8, Alg. (11.2)]. More precisely, the main idea is to determine binomial basis of Newton's formula, in terms of variants of difference operators, for example

$$\left\{ \begin{array}{l} f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^{[k]}}{k!} \delta^k f(a) = \sum_{k=0}^{\infty} \frac{x-a}{k} \binom{x-a+\frac{k}{2}-1}{k-1} \delta^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)_k}{k!} \Delta^k f(a) = \sum_{k=0}^{\infty} \binom{x-a}{k} \Delta^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^{(k)}}{k!} \nabla^k f(a) = \sum_{k=0}^{\infty} \binom{x-a+k-1}{k} \nabla^k f(a), \\ f(x) = \sum_{k=0}^{\infty} \frac{(x-a)^k}{k!} \frac{d^k f(a)}{dx^k}. \end{array} \right.$$

We may observe that each variation of Newton's formula above has its binomial basis

$$\left\{ \begin{array}{l} \frac{(x)_n}{n!} = \frac{1}{n!} x(x-1)(x-2) \cdots (x-n+1) = \binom{x}{n}, \\ \frac{x^{(n)}}{n!} = \frac{1}{n!} x(x+1)(x+2) \cdots (x+n-1) = \binom{x+n-1}{n}, \\ \frac{x^{[n]}}{n!} = \frac{1}{n!} \left(n + \frac{k}{2} - 1\right) \left(n + \frac{k}{2} - 2\right) \cdots \left(n - \frac{k}{2} + 1\right) = \frac{x}{n} \binom{x+\frac{n}{2}-1}{n-1}. \end{array} \right.$$

This allows us to successfully combine Newton's formula with hockey-stick type identity, to find closed forms for sums of powers of integers. Thus, we focus on closed forms of powers sums, by utilizing central Newton's formula, and hockey-stick identity (1.8).

3. MAIN RESULTS

We start our discussion from Lemma (1.5), which yields backward Newton's interpolation formula for powers. Therefore, by applying Lemma (1.6) to Newton's interpolation formula (1.5), for non-negative integers n, m and an arbitrary integer t yields

$$\begin{aligned} n^m &= \frac{(n-t)^{[0]}}{0!} \delta^0 t^m + \sum_{k=1}^m \frac{n-t}{k} \binom{n+t+\frac{k}{2}-1}{k-1} \delta^k t^m \\ &= t^m + \sum_{k=1}^m (n-t) \binom{n-t+\frac{k}{2}-1}{k-1} \frac{\delta^k t^m}{k}. \end{aligned}$$

Note that we isolate the term $\frac{(n-t)^{[0]}}{0!} \delta^0 t^m$ to avoid division by zero. By expanding the brackets, we have

$$n^m = t^m + \sum_{k=1}^m \frac{\delta^k t^m}{k} \left[n \binom{n-t+\frac{k}{2}-1}{k-1} - t \binom{n-t+\frac{k}{2}-1}{k-1} \right].$$

Now we notice that sum above has an additional factor n in its binomial basis $n \binom{n-t+\frac{k}{2}-1}{k-1}$, preventing us from using centered hockey-stick identity (1.8) for closed form. Thus, consider the following Lemma

Lemma 3.1 (Binomial decomposition). *For non-negative integers r, n, m*

$$n \binom{n+r}{m} = (m+1) \binom{n+r}{m+1} - (r-m) \binom{n+r}{m}.$$

Proof. By expanding the brackets yields,

$$n \binom{n+r}{m} = m \binom{n+r}{m+1} + \binom{n+r}{m+1} - r \binom{n+r}{m} + m \binom{n+r}{m}.$$

Recall the extraction property of binomial coefficients $\binom{n}{k+1} = \frac{n-k}{k+1} \binom{n}{k}$. Therefore, by extraction property

$$\binom{n+r}{m+1} = \frac{n+r-m}{m+1} \binom{n+r}{m}.$$

Thus,

$$n \binom{n+r}{m} = m \frac{n+r-m}{m+1} \binom{n+r}{m} + \frac{n+r-m}{m+1} \binom{n+r}{m} - r \binom{n+r}{m} + m \binom{n+r}{m}.$$

By factoring out binomial coefficient $\binom{n+r}{m}$ yields

$$\begin{aligned} n \binom{n+r}{m} &= \binom{n+r}{m} \left[m \frac{n+r-m}{m+1} + \frac{n+r-m}{m+1} - r + m \right] \\ &= \binom{n+r}{m} \left[(m+1) \frac{n+r-m}{m+1} - r + m \right] \\ &= \binom{n+r}{m} n. \end{aligned}$$

This completes the proof. □

Hence, by setting $r \rightarrow -t + \frac{k}{2} - 1$, and $m \rightarrow k - 1$ into Lemma (3.1) gives

$$n \binom{n-t+\frac{k}{2}-1}{k-1} = k \binom{n-t+\frac{k}{2}-1}{k} + \left[t + \frac{k}{2} \right] \binom{n-t+\frac{k}{2}-1}{k-1}.$$

Because, by binomial decomposition (3.1), we have

$$\begin{aligned} n \binom{n-t+\frac{k}{2}-1}{k-1} &= k \binom{n-t+\frac{k}{2}-1}{k-1+1} - \left[-t + \frac{k}{2} - 1 - (k-1) \right] \binom{n-t+\frac{k}{2}-1}{k-1} \\ &= k \binom{n-t+\frac{k}{2}-1}{k} - \left[-t - \frac{k}{2} \right] \binom{n-t+\frac{k}{2}-1}{k-1} \\ &= k \binom{n-t+\frac{k}{2}-1}{k} + \left[t + \frac{k}{2} \right] \binom{n-t+\frac{k}{2}-1}{k-1}. \end{aligned}$$

Thus,

$$n^m = t^m + \sum_{k=1}^m \frac{\delta^k t^m}{k} \left[k \binom{n-t+\frac{k}{2}-1}{k} + \left[t + \frac{k}{2} \right] \binom{n-t+\frac{k}{2}-1}{k-1} - t \binom{n-t+\frac{k}{2}-1}{k-1} \right].$$

By collapsing the terms $t \binom{n-t+\frac{k}{2}-1}{k-1}$, and by simplifying the factors k yields

$$n^m = t^m + \sum_{k=1}^m \left[\binom{n-t+\frac{k}{2}-1}{k} + \frac{1}{2} \binom{n-t+\frac{k}{2}-1}{k-1} \right] \delta^k t^m.$$

We move the term t^m under the summation, thus

$$n^m = \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}-1}{k} + \frac{1}{2} \binom{n-t+\frac{k}{2}-1}{k-1} \right] \delta^k t^m.$$

Now, we simplify the binomial basis $\binom{n-t+\frac{k}{2}-1}{k} + \frac{1}{2}\binom{n-t+\frac{k}{2}-1}{k-1}$ to get rid of fractions inside of it. Let $a = n - t + \frac{k}{2} - 1$, then

$$\binom{a}{k} + \frac{1}{2}\binom{a}{k-1} = \frac{1}{2}\left[2\binom{a}{k} + \binom{a}{k-1}\right] = \frac{1}{2}\left[\binom{a}{k} + \binom{a+1}{k}\right].$$

Therefore,

$$n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}-1}{k} + \binom{n-t+\frac{k}{2}}{k-1} \right] \delta^k t^m.$$

Note that we do not bother ourselves about fractions in upper index of binomial coefficients $\binom{n-t+\frac{k}{2}-1}{k}$, and $\binom{n-t+\frac{k}{2}}{k-1}$, because binomial coefficient $\binom{n}{k}$ is a polynomial in n . The well-known identity in unsigned Stirling numbers of the first kind $[n]_k$ shows it explicitly

$$\binom{n}{k} = \sum_{j=0}^k \frac{(-1)^{k-j}}{k!} [k]_j n^j.$$

For example,

Example 3.2. For integer $0 \leq t \leq 3$, we have

$$\begin{aligned} t = 0 : \quad n^3 &= \left(1 + \binom{n}{-1}\right) \cdot 0 + \frac{1}{2}(1 + 2n) \cdot \frac{1}{4} + \frac{1}{2}(2 + n + n^2) \cdot 0 \\ &\quad + \frac{1}{48}(21 + 46n + 12n^2 + 8n^3) \cdot 6, \\ t = 1 : \quad n^3 &= \left(1 + \binom{n-1}{-1}\right) \cdot 1 + \frac{1}{2}(-1 + 2n) \cdot \frac{13}{4} + \frac{1}{2}(2 - n + n^2) \cdot 6 \\ &\quad + \frac{1}{48}(-21 + 46n - 12n^2 + 8n^3) \cdot 6, \\ t = 2 : \quad n^3 &= \left(1 + \binom{n-2}{-1}\right) \cdot 8 + \frac{1}{2}(-3 + 2n) \cdot \frac{49}{4} + \frac{1}{2}(4 - 3n + n^2) \cdot 12 \\ &\quad + \frac{1}{48}(-87 + 94n - 36n^2 + 8n^3) \cdot 6, \\ t = 3 : \quad n^3 &= \left(1 + \binom{n-3}{-1}\right) \cdot 27 + \frac{1}{2}(-5 + 2n) \cdot \frac{109}{4} + \frac{1}{2}(8 - 5n + n^2) \cdot 18 \\ &\quad + \frac{1}{48}(-225 + 190n - 60n^2 + 8n^3) \cdot 6. \end{aligned}$$

Hence, we can utilize centered hockey-stick identity (1.8) for closed formula for sums of powers, because

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\sum_{j=1}^n \binom{j-t+\frac{k}{2}-1}{k} + \sum_{j=1}^n \binom{j-t+\frac{k}{2}}{k-1} \right] \delta^k t^m.$$

Therefore, by Lemma (1.8)

$$\begin{cases} \sum_{j=1}^n \binom{j-t+\frac{k}{2}-1}{k} &= \binom{n-t+\frac{k}{2}+0}{k+1} - \binom{0-t+\frac{k}{2}}{k+1}, \\ \sum_{j=1}^n \binom{j-t+\frac{k}{2}}{k-1} &= \binom{n-t+\frac{k}{2}+1}{k+1} - \binom{1-t+\frac{k}{2}}{k+1}. \end{cases}$$

Thus, closed formula for ordinary sums of powers follows

Proposition 3.3 (Ordinary sums of powers). *For non-negative integers n, m , and an arbitrary integer t*

$$\Sigma^1 n^m = \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+0}{k+1} + \binom{n-t+\frac{k}{2}+1}{k+1} - \binom{0-t+\frac{k}{2}}{k+1} - \binom{1-t+\frac{k}{2}}{k+1} \right] \delta^k t^m.$$

For example,

Example 3.4. *For integer $0 \leq t \leq 4$, we have*

$$\begin{aligned} t = 0 : \quad \Sigma^1 n^3 &= 2n \cdot 0 + (n + n^2) \cdot \frac{1}{4} + \frac{1}{6}(n + 3n^2 + 2n^3) \cdot 0 \\ &\quad + \frac{1}{24}(-n + n^2 + 4n^3 + 2n^4) \cdot 6, \\ t = 1 : \quad \Sigma^1 n^3 &= 2n \cdot 1 + (-n + n^2) \cdot \frac{13}{4} + \frac{1}{6}(n - 3n^2 + 2n^3) \cdot 6 \\ &\quad + \frac{1}{24}(n + n^2 - 4n^3 + 2n^4) \cdot 6, \\ t = 2 : \quad \Sigma^1 n^3 &= 2n \cdot 8 + (-3n + n^2) \cdot \frac{49}{4} + \frac{1}{6}(13n - 9n^2 + 2n^3) \cdot 12 \\ &\quad + \frac{1}{24}(-21n + 25n^2 - 12n^3 + 2n^4) \cdot 6, \\ t = 3 : \quad \Sigma^1 n^3 &= 2n \cdot 27 + (-5n + n^2) \cdot \frac{109}{4} + \frac{1}{6}(37n - 15n^2 + 2n^3) \cdot 18 \\ &\quad + \frac{1}{24}(-115n + 73n^2 - 20n^3 + 2n^4) \cdot 6, \\ t = 4 : \quad \Sigma^1 n^3 &= 2n \cdot 64 + (-7n + n^2) \cdot \frac{193}{4} + \frac{1}{6}(73n - 21n^2 + 2n^3) \cdot 24 \\ &\quad + \frac{1}{24}(-329n + 145n^2 - 28n^3 + 2n^4) \cdot 6. \end{aligned}$$

Similarly, by setting $r = 0$, and $r = 1$ into Lemma (1.8), we have

$$\begin{cases} \sum_{j=1}^n \binom{j-t+\frac{k}{2}+0}{k+1} &= \binom{n-t+\frac{k}{2}+1}{k+2} - \binom{1-t+\frac{k}{2}}{k+2}, \\ \sum_{j=1}^n \binom{j-t+\frac{k}{2}+1}{k+1} &= \binom{n-t+\frac{k}{2}+2}{k+2} - \binom{2-t+\frac{k}{2}}{k+2}. \end{cases}$$

Hence,

Proposition 3.5 (Double sums of powers). *For non-negative integers n, m , and an arbitrary integer t*

$$\begin{aligned} \Sigma^2 n^m &= \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+1}{k+2} + \binom{n-t+\frac{k}{2}+2}{k+2} \right. \\ &\quad - \left(\binom{0-t+\frac{k}{2}}{k+1} + \binom{1-t+\frac{k}{2}}{k+1} \right) \Sigma^1 n^0 \\ &\quad \left. - \left(\binom{1-t+\frac{k}{2}}{k+2} + \binom{2-t+\frac{k}{2}}{k+2} \right) \Sigma^0 n^0 \right] \delta^k t^m. \end{aligned}$$

For example,

Example 3.6. *For integer $0 \leq t \leq 2$, we have*

$$\begin{aligned} t = 0 : \quad \Sigma^1 n^3 &= \frac{1}{2}(n + n^2) \cdot 0 + \frac{1}{48}(-3 - 2n + 12n^2 + 8n^3) \cdot \frac{1}{4} \\ &\quad + \frac{1}{24}(-2n - n^2 + 2n^3 + n^4) \cdot 0 \\ &\quad + \frac{1}{3840}(45 + 18n - 200n^2 - 80n^3 + 80n^4 + 32n^5) \cdot 6, \\ t = 1 : \quad \Sigma^1 n^3 &= \frac{1}{2}(-n + n^2) \cdot 1 + \frac{1}{48}(3 - 2n - 12n^2 + 8n^3) \cdot \frac{13}{4} \\ &\quad + \frac{1}{24}(2n - n^2 - 2n^3 + n^4) \cdot 6 \\ &\quad + \frac{1}{3840}(-45 + 18n + 200n^2 - 80n^3 - 80n^4 + 32n^5) \cdot 6, \\ t = 2 : \quad \Sigma^1 n^3 &= \frac{1}{2}(2 - 3n + n^2) \cdot 8 + \frac{1}{48}(-15 + 46n - 36n^2 + 8n^3) \cdot \frac{49}{4} \\ &\quad + \frac{1}{24}(-6n + 11n^2 - 6n^3 + n^4) \cdot 12 \\ &\quad + \frac{1}{3840}(105 - 142n - 360n^2 + 560n^3 - 240n^4 + 32n^5) \cdot 6. \end{aligned}$$

Thus, in general

Theorem 3.7 (Multifold sums of powers). *For non-negative integers r, n, m , and an arbitrary integer t*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+r-1}{k+r} + \binom{n-t+\frac{k}{2}+r}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left(\binom{s-t+\frac{k}{2}}{k+s+1} + \binom{s-t+\frac{k}{2}+1}{k+s+1} \right) \Sigma^{r-1-s} n^0 \right] \delta^k t^m. \end{aligned}$$

By Lemma (1.9) pure binomial form follows

Proposition 3.8 (Multifold Binomial sums of powers). *For non-negative integers r, n, m , and an arbitrary integer t*

$$\begin{aligned} \Sigma^r n^m = & \frac{1}{2} \sum_{k=0}^m \left[\binom{n-t+\frac{k}{2}+r-1}{k+r} + \binom{n-t+\frac{k}{2}+r}{k+r} \right. \\ & \left. - \sum_{s=0}^{r-1} \left(\binom{s-t+\frac{k}{2}}{k+s+1} + \binom{s-t+\frac{k}{2}+1}{k+s+1} \right) \binom{r-s+n-2}{r-s-1} \right] \delta^k t^m. \end{aligned}$$

Therefore, we have successfully derived multifold formulas for sums of powers, in closed form. The work [9] provides implementation of algorithm proposed in [8, Alg. (11.2)], in terms of forward finite differences Δ , and Newton's formula. Similarly, the work [10] discusses multifold formulas for sums of powers in terms of backward differences ∇ , and Newton's formula.

4. CONCLUSIONS

In this manuscript, we derive closed formulas for multifold sums of powers of integers by combining the central Newton interpolation formula with hockey-stick identities for binomial coefficients. We further obtain representations of multifold sums of powers in terms of Stirling numbers of the second kind and Eulerian numbers. Finally, we provide Wolfram Mathematica programs for the efficient verification of the derived identities.

REFERENCES

- [1] Knuth, Donald E. Johann Faulhaber and sums of powers. *Mathematics of Computation*, 61(203):277–294, 1993. <https://arxiv.org/abs/math/9207222>.

- [2] J. F. Steffensen. On the definition of the central factorial. *Journal of the Institute of Actuaries (1886-1994)*, 64(2):165–168, 1933. <https://www.jstor.org/stable/41137516>.
- [3] Paul Leo Butzer, K. Schmidt, E.L. Stark, and L. Vogt. Central factorial numbers; their main properties and some applications. *Numerical Functional Analysis and Optimization*, 10(5-6):419–488, 1989. <https://doi.org/10.1080/01630568908816313>.
- [4] Steffensen, Johan Frederik. *Interpolation*. Williams & Wilkins, 1927. <https://www.amazon.com/-/de/Interpolation-Second-Dover-Books-Mathematics-ebook/dp/B00GHQVON8>.
- [5] John Riordan. *Combinatorial identities*, volume 217. Wiley New York, 1968. <https://www.amazon.com/-/de/Combinatorial-Identities-Probability-Mathematical-Statistics/dp/0471722758>.
- [6] Graham, Ronald L. and Knuth, Donald E. and Patashnik, Oren. *Concrete mathematics: A foundation for computer science (second edition)*. Addison-Wesley Publishing Company, Inc., 1994. <https://archive.org/details/concrete-mathematics>.
- [7] Donald E. Knuth. Two notes on notation, 1992. <https://arxiv.org/abs/math/9205211>.
- [8] Petro Kolosov. Sums of powers of integers: A complete framework for closed formulas, June 2026. <https://doi.org/10.5281/zenodo.20548019>.
- [9] Petro Kolosov. Newton's interpolation formula and sums of powers, December 2025. <https://doi.org/10.5281/zenodo.18040979>.
- [10] Petro Kolosov. Sums of powers via backward finite differences and Newton's formula, January 2026. <https://doi.org/10.5281/zenodo.18118011>.

MATHEMATICA PROGRAMS

Use this [GitHub](#) repository to validate the results using Mathematica programs.

Mathematica Function	Validates / Prints
<code>ValidateOrdinarySumsOfPowers.txt</code>	Validates Prop. (3.3)
<code>ValidateDoubleSumsOfPowers.txt</code>	Validates Prop. (3.5)
<code>ValidateMultifoldSumsOfPowers.txt</code>	Validates Thm. (3.7)
<code>ValidateMultifoldBinomialSumsOfPowers.txt</code>	Validates Prop. (3.8)

Metadata

- **Initial release date:** January 3, 2026.
- **Current release date:** July 1, 2026.
- **Version:** Local-0.1.0
- **MSC2010:** 05A19, 05A10, 11B73, 11B83 .
- **Keywords:** Sums of powers, Newton's interpolation formula, Finite differences, Binomial coefficients, Faulhaber's formula, Bernoulli numbers, Bernoulli polynomials, Interpolation, Discrete convolution, Combinatorics, Polynomial identities, Central factorial numbers, Stirling numbers, Eulerian numbers, Worpitzky identity, Pascal's triangle, OEIS .
- **License:** This work is licensed under a [CC BY 4.0 License](#).
- **DOI:** <https://doi.org/10.5281/zenodo.18096789>
- **Web Version:** kolosovpetro.github.io/sums-of-powers-central-differences/
- **Sources:** [github/kolosovpetro/SumsOfPowersCentralDifferencesNewtonFormula](https://github.com/kolosovpetro/SumsOfPowersCentralDifferencesNewtonFormula)
- **ORCID:** [0000-0002-6544-8880](https://orcid.org/0000-0002-6544-8880)
- **Email:** kolosovp94@gmail.com

DEVOPS ENGINEER

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>