LATEX TEMPLATE FOR GITHUB

PETRO KOLOSOV

Abstract. Your abstract here.

Contents

1. Introduction 1
1.1. Rising factorials 4
1.2. Central factorials 4
1.3. Derivatives 4
Conclusions 5
Acknowledgements 5
References 5
Mathematica 6

1. Introduction

Include some references [1, 2, 3]. Lorem Ipsum is simply dummy text of the printing and typesetting industry. Lorem Ipsum has been the industry's standard dummy text ever since the 1500s, when an unknown printer took a galley of type and scrambled it to make a type

Date: November 5, 2026.

2010 *Mathematics Subject Classification.* 05A19, 05A10, 41A15, 11B68, 11B73, 11B83.

Key words and phrases. Sums of powers, Newton's interpolation formula, Finite differences, Binomial coefficients, Faulhaber's formula, Bernoulli numbers, Bernoulli polynomials, Interpolation, Approximation, Discrete convolution, Combinatorics, Polynomial identities, Central factorial numbers, Stirling numbers, Eulerian numbers, Worpitzky identity, Pascal's triangle, OEIS.

DOI: https://doi.org/10.5281/zenodo.18040979

specimen book. It has survived not only five centuries, but also the leap into electronic typesetting, remaining essentially unchanged. It was popularised in the 1960s with the release of Letraset sheets containing Lorem Ipsum passages, and more recently with desktop publishing software like Aldus PageMaker including versions of Lorem Ipsum.

Image example




Figure 1. Image example (from caption).

m/r	0	1	2	3	4	5	6	7
0	1							
1	1	6						
2	1	0	30					
3	1	-14	0	140				
4	1	-120	0	0	630			
5	1	-1386	660	0	0	2772		
6	1	-21840	18018	0	0	0	12012	
7	1	-450054	491400	-60060	0	0	0	51480

Table 1. Coefficients $\mathbf{A}_{m,r}$. See OEIS sequences [4, 5].

And for any natural m we have polynomial identity

$$x^m = \sum_{k=1}^{m} T(m,k) x^{[k]} \tag{1}$$

where $x^{[k]}$ denotes central factorial defined by

$$x^{[n]} = x\left(x + \frac{n}{2} - 1\right)_{n-1}$$

where $(n)_k = n(n-1)(n-2)\cdots(n-k+1)$ denotes falling factorial in Knuth's notation. In particular,

$$x^{[n]} = x\left(x + \frac{n}{2} - 1\right)\left(x + \frac{n}{2} - 1\right)\cdots\left(x + \frac{n}{2} - n - 1\right) = x\prod_{k=1}^{n-1}\left(x + \frac{n}{2} - k\right) \tag{2}$$

This is an equation reference (1).

Continuing similarly, we are able to derive the formula for multifold sums of powers, which is

Theorem 1.1 (Multifold sums of powers via Newton's series)**.** *For non-negative integers r, n, m and an arbitrary integer t*

$$\Sigma^r\, n^m = \sum_{j=0}^{m} \Delta^j t^m \left[\left(\sum_{s=1}^{r} (-1)^{j+s-1} \binom{j+t-1}{j+s} \Sigma^{r-s}\, n^0\right) + \binom{n-t+r}{j+r}\right]$$

Proof. By Newton's series for power and repeated applications of the segmented hockey stick identity. □

Proposition 1.2 (Falling factorial)**.**

$$(x)_n = x(x-1)(x-2)(x-3)\cdots(x-n+1) = \prod_{k=0}^{n-1}(x-k)$$

Proposition 1.3.

$$\frac{(x)_n}{n!} = \binom{x}{n}$$

1.1. **Rising factorials.**

Proposition 1.4 (Rising factorial)**.**

$$x^{(n)} = x(x+1)(x+2)(x+3)\cdots(x+n-1) = \prod_{k=0}^{n-1}(x+k)$$

Proposition 1.5.

$$\frac{x^{(n)}}{n!} = \binom{x+n-1}{n}$$

1.2. **Central factorials.**

Lemma 1.6 (Central factorial)**.**

$$n^{[k]} = n\left(n+\frac{k}{2}-1\right)\left(n+\frac{k}{2}-2\right)\cdots\left(n-\frac{k}{2}+1\right) = n\prod_{j=1}^{k-1}\left(n+\frac{k}{2}-j\right)$$

Proposition 1.7.

$$n^{[k]} = n\left(n+\frac{k}{2}-1\right)_{k-1}$$

1.3. **Derivatives.**

$$\frac{\mathrm{d}x}{\mathrm{d}y} = \frac{f(x+h)-f(x)}{h}$$

$$\frac{\mathrm{d}^3x}{\mathrm{d}y^3} = \frac{f(x+h)-f(x)}{h}$$

Conclusions

Conclusions of your manuscript.

Here is an itemize list with adjusted margins

- Conclusion 1 ....
- Conclusion 2 ....
- Conclusion 3 ....

Acknowledgements

The author is grateful to [Full Name] for his valuable contribution [contribution] about the fact that [interesting claim].

References

[1] Petro Kolosov. Finding the derivative of polynomials via double limit. *GitHub*, 2024.

[2] Alekseyev, Max. MathOverflow answer 297916/113033. https://mathoverflow.net/a/297916/113033, 2018.

[3] Petro Kolosov. The coefficients U(m, l, k), m = 3 defined by the polynomial identity. https://oeis.org/A316387, 2018.

[4] Petro Kolosov. Entry A302971 in The On-Line Encyclopedia of Integer Sequences. https://oeis.org/A302971, 2018.

[5] Petro Kolosov. Entry A304042 in The On-Line Encyclopedia of Integer Sequences. https://oeis.org/A304042, 2018.

Mathematica

Use this GitHub repository to validate the results using Mathematica programs.

Mathematica Function	**Validates / Prints**
`ValidateOrdinarySumsOfPowers.txt`	Validates Prop. (**??**)
`ValidateDoubleSumsOfPowers.txt`	Validates Prop. (**??**)
`ValidateMultifoldSumsOfPowers.txt`	Validates Thm. (**??**)
`ValidateMultifoldBinomialSumsOfPowers.txt`	Validates Prop. (**??**)
`ValidateOrdinarySumsOfPowersAtZero.txt`	Validates Prop. (**??**)
`ValidateDoubleSumsOfPowersAtZero.txt`	Validates Prop. (**??**)
`ValidateMultifoldSumsOfPowersAtZero.txt`	Validates Thm. (**??**)
`ValidateMultifoldSumsOfPowersAtZeroBinomialForm.txt`	Validates Thm. (**??**)

Metadata

- **Initial release date:** December 24, 2025.
- **Current release date:** November 5, 2026.
- **Version:** `2.2.0+main.eb7888f`
- **MSC2010:** 05A19, 05A10, 41A15, 11B68, 11B73, 11B83
- **Keywords:** Sums of powers, Newton's interpolation formula, Finite differences, Binomial coefficients, Faulhaber's formula, Bernoulli numbers, Bernoulli polynomials, Interpolation, Approximation, Discrete convolution, Combinatorics, Polynomial identities, Central factorial numbers, Stirling numbers, Eulerian numbers, Worpitzky identity, Pascal's triangle, OEIS

- **License:** This work is licensed under a CC BY 4.0 License.

- **DOI:** https://doi.org/10.5281/zenodo.18040979
- **Web Version:** kolosovpetro.github.io/math/test
- **Sources:** github.com/kolosovpetro/github-latex-template

- **ORCID:** 0000-0002-6544-8880
- **Email:** kolosovp94@gmail.com

DevOps Engineer

Email address: kolosovp94@gmail.com

URL: https://kolosovpetro.github.io