

A large blue circular graphic on the left side of the slide, partially cut off by the edge.

# Statistics for Data Science

## Hypothesis Testing

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# Agenda - Hypothesis Testing

1. Hypothesis Testing
  - a. Introduction
  - b. Hypothesis Formulation
2. Basic concepts of Hypothesis Testing
  - a. Importance of null
  - b. Importance of test statistic
  - c. Type I and Type 2 errors
  - d. Hypothesis testing template
3. Performing a Hypothesis Test
  - a. Some key ideas
  - b. Assumptions
  - c. Critical point
  - d. Rejection region approach
  - e. p-value approach
4. One-Tailed and Two-Tailed Tests
5. Confidence Interval and Hypothesis Test

# Real World Problem

Suppose you are a quality analyst at a bulb manufacturing company and analyze the reliability of bulbs. Historically, 70% of the bulbs pass the reliability test.


Now, a slightly altered manufacturing process(B) has been introduced to produce the bulbs.

Can you conclude whether the new process improves the reliability of the bulbs or not by checking the number of reliable bulbs in a sample?

# Gathering evidence for statistical Inference

We selected a random sample of 100 bulbs out of which 73 are reliable. Does this provide strong evidence that the new manufacturing process is more reliable?

If the new manufacturing process was only as good as the current process - What is the probability of getting 73 or more reliable bulbs in a sample of 100 bulbs?




The probability of getting 73 or more reliable bulbs in a sample of 100 bulbs is  $\sim 0.30$ .



Thus, there is no strong evidence that the new process improves reliability

# Gathering evidence for statistical Inference

A similar experiment was run with yet another manufacturing process (C). A sample of 100 bulbs produced using this process had 81 reliable bulbs.



The probability of getting 81 or more reliable bulbs in a sample of 100 bulbs is  $\sim 0.01$ .



Thus, there is strong evidence that the new process improves reliability

# Why Hypothesis?

## Estimation

The problem of estimation is considered, when there is no previous knowledge of the population parameter. The problem is simpler in that case. A random sample is taken, a sample statistic is computed and an appropriate point and interval estimate is suggested.

## Hypothesis Testing

Often the interest is not in the numerical value of the point estimate of the parameter, but in knowing the plausibility of a hypothesis about the population parameter by using sample data. Estimation is not enough to arrive at a conclusion in such cases.

# What is Hypothesis?

Often we are interested in population parameter(s)



**A hypothesis is a conjecture about the population parameter(s)**



For example, a bulb manufacturing company is interested in knowing whether the new manufacturing process improves reliability of the bulbs.



The objective of the Hypothesis Testing is to SET a value for the parameter(s) and perform a statistical TEST to see whether that value is tenable in the light of the evidence gathered from the sample.

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# Overview of Applications

## Applications of Hypothesis Testing

```
graph TD; A[Applications of Hypothesis Testing] --> B[Testing Research Hypotheses]; A --> C[Testing the validity of a claim]; A --> D[Testing the business decisions]; B --> E[e.g. a new automobile system increases the mean mpg performance]; C --> F[e.g. a manufacturer claims that 1L soft drink bottles are filled with an average of at least 0.99L]; D --> G[e.g. new online ad has resulted in higher online conversion rates for an E-commerce website];
```

**Testing Research Hypotheses**

e.g. a new automobile system increases the mean mpg performance

**Testing the validity of a claim**

e.g. a manufacturer claims that 1L soft drink bottles are filled with an average of at least 0.99L

**Testing the business decisions**

e.g. new online ad has resulted in higher online conversion rates for an E-commerce website



# Stating the Hypothesis

**Null and Alternative Hypotheses - Two mutually exclusive statements about the population parameter(s)**

```
graph TD; A[Null and Alternative Hypotheses - Two mutually exclusive statements about the population parameter(s)] --> B[Null Hypothesis (H0)]; A --> C[Alternative Hypothesis (Ha)];
```

## **Null Hypothesis ( $H_0$ )**

The presumed current state of the matter or status quo.

E.g. The new process for manufacturing bulbs does not improve reliability.

## **Alternative Hypothesis ( $H_a$ )**

The rival opinion or research hypothesis or an improvement target.

E.g. The new process for manufacturing bulbs improves reliability.

## Null & Alternative Formulation : Example

Mean length of lumber is specified to be 8.5m for a certain building project. A construction engineer wants to make sure that the shipments she received adhere to that specification.



The population parameter about which the hypothesis will be formed is **population mean  $\mu$** .



The hypotheses are

$$H_0 : \mu = 8.5$$

$$H_a : \mu \neq 8.5$$

## Null & Alternative Formulation : Example

There is a belief that 20% of men on business travel abroad brings a significant other with them. A chain hotel claims that number is too low.



The population parameter about which the hypothesis will be formed is **population proportion  $\pi$** .



The hypotheses are

$$H_0 : \pi = 0.2$$

$$H_a : \pi > 0.2$$

# Tips to formulate Null & Alternative

Am I testing a status quo that already exists?



Null Hypothesis



Negation of the research question



Always contains equality ( $=$ ,  $\geq$ ,  $\leq$ )

Am I testing an assumption or claim that is beyond what I know?



Alternate Hypothesis



Research question to be proven



Doesn't contain equality ( $\neq$ ,  $>$ ,  $<$ )



# Basic Concepts of Hypothesis Testing

# Importance of Null

Null hypothesis is assumed to be true unless reasonably strong evidence to the contrary is found.

Based on a random sample a decision is made whether there exists reasonably strong evidence against the null hypothesis.

**Evidence is strong** (satisfies the predetermined decision rule)



**Reject the null hypothesis**  
in favour of alternative hypothesis

**Evidence is not strong** (does not satisfy the predetermined decision rule)



**Fail to reject the null hypothesis**  
in favour of alternative hypothesis

# Importance of Test Statistic

The test statistic is calculated from the sample data and tested against the predetermined Decision Rule.

The test statistic is a random variable that follows a standard distribution such as Normal, T, F, Chi-square etc. Sometimes the tests are named after the test statistic


Since hypothesis testing is done on the basis of sampling distribution, the decisions made are probabilistic.

Hence, it is very important to understand the errors associated with hypothesis testing.

# Type I and Type II Error



# Type I and Type II Errors



# Type I and Type II Errors : Example

**Null Hypothesis:** The patient doesn't have cancer

**Alternate Hypothesis:** The patient has cancer

▶ **Type I error (false positive):** “The patient doesn't have cancer but doctors says she does”

▶ **Type II error (false negative):** “The patient does have cancer but report says she doesn't”



# Template for Hypothesis Testing

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# Hypothesis Testing Template

1	Identify the key question	<i>What is the research question that you are trying to answer?</i>
2	Establish the hypotheses	<i>What is the metric of interest? Define the Null and Alternate Hypothesis.</i>
3	Understand and prepare data	<i>What data do you have? Do you understand what it means? Can it be used directly?</i>
4	Identify the right test	<i>Choose the method for testing based on the last three points</i>
5	Check the assumptions	<i>Ensure that data satisfies the assumption for the test.</i>
6	Perform the test	<i>Get to conclusion based on the results (p-value)</i>



# Performing a hypothesis test

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# Some key ideas first

Level of  
Significance ( $\alpha$ )



- Probability of rejecting the null hypothesis when it is true
- Fixed before the hypothesis test.

p-value



- Probability of observing test statistic or more extreme results than the computed test statistic, under the null hypothesis.
- Depends on the sample data. Alpha is pre-fixed but p-value depends on the value of the test statistic

Acceptance or  
Rejection Region



- The total area under the distribution curve of the test statistic is partitioned into acceptance and rejection region
- Reject the null hypothesis when the test statistic lies in the rejection region, Else we fail to reject it

# Let's start simple

Consider the following questions in hypothesis testing

What are the null and alternative hypotheses?

What is an appropriate test statistic?

What is preset level of significance?

How to check whether the data is giving significant evidence against the null hypothesis or not?

Let's see an example and understand the significance of the above questions



For simplicity, we will assume that the population standard deviation is known and the sample size is more than 30.

## Example

It is known from experience that for a certain E-commerce company the mean delivery time of the products is 5 days with a standard deviation of 1.3 days.

The new customer service manager of the company is afraid that the company is slipping and collects a random sample of 45 orders. The mean delivery time of these samples comes out to be 5.25 days.

Is there enough statistical evidence for the manager's apprehension that the mean delivery time of products is greater than 5 days.

**This is clearly a one-tailed test, concerning population mean  $\mu$ , the mean delivery time of products.**

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.



# First test - z-test for One Mean


Significance of the test	Assumptions	Test Statistic Distribution
Test for population mean $H_0 : \mu = \mu_0$	<ul style="list-style-type: none"><li>• Continuous data</li><li>• Normally distributed population or sample size <math>&gt; 30</math></li><li>• Known population standard deviation <math>\sigma</math></li><li>• Random sampling from the population</li></ul>	Standard Normal distribution



# One-tailed and Two-tailed Tests

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# One-tailed and Two-tailed Tests




Choice of One tailed vs Two tailed depends on the nature of the problem, not on the sample data!


# Difference between One-tailed and Two-tailed Tests

Test statistic value **does not change** for two-tailed or one-tailed test.

Only the critical value(s) / p-value associated with the test statistic changes



The difference is not tested on this side and the hypothesis test has greater power on the other side.



The difference is tested on both the sides.

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.



# Connecting the dots with Confidence Intervals

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.

# Confidence Interval vs Hypothesis Testing

Suppose we calculate the  $(100 - 5)\%$  confidence interval for the mean

We also conduct the Z-test for the mean with a 5% significance level.

The hypotheses of the Z-test are

$$H_0 : \mu = \mu_0 \text{ against } H_a : \mu \neq \mu_0$$

Is there any relationship between the estimated confidence interval and the hypothesis test?

The confidence interval contains all values of  $\mu_0$  for which the null hypothesis will not be rejected.

This file is meant for personal use by collinsbenda360@gmail.com only.  
Sharing or publishing the contents in part or full is liable for legal action.