

Unit - 2

Set Theory

Set: A collection of well-defined objects is called a set.

Elements of a set are usually denoted by lower-case letters while sets are denoted by capital letters

Ex: The vowels

$$V = \{a, e, i, o, u\}$$

Set of natural numbers

$$N = \{1, 2, 3, 4, \dots\}$$

Set of whole numbers

$$W = \{0, 1, 2, 3, \dots\}$$

There are 2 types of set

Notations

1) Roster form

2) Set builder form

1) Roster form

It is the list of all elements of the set.

2) Set-builder form

All the elements of the set must have same property.

Ex. $S = \{1, 3, 5, 7, \dots\}$
All odd numbers.

$S = \{x/x \text{ is an odd number}\}$

cardinality of set:

The no. of elements present in the set. Denoted as $|S|$.

Ex: 1) Let 'S' be the set of odd +ve numbers less than 10.

$S = \{1, 3, 5, 7, 9\}$

the cardinality of set is

$$|S| = 5.$$

2) Let 'A' be the set of letters of English alphabets. Then cardinality of A is $|A| = 26$.

Subset [⊆]

If every element in set A is also the element of set B, then A is called subset of B.

It is denoted by $A \subseteq B$

Ex:- $A = \{1, 2, 3\}$, $B = \{1, 2, 3, 4, 5\}$

$A \subseteq B$

But $B \not\subseteq A$

Note:- Null set is a subset

of every set.

Equality of sets

TWO sets A & B are equal if and only if $A \subseteq B$ & $B \subseteq A$.

Ex:- $A = \{1, 2, 3\}$, $B = \{1, 2, 3\}$

$\therefore A \subseteq B$ & $B \subseteq A$

\therefore We can say that $A = B$.

Types of sets

Empty set or void set or null set

$$\phi = \{\} \text{ or } \emptyset$$

No. elements in the set.

Finite set: A set with countable number of elements.

$$\text{Ex: } A = \{1, 2, 3\}$$

Infinite set: A set with uncountable number of elements.

$$\text{Ex: } A = \{1, 2, 3, \dots\}$$

Singleton set: A set with only one element. Ex: $A = \{1\}$

Proper subset :- $[P \subset J]$

$A \subset B$ $\Leftrightarrow A \neq B$ & A is a subset of B .

If A is

Proper subset :- $[S \subset A]$ subset of A

If $A \subset B$ and $B \neq A$ then

$A \subset B$

$[S \subset A]$ subset of A

$$\text{Ex: } A = \{1, 2, 3\}, B = \{1, 2, 3, 4\}$$

$\therefore A \subset B$ & $B \neq A$ if

$\therefore A \subset B$



Power set: It is the set of all subsets of A.

Denoted by $P(A)$.

Ex: If $A = \{1, 2\} \rightarrow 2^2 \Rightarrow 2^2 = 4$

then $P(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$

Universal set: Example problem.

It is the union of all the sets denoted by \cup (or \in).

Ex: If $A = \{1, 2\}, B = \{3, 4\}, C = \{a, b\}$

then $U = \{1, 2, 3, 4, a, b\}$

Set operations

Union $[A \cup B]$

All elements in A or B.

Intersection $[A \cap B]$

and common elements

Difference $[A - B]$

Elements in A but not

in B. $A - B \neq B - A$

complement (A^c or A'):

Elements in universal set U but not in A . [$A^c = U - A$]

Symmetric difference:

$$A \Delta B = (A - B) \cup (B - A)$$

Ex:- Let $A = \{1, 2, 3, 4\}$, $B = \{3, 4, 5, 6\}$

$$A \cup B = \{1, 2, 3, 4, 5, 6\}$$

$$A \cap B = \{3, 4\}$$

$$A - B = \{1, 2\}$$

$$B - A = \{5, 6\}$$

$$A \Delta B = (A - B) \cup (B - A)$$

$$A \Delta B = \{1, 2, 5, 6\}$$


A^c = If $U = \{1, 2, 3, 4, 5\}$, $A = \{1, 3, 5\}$

$$A^c = U - A = \{2, 4\}$$


Venn Diagrams

Visual representation of sets using circles inside a rectangle [the universal set]


$$A \cup B =$$




$$A \cap B =$$




$$A - B =$$




$$B - A =$$



$$A \Delta B = (A - B) \cup (B - A)$$



$$=$$



Laws of Set Theory

1) Identity law

$$A \cup \emptyset = A \quad A \cap U = A$$

2) Idempotent law

$$A \cup A = A \quad A \cap A = A$$

3) Commutative law

$$A \cup B = B \cup A \quad A \cap B = B \cap A$$

Wrong Ans. state

4) Associative law

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

5) Distributive law

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$(A \cap B) \cup C$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

6) Absorption law

$$A \cup (A \cap B) = A$$

$$A \cap (A \cup B) = A$$

(Ans.) Ans. base case Ans. Ans.



7) De-morgan's law

$$(A \cup B)' = A' \cap B'$$

$$(A \cap B)' = A' \cup B'$$

8) complement law

$$(A')' = A$$

$$A \cup A' = U, \text{ and } A \cap A' = \emptyset$$

State and prove

Distributive laws.

Def:- Let 'x' is an arbitrary element of $A \cup (B \cap C)$

Let $x \in A \cup (B \cap C)$

$\Leftrightarrow x \in A$

$\Leftrightarrow (x \in A \text{ or } x \in (B \cap C))$

$\Leftrightarrow x \in A \text{ (or) } (x \in B \text{ and } x \in C)$

$\Leftrightarrow (x \in A \text{ or } x \in B) \text{ and}$

$(x \in A \text{ or } x \in C)$

$\Leftrightarrow x \in (A \cup B) \text{ and } x \in (A \cup C)$

$$\text{Q) } x \in (A \cup B) \cap (A \cup C)$$

$$\therefore A \cup (B \cap C) = (A \cup B) \cap (A \cup C) //$$

Prove De-morgan's law.

$$(A \cup B)' = A' \cap B' \text{ for } \forall x$$

$$\Rightarrow x \in (A \cup B) \text{ has two cases}$$

$$x \in A \text{ or } x \in B \text{ has two cases}$$

$$\Rightarrow x \in (A \cap B)' \text{ has two cases}$$

$$x \in A \text{ and } x \in B \text{ has two cases}$$

$$\Rightarrow x \notin A \cup B$$

$$\Rightarrow x \notin A \text{ and } x \notin B \text{ has two cases}$$


$$\Rightarrow x \notin (A \cup B)' \text{ has two cases}$$

$$x \notin A \text{ or } x \notin B$$

$$\text{example has two cases}$$

Ex. Let $U = \{1, 2, 3, 4, 5\}, A = \{2, 4\}$

$$A^c = U - A = \{1, 3, 5\}$$



$x \in A^c$ thus $x \notin A$



Prove De-morgan's law

$$(A \cup B)' = A' \cap B'$$

Sol:

L.H.S :- Let $x \in (A \cup B)'$

$$\Leftrightarrow x \notin (A \cup B)$$

$$\Leftrightarrow x \notin A \text{ and } x \notin B$$

$$\Leftrightarrow x \in A' \text{ and } x \in B'$$

$$\Leftrightarrow x \in (A' \cap B') \Rightarrow \text{R.H.S.}$$

$$\therefore (A \cup B)' = (A' \cap B')$$

S.T

$$A - (A \cap B) = A - B$$

Sol:

$$\text{let } x \in A - (A \cap B)$$

$$\Leftrightarrow x \in A \text{ and } x \notin A \cap B$$

$$\Leftrightarrow x \in A \text{ and } \{x \notin A \text{ or } x \notin B\}$$

$$\Leftrightarrow x \in A \text{ and } x \notin A \cap B$$

$$\Leftrightarrow x \in A \text{ and } x \notin B$$

$$\Leftrightarrow x \in A - B$$

$$\Leftrightarrow x \in A - B$$



If $A = \{2, 3, 4\}$, $B = \{1, 2\}$, $C = \{4, 5, 6\}$

find $A+B$, $B+C$, $A+B+C$

Sol: $A+B = \{2, 3, 4\} \cup \{1, 2\} = \{1, 2, 3, 4\}$

$+ & A$ are equal

$$\therefore A \Delta B = (A-B) \cup (B-A)$$

(i) $[A+B = (A-B) \cup (B-A)]$

$$\therefore A-B = \{3, 4\}$$

$$B-A = \{1\}$$

$$A+B = \{3, 4\} \cup \{1\} = \{1, 3, 4\}$$

(ii) $B+C$

$$\therefore [B+C = (B-C) \cup (C-B)]$$

$$\Rightarrow B-C = \{1, 2\}, C-B = \{4, 5, 6\}$$

$$\therefore B+C = \{1, 2\} \cup \{4, 5, 6\} = \{1, 2, 4, 5, 6\}$$

(iii) $A+B+C$

$$\Rightarrow (A-B) \cup (B-A) \cup (B-C) \cup (C-B)$$

$$(ii) \quad A + B + B + C$$

$$\text{Sol: } = ((A+B) - (B+C)) \cup ((B+C) - (A+B))$$

$$\Rightarrow [\{1, 3, 4\} - \{1, 2, 4, 5, 6\}] \cup [\{1, 2, 4, 5, 6\} - \{1, 3, 4\}]$$

$$\Rightarrow \{3\} \cup \{2, 5, 6\}$$

$\{3\} = S - A$

$$\Rightarrow \{2, 3, 5, 6\}$$

$\{2, 3, 5, 6\} = (S - A) - (S - B) = B - A$

$$(iv) \quad \underline{A + B + C}$$

$$\text{Sol: } \Rightarrow ((A+B)-C) \cup (C-(A+B))$$

$$\Rightarrow [\{1, 3, 4\} - \{4, 5, 6\}] \cup [\{4, 5, 6\} - \{1, 3, 4\}]$$

$$\Rightarrow \{1, 3\} \cup \{5, 6\}$$

$\{1, 3\} = S - A$

$$\Rightarrow \{1, 3, 5, 6\}$$

$\{1, 3, 5, 6\} = S - A - B$

$$\{1, 3, 5, 6\} = \{1, 3, 5\} \cup \{6\} = S - A - B$$



cartesian product

If A and B are 2 sets, then
cartesian product is

$$A \times B = \{(a, b) | a \in A, b \in B\}$$

e.g. If $A = \{1, 2, 3\}$, $B = \{4, 5\}$

then $A \times B$

$$= \{(1, 4), (1, 5), (2, 4), (2, 5), (3, 4), (3, 5)\}$$

$$B \times A = \{(4, 1), (4, 2), (4, 3), (5, 1), (5, 2), (5, 3)\}.$$

$$\therefore A \times B \neq B \times A$$

If $A = \{2, 4, 6, 8\}$, $B = \{1, 3, 5, 7, 9\}$

Find the cartesian product

$$A \times B \text{ and } B \times A$$

Sol:- $A \times B = \{(2, 1), (2, 3), (2, 5), (2, 7), (2, 9), (4, 1), (4, 3), (4, 5), (4, 7), (4, 9), (6, 1), (6, 3), (6, 5), (6, 7), (6, 9), (8, 1), (8, 3), (8, 5), (8, 7), (8, 9)\}$

$$B \times A = \{ \text{ } \} \quad \text{Taking first element}$$

$$A = \{ 2, 4, 6, 8 \}, B = \{ 1, 3, 5, 7, 9 \}$$

$$B \times A = \{ (1, 2), (1, 4), (1, 6), (1, 8) \}$$

$$(3, 2), (3, 4), (3, 6), (3, 8), (5, 2)$$

$$(5, 4), (5, 6), (5, 8), (7, 2), (7, 4)$$

$$(7, 6), (7, 8), (9, 2), (9, 4),$$

$$(9, 6), (9, 8) \}$$

next
3cm



Relation

Any subset of $A \times B$ is called a relation.

Ex: If $A \times B = \{ (a, 1), (a, 2), (a, 3) \}$

$\{ (a, 1), (a, 2) \}$ is a relation

because it is subset of $A \times B$.

(P, Q) (F, S) (C, P) (S, C) (Q, S) (P, S)

(P, S) (F, S) (C, S) (Q, S) (P, Q)

{(F, S), (F, Q), (C, S), (C, Q), (P, S), (P, Q)}

Domain and Range of Relations

Domain of $R = \{x / \forall (x, y) \in R\}$

Range of $R = \{y / \forall (x, y) \in R\}$

e.g.: If $R = \{(1, 2), (2, 4), (2, 7)\}$

Domain = $\{1, 2\}$, Range = $\{2, 4, 7\}$.

for sets $P = \{a, b, c, d\}$, $Q = \{e, f, g\}$

find $P \cup Q$, $P - Q$, $Q - P$, $P \cap Q$

Sol.: $P \cup Q = \{a, b, c, d, e, f, g\}$

$P - Q = \{a, b\}$

$Q - P = \{e, f, g\}$

$P \cap Q = \{c, d\}$ i.e.

Intersection (संतुष्टिमयीकरण)

A student with 4 subjects (Math, Science, English, Social)

Intersection of both, i.e.

A student with 3 subjects (Math, Science, English)

i.e. 3 subjects

Types of Relations

- 1) Reflexive Relation
- 2) Irreflexive
- 3) Symmetric
- 4) Anti-symmetric
- 5) Transitive

1) Reflexive Relation

R is reflexive if $\forall a \in A$ such that

$$(a,a) \in R \quad \forall a \in A \quad R = \{ \}$$

2) Irreflexive

If $(a,a) \notin R \quad \forall a \in A$

3) Symmetric

If $(a,b) \in R$ then $(b,a) \in R$.

4) Anti-symmetric

If $(a,b) \in R, (b,a) \in R$

then $a = b$.

5) Transitive:

If $(a,b) \in R$, $(b,c) \in R$ then

$(a,c) \in R$.

Ex If $X = \{1, 2, 3\}$

then ordered pairs

$$R = \{(1,1), (1,2), (1,3), (2,1), (2,2), (2,3), (3,1), (3,2), (3,3)\}$$

$\therefore (1,1) \in R \rightarrow$ Reflexive.

$\Rightarrow (1,1), (2,2), (3,3) \in R$ then
irreflexive.

\Rightarrow let $X = \{a, b, c, d\}$

$$R = \{(a,a), (b,b), (c,c), (d,d)\}$$

It is symmetric

$(x,y), (y,z) \in R$ only if $x = y$ *

$$\{(2,3), (3,2), (2,2)\} \times$$

$$\{(1,1), (2,2), (3,3)\}$$

Antisymmetric

$$\{(1,1), (2,2), (1,3)\} \times \{(3,1)\} \times$$

\Rightarrow Transitive:

$$\text{If } A = \{1, 2, 3\} \text{ (domain)} \quad \begin{array}{c} 1, 2 \\ \hline 2 \\ 1, 3 \end{array}$$

$$R = \{(1, 2), (2, 3), (1, 3)\}$$

$$\text{If } A = \{1, 2, 3, 4\} \quad \begin{array}{c} \text{Each } f = x + 1 \\ \text{eg. } 1+1=2, 2+1=3, 3+1=4 \end{array}$$

$$R = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 3), (4, 4)\}$$

then find if relation is

reflexive, symmetric

anti-symmetric and transitive,

Given: $A = \{1, 2, 3, 4\}$

$$R = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 3), (4, 4)\}$$

$\therefore \{(1, 1), (2, 2), (3, 3), (4, 4)\} \in R$

It is reflexive.

(ii) Symmetric

$$\therefore \{(1, 2), (2, 1)\} \in R$$

It is symmetric

non-antisymmetric

as $(1, 2) \in R$ but $(2, 1) \notin R$



(iii) Anti-symmetric

$(2,1) \notin R$.

It is not anti-symmetric

(iv) transitive

$\therefore \{(1,2), (2,1), (1,1)\} \subset R$

\therefore It is transitive.

2) Let $A = \{1, 2, 3, 4\}$.

$R = \{(1,1), (1,2), (2,3), (1,3), (4,4)\}$

Sol.: Not reflexive

$\{(1,1), (2,2), (3,3), (4,4)\} \notin R$

Not symmetric - $(1,2) \in R$ but $(2,1) \notin R$

$\{(1,2), (2,1)\} \subset R$ but $(2,1) \notin R$

It is Anti-symmetric.

It is transitive.

Participate in a T.Q.

Ques. 1. If R is a relation

of $A \times A$, then R is said to be

binary relation on A .



Equivalence Relation

A relation is said to be equivalence if it is

(i) reflexive.

(ii) symmetric

(iii) transitive

Poset

A relation is said to be poset if it is

(i) Reflexive

(ii) Antisymmetric

(iii) Transitive

Ex If $A = \{1, 2, 3, 4\}$

$R = \{(1, 1), (1, 4), (2, 2), (2, 3),$

$(4, 1), (4, 4), (3, 2), (3, 3)\}$

P.T R is an equivalence

relation. $\therefore (a, a) \in R$

Ldt. $\{(1, 1), (2, 2), (3, 3), (4, 4)\} \subset R$

$\therefore R$ is reflexive.

~~R is symmetric~~ for relation R.
for each $(x, y) \in R$ we
have $(y, x) \in R$.

$$(1, 4) \in R \rightarrow (4, 1) \in R$$

$$(2, 3) \in R \rightarrow (3, 2) \in R$$

$\therefore R$ is symmetric

$\therefore R$ is transitive.

$\Rightarrow R$ is reflexive.
 $\forall (x, y) \in R$, if $(y, z) \in R$ we
have $(x, z) \in R$.

$$\begin{array}{c} 2 \rightarrow 3 \\ (1, 4) \\ (4, 1) \\ \hline (1, 1) \in R \end{array}$$

$2 \rightarrow 3$

$3 \rightarrow 2$

$(2, 2) \in R$

$\therefore R$ is reflexive, symmetric,
but not transitive.

$\therefore R$ is equivalence relation.

Exercise

Given $A = \{1, 2, 3, 4\}$

$R_1 = \{(1, 2), (2, 1), (3, 4), (4, 3)\}$

$R_2 = \{(1, 2), (2, 1), (3, 3), (4, 4)\}$

using binary relation R ,

Matrix representation of relation

A relation can be represented by a matrix called Relation matrix and it is defined by $M_R = \{m_{ij}\}$

$$M_R = [m_{ij}]$$

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$

Ex: Let $A = \{1, 2, 3, 4\}$ and $R = \{(x, y) | x > y\}$ Find the matrix relation and directed graph of R .



Sol: Given: $A = \{1, 2, 3, 4\}$

$R = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (2, 4), (3, 1), (3, 2), (3, 3), (3, 4), (4, 1), (4, 2), (4, 3), (4, 4)\} \rightarrow \text{ordered pairs}$

\therefore Given $R = \{(x, y) | x > y\}$. Hence
 $\Rightarrow \{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2),$
 $(4, 3)\}$

Now writing 2 3 4
 $\therefore M_R = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix}$

Di-graph [Directed graph]



\therefore Game and activity

2) If $A = \{1, 2, 3, 4\}$ defined by

$$R = \{(1, 1), (2, 2), (3, 3), (4, 4), (4, 3), (4, 2), (4, 1), (3, 2), (3, 1)\}$$

Find


the matrix relation and
directed graph.

Sol.: Given: $A = \{1, 2, 3, 4\}$

R is given.

$$M_R = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

Directed graph



$\left. \begin{array}{l} 1 \text{ and } 4 \text{ divisible by } 2 \\ 2 \text{ divides } 4 \end{array} \right\} 2 \text{ divides } 4 \text{ or not?} \quad 4 \mid 2$

Let $X = \{1, 2, 3, 4, 5, 6, 7\}$ and

$R = \{(x, y) | x-y \text{ is divisible by } 3\}$

in X . S.T 'R' is an equivalence relation

Sol: Given: $X = \{1, 2, 3, 4, 5, 6, 7\}$

$R = \{(x, y) | x-y \text{ is divisible by } 3\}$

Reflexive

$\forall x \in X$, we have $x-x=0$

is divisible by 3

$\therefore (x, x) \in R \quad \forall x \in X$.

$\therefore R$ is reflexive.

Symmetrie:

Let $x, y \in X$ and $(x, y) \in R$

then $x - y$ is divisible by 3

$\Rightarrow x - (x - y)$ is also divisible by 3

$\Rightarrow y - x$ is also divisible by 3.

$\therefore (y, x) \in R$

$\therefore (x, y) \in R \rightarrow (y, x) \in R$

$\therefore R$ is symmetric.

Transitive:

Let $x, y, z \in X$ & $(x, y) \in R$, $(y, z) \in R$

$(y, z) \in R$ then

$$\frac{x-y}{3} = k_1 \quad \& \quad y-z = 3k_2$$

$$\Rightarrow x-y + y-z = 3k_1 + 3k_2$$

$$\Rightarrow x-z = 3(k_1 + k_2)$$

divisible by 3

$$\Rightarrow (x, z) \in R$$

$\therefore R$ is Transitive

\therefore It is equivalence

composition of Relations

If $A = \{1, 2, 3, 4\}$ and R and S are relations on a set A defined by

$$R = \{(1, 2), (1, 3), (2, 4), (4, 4)\}$$

$$S = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 3), (2, 4)\}$$

find $ROS, SOR, ROR, SOS, RO(SOR)$,

$$(ROS)_{DR}, (ROR)_{OR}, (SOR)_{OS},$$

$$R \cup S, R \cap S, S^{-1}, \text{Dom}(R), \text{Range}(R),$$

$$\text{Dom}(S), \text{Range}(S), \text{Dom}(R \cup S),$$

$$\text{Range}(R \cap S)$$

$$\therefore ROS$$

$$\Rightarrow \{(1, 3), (1, 4)\}$$

$$SOR$$

$$\Rightarrow \{(1, 2), (1, 3), (1, 4), (2, 4)\}$$

$$R^2 = ROR$$

$$\Rightarrow \{(1, 4), (2, 4)\}$$

$$S^2 = SOS$$

$$\Rightarrow \{(1, 2), (1, 3), (1, 4)\}$$

$$R_0(S_0 R) = \{(1, 4)\}$$

$$(R_0 S) O R = \{(1, 4)\}$$

$$(R_0 R) O R = \{(1, 4), (2, 4)\}$$

$$(S_0 S) O S = \{(1, 3), (1, 4)\}$$

$$R \cup S = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (4, 4)\}$$

$$R \cap S = \{(2, 3), (2, 4), (4, 4)\}$$

$$R \circ S = \{(1, 2), (1, 3), (2, 4)\}$$

$$S^T = \{(1, 1), (2, 1), (3, 1), (4, 1), (3, 2), (4, 2)\}$$

$$S^{-1} = \{(1, 1), (2, 1), (3, 1), (4, 1), (3, 2), (4, 2)\}$$

$$D(R) = \{1, 2, 4\}$$

$$R(R) = \{2, 3, 4\}$$

$$D(S) = \{1, 2\}$$

$$R(S) = \{1, 2, 3, 4\}$$

$$D(R \cup S) = \{1, 2, 4\}$$

$$R(R \cap S) = \{2, 3, 4\}$$

$$S' = A - S$$

$$\Rightarrow (A \times A) - S$$

$$A = \{1, 2, 3, 4\}$$

$$A = \{1, 2, 3, 4\}$$

$$\therefore A \times A = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (2, 2), (2, 3), (2, 4), (3, 1), (3, 2), (3, 3), (3, 4), (4, 1), (4, 2), (4, 3), (4, 4)\}$$

$$\therefore (A \times A) - S \Rightarrow$$

$$S = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 3), (2, 4)\}$$

$$\therefore (A \times A) - S \Rightarrow \{(2, 1), (2, 2), (3, 1), (3, 2), (3, 4), (3, 3), (4, 4), (4, 1), (4, 2), (4, 3)\}$$

H.W.: Let $A = \{a, b, c, d\}$ defined by.

$$R = \{(a, a), (a, c), (c, b), (c, d), (d, b)\}$$

$$S = \{(b, a), (c, c), (c, d), (d, a)\}$$

Find (i) $R \circ S$

(i) $S \circ R$

(ii) $R \circ R$

congruence modulo m

for any positive integers x, y, m
if m divides $x - y$ then x and y are said to be congruent
modulo m and is written as

$$x \equiv y \pmod{m}$$

Ex. $5 \equiv 3 \pmod{2}$

$$14 \equiv 2 \pmod{3}$$

P.T the congruence relation is an equivalence relation.

(i) Reflexive

As $x - x = 0$ is divisible by m

$$\text{i.e., } x \equiv x \pmod{m}$$

(ii) Symmetric

$$\therefore x \equiv y \pmod{m}$$

$$m/x-y$$

similarly $\Rightarrow m/y-x$ also

Ex. $3 \text{ divides } 7-1$
 $3/7-1 \Rightarrow 3/6$



$y \equiv z \pmod{m}$

∴ It is symmetric.

Transitive:

(ii) $\therefore a \equiv y \pmod{m}, y \equiv z \pmod{m}$

$m/a-y$ and $m/y-z$.

$$\Rightarrow m/a-y+y-z \Rightarrow m/a-z$$

$$\Rightarrow a \equiv z \pmod{m}.$$

∴ congruence modulo m is
reflexive, symmetric and
transitive.

hence it is equivalence

relation.

QUESTION 10

ANSWER



Scanned with OKEN Scanner

So the relation "greater than or equal to" is a partial ordering on the set of integers.

Let: let \mathbb{Z}^+ be the set of all positive integers

$$\text{Given } R = \{(x, y) | x \geq y\} \text{ on } \mathbb{Z}^+$$

(i) Reflexive:

$\forall x \in \mathbb{Z}^+$, we have $x \geq x$.

$\therefore (x, x) \in R \quad \forall x \in \mathbb{Z}^+$

$\therefore R$ is reflexive.

(ii) Anti-symmetric

$\forall x, y \in \mathbb{Z}^+$, clearly $x \geq y$

and $y \geq x \Rightarrow x = y$

i.e. $(x, y) \in R \Rightarrow (y, x) \notin R$

then $x = y$

$\therefore R$ is Anti-symmetric.

(iii) Transitive

$\forall x, y, z \in \mathbb{Z}^+$, if $x \geq y$ & $y \geq z$
then $x \geq z$

i.e., $(x, y) \in R$ & $(y, z) \in R$ then
 $(x, z) \in R$

hence, (\mathbb{Z}^+, \leq) is a POSET

Partial ordering set [POSET]

A set P with a partial ordering \leq is called a partially ordered set (or) POSET.

It is denoted by (P, \leq) .

Hasse-Diagram

A partial ordering (\leq) on a set 'P' can be represented by means of a diagram known as Hasse diagrams of (P, \leq) .

→ In Hasse-diagram, each element is represented by a small circle or a dot.

→ we do not put arrows on edges and we do not draw self-loops at vertices

→ There will be no transitive edges

✓ A · B ⊂

..... B ⊂ x



→ Diagram will be from bottom to top.

→ There will be no arrow heads.

To draw the Hasse-diagram


Q) Let $X = \{2, 3, 6, 12, 24, 36\}$ and \leq be the relation "divides", Draw the Hasse-diagram of (X, \leq) .

Q) Draw the Hasse-diagram for the divisibility relation on

defined on set $A = \{2, 3, 6, 12, 24, 36\}$


Ans: $R = \{(2, 2), (2, 6), (2, 12), (2, 24), (2, 36), (3, 3), (3, 6), (3, 12), (3, 24), (3, 36), (6, 6), (6, 12), (6, 24), (6, 36), (12, 12), (12, 24), (12, 36), (24, 24), (24, 36), (36, 36)\}$

Hasse-diagram




$\therefore 2$ doesn't divide 3
so 2 & 3 are represented with
separate dots. $\boxed{2 \cdot \cdot 3}$

$\rightarrow \because 2$ divides 6
 $\therefore 2$ divides 6 also, so 6
& 3 are in the middle of 2 & 3
in the middle of 2 & 3



$\rightarrow \because 6$ divides 12
so 6 to 12 one edge.




~~$\# 2$ & 3 also divides 12,
but they are transitive edges~~

$$\begin{array}{r} 2, 6 \\ 6, 12 \\ \hline 2, 12 \end{array} \quad \begin{array}{r} 3, 6 \\ 6, 12 \\ \hline 3, 12 \end{array}$$

transitive edges

$\rightarrow \because 12$ divides 24
12 also divides 36
so as edge from 12 to 24 & 36




∴ Draw Hasse-diagram for
 $\{1, 2, 3, 4, 6, 9\}$ for divisibility

Sol: $R = \{(1,1), (1,2), (1,3), (1,4),$
 $(1,6), (1,9), (2,2), (2,4), (2,6),$
 $(3,3), (3,6), (3,9), (4,4), (6,6),$
 $(9,9)\}$

Hasse-diagram

$\{1, 2, 3, 4, 6, 9\}$




1 divides 2

1 divides 3 also.

Info with 30 edges.

2 divides 4


so 2 to 4 edge




2 divides 6.

3 also divides 6

so edges from
2 & 3 to 6



3 divides 9
so 3 to 9 edges



→ Draw the Hasse-diagram for the relation R on $A = \{1, 2, 3, 4, 5\}$. whose relation matrix is given

below.


	①	②	③	④	⑤
1	1	0	1	1	1
2	0	1	1	1	1
3	0	0	1	1	1
4	0	0	0	1	0
5	0	0	0	0	1

Sol: let $R = \{(1, 1), (1, 3), (1, 4), (1, 5),$

$(2, 2), (2, 3), (2, 4), (2, 5), (3, 3),$

$(3, 4), (3, 5), (4, 4), (5, 5)\}$

Hasse-diagram for $\{1, 2, 3, 4, 5\}$



$1 \leq 2$ [But not in relation]

so no edge for 1 to 2

$$\therefore 1 \leq 3$$

$$2 \text{ also } 2 \leq 3$$

so edge from


$$1, 2 \text{ to } 3.$$



Also $3 \leq 4$ and so on

so $3 \leq 5$


so edges from 3 to 4 & 5



Draw Hasse diagrams for

$\{1, 3, 5, 9, 15, 45\}$ for divisibility


So $\therefore R = \{(1, 1), (1, 3), (1, 5), (1, 9), (1, 15), (1, 45), (3, 3), (3, 9), (3, 15), (3, 45), (5, 5), (5, 15), (5, 45), (9, 9), (9, 45), (15, 15), (15, 45), (45, 45)\}$



Applications in康定

Hasse-Diagram

$\{1, 3, 5, 9, 15, 45\}$



$\therefore 1$ divides 3


1 also divides 5

so edge from 1 to 3 + 5



3 doesn't divide 9


but, 3 divides 9, so 3 to 9 → edge



9 doesn't divide 15

but 3 & 5 divide 15


so 3 + 5 to 15 → edge



both 15 + 9 divides 45


so edge from 15, 9 to 45

→ no transitive edges
should be drawn




Q) Let $A = \{1, 2, 3, 4, 6, 8, 12, 24, 48\}$

Sol:



(or)
neatly
→



Properties present in
poset of a set P are

Q) Draw Hasse diagram of poset

$\{1, 2, 3, 4, 5\}$ for \leq




1. \leq is reflexive & transitive
2. \leq is antisymmetric
3. Every two elements are comparable
4. Every two elements are comparable
5. Every two elements are comparable
6. Every two elements are comparable
7. Every two elements are comparable
8. Every two elements are comparable
9. Every two elements are comparable
10. Every two elements are comparable
11. Every two elements are comparable
12. Every two elements are comparable
13. Every two elements are comparable
14. Every two elements are comparable
15. Every two elements are comparable
16. Every two elements are comparable
17. Every two elements are comparable
18. Every two elements are comparable
19. Every two elements are comparable
20. Every two elements are comparable
21. Every two elements are comparable
22. Every two elements are comparable
23. Every two elements are comparable
24. Every two elements are comparable
25. Every two elements are comparable
26. Every two elements are comparable
27. Every two elements are comparable
28. Every two elements are comparable
29. Every two elements are comparable
30. Every two elements are comparable
31. Every two elements are comparable
32. Every two elements are comparable
33. Every two elements are comparable
34. Every two elements are comparable
35. Every two elements are comparable
36. Every two elements are comparable
37. Every two elements are comparable
38. Every two elements are comparable
39. Every two elements are comparable
40. Every two elements are comparable
41. Every two elements are comparable
42. Every two elements are comparable
43. Every two elements are comparable
44. Every two elements are comparable
45. Every two elements are comparable
46. Every two elements are comparable
47. Every two elements are comparable
48. Every two elements are comparable
49. Every two elements are comparable
50. Every two elements are comparable
51. Every two elements are comparable
52. Every two elements are comparable
53. Every two elements are comparable
54. Every two elements are comparable
55. Every two elements are comparable
56. Every two elements are comparable
57. Every two elements are comparable
58. Every two elements are comparable
59. Every two elements are comparable
60. Every two elements are comparable
61. Every two elements are comparable
62. Every two elements are comparable
63. Every two elements are comparable
64. Every two elements are comparable
65. Every two elements are comparable
66. Every two elements are comparable
67. Every two elements are comparable
68. Every two elements are comparable
69. Every two elements are comparable
70. Every two elements are comparable
71. Every two elements are comparable
72. Every two elements are comparable
73. Every two elements are comparable
74. Every two elements are comparable
75. Every two elements are comparable
76. Every two elements are comparable
77. Every two elements are comparable
78. Every two elements are comparable
79. Every two elements are comparable
80. Every two elements are comparable
81. Every two elements are comparable
82. Every two elements are comparable
83. Every two elements are comparable
84. Every two elements are comparable
85. Every two elements are comparable
86. Every two elements are comparable
87. Every two elements are comparable
88. Every two elements are comparable
89. Every two elements are comparable
90. Every two elements are comparable
91. Every two elements are comparable
92. Every two elements are comparable
93. Every two elements are comparable
94. Every two elements are comparable
95. Every two elements are comparable
96. Every two elements are comparable
97. Every two elements are comparable
98. Every two elements are comparable
99. Every two elements are comparable
100. Every two elements are comparable

Totally ordered set.

→ A partial ordering R on a set
 $A = \{1, 2, 3, 4\}$ is represented by the
following digraph. Draw the
Hasse-diagrams of (R, \leq) . $\rightarrow \leq = 1$

sol : $R = \{(1, 1), (1, 2), (1, 3), (1, 4),$
 $(2, 2), (2, 3), (2, 4), (3, 3), (4, 4)\}$.


Digraph



Hasse-Diagram

divides

$$\{1, 2, 3, 4\}, 1$$




→ Draw hasse-diagram for
divisors of 36 i.e., (D_{36}, \mid)

Sol: $A = \{1, 2, 3, 4, 6, 9, 12, 18, 36\}$

$R = \{(x, y) \mid x \mid y\}$


$A = \{1, 2, 3, 4, 6, 9, 12, 18, 36\}$



→ Draw H-D for. divisibility

for $A = \{1, 2, 3, 5, 6, 15, 30\}$


Sol:-




H-D for $\{1, 2, 3, 6, 9, 18\}, 1$

$\{1, 2, 3, 6, 9, 18\}$

soln




$\{1, 2, 3, 4, 5, 10\}$



D(10)

let $A = \{1, 2, 5, 10\}$




D(15), D(32), D(45).

D100

Let $A = \{1, 2, 4, 5, 10, 20, 25, 50, 100\}$

Hasse-Diagram



(a) a
partial order



Lattice

A lattice is a partially ordered set (POSET), denoted as (L, \leq) in which every pair of elements such as $(a, b) \in L$ has a least upper bound (LUB) & greatest lower bound (GLB).

LUB: It is also called as supremum, join, sum of elements
→ It is denoted by $a \vee b$.


$$\text{LUB}(a, b) = a \vee b \quad (\text{or}) \quad a \text{ joins } b \\ \Rightarrow a \oplus b$$

GLB: It is also called as infimum, meet, product of elements
→ It is denoted as $a \wedge b$.

$$\text{GLB}(a, b) = a \wedge b, \quad (\text{or}) \quad a \text{ meets } b \\ \Rightarrow a \ast b$$

It is a structure having two binary operations \wedge and \vee such that each one with respect to the other is commutative, associative and has identity element.

Ex:- Define whether POSET defined by Hasse diagram is lattice or not.




LUB table

\vee	1	2	3	6
1	1	2	3	6
2	2	2	6	6
3	3	6	3	6
6	6	6	6	6

GLB table

\wedge	1	2	3	6
1	1	1	1	1
2	1	2	1	2
3	1	1	3	3
6	1	2	3	6

\therefore Every pair of elements in the given Hasse-diagram has LUB & GLB
 \therefore It is a lattice.



LUB table

a	a	b	c	d	e	f
a	a	c	c	d	e	f
b	c	b	c	d	e	f
c	c	c	c	d	e	f
d	d	d	d	d	e	f
e	e	e	e	e	e	-
f	f	f	f	f	-	f

GLB table

1	a	b	c	d	e	f
a	a	-	a	a	a	a
b	-	b	b	b	b	b
c	a	b	c	c	c	c
d	a	b	c	d	d	d
e	a	b	c	d	e	d
f	a	b	c	d	d	f

$\therefore \text{LUB}\{e, f\} \& \text{GLB}\{a, b\}$ don't exist, the given poset is not a lattice.

Some properties of lattices

If L is any lattice then

$$\forall a, b, c \in L$$

$$(i) a * a = a, a \oplus a = a \quad \text{Idempotent}$$

$$(ii) a * b = b * a, a \oplus b = b \oplus a$$

↓
commutative.

$$(iii) a * (b * c) = (a * b) * c$$

$$a \oplus (b \oplus c) = (a \oplus b) \oplus c$$

↓
Associative.

$$(iv) a * (b \oplus c) = (a * b) \oplus (a * c)$$

$$(a * b) \oplus c = (a \oplus c) * (b \oplus c)$$

↓
Distributive.

$$(v) a * (a \oplus b) = a$$

$$a \oplus (a * b) = a$$

↓
Absorption

Duality law

The dual of any statement can be obtained by interchanging * and \oplus .


Ex:- 1) The dual of

$$a * (b \oplus c) \text{ is } a \oplus (b * c)$$

2) The dual of

$$(a \wedge b) \vee c \text{ is } (a \vee b) \wedge c.$$

Types of lattices:



A Hasse diagram for a lattice

It is formed with binary operations (join). The join operation

is denoted by \wedge and the meet operation

is denoted by \vee .

Distributive lattice

A lattice (L, \leq) is said to be distributive lattice if and only if it satisfies the following properties.

$$(i) a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

$$(ii) a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

where $a, b, c \in L$.


Ex : Verify the following

hasse-diagrams (or) lattices

are distributive lattice

or not.

i)



Sol : It is said to be distributive lattice if it satisfies the following properties

$$(i) a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

Let $a=1, b=2, c=3$

$$\Rightarrow 1 \vee (2 \wedge 3) = (1 \vee 2) \wedge (1 \vee 3)$$

$$\Rightarrow 1 \vee (2 \wedge 3) = 2 \wedge 3$$

$$\Rightarrow 1 = 1 \quad \therefore [L.H.S = R.H.S]$$

$$(ii) a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

$$\therefore a=1, b=2, c=3$$

$$\Rightarrow 1 \wedge (2 \vee 3) = (1 \wedge 2) \vee (1 \wedge 3)$$

$$\Rightarrow 1 \wedge 4 = 1 \vee 1$$


$$\therefore L.H.S = R.H.S$$

\therefore It satisfied the 2 properties

\therefore The given base-diagram
is a distributive lattice.



∴ The given base-diagram is a distributive lattice.



$$(i) a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

$$\text{Let } a=1, b=2, c=3$$

$$\Rightarrow 1 \vee (2 \wedge 3) = (1 \vee 2) \wedge (1 \vee 3)$$

$$\Rightarrow 1 \vee (0) = 1 \wedge 2$$

$$\Rightarrow 1 \neq 1 \quad [\text{Satisfied}]$$

$$(ii) a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

$$a=1, b=2, c=3$$


$$\Rightarrow 1 \wedge (2 \vee 3) = (1 \wedge 2) \vee (1 \wedge 3)$$

$$\Rightarrow 1 \wedge (0) = 1 \neq 0$$

$$\Rightarrow 1 \neq 2 \quad [\text{L.H.S} \neq \text{R.H.S}]$$

\therefore It is not a distributive lattice.

37



$$(i) a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

$$\text{Let } a = a, b = b, c = c$$

$$\Rightarrow a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

$$\Rightarrow a \vee d = e \wedge e$$

$$\Rightarrow a \neq e$$

$$(ii) a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

$$\Rightarrow a \wedge (e) = d \vee d$$

$$\Rightarrow a \neq d$$

Hardly
 ★ must have exactly one
 complement for each element.



Then it becomes

distributive lattice

Verify that (D_{16}, \leq) is a lattice or not?

Sol.: $D_{16} = \{1, 2, 4, 8, 16\}$

Hasse-Diagram,



LUB(LV)	1	2	4	8	16
1	1	2	4	8	16
2	2	2	4	8	16
4	4	4	4	8	16
8	8	8	8	8	16
16	16	16	16	16	16

GLB	1	2	4	8	16
1	1	1	1	1	1
2	1	2	2	2	2
4	1	2	4	4	4
8	1	2	4	8	8
16	1	2	4	8	16


Every pair of elements in D_{16} has LUB & GLB
 $\therefore (D_{16}, \leq)$ is a lattice

Bounds of a Lattice

The greatest element and the least element in a lattice are denoted by 1 & 0 respectively and they are called as bounds of the lattice.

Bounded lattice

A lattice (L, \leq) is bounded if it has the greatest element (1) and the least element (0).



complete lattice

If every subset of L has a join and meet, then the lattice (L, \leq) is called a complete lattice.

complement of an element.

Let $a \in L$ be an element in a lattice $(L, *, \oplus)$ with bounds 0 and 1 .

If $\exists b \in L$ such that $a * b = 0$ and $a \oplus b = 1$ then

b is said to be the complement of a . A lattice where each element has a complement is called a complemented lattice.

If each element has a complement in a lattice, then the lattice is said to be complemented lattice.

~~→ complement is not unique.~~

Need to get

$$\begin{aligned} [a * b = 1] &= b * \\ [a \oplus b = 0] &= b \oplus \end{aligned}$$

$$\begin{array}{l} \oplus \rightarrow v \\ * \rightarrow 1 \end{array}$$

so a is complement of b if b is complement of a .

Now we have to prove a is complement of b .


So $b * a = 0$ and $b \oplus a = 1$.

Now we have to prove $a * b = 0$ and $a \oplus b = 1$.



Q. T the given lattice is a complemented lattice or not.

1)



Need to get $\begin{cases} a \vee b = 1 \\ a \wedge b = 0 \end{cases}$.

Here $c = 1, a = 0$

Any way $c \vee a = c = 1$

$c \wedge a = a = 0$.

$\therefore c$ is complement of a

& a is complement of c

Next $b \& d$.

$$b \vee d = c = 1$$


$$b \wedge d = a = 0$$

$\therefore b$ is complement of d

d is complement of b

\therefore In the given Hasse-diagram
for each & every element, there
is a complement

\therefore It is a complemented lattice.



$$a \vee d = a = 1$$

$$a \wedge d = d = 0$$

so $a \wedge c = d$
 $d \vee c = a = 1$

Next for $b \wedge c$

$$b \vee c = a = 1$$

$$b \wedge c = d = 0$$

$$\therefore b \wedge c = c$$

$$c \wedge b = b$$

\therefore All the elements have complements.

\therefore It is a complemented lattice.

$a = 1$ is the top element

$d = 0$ is the bottom element

b, c are the middle elements

$b \wedge c = d$ is the meet operation

$b \vee c = a$ is the join operation


$b \wedge a = b$ is the identity for meet

$b \vee a = a$ is the identity for join

$b \wedge b = b$ is the idempotent law for meet

$b \vee b = a$ is the idempotent law for join

37



Sol: one element may have
any no. of complements.

$$\text{Anyway } 1 \vee 0 = 1 \quad 1 \wedge 0 = 0$$

$$1^c = 0$$

$$0^c = 1$$

for a & b

$$a \vee b = 1$$

$$a \wedge b = 0$$

$$a^c = b$$

$$b^c = a$$

for b & c

$$b \vee c = 1$$

$$b \wedge c = 0$$

$$b^c = c$$

$$c^c = b$$

c & d

$$c \vee d = 1$$

$$c \wedge d = 0$$

$$c^c = d$$

$$d^c = c$$

for a & d

$$a \vee d = 1$$

$$a \wedge d = 0$$

$$a^c = d$$

$$d^c = a$$

So one may have any no. of elements.

$$a^c \rightarrow d, b, c$$

$$b^c \rightarrow c, d, a$$

$$c^c \rightarrow b, a, d$$

$$d^c \rightarrow a, c, b$$

$$b.d.$$

$$b \vee d = 1$$

$$b \wedge d = 0$$

$$b^c = d$$

$$d^c = b$$

$$b \vee a = 1$$


$$b \wedge a = 0$$

$$b^c = a$$

$$a^c = b$$

\therefore All elements are having complements

\therefore It is a complemented lattice.



$$\therefore c^c = b, o^c = 1$$

check for a.

Suppose a, d

$$a \vee d = d \neq 1$$

$$a \wedge d = a \neq 0$$

$$\begin{cases} a \vee b = d \neq 1 \\ a \wedge b = 0 \\ a^c \neq b \\ b^c \neq a \end{cases}$$

$$\begin{cases} a \vee d = d \neq 1 \\ a \wedge d = 0 \\ a^c \neq b \\ b^c \neq a \end{cases}$$

Suppose a, c

$$\begin{cases} a \vee c = c \neq 1 \\ a \wedge c = a \neq 0 \end{cases}$$

$$\begin{cases} a^c = c \\ c^c \neq a \end{cases}$$



a. doesn't have complement

Even if one element doesn't have a complement

It is not a complemented lattice


~~lattice~~ doesn't have a complemented lattice

If for b, d

$$b \vee d = d \neq 1$$

$$b \wedge d = d \neq 0$$

5)



$$\begin{aligned}b^c &= c, c^c \\b \wedge c &= 1 \\b \vee c &= 0\end{aligned}$$

$$\text{because } a \vee g = 1, a \wedge g = 0$$

Suppose e, f

$$\begin{aligned}e \vee f &= f \neq 1 & e^c &\neq f \\e \wedge f &= e \neq 0 & f^c &\neq e\end{aligned}$$

Suppose e, d

$$\begin{aligned}e \vee d &= d \neq 1 & e^c &\neq d \\e \wedge d &= e \neq 0 & d^c &\neq e\end{aligned}$$

Suppose e, a

$$eva = e \neq 0$$

$$ena = a = 0$$

$\therefore e$ doesn't have even one complement

\therefore It is not a complemented

Lattice

Functions

Let X and Y be any 2 sets
A relation f from X to Y is
called a function if for every
 $x \in X$ there is a unique $y \in Y$.

such that $(x, y) \in f$


Ex: let $X = \{1, 5, P, Jack\}$,

$Y = \{2, 5, 7, q, Jill\}$.

& $f = \{(1, 2), (5, 7), (P, q), (Jack, q)\}$

clearly every element has
unique image

$\therefore f$ is a function.



domain of $f = \{1, 5, P, \text{Jack}\}$

Range of $f = \{2, 5, 7, \text{Jill}\}$

Codomain of $f = \{2, 5, 7, 9, \text{Jill}\}$

→ If an element x in X is mapped to y in Y , then y is said to be image of x and x is said to be pre-image of y .

Ex: If the function f is defined by $f(x) = x^2 + 1$ on the set $\{-2, -1, 0, 1, 2\}$ find the range of f .

Sol: Given $f(x) = x^2 + 1$ for $x \in \{-2, -1, 0, 1, 2\}$

$$f(-2) = (-2)^2 + 1 = 5$$

$$f(-1) = (-1)^2 + 1 = 2$$

$$f(0) = 0^2 + 1 = 1$$

$$f(1) = 1^2 + 1 = 2$$

$$f(2) = 2^2 + 1 = 5$$

∴ Range of $f = \{1, 2, 5\}$


Types of functions

1) one-to-one [Injective function]


A function $f: A \rightarrow B$ is called one-to-one function if distinct elements of set A are mapped to the distinct elements of set B.

\Rightarrow Two elements in set A can't be mapped to the same element of set B.


Ex. 1)



2)






3)




Onto function [surjective]

A function $f: A \rightarrow B$ is onto function if every element of Set B will be an image of some element of set A.




It is onto

but not one-one



is injective

but not surjective



not one-one
and onto.

Bijective function : A mapping $f: X \rightarrow Y$ is said to be bijective if it is both one-to-one and onto.

S.T. the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 3x^3 + 5 \quad \forall x \in \mathbb{R}$ is a bijection.

Q: Let $x, y \in \mathbb{R}$. Then
∴ bijection means both one-one & onto

one-one

$$\begin{aligned} f(x) &= f(y) \\ (f(x) - 3x^3 + 5) &= (f(y) - 3y^3 + 5) \end{aligned}$$

$$\begin{aligned} \Rightarrow 3x^3 + 5 &= 3y^3 + 5 \\ \Rightarrow 3x^3 &= 3y^3 \\ \Rightarrow x^3 &= y^3 \Rightarrow x = y \\ \therefore f \text{ is one-one} & \end{aligned}$$

onto

Let $f(x) = y \quad \forall x \in \mathbb{R}$

$$\Rightarrow 3x^3 + 5 = y$$

$$\Rightarrow 3x^3 = y - 5$$

$$\Rightarrow x^3 = \frac{y-5}{3} \Rightarrow x = \left(\frac{y-5}{3}\right)^{\frac{1}{3}} \in \mathbb{R}$$

$$\text{Ex: } f(x) = 3x^3 + 5 \xrightarrow{\text{Eqn 1}} \text{is it onto?}$$

$$\therefore \text{Sub } x = (\frac{y-5}{3})^{1/3} \text{ in Eqn 1}$$

$$\Rightarrow \text{Eqn 1} \rightarrow 3\left(\left(\frac{y-5}{3}\right)^{1/3}\right)^3 + 5$$

$$\Rightarrow y - 5 + 5 = y$$

\therefore We got $f(x) = y$

$\therefore f$ is onto.

$\therefore f$ is one-one and onto

$\therefore f$ is a bijection

Inverse function

A function $f: x \rightarrow y$ is invertible if $f: x \rightarrow y$ is bijective.

Ex: Let $f: R \rightarrow R$ be given by
 $f(x) = x^3 - 2$. Find f^{-1} if it exists

Sol: One-one $\Leftrightarrow f(x) = f(y)$

$$x^3 - 2 = y^3 - 2$$

$$x^3 = y^3 \Rightarrow x = y$$



$\therefore f$ is one-one.

Let $f(x) = y$

$$\Rightarrow x^3 - 2 = y \Rightarrow x^3 = y + 2$$
$$\Rightarrow [x = (y+2)^{1/3}] \rightarrow \textcircled{1}$$

$\therefore f(x) = y \Rightarrow x^3 - 2 = y \in \mathbb{R}$

$$\Rightarrow [(y+2)^{1/3}]^3 - 2$$
$$\Rightarrow y + 2 - 2 = y$$

\therefore got $f(x) = y$

$\therefore f$ is onto.

not needed

To get Inverse

$$\therefore f(x) = y$$
$$\Rightarrow x = f^{-1}(y)$$
$$\Rightarrow (y+2)^{1/3} = f^{-1}(y) \quad \text{[from } \textcircled{1} \text{]}$$

\Rightarrow Similarly

$$f^{-1}(x) = (x+2)^{1/3}/k$$

composition of functions

Let $f: x \rightarrow y$ and $g: y \rightarrow z$ be
 functions, then the composition
 of $g \circ f$ is the function from x to z
 defined by $(g \circ f)(x) = g(f(x))$.

$$(g \circ f)(x) = g(f(x)) \quad \forall x \in \mathbb{R}$$

Ex. Let $f(x) = x+2$, $g(x) = 3x^2 + 4$
 $\& h(x) = x^2 + 1$ [for $x \in \mathbb{R}$. Find
 $gof, fog, fof, gog, fo(h), hog, hof,$
 $\& fo(goh), (fog)o(h)$. Verify
that $fo(goh) = (fog)o(h)$

$$\text{Sol.: } \text{Given: } f(x) = x + 2, g(x) = 3x^2 + 4, h(x) = x^2 + 1$$

$$\begin{aligned}
 & \text{(ii) } (g \circ f)(x) \\
 & \Rightarrow g[f(x)]. \Rightarrow g[x+2] \\
 & \Rightarrow 3(x+2)^2 + 4 \\
 & \Rightarrow 3[x^2 + 4x + 4] + 4 \quad [\because g(x) = 3x^2 + 4] \\
 & \Rightarrow 3x^2 + 12x + 12 + 4 \\
 & \Rightarrow 3x^2 + 12x + 16 \quad \text{Ans.}
 \end{aligned}$$

$$\text{(iii) } f \circ g(x) = f[g(x)]$$

$$= f[3x^2 + 4]$$

$$\text{but } \Rightarrow (3x^2 + 4) + 2 \Rightarrow 3x^2 + 6$$

$$\text{(iv) } f \circ f(x) = f[f(x)]$$

$$= f[x+2]$$

$$\text{but } \Rightarrow (x+2) + 2 \Rightarrow x+4$$

$$\text{(v) } g \circ g(x) = g[g(x)]$$

$$\text{but } \Rightarrow g[3x^2 + 4]$$

$$\text{but } \Rightarrow 3(3x^2 + 4)^2 + 4$$

$$= 3[9x^4 + 16 + 24x^2] + 4$$

$$\Rightarrow 27x^4 + 48 + 72x^2 + 4$$

$$\Rightarrow 27x^4 + 72x^2 + 52$$

$$\text{(vi) } f \circ h(x) = f[h(x)]$$
~~$$= f[x^2 + 1]$$~~

$$= (x+2)^2 + 1$$

$$\Rightarrow x^2 + 4 + 4x + 1$$

$$\Rightarrow x^2 + 4x + 5$$

$\therefore f(x) = x+2$

$f(x^2+1) = (x^2+1)+2$

$= x^2+1+2$

$$\text{(v)} f \circ h(x) = f[h(x)] \\ = f[x^2 + 1]$$

$$= \therefore f(x) = x + 2$$

$$f(x^2 + 1) = (x^2 + 1) + 2$$

$$\Rightarrow x^2 + 3$$

$$\text{(vi)} h \circ g(x) = h[g(x)]$$

$$= h[3x^2 + 4]$$

$$= (3x^2 + 4)^2 + 1 \quad [\because h(x) = x^2 + 1]$$

$$= 9x^4 + 16 + 24x^2 + 1$$

$$\Rightarrow 9x^4 + 24x^2 + 17$$

$$\text{(vii)} h \circ f(x) = h[f(x)]$$

$$= h[x + 2]$$

$$= (x + 2)^2 + 1 \quad [\because h(x) = x^2 + 1]$$

$$= x^2 + 4 + 4x + 1$$

$$= x^2 + 4x + 5$$

$$\text{(viii)} f \circ (g \circ h)(x) = f[g[h(x)]]$$

$$= f[g[x^2 + 1]]$$

$$= f[3(x^2 + 1)^2 + 4]$$

$$= f[3(x^4 + 1 + 2x^2) + 4]$$

$$= f[3x^4 + 3 + 6x^2 + 4]$$

$$\Rightarrow f[3x^4 + 6x^2 + 7]$$

$$\Rightarrow (3x^4 + 6x^2 + 7) + 2 \quad \{ \because f(x) = x + 2 \}$$

$$\Rightarrow \underline{3x^4 + 6x^2 + 9} //$$

3) $(f \circ g) \circ h(x) \Rightarrow f[g[h(x)]]$

$(f \circ g) \circ h$

$$\Rightarrow f[g(x)] = f[g(x)]$$

$$= f[3x^2 + 4]$$

$$= (3x^2 + 4) + 2$$

$$\{ f[g(x)] = 3x^2 + 6 \}$$

$(f \circ g) \circ h(x)$

$\Rightarrow (f \circ g)h(x)$

$$\Rightarrow (f \circ g)(x^2 + 1)$$

$$\Rightarrow 3(x^2 + 1)^2 + 6$$

$$\Rightarrow 3[x^4 + 1 + 2x^2] + 6$$

$$\Rightarrow 3x^4 + 3 + 6x^2 + 6$$

$$\Rightarrow \underline{3x^4 + 6x^2 + 9} //$$

$\therefore f \circ (g \circ h)(x) = (f \circ g) \circ h(x)$

S.T $f(x) = 3 - 4x$ is bijective, $f: \mathbb{R} \rightarrow \mathbb{R}$

$$f(x) = 3 - 4x$$

one-one

$$\text{let } f(x_1) = f(x_2)$$

$$3 - 4x_1 = 3 - 4x_2$$

$$\cancel{3} - 4x_1 = -4x_2$$

$$\Rightarrow x_1 = x_2$$

f is one-one

onto

$$\text{let } f(x) = y$$

$$\cancel{3} - 4x = y$$

$$\cancel{3} - 4x = y - 3$$

$$\cancel{x} = \frac{y - 3}{-4}$$

$$x = \frac{-(3-y)}{-4} = \frac{3-y}{4} \in \mathbb{R}$$

$\therefore f$ is onto.

$\therefore f$ is bijective

$f(x) = 2x + 1 \forall x \in \mathbb{R}$ is also bijective

$$\text{onto} \rightarrow \left\{ x = \frac{y-1}{2} \in \mathbb{R} \right\}$$