UPSC CSE 2027 · GS FOUNDATION

Geography

Geomorphology · Climatology · Oceanography · Indian geography · Human geography · Economic geography

Personal Study Edition

Built from the Sarrthi IAS GS Foundation Geography handouts and mind maps.

Copyright belongs to the teacher and institute; keep this private.

How to use this book

Every chapter opens with the **Exam lens** and plain-language key terms, then the full notes in handout order with PYQ tags, alerts, mnemonics and diagrams, and ends with a recall sheet. The practice quizzes, flashcards and question banks live on the Study Shelf.

Priority tags: **H** = two or more PYQs map to the chapter; **M** = one PYQ or a partial fit; **L** = none quoted in the material.

The wide blank column on the right of every page, and the binding margin on the left, are kept free for your own current affairs, data and value additions.

Reign-year codes (for example Babur 26-30) are kept exactly as the teacher writes them.

Contents

1 Geomorphology 4
2 Climatology 118
3 Oceanography 198
4 Indian Geography 245
5 Human Geography 370
6 Economic Geography 394
7 Mind maps · Geomorphology 484
8 Mind maps · Climatology 513
9 Mind maps · Oceanography 527
10 Mind maps · Indian Geography 542
11 Mind maps · Human Geography 562
12 Mind maps · Economic Geography 567

GEOGRAPHY · GS FOUNDATION · SUBJECT 1

Geomorphology

Handout + mind maps: no class slides | Handout pages 67

HIGH · 19 Prelims, 2 Mains

The whole Geomorphology handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 36 mind-map pages are in Chapter 7.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: HIGH for Prelims.** 19 Prelims PYQs are printed in this part of the handout (15 with a year: 2011, 2012×2, 2013×2, 2014, 2015, 2018, 2019, 2022, 2023, 2024×3, 2025; 4 with no year printed). Mains quoted: **Juno mission (2017, 10m)** and **primary rocks (2022, 10m)**.
- **Hot clusters:** Earth–Sun geometry (pole star, day length, 21 June, time zones) · solar storm effects · P vs S waves · weathering by rain · drift evidences · volcanic products · chalk vs clay.
- **No teacher ink in this PDF**, so the priority tag comes only from PYQ counts in the handout.
- **Prepare:** *Prelims* – who/when/where of theories, layers and discontinuities, wave properties, landform identification. *Mains* – drift → plate tectonics reasoning, rock types and economic value, landforms by agent (river, ice, sea, wind, groundwater).

1 · UNIVERSE

THEORIES OF ORIGIN OF THE SOLAR SYSTEM

Gaseous Hypothesis:

- Given by **Kant**



- Large particles of **primordial matter** were **scattered** in the universe.
- They started **colliding against each other** .
- **Collision resulted in the generation of heat.**
- Heat changed the primordial matter from **solid to liquid and from liquid to gaseous** state, **creating a nebula** .
- **Nebula started rotating** around its axis, **creating a bulge** in the center of the gaseous mass.



- When this bulge increased in size, the **rings started forming** one by one and were **separated from the middle part** of the nebula.



- **Residual central mass became the sun** and the rest of the **rings became the planets** .
- **How did the moons form?**



- Repetition of the **same process at the planet level.**
- Rings were separated from the newly formed planets.
- Material of each ring condensed to form satellites of the concerned planets.

- **Drawback:**
  - Not explained the **source of the primordial matter** .
  - According to the science of the law of motion, the collision of particles doesn't generate **rotatory motion** .

Nebular Hypothesis:

- 1796.
- By **Laplace**
- Modified version of Kant's theory
- **Postulates:**
  - A **huge and hot gaseous nebula** is present in the space. Thus, he solved the problem of heat of the nebula through this assumption.
  - From the very beginning, a **huge and hot nebula was rotating** (spinning) on its axis.
  - A huge and hot gaseous matter called a nebula existed in space which was continuously rotating on its axis.
  - Nebula was **losing heat** from its outer surface due to the **process of radiation** and was thus **reducing in size and volume** due to contraction.
  - As its size reduced, its **velocity of rotatory motion began to increase** .



- With increased velocity, the **rings started detaching** .
- Each **ring condensed** at a point in the form of a gaseous mass which started **rotating around the nebula** .
- **Gaseous mass** later cooled and formed the **planets** .
- The **remaining part** of the nebula thus became the **sun** .
- **Limitations:**
  - Did not explain the source of **origin of the nebula** .
  - If the planets have been formed from the rotating nebula, then the residual part of the nebula i.e., the **Sun should rotate at the highest speed** due to the decrease in its size, but **it is not so** .

Chamberlin–Moulton planetesimal hypothesis

- According to this hypothesis, a **star passed close to the young Sun, causing tidal bulges and solar prominences that led to repeated ejections of material from the Sun.**



- **Gravitational effects** from the passing star created **spiral arms** of ejected material, with some of it **remaining in orbit** .
- This material **cooled and condensed** into small bodies called **planetesimals** and larger **protoplanets** .
- Over time **, collisions among** these objects formed the **planets and their moons, with comets and asteroids** as leftover debris.

Accretion Model

- By **Otto Schmidt (1944) and Viktor Safronov (1969).**
- **Constant accretion** of particles convert them from a **nebula to planets** eventually.

- Nebula → Dust grains → Boulders → Planetesimals → Planets



Big Bang Hypothesis:

- **Georges Lemaitre** proposed in 1920s.
- He theorized that the **universe began from a single primordial atom.**
- The idea received major boosts from Edwin Hubble's observations that galaxies are speeding away from us in all directions, as well as from the 1960s discovery of cosmic microwave radiation—interpreted as echoes of the big bang—by Arno Penzias and Robert Wilson.
- Theory was finally validated in 1972.



- All matter in the universe existed in the form of **dense and huge primordial matter.**
- **A violent explosion** took place in this primordial matter.
- Due to this, the **dust particles** present in this matter were **scattered in the universe.**
- Over several million years, slightly **denser regions become gravitationally attracted** to each other.
- They grew even **denser, forming gas clouds, stars, galaxies** , and other astronomical structures.

PROOF:



(Hotter the body, shorter the wavelength. Why? More energy = faster particle motion → higher frequency waves = shorter wavelength)



- **Hubble's Law (Hubble-Lemaitre law):** galaxies farther away from us are moving away at faster speeds, indicating the **expansion of the universe.**
  - Basis of Hubble's law is **redshift** , which is the **elongation of light waves** to longer wavelengths, often **toward the red end** of the spectrum, caused by an **object moving away from the observer** , commonly observed in distant galaxies due to the expansion of the universe.
- **Cosmic Microwave Background Emissions:**
  - Discovered by American physicists **Arno Penzias and Robert Wilson,** who detected it in **1965** .
  - It is the faint glow of electromagnetic radiation that fills the universe. It is the oldest light in the universe, dating back to about 380,000 years after the Big Bang.
  - In the early universe, it was so hot that matter and radiation were constantly interacting — a plasma of electrons and nuclei. There were no atoms due t the heat.
  - When the universe cooled enough (~3000 K), electrons combined with protons to form neutral atoms — a process called recombination.
  - At that point, photons could travel freely — this is the moment the CMB was released.
  - **CMB is like a baby photo of the universe** , showing us how it looked just after birth. Its existence, temperature, uniformity, and small irregularities are **exactly what**

the Big Bang Theory predicts , and no other theory explains it so well.

⚡ **Quick revise** Origin theories – "who said what" Fold ▲

Theory	By	Core idea	Drawback (handout)
Gaseous hypothesis	**Kant**	Colliding particles → heat → **nebula**; rotating nebula threw **rings** → planets; same process → moons	Source of primordial matter unexplained; collisions can't give rotation
Nebular hypothesis	**Laplace, 1796** (modified Kant)	Hot rotating nebula **cooled, shrank, spun faster**, rings detached → planets; rest = Sun	Source of nebula unexplained; Sun should spin fastest but doesn't
Planetesimal	**Chamberlin–Moulton**	A **passing star** pulled material off the young Sun → **planetesimals** → protoplanets → planets	–
Accretion model	**Otto Schmidt 1944**, **Safronov 1969**	Nebula → dust grains → boulders → planetesimals → planets	–
Big Bang	**Lemaître** (1920s), "primordial atom"	Dense primordial matter exploded; slightly denser regions pulled together → gas clouds, stars, galaxies	*Validated 1972 (handout)*

👨‍👩‍👧 **Who-is-who:** *K*ant = *K*ollision, *L*aplace = *L*oses heat (cools & shrinks, spins faster). Schmidt–Safronov = **S**mall **S**teps (dust → planet).

⚡ **Quick revise** Proofs of the Big Bang Fold ▲

Proofs of the Big Bang: Hubble's law – farther galaxies recede faster (basis: **redshift**). **CMB** – found by **Penzias & Wilson, 1965**; oldest light, **~380,000 yrs** after Big Bang; at ~3000 K electrons joined protons (**recombination**) so photons flew free – "a baby photo of the universe". Universe age **13.8 bn yrs**.

2 · DARK ENERGY & DARK MATTER

- **Dark energy and dark matter** are two mysterious components that together make up about **95% of the universe** , yet they remain largely unexplained.

- **Dark matter** , which accounts for roughly **27%** , does not emit, absorb, or reflect light, but its presence is inferred from its **gravitational effects** , such as the rotation of galaxies and gravitational lensing (bending of light as it passes from near a massive celestial object).
- It acts as an invisible scaffold, holding galaxies together.
- On the other hand, **dark energy** , making up about **68%** , is thought to be responsible for the **accelerating expansion of the universe** .
- Unlike dark matter, dark energy is not associated with matter or gravity pulling things together, but rather with a **repulsive force** that pushes space itself apart.
- Though both are invisible and not directly detectable, their effects are crucial in explaining the structure, behavior, and fate of the cosmos.

⚡ **Quick revise** Dark side Fold ▲

Dark side: dark matter ≈ **27%** (gravity glue, lensing); dark energy ≈ **68%** (pushes space apart, accelerating expansion); together ≈ 95%.

3 · GALAXIES

- **Formation**

  - **After the Big Bang** (13.8 billion years ago): Initial fluctuations in the density of matter led to the congregation of material at those places. The role of Dark matter’s gravity is considered here.
  - **First Galaxies (about 1 billion years post-Big Bang)** : Gas clouds collapse under gravity, forming the first stars and galaxies.
  - **Galaxy Evolution (1 billion years to present):** Galaxies grow and evolve through mergers and interactions.

- **Types of Galaxies**
  - **Spiral Galaxies:** These galaxies feature a **flat, rotating disk with spiral arms winding outward from a central bulge** . The **Milky Way** is a prime example of a spiral galaxy. They are rich in gas and dust, facilitating **active star formation, particularly in the spiral arms.**
  - **Elliptical Galaxies:** Ranging from nearly spherical to elongated shapes, elliptical galaxies **have a smooth, featureless light profile** . They contain **older, cooler stars and possess minimal gas and dust** , resulting **in low rates of new star formation.**

  - **Lenticular Galaxies** : These are **intermediate between spiral and elliptical** galaxies, characterized by a **central bulge and a disk but lacking** prominent **spiral arms** .

They have little interstellar matter, leading to **minimal star formation** activity.

  - **Irregular Galaxies:** Lacking a distinct shape, irregular galaxies often **appear chaotic** , possibly due to gravitational interactions or collisions with other galaxies. They typically **contain abundant gas and dust, with active regions** of star formation.

- **Active Galactic Nuclei: highly energetic regions** at the **centers of some galaxies** , powered by **supermassive black holes** accreting matter. These regions emit enormous amounts of radiation across the electromagnetic spectrum and are categorized based on their observational characteristics.
- **Active galaxies** make up about **10% of known galaxies** and have centers that are over 100 times brighter than the combined light of their stars. This intense energy is powered by supermassive black holes surrounded by accretion disks of infalling gas and dust, which emit radiation from infrared to X-rays. Some of this material is also ejected as jets near the speed of light.
- **Milky way** is **not active.**
- **Types of galaxies on basis of activity:**

- **Seyfert Galaxies Most common and lowest energy** active galaxies. **Appear normal in visible light but emit strongly in infrared and sometimes X-rays.** Usually radio-quiet, though **some produce jets** .
- **Quasars Brightest and most energetic** type of active galaxy. **Can outshine entire galaxies** and are visible across great cosmic distances. **Powered by massive accretion activity** ; may be episodic, lasting ~10 million years. Useful for studying the early universe and black hole growth.
- **Blazars** A type of active galaxy whose **jets point almost directly at Earth** . Appear **extremely bright** and are **sources of high-energy particles like neutrinos.** Give insight into the extreme environments near black holes.

⚡ **Quick revise** Galaxies & stars Fold ▲

- **Types:** Spiral (Milky Way) · Elliptical (old, cool stars, little gas) · Lenticular (bulge + disk, no arms) · Irregular (chaotic, gas-rich).
- **Active galaxies** (~10%; supermassive black hole + accretion disk; **Milky Way is not active**): **Seyfert** (most common, lowest energy) · **Quasar** (brightest) · **Blazar** (jet points at Earth).
- **Populations: Pop III** – hypothetical first stars, no heavy metals · **Pop II** – oldest observed (5–10 bya), halo/globular clusters · **Pop I** – younger, metal-rich (4–5 bya), disk & arms, **our Sun**.

4 · STARS

- Their formation after the Big Bang within the galaxies.
  - Approximately 380,000 years post-Big Bang, the universe cools enough for electrons to combine with protons and form neutral atoms, releasing the CMB.
  - **Formation of Gas Clouds (100 million to 200 million years post-Big Bang)** : Gravity causes hydrogen and helium gas to clump together, forming large clouds.
  - **Protostar Formation (200 million to 400 million years post-Big Bang):** These gas clouds collapse under gravity, increasing temperature and pressure, forming protostars.
  - **Nuclear Fusion Ignition (200 million to 400 million years post-Big Bang):** When core temperatures are high enough, nuclear fusion ignites, and stars are born.
- **Lifecycle of stars:**
  - **Stellar Nebula** : Stars form from collapsing clouds of gas and dust.
  - **Protostar:** As the nebula collapses, a protostar forms and heats up.
  - **Groups of stars that have newly formed from molecular clouds** are commonly referred to as **stellar clusters** .
  - **Molecular clouds that contain many of these clusters** are known as **stellar nurseries** .
  - **Main Sequence:** Star enters a stable phase, fusing hydrogen into helium.
    - **Low-Mass Stars (mass < 8 times that of Sun):**

- **Red Giant:** Exhausts hydrogen, expands, and cools.
- **Planetary Nebula:** Outer layers shed.
- **White Dwarf:** Core remains, cools over time. It may convert into a black dwarf over time, but we do not have any solid proof of this as of now.
    - **High-Mass Stars (mass > 8 times that of the sun): Red Supergiant:** Similar process but on a larger scale.
- **Supernova:** Catastrophic explosion after iron buildup in core.
- **Neutron Star/Black Hole:** The remnant core collapses into a neutron star or black hole, depending on mass. If it's between 8-20/25 times the mass of the sun, we get Neutron stars. If it's more, we get Black Holes.
- **Chandrashekhar Limit:** If after the shedding of the outer layers, due to supernova, or without it, the mass of the remaining core is less than **1.44 times the mass of the sun** , we get a white dwarf. Otherwise, we get a neutron star (1.44-3) or a black hole (>3).

Population 3, 2 and 1 stars

- Stars are classified into populations based on their metallicity, which refers to the abundance of elements heavier than hydrogen and helium.
- **Population III Stars (Hypothetical):** These are theorized to be the **very first stars** born in the universe, hundreds of millions of years after the Big Bang.
    - **Extremely Metal-Poor:** They are believed to have been almost entirely composed of hydrogen and helium, with virtually no heavier elements because those elements hadn't been forged in abundance yet.
    - **Massive and Short-Lived:** Due to their large size, they are thought to have been incredibly massive and short-lived, burning out rapidly in a few million years.
    - Direct evidence for their existence has not yet been found.
    - Do not have a planetary system.
- **Population II Stars (Oldest Observed; 5-10 bya):** These are the oldest stars still observable in the Milky Way galaxy.
    - **Metal-Poor:** They contain very little of the heavier elements compared to younger stars.
    - **Location:** Population II stars are typically found in the galactic halo (spherical region surrounding the disk) and globular clusters (dense collections of millions of old stars).
    - **Slower Evolution:** Due to their lower metallicity, they evolve more slowly and have longer lifespans than stars with higher metal content.
- **Population I Stars (Younger and Richer in Metals; 4-5 bya)** : These are the younger stars, including our Sun.



- **Metal-Rich:** They contain a higher proportion of heavier elements compared to Population II stars. This enrichment is due to previous generations of stars that lived, died, and scattered these elements into the interstellar medium.
- **Location:** Population I stars are found in the galactic disk (flattened disc where most stars and gas reside) and spiral arms of the Milky Way.
- **Faster Evolution:** Their higher metallicity can lead to faster stellar evolution and shorter lifespans compared to Population II stars.

Neutron Stars

- Neutron stars are extremely dense stellar remnants formed from the collapsed core of **massive stars (about 8-25 times the mass of the Sun)** after a supernova explosion.
- They are **composed almost entirely of neutrons** and have incredibly **strong gravitational and magnetic fields.**
- Sometimes they can accrete mass from their surroundings to collapse into black holes.
- **Pulsars: Rapidly rotating neutron stars** that emit beams of electromagnetic radiation from their magnetic poles, observed as regular pulses.

Black Holes

- Black holes are regions of spacetime with **gravitational pull** so **strong** that nothing, **not even light, can escape** from them. (For reference, Earth's escape velocity is 11.8 m/s.)
- They form from the remnants of massive stars (more than 25 times the mass of the Sun) after they undergo a supernova explosion and collapse under their own gravity.



- **Event Horizon:** The **boundary** around a black hole beyond which nothing can escape.
- **Accretion Disks:** Disks of **gas and dust spiraling into black holes** , emitting intense radiation due to friction and heating.
- Accretion Disk, then event horizon, then the void.
- **Gravitational Waves: Ripples in spacetime** produced by the acceleration of massive objects like merging black holes, detectable by observatories like **LIGO** (Laser Interferometer Gravitational-Wave Observatory; Nobel prize in 2017) **and Virgo** .



- **Spaghettification:** Spaghettification is the process where **objects are stretched and torn apart** by **intense tidal forces** as they approach a black hole.

- **Hawking Radiation:** Hawking radiation is the **theoretical emission** of particles from black holes due to quantum effects near the event horizon, leading to **gradual black hole evaporation** .

⚡ **Quick revise** Galaxies & stars Fold ▲

⭐ **Star fate – "Small stars retire quietly, big stars quit with a bang"**

Mass	Path	End
< 8 × Sun	Main sequence → Red giant → Planetary nebula	**White dwarf**
> 8 × Sun	Red supergiant → **Supernova** (iron core)	**Neutron star** (8–20/25×) or **Black hole** (>25×)

Chandrasekhar limit 1.44 – 3: core < **1.44** = white dwarf; **1.44–3** = neutron star; **> 3** = black hole. *Pulsar = spinning neutron star ("lighthouse").* Black-hole words: event horizon, accretion disk, **spaghettification**, **Hawking radiation**, gravitational waves (**LIGO, Nobel 2017**).

5 · SOLAR SYSTEM

Components of the Solar System

- **The Sun:** The **central star** of the solar system, a massive ball of hydrogen and helium undergoing nuclear fusion.
- **Planets: Major celestial bodies orbiting the Sun** , classified as terrestrial (Mercury, Venus, Earth, Mars) and gas giants (Jupiter, Saturn) or ice giants (Uranus, Neptune).
- In recent years, **astronomers have proposed** the existence of a ninth planet, nicknamed " **Planet Nine** ," or " **Planet X,** " to **explain the unusual orbital patterns of some objects in the Kuiper Belt** , a region beyond Neptune. If Planet Nine exists, it is **hypothesized to be a gas or ice giant** , potentially **5-7 times Earth's mass** , and located **very far from the Sun** , possibly 200 times farther than Earth.
- **Dwarf Planets:** Celestial bodies **orbiting the Sun** that are **spherical but have not cleared their orbital path.** The solar system has 5 known dwarf planets: **Pluto (the biggest), Ceres (the biggest object in the asteroid belt), Eris, Haumea, and Makemake (the newest).** What are the **criteria** for calling a body a dwarf planet?
  - **In orbit around the Sun** : The object must be gravitationally bound to the Sun and travel in an orbit around it.
  - **Massive enough for hydrostatic equilibrium (nearly round shape)** : The object's own gravity must be strong enough to pull it into a roughly round shape, overcoming the forces of its rigid body. This is in contrast to asteroids, which are irregularly shaped due to their weaker gravity.
  - **Not a satellite (moon) of another object:** The object cannot be gravitationally bound to another planet or large object and act as its moon.
  - **Has not cleared the neighborhood around its orbit:** This is the key distinction between a dwarf planet and a full-fledged planet. The object's gravity must not be

strong enough to have cleared the path around its orbit of other objects of comparable size. (Orbit of Pluto and Neptune intersects.)

- **Moons: Natural satellites** orbiting planets and dwarf planets. Moons in our Solar system: 0 for Mercury & Venus (has a Quasi-satellite called Zoozve), 1 for Earth; 2 for Mars; 101 at Jupiter; 274 at Saturn; 28 at Uranus; 16 at Neptune; and five for dwarf planet Pluto. Jupiter has the biggest moon in our solar system, Ganymede.
- **Quasi-Moons:** Quasi-moons are **asteroids that orbit the Sun while staying close to a planet** . A quasi-satellite's orbit is usually more oblong and less stable than the planet's orbit. In time, the shape of a quasi-satellite's orbit may change, and it may move away from the planet.
- **Kuiper Belt:** A **region beyond Neptune** filled with icy bodies and dwarf planets. In addition to rock and water ice, objects in the Kuiper Belt **also contain a variety of other frozen compounds like ammonia and methane.** Distance from the sun is about 30-50 AU (1 Astronomical Unit = 149.6 million km).
  - The Kuiper Belt is **gradually wearing down** as objects within it occasionally collide, breaking into smaller fragments, which can include comets and dust that gets carried out of the solar system by solar wind.
- **Oort Cloud:** A distant **spherical shell of icy objects surrounding the solar system** , believed to be the source of long-period comets.



  - The Oort Cloud is believed to begin around **2,000 to 5,000 AU** from the Sun, while its outer boundary is estimated to lie between **10,000 and 100,000 AU** away.
- **Asteroids: Small rocky bodies** , **primarily** found in the asteroid belt **between Mars and Jupiter** . Asteroids vary in size, from **Vesta** , the **largest** at approximately **329 miles (530 kilometers)** wide, to much smaller objects measuring under **33 feet (10 meters)** . Altogether, the **combined mass of all asteroids** is still **less than the mass of Earth's Moon** .
  - **Trojans:** These **asteroids share an orbit with a larger planet, but do not collide with it** because they **gather around** two special places in the orbit (called the **L4 and L5 Lagrangian points**).
  - **Lagrangian points** are **positions in space where the gravitational forces of two large bodies** , like the Earth and Sun, **balance the centripetal force felt by a smaller object** . These points allow the object to remain relatively stable with respect to the two larger bodies.
- **Comets: Icy bodies that release gas and dust** , forming a **glowing coma and tail** when near the Sun, originating from the Kuiper Belt or Oort Cloud.



- **Meteoroids:** Small particles from comets or asteroids that travel through space. Some meteoroids are rocky, while others are metallic or a combination of rock and metal.
- **Meteor:** When a **meteoroid enters Earth's atmosphere** , it heats up due to friction with air molecules. This burning object, streaking across the sky, is what we call a meteor, commonly known as a "shooting star." Most meteors are completely vaporized by the heat before reaching the ground.
- **Meteorite:** If a **meteoroid is large and sturdy enough to survive its fiery passage** through the atmosphere and impact Earth's surface, it becomes a meteorite. Meteorites can be of various sizes, from a pebble to a massive rock. Of more than 50,000 meteorites found on Earth, 99.8% come from asteroids. The remaining small fraction (0.2%) of meteorites is split roughly equally between meteorites from Mars and the Moon.
- **Heliosphere** : The **bubble of solar wind** surrounding the solar system.
- **Termination Shock** : Where the **solar wind slows from supersonic to subsonic speeds** . This slowing down is due to the solar wind particles encountering the denser interstellar medium's pressure.
- **Heliosheath** : The **turbulent region beyond the termination shock** . In this region, the solar wind's velocity decreases further, and the magnetic field becomes more irregular. The heliosheath acts as the outer layer of the heliosphere and is a turbulent region where the solar wind's influence weakens.
- **Heliopause** : The boundary where **solar wind pressure balances with interstellar pressure** . It marks the outer edge of the heliosphere. Beyond the heliopause, the interstellar medium predominates, and the influence of the solar wind ceases. The heliopause is considered the final frontier of the solar wind's reach, marking the transition between the heliosphere and interstellar space.
- **Bow Shock** : The **shockwave formed as the heliosphere moves through the interstellar medium** . It is analogous to the shock wave that forms in front of a moving boat or a supersonic aircraft. As the heliosphere plows through the interstellar medium, it creates this shockwave, where the interstellar medium is abruptly heated and slowed down due to the presence of the heliosphere. However, the existence of such a region has been contested by the data received from the Voyager 1 and 2 missions by NASA.

6 · PYQ

Q.1) The term 'Goldilocks Zone' is often seen in the news in the context of (2015)

(a) the limits of habitable zone above the surface of the Earth

(b) regions inside the Earth where shale gas is available

(c) search for the Earth-like planets in outer space

(d) search for meteorites containing precious metals

Answer - C

Explanation:

- Goldilocks zone is the distance from the star, where the temperature is such that water can stay in Liquid form. Earth lies in the Goldilocks Zone of the Sun. If we wish to find life form similar to that found on the Earth, then we need to search for the planets lying in the Goldilocks zone of their respective stars.

Q.2) What is the difference between asteroids and comets? (2011)

1. Asteroids are small rocky planetoids, while comets are formed of frozen gases held together by a rocky and metallic material.

2. Asteroids are found mostly between the orbits of Jupiter and Mars, while comets are found mostly between Venus and mercury.

3. Comets show a perceptible glowing tail, while asteroids do not.

Which of the statements given above is/are correct ?

- 1 and 2 only
- 1 and 3 only
- 3 only
- 1, 2 and 3

Answer. B

Explanation

- The main difference between asteroids and comets is their composition, as in, what they are made of Asteroids are made up of metals and rocky material, while comets are made up of ice, dust and rocky material.
- Both asteroids and comets were formed early in the history of the solar system about 4.5 billion years ago.
- Asteroids formed much closer to the Sun, where it was too warm for ice to remain solid. Comets formed farther from the Sun where ice would not melt. Comets which approach the Sun lose material with each orbit because some of their ice melts and evaporates to form a tail.
- Asteroids are a series of very small planets or fragments of planets lying between the orbits of Mars and Jupiter.
- They revolve around the Sun in the same way as planets, unlike comets.

⚡ **Quick revise** Asteroid vs comet (PYQ 2011) Fold ▲

Asteroid vs comet (PYQ 2011): asteroids = rock/metal, between Mars and Jupiter, orbit like planets; comets = ice + dust, show a glowing **coma and tail** near the Sun. (Statement "comets between Venus and Mercury" is wrong.)

7 · PLANETS

Feature	Inner Planets	Outer
Number	4 (Mercury, Venus, Earth, Mars)	4 (Neptu
Composition	Primarily rocky (iron, silicon)	Prima heliur
Size	Smaller diameter and mass	Signif mass
Density	Higher density	Lowe
Surface	Solid surfaces with craters, volcanoes, and canyons (except Venus which has a thick atmosphere)	No sc comp heliur
Magnetic Field	Weaker magnetic fields (except Earth)	Stron
Rings	No rings	Most and ic 2, Ura
Moons	Fewer moons (Earth: 1, Mars: 2)	Many 274,
Temperature	Wider range of temperatures, with hot days and cold nights (except Venus with a constant hot surface)	Colde distar
Formation	Closer to the Sun, formed from denser material in the early solar system	Farthe leftov
Orbit	Shorter orbital periods (Mercury: 88 days, Earth: 365 days, Mars: 687 days)	Longe years years

- **Venus and Uranus rotate from East to West.**

- **The Axis of Uranus is parallel to its orbit.**
- The **Giant and the Little red spots** are a defining feature of Jupiter. They are huge storms. The former is 16000 km in diameter while the latter is 3500 km in diameter.
- Atmosphere of different Planets:
  - **Mercury:** Very thin, most of it is lost due to solar storms
  - **Venus** : thick atmosphere, mainly made of CO2, making the planet super-hot due to the greenhouse effect.
  - **Mars:** Very thin, rich in CO2
  - **Jupiter and Saturn:** Hydrogen and Helium with the addition of some sulphurous and carbon gases.
  - **Uranus and Neptune** : Mainly Methane and Ammonia

⚡ **Quick revise** Solar system – numbers worth holding Fold ▲

Item	Fact
Planets	**Terrestrial:** Mercury, Venus, Earth, Mars · **Gas giants:** Jupiter, Saturn · **Ice giants:** Uranus, Neptune
Dwarf planets (5)	**Pluto** (biggest), **Ceres** (biggest in asteroid belt), Eris, Haumea, **Makemake** (newest). Criteria: orbits Sun · round · not a moon · **has not cleared its orbit**
Moons	Mercury **0**, Venus **0** · Earth **1** · Mars **2** · Jupiter **101** · Saturn **274** · Uranus **28** · Neptune **16** · Pluto **5**. Biggest moon: **Ganymede** (Jupiter)
Kuiper belt	Beyond Neptune, **30–50 AU**; icy bodies; source of comets (1 AU = 149.6 m km)
Oort cloud	Spherical shell; starts **2,000–5,000 AU**, outer edge 10,000–100,000 AU; **long-period comets**
Asteroids	Belt **between Mars & Jupiter**; **Vesta** largest (~530 km); **Trojans** sit at L4/L5 Lagrange points
Meteors	Meteoroid (in space) → **meteor** (burns in air) → **meteorite** (hits ground). 99.8% of meteorites are from asteroids; 0.2% Mars + Moon
Heliosphere	Termination shock → heliosheath → **heliopause** (solar wind = interstellar pressure) → bow shock (existence contested by Voyager data)

> **Handout alert:** Saturn's moons are printed as **274** in the moons list but **74** in the inner/outer planets table → use **274**. "Earth's escape velocity 11.8 m/s" – unit looks like a typo; don't memorise it. *Planet Nine* is only hypothesised (5–7 × Earth mass).

Inner vs outer planets: inner = rocky, dense, small, few/no moons, weak magnetic field, short orbits (Mercury 88 d, Earth 365, Mars 687) · outer = gaseous, big, many moons, rings, strong fields, long orbits (Jupiter 12 y, Saturn 29, Uranus 84, Neptune 165). **Venus and Uranus rotate East→West**; Uranus's axis is parallel to its orbit. Jupiter's **Great Red Spot 16,000 km**, Little Red Spot 3,500 km (storms). Atmospheres:

Venus = CO_2, super-hot (greenhouse); Mars thin CO_2; Jupiter/Saturn H + He; Uranus/Neptune methane + ammonia.

8 · THE SUN

- **Composition:** 71% Hydrogen; 27.1% Helium; 0.97% Oxygen; 0.4% Carbon, trace amount of other elements.
- **Interior:**
  - **Core: Nuclear fusion** takes place here. The temperature here is over 15 million degrees C.
  - **Radiative Zone** : Region in which **energy transfer** is **through thermal radiation** instead of convection. (Radiation: emission or transmission of energy in the form of electromagnetic waves or particles.)
  - **Convective Zone:** The **convection cells move in this zone** for energy transportation. (Convection: transfer of heat through the movement of a fluid (such as air or water) caused by differences in temperature and density.)
- The Sun is made up of hot gases, so it **lacks any "surface"** like earth.





- Instead, as we move from the Sun’s outer atmosphere toward the Sun's core, the gas gets denser and denser.
- **Atmosphere:**
  - **Photosphere:**
    - Sun’s outer atmosphere is transparent, but when we try to look through the atmosphere deeper into the Sun, our view is blocked.
    - The photosphere is the **layer that begins where the Sun starts to look opaque**





  - **Chromosphere:** layer **above the photosphere.** Outer chromosphere is about 10,000 degrees C.



  - **Corona: Layer of hot plasma** which extends millions of kilometers into the space from the surface of the chromosphere and has an irregular shape.
- **Coronal Mass Ejection:**
  - Large **expulsions of plasma and magnetic field from the Sun’s corona** .
  - 15-18 hours to reach Earth. Can be slower.
  - Expand as they move away from the sun.

- **Solar Winds:**



  - Solar wind is a **stream of charged particles released from the upper atmosphere of the Sun** (Corona).
  - **Solar wind is created by the outward expansion of plasma.**
  - Plasma is continually heated to the point that the Sun's gravity can't hold it down.
- **Solar flares:**
  - Tremendous **explosions due to the release of energy in twisted magnetic fields** (Like when rubber bands come back to their original size.)
  - It causes a sudden flash of increased brightness on the Sun.
  - Can be, but **not always, associated with CMEs** .
  - Release **X-rays & UV radiations** .
  - Flares are categorized according to their strength (energy output). The smallest ones are **A-class, followed by B, C, M and X, the largest.**
- **Geomagnetic Storms:**
  - When Earth is in the path of CME.
  - **Impacts:**
    - **Technological Systems:**
- Satellite malfunction: In 2022, a minor geomagnetic storm caused the failure of 40 Starlink satellites.
- Navigation errors: GPS signal degradation affects civil aviation and defence navigation.
    - **Power Infrastructure:**
- Grid failure risks: The 1989 Quebec blackout exemplifies how CMEs induce geomagnetically induced currents (GICs), damaging transformers.
    - **Aviation and Space:**

- Polar flights face increased radiation exposure, forcing rerouting.
- Astronauts aboard the ISS require radiation shielding during intense solar events.
    - **Military and Strategic Systems:**
- Ballistic missile early warning systems and radar operations can be temporarily disabled.
    - **Auroras**
- **Auroras:**
    - Aurora (polar light) is a natural phenomenon which is characterised by a display of lights in the **Polar region** skies.
    - **Solar wind has charged protons and electrons that excite atmospheric Oxygen and Nitrogen.**



    - **When Oxygen and Nitrogen return from excited states to the ground state, they release Photons, which are displayed as the Aurora lights.**
    - Northern Hemisphere: **Aurora Borealis**
    - Southern Hemisphere: **Aurora Australis**
    - **Why are auroras only on the poles?**
        - Earth is a magnet. **Currently, the North and the South Poles of this magnet are near our geographical North and South Poles.**
        - The charged particles get attracted towards the magnetic Poles.
        - This causes auroras at the geographical north and south poles.
- **Van Allen Radiation Belt:** The Van Allen radiation belts consist of **charged particles trapped by Earth's magnetic field** , forming doughnut-shaped regions around the planet. They protect Earth from solar radiation but can pose risks to spacecraft passing through them.

- The **inner and outer radiation belts** are two distinct regions within the Van Allen radiation belts surrounding Earth. The **inner belt is composed mainly of protons** and extends from about 1,000 to 6,000 kilometers above the Earth's surface, while the **outer belt consists mostly of electrons** and extends from about 13,000 to 60,000 kilometers above the surface.



- **Sunspots** : **darker (cooler) areas** on the solar surface.
- **Solar Maxima & Solar Minima:**
  - Solar minima and maxima are the **two extremes of the Sun's 11-year and 400-year activity cycles** .
  - Solar maximum: The **highest solar activity** is observed during this period. The Sun has many sunspots and eruptions of solar flares. We also receive the **highest insolation** during Solar Maxima, because the Sun emits more energy due to increased sunspot and solar flare activity, boosting total solar irradiance reaching Earth. Right now sun is in the **Maxima Phase.**
  - Solar minimum is the period of **least solar activity** in the 11-year solar cycle.
- **Goldilocks Zone:**



  - **Distance from the star where a planet’s atmospheric conditions are such that water remains liquid.**
  - Distance from the star where the Goldilocks zone exists depends on the size and the strength of the star.
  - E.g. Earth; Kepler-186f.

⚡ **Quick revise** The Sun Fold ▲

> **Sun = a samosa** (inside → out): **Core** = hot filling, fusion, **> 15 million °C** → **Radiative zone** = mashed potato (energy by radiation) → **Convective zone** = bubbling oil (convection cells) → **Photosphere** = golden crust (where Sun turns opaque; no real "surface") → **Chromosphere** = oil sheen (outer part ~**10,000 °C**) → **Corona** = steam (hot plasma, millions of km).

Composition: **71% H, 27.1% He, 0.97% O, 0.4% C**.

Space-weather term	What it is	Remember
Solar wind	Charged particles streaming from the corona	plasma too hot for gravity to hold
Solar flare	Explosion from twisted magnetic fields; X-rays + UV	classes **A, B, C, M, X** (X biggest); *may or may not* come with CME
CME	Plasma + magnetic field blobs from corona	**15–18 h** to reach Earth
Geomagnetic storm	Earth in path of a CME	Starlink **40 satellites** lost (2022); **Quebec blackout 1989**; GPS errors; polar flight rerouting; radar/missile-warning disruption
Aurora	Solar-wind particles excite O and N → photons	**Borealis** (N), **Australis** (S); at poles because the magnetic poles are near them
Van Allen belts	Charged particles trapped by Earth's field	inner: **protons**, 1,000–6,000 km · outer: **electrons**, 13,000–60,000 km
Sunspots	Cooler, darker patches	**11-year** (and 400-year) cycle; now in **Maxima**

PYQ 2022 solar storm → answer (c): *possible* effects = GPS/navigation failure, power-grid damage, **intense auroras**, satellite orbit disturbance, polar short-wave radio disruption. **Not** tsunamis (need a vertical water shift) and **not** forest fires (flares don't raise Earth's temperature). Handout note: the May 2024 storm showed auroras as far south as **Leh**.

Goldilocks zone (PYQ 2015, ans c): distance from a star where **water stays liquid** – used in the **search for Earth-like planets** (Earth; Kepler-186f).

9 · EVOLUTION OF EARTH

Phases:

- **Crust formation:**
  - **Light substances floated up from deep inside** , and then they changed into hard rocks after cooling.
  - As the earth's interior continued to cool, it contracted, and the outer crust wrinkled forming ridges and basins.
- **Atmosphere Formation:**
  - **Degassing:** lighter substances swarmed up above the earth's surface forming an atmosphere of gases.
  - **Happened in 3 phases:**
    - **Early Atmosphere was rich in Hydrogen and Helium** . These gases were eventually **stripped away by solar wind.**
    - Stage 2: The **volcanic activities** on partially solidified Earth, led to the release of gases like **Co2, Nitrogen Oxides, Sulphur Oxides and water vapour** . The water vapour released formed clouds and led to the formation of oceans through rainfall.
    - Stage 3: **As life evolved** on Earth, the configuration of the atmosphere changed, and the concentration of **Oxygen** started to increase.



- **Ocean Formation:**
  - Atmospheric Gases formed the Clouds, which cause **heavy rains for thousands of year**



  - This **water was collected in basins, forming oceans.**
  - Then **life started in the ocean** .
- **Life on Earth:**
  - **4.5 Billion Years Ago:** Earth forms, hot and bombarded by asteroids.
  - **Over the next billion years:** As Earth cools, conditions become favorable for **complex molecules** to form in the oceans.
  - **Around 3.8 Billion Years Ago: Simple life emerges** , possibly near deep-sea vents, utilizing chemicals for energy (not sunlight). These were likely **single-celled**

organisms (blue-green algae) with basic building blocks of life.

- **Proof: Chemical Biogenesis or the Miller-Urey experiment, conducted in 1953** by Stanley Miller and Harold Urey. It proved that in plausible early Earth conditions, organic molecules important for life could arise naturally. Inorganic molecules could convert to organic ones. This can lead to the creation of necessary amino acids, essential for the creation of single-celled organisms that could divide.
- **LUCA: Last Universal Common Ancestor.** Might have existed about **4.2 billion years** ago.

⚡ **Quick revise** Crust → atmosphere → ocean → life Fold ▲

- **Crust → atmosphere → ocean → life.** Atmosphere in **3 stages:** (1) H & He, blown away by solar wind → (2) **degassing**: volcanoes give CO_2, N & S oxides, water vapour → clouds → rain → **oceans** → (3) life raises **oxygen** (photosynthesis).
- **Timeline:** Earth **4.5 bya** · simple life **3.8 bya** near deep-sea vents (blue-green algae) · **LUCA ~4.2 bya** (as printed) · **Miller–Urey 1953** (organic molecules from inorganic).
- **PYQ 2018:** Statements 1 (magnetic field reverses every few hundred thousand yrs) ✔ and 3 (life modified early atmosphere) ✔ are right; "54% oxygen, no CO_2 at birth" ✘. → **1 & 3**. (Early air: water vapour, CO_2, N_2, no O_2.)
- **PYQ 2012:** elements behind origin of life = **Carbon, Hydrogen, Nitrogen**.

10 · EARTH FACTS:

- **Titled on Axis** : **23.5 degrees** from the vertical. (**towards Pole/North Star**)



- According to a study, between 1993 and 2010, excessive groundwater extraction, particularly in regions like northwestern India and western North America, caused Earth's rotational pole to drift approximately 80 centimeters eastward (0.00000719 degrees).

- Melting of ice caps in higher latitudes, and migration of water to lower latitudes, is also causing changes in Earth's axis.
- **1 rotation: 23 hrs 56 mins 4.1 seconds** (sidereal day).
- **Speed of rotation is reducing** due to:
  - Wind and wave friction.
  - Movement of moon away from us.
  - Slowdown of the Core, and its movement in opposite direction.
- **Looking down at the North Pole from space, Earth is rotating in a counterclockwise direction.**
- Revolution in elliptical orbit. **365 days, 5 hours and 48 minutes** (sidereal year) **.**
- **Aphelion:** Point of the Earth's orbit that is **farthest away from the Sun** .
- **Perihelion:** Point of the Earth's orbit that is **nearest to the Sun** .

Geomagnetism

- Paleomagnetic studies of rocks and ocean sediment have demonstrated that the **orientation of the earth's magnetic field has frequently alternated over geologic time** .
- Flips every 2-3 lakh years.
- Periods of "normal" polarity (i.e., when the north-seeking end of the compass needle points toward the present north magnetic pole, as it does today) have alternated with periods of "reversed" polarity (when the north-seeking end of the compass needle points southward).
- Currently, there has been an accelerated movement of Earth's magnetic North Pole towards Russia. The magnetic North Pole has been drifting from northern Canada into the Arctic Ocean and towards Siberia since the 1990s, with its speed varying over time
- Importance of Earth's magnetic Field:
  - Protects Earth from harmful solar wind and charged particles.
  - Prevents excessive atmospheric loss into space.
  - Creates the magnetosphere around Earth, which contains the Van Allen Belts. (Magnetic field = force; magnetosphere = zone of influence of that force)
  - Causes auroras (Northern and Southern Lights) near polar regions.
  - Helps in navigation using magnetic compasses.
  - Important for migratory animals and birds for orientation.
  - Protects satellites, communication systems, and power grids from solar storms.
  - Maintains conditions suitable for life on Earth.

⚡ **Quick revise** Earth – evolution, motion, Moon, time Fold ▲

Earth fact	Value
Axial tilt	**23.5°**
Rotation (sidereal day)	**23 h 56 m 4.1 s**, West→East (counter-clockwise from above the North Pole); slowing (wind/wave friction, Moon receding, core slowdown)

Earth fact	Value
Revolution (sidereal year)	**365 d 5 h 48 m**; **aphelion** farthest, **perihelion** nearest
Pole drift	Groundwater pumping (NW India, W North America) shifted the pole ~**80 cm east** (1993–2010); melting ice moving water to low latitudes also shifts the axis
Magnetic field	Flips every **2–3 lakh years**; magnetic north drifting from Canada → Arctic → **Siberia** since the 1990s

Why the magnetic field matters: blocks solar wind, stops atmospheric loss, creates the **magnetosphere** (holds Van Allen belts), causes auroras, helps compasses, guides migrating animals, protects satellites/grids.

Day/night & seasons – PYQs 2012, 2013, 2019, 2024: - Pole star lies due North: villager 5 km **east** keeps the pole star **on his left** (2012). - Variation of day length with season = **revolution on a tilted axis + latitude** (ans B, 2013). With no tilt, no seasons and equal days. - **21 June:** Sun does **not set at the Arctic Circle** (2019, ans A). >12 h daylight on 21 June at the **Tropic of Cancer and Arctic Circle** (2024, ans d: 2 and 4). Equator gets overhead Sun only at the **equinoxes**.

11 · MOON



- The moon is **Earth's only natural satellite.**
- It orbits the Earth at an **average distance of 384,400 km.**
- The moon is **a bit more than one-fourth (27%) the size of Earth.**
- Its gravity influences Earth's tides.
- **Size** : **27%** of the Earth.
- **Perigee** : The point on the Moon's orbit **closest to the Earth** .
- **Apogee:** The point on the Moon's orbit **farthest away from the Earth** .
- **Tidally locked to the Earth:**
  - This means that the moon **always shows one face to the Earth** .

- Earth’s gravitational interaction slowed down the moon’s rotation until it became tidally locked.

- **Phases of the Moon**



- **Creation:**
  - **Fission Theory, George Darwin (1838):** The Moon once formed part of the **rapidly spinning Earth** , which **flung off a large chunk due to the centrifugal force,** creating the Moon. This explains the Moon's receding path from Earth and Earth's slightly bulging equatorial region. Not accepted anymore.
  - **Giant Impact Hypothesis (Big Splash): collision between the proto-Earth and a Mars-sized planet (called Theia)** , around **4.5 bn years ago** created the Moon.
  - **Binary Exchange Capture Theory (Proposed):** The moon was not born out of Earth, but was infact part of a binary system of celestial objects. When they came close to the Earth, earth captured one and expelled away the other part of the binary system.



⚡ **Quick revise** Earth – evolution, motion, Moon, time Fold ▲

🌙 **Moon quick-fire:** 384,400 km · 27% of Earth's size · **tidally locked** (same face always) · **Perigee** = **P**ear-close, **Apogee** = **A**way. **Origins:** Fission (George Darwin) – rejected · **Giant Impact** (Theia, Mars-sized, 4.5 bya) – accepted · Binary-exchange capture – proposed.

12 · ECLIPSES:

- **Lunar Eclipse:** Earth between the Sun and the Moon



  - **Umbra: darkest region** of a shadow
  - **Penumbra:** region in which **only a portion of the light source is blocked** by the occluding body.
  - **Blood Moon: When the whole of the lunar disc enters the Earth's umbra** , the Moon appears reddish.
  - **Blue Moon:** Not related to the eclipses. When we get **2 full moons in a month, the 2 nd one is called Blue Moon.**
- **Solar Eclipse: Moon Between** Earth and the Sun.
  - The moon blocks the light of the sun from reaching the Earth's surface and casts a shadow (Umbra) on the Earth.
  - **Annular Solar Eclipse:** When the **Moon** passes directly between the Earth and the Sun but **does not completely cover the Sun's disk** , as the Moon is farther away from Earth, making its shadow on Earth smaller.





- **Total Solar Eclipse:** When the **moon completely covers the sun** , as it is closer to the Earth. It also leads to the **Diamond ring effect** (As the Moon moves across the face of the Sun, a point comes when only a single bright spot of sunlight remains visible, looking much like a shining diamond.)



- **Partial Solar eclipse can be due to 2 reasons:**
  - When sun, moon and the earth are **not in a straight line** , only a penumbra region is formed, causing partial eclipse.
  - When they are in a **straight line, some regions see a full solar eclipse (umbra) while some see a partial one (penumbra).** All Lunar eclipses are Full Moon days but not all Full Moon days are Lunar Eclipses. All Solar Eclipses are New Moon days but not all New moon days are solar eclipse days. This is because the moon’s orbital plane is titled at 5 degrees compared to Earth-Sun orbital plane.



Latitudes and Longitudes:

- **Imaginary lines** created to make it easy to locate places on the Earth.



- **Latitudes: Parallel lines** running in **east-west directions** are latitudes.
- **Longitudes:** Lines running **North to South** and **meeting** each other **at the poles** are the longitudes.
- **Climatic Zones**
- **Time Zones**
  - Earth takes **4 minutes to rotate by 1 degree longitude**
  - **Earth rotates from West to East,** that's why **Eastern locations experience the sun before west** , and are thus ahead in time compared to western locations.
  - The **International Date Line exists to maintain a standard calendar day** globally by marking the point where the date changes by one day when crossing time zones.



⚡ **Quick revise** Earth – evolution, motion, Moon, time Fold ▲

Time. 4 minutes per 1° of longitude; Earth rotates W→E so east is ahead; **International Date Line** keeps the calendar day consistent.

List of Countries with the most timezones:

Rank	Country	Time Zones
1	France	12
2	Russia	11
2	USA	11
3	Antarctica*	9
3	Australia	9
3	UK	9
4	Canada	6
5	Denmark	5

⚡ **Quick revise** Eclipses Fold ▲

Eclipses. *Lunar* = Earth between Sun and Moon (**Blood Moon** = whole Moon inside umbra; **Blue Moon** = 2nd full moon in a month, *no relation to eclipses*). *Solar* = Moon between Sun and Earth; **Annular** (Moon farther → ring), **Total** (Moon closer; diamond-ring effect), **Partial**

(penumbra only). Not every full/new moon gives an eclipse because the Moon's orbit is tilted **5°**.

⚡ **Quick revise** Earth – evolution, motion, Moon, time Fold ▲

🕐 **Most time zones:** France **12** · Russia & USA **11** ("two elevens – a cricket team each") · **A-A-U** = Antarctica, Australia, UK **9** · Canada **6** · Denmark **5**.

13 · PYQ

Q.1) If a major solar storm (solar flare) reaches the Earth, which of the following are the possible effects on the Earth ? (2022)

2. GPS and navigation systems could fail.

3. Tsunamis could occur at equatorial regions.

4. Power grids could be damaged.

5. Intense auroras could occur over much of the Earth.

6. Forest fires could take place over much of the planet.

7. Orbits of the satellites could be disturbed.

8. Shortwave radio communication of the aircraft flying over polar regions could be interrupted.

Select the correct answer using the code given below:

(a) 1, 2, 4 and 5 only

(b) 2, 3, 5, 6 and 7 only

(c) 1, 3, 4, 6 and 7 only

(d) 1, 2, 3, 4, 5, 6 and 7

Answer - C

Explanation: (e) Solar storms can play a havoc on the electrical and electronic systems of the Earth, affecting power systems, satellite communication, nd can cause intense auroras. In May 2024 one such intense solar storm caused the auroras to be seen as far away from the pole as Leh in India. (f) Tsunamis require vertical shift of water which is not caused by the flares. These flare are not related to increased temperature on earth, so they do not cause forest fires.

Q.2) Which one of the following sets of elements was primarily responsible for the origin of life on the Earth? (2012)

- Hydrogen, Oxygen, Sodium
- Carbon, Hydrogen, Nitrogen
- Oxygen, Calcium, Phosphorus
- Carbon, Hydrogen, Potassium

Answer. B

Explanation The correct answer to this question, as most of you know, is Carbon, Hydrogen and Nitrogen.

- In fact, the early biochemistry was not simply the reactions of Carbon, Hydrogen, Nitrogen and Oxygen was a bit far from it, was CO2, N and water resulting in the organic acids such as oxalic acid.
- The infrared part of solar radiation. The increasing amount of carbon dioxide in the air is slowly raising the temperature of the atmosphere because it absorbs the infrared part of the solar radiation. Only water vapour has the ability to absorb both incoming UV and outgoing infrared radiation.

Q.3) A person stood alone in a desert on a dark night and wanted to reach his village which was situated 5 km east of the point where he was standing. He had no instruments to find the direction but he located the polestar. The most convenient way now to reach his village is to walk in the: (2012)

- direction facing the polestar
- a direction opposite to the polestar
- direction keeping the polestar to his left
- direction keeping the polestar to his right

Answer. C

Explanation

- The correct answer is direction, keeping the pole star to his left. In ancient times, people used to determine directions during the night with the help of stars. The North star indicates the north direction. It is also called the Pole Star.

Q.4) Variations in the length of daytime and nighttime from season to season are due to - (2013)

4. The earth’s rotation on its axis

5. Revolution of the earth on a tilted axis

6. Latitudinal position of the place

7. The earth’s revolution around the sun in an elliptical manner

Answer. B

Explanation Had the axis of the Earth been perfectly perpendicular to the plane of the ecliptic, the sun rays would have been vertical over the equator all year round and the

amount of energy received at any place would have remained constant throughout the year; hence no change of seasons.

- Also in that situation, the days and nights all over the earth would have been exactly equal throughout the year as it happens at the time of equinoxes.
- The change between day and night is caused by the rotation of the Earth on its axis. ... The changing length of days and nights depends on where you are on Earth and the time of year. Also, daylight hours are affected by the tilt of the Earth's axis and its path around the sun.
- As the Earth moves around the Sun, the length of the day changes. ... This is all caused by the 23.5-degree tilt of the Earth's axis as it travels around the sun. In the Northern Hemisphere, days are longest at the time of the summer solstice in June, and the shortest days are at the winter solstice in December.

Q.5) Consider the following statements:

- The Earth's magnetic field has reversed every few hundred thousand years.
- When the Earth was created more than 4000 million years ago, there was 54% oxygen and no carbon dioxide.
- When living organisms originated, they modified the early atmosphere of the Earth.

Which of the statements given above is/are correct? (2018)

- 1 only
- 2 and 3 only
- 1 and 3 only
- 1, 2 and 3

Answer. C

Explanation

- A geomagnetic reversal is a change in a planet's magnetic field such that the positions of magnetic north and magnetic south are interchanged. The Earth's field has alternated between periods of normal polarity, in which the predominant direction of the field was the same as the present direction, and reverse polarity, in which it was the opposite. These periods are called chrons.
- Reversal occurrences are statistically random. There have been 183 reversals over the last 83 million years.
- The Earth formed approximately 4.5 billion years ago, along with the other seven planets in the solar system.
- As the Earth cooled, a primitive atmosphere was created by the out-gassing of early volcanoes.
- The early atmosphere contained no oxygen and would have been toxic to human beings, as well as most other life on Earth today.
- Based on an analysis of gases vented by modern volcanoes, it seems likely that this early atmosphere consisted mostly of water vapour (H_2O) and carbon dioxide (CO_2) and nitrogen gas (N_2).
- In the whole process of evolution, the living organism changed the chemical composition of the Atmosphere. For example, oxygen came from photosynthesis.

Q.6) On 21st June, the Sun (2019)

- does not set below the horizon at the Arctic Circle
- does not set below the horizon at Antarctic Circle
- shines vertically overhead at noon on the Equator
- shines vertically overhead at the Tropic of Capricorn

Answer. A

Explanation

- The Sun is directly overhead at "high-noon" on the equator twice per year, at the two equinoxes. On the Arctic Circle, the Sun does not set at all on the Summer Solstice which occurs on 21st June.
- On that one day, the Sun traces a complete circle just above the horizon as the Earth rotates.

Q.7) On June 21 every year, which of the following latitude(s) experience(s) a sunlight of more than 12 hours? (2024)

- Equator
- Tropic of Cancer
- Tropic of Capricorn
- Arctic Circle

Select the correct answer using the code given below:

(a) 1 only

(b) 2 only

(c) 3 and 4

(d) 2 and 4

Answer: (d) 2 and 4

Q.8) Consider the following statements: Statement I: Scientific studies suggest that a shift is taking place in the Earth's rotation and axis. **Statement II:** Solar flares and associated coronal mass ejections bombarded the Earth's outermost atmosphere with tremendous amount of energy. **Statement III:** As the Earth's polar ice melts, the water tends to move towards the equator.

Which one of the following is correct in respect of the above statements?

(a) Both Statement II and Statement III are correct and both of them explain Statement I

(b) Both Statement II and Statement III are correct but only one of them explains Statement I

(c) Only one of the Statements II and III is correct and that explains Statement I

(d) Neither Statement II nor Statement III is correct

Answer: (b) Statement I: Correct Scientific studies, including those from NASA and peer-reviewed journals, confirm that Earth's axis and rotation are gradually shifting due to

redistribution of mass (e.g., melting glaciers and sea-level rise). **Statement II:** Correct, but does not explain Statement I Solar flares and coronal mass ejections (CMEs) do indeed impact the Earth's magnetosphere and upper atmosphere. However, they do not significantly affect the Earth's axial tilt or rotation. **Statement III:** Correct and explains Statement I Melting polar ice (especially in Greenland and Antarctica) leads to mass redistribution toward oceans, which contributes to a shift in the Earth's rotational axis (a phenomenon called true polar wander).

Q.9) Consider the following statements: Statement I: Anadyr in Siberia and Nome in Alaska are a few kilometers from each other, but when people are waking up and getting set for breakfast in these cities, it would be different days. **Statement II:** When it is Monday in Anadyr, it is Tuesday in Nome.

Which of the statements given above is/are correct?

(a) I only

(b) II only

(c) Both I and II

(d) Neither I nor II

Answer: (a) I only Statement I: Correct Anadyr is in the Chukotka Autonomous Okrug, Russia (Siberia). Nome is in western Alaska, USA. They are about 800 km apart, separated by the Bering Strait. Time Difference: Anadyr uses UTC+12. Nome uses UTC−9 (Alaska Standard Time). Statement II: Incorrect When it is Monday in Anadyr, it is not Tuesday in Nome : it is actually Sunday.

Q.10) Consider the following countries: 1.United Kingdom 2.Denmark 3.New Zealand 4.Australia 5.Brazil

How many of the above countries have more than four time zones?

(a) All the five

(b) Only four

(c) Only three

(d) Only two

Answer: (b) Only four United Kingdom

- Time zones:
  - Mainland: GMT or BST (1 zone)
  - Overseas territories include Bermuda, Falklands, Gibraltar, etc.
- Total time zones: Up to 9, including overseas territories. Denmark : More than 4 Time Zones
- Mainland: UTC+1 (CET)
- Greenland:
  - Danmarkshavn : UTC+0
  - Ittoqqortoormiit : UTC−1

- Nuuk (Capital) : UTC−3
- Thule : UTC−4

- Total: 5 Time Zones New Zealand : More than 4 Time Zones
- Mainland: NZST/NZDT (UTC+12/+13)
- Territories:
  - Tokelau : UTC+13
  - Cook Islands : UTC−10
  - Niue : UTC−11
- Total: 5 Time Zones Australia : More than 4 Time Zones
- Mainland:
  - AWST (UTC+8), ACST (UTC+9:30), AEST (UTC+10)
- Territories:
  - Lord Howe Island : UTC+10:30
  - Norfolk Island : UTC+11
  - Cocos Islands : UTC+6:30
  - Christmas Island : UTC+7
- Total: 6+

Brazil : Only 4 Time Zones

- Historically had 4 zones: UTC−2 to UTC−5
- Now has 3 official zones, but still only 4 recognized historically
- Not more than 4

⚡ **Quick revise** PYQs with no year printed Fold ▲

PYQs with no year printed: (a) *Anadyr vs Nome* → only Statement I right (different days at breakfast; but Monday in Anadyr is **Sunday** in Nome, not Tuesday). (b) *UK, Denmark, NZ, Australia, Brazil* – **four** have more than four zones (Denmark 5, NZ 5, Australia 6+, UK 9 with territories; Brazil only 4). (c) *Shift in Earth's rotation/axis* → answer (b): melting polar ice moving water toward the equator explains it; solar flares are true but don't explain it.

14 · INTERIOR OF THE EARTH

Why do we need to know about this?

- Understanding the earth's interior is essential to **understanding the nature of changes that take place over and below the earth's surface.**
- To understand **geophysical phenomena** like volcanism, earthquakes, etc.
- To understand the **internal structure of various objects of the solar system.**
- To understand the **evolution and present composition of the atmosphere.**

- Future **deep-sea mineral exploration** etc.

Sources:

- **Direct Sources:** Facilitate direct observation of what's under the surface.
  - **Deep Earth mining and Drilling** (deepest mine: Mponeng gold mine and TauTona gold mine in South Africa at 3.6km; deepest drilling 12km in Kola peninsula, Russia).
  - **Volcanic Eruptions** : Bring the material inside the Earth to the surface, giving us an insight into what is present under the Earth.
- **Indirect Sources** : Help us statistically extrapolate what's inside the Earth.
  - **Meteors and other celestial bodies** (other planets, moons, etc): Similar formation Process as Earth. So, the internal structure of these bodies will be similar to that of the Earth. Helps in expanding our imagination as to how Earth would look from the inside.



  - **Gravitation** : Gravity value is not the same everywhere, and this is known as a gravitational anomaly, helps us identify material distribution inside the earth. Wherever heavier materials have accumulated, gravitational force will be higher as it is directly proportional to the mass of the bodies in question.
  - **Shifts in our magnetic fields:** Our magnetic field constantly keeps moving. It reverses in approximately 200000 to 300000 years. It is the material inside the earth that is responsible for the magnetic field. This reversal gives us the idea that the material inside the earth can move.
  - **Earthquake waves:** Body waves that are generated during earthquakes are by far the best indirect source of information. (Details in the earthquake section.)
- **Other minor sources and observations:**
  - **Volcanic eruptions and existence of hot springs, geysers** etc. point to an interior which is very hot.
  - The high temperatures are attributed to automatic disintegration of the radioactive substances.
  - Gravitation and the diameter of the earth helps in estimating pressures deep inside

LAYERS OF EARTH

- Earth's layers are identified by studying various direct and indirect sources.



- The structure of the earth's interior is made up of several **concentric layers.**
- Broadly based on chemical properties, three layers of earth can be identified— crust, mantle, and core.

Crust: 30-50 km

- The outer thin layer of the Earth.
- It has 2 parts: the **denser oceanic (basaltic) and the lighter continental (granitic) crust.**
- The **outer covering** of the crust is of **sedimentary material (granitic rocks)** and **below** that lie **crystalline, igneous and metamorphic rocks** which are acidic in nature.
- The lower layer of the crust consists of basaltic and ultra-basic rocks.
- **Oceanic crust (SiMa) denser and thinner** (5-30km)



- **Continental Crust (SiAl) lighter and thicker** (50-70km).
- The **continental crust is thicker** in the areas of **major mountain systems** . It is as much as 70 -100 km thick in the Himalayan region.
- The continental and the oceanic crusts are **separated by the Conrad discontinuity.**
- It forms **0.5-1.0 per cent of the earth's volume.**

Mantle:

- **Mohorovicic Discontinuity to 2900km (Gutenberg Discontinuity)** .
- It forms **83% of Earth's volume** .
- **Denser than crust** (upper is 2.9-3.3g/cm3; lower is 3.3-5.7g/cm3; average density 4.5g/cm3).
- The **crust and the uppermost part of the mantle** are called **LITHOSPHERE** . Its thickness ranges from **10-200 km** . It is **Brittle** in nature.

- Below the Lithosphere lies the **ASTHENOSPHERE, till 400 km depth** . It is a **semi-solid molten** layer, entirely within the mantle. This layer is the source of Magma.
- The lower mantle extends beyond the asthenosphere. It is in solid state.
- The **upper and the lower mantle are separated by Repetti Discontinuity.**
- The mantle and the core are separated by Gutenberg Discontinuity.

Core: 2900-6400km

- Lies between **2900 km and 6400 km** below the earth's surface.
- It accounts for **16 per cent** of the **earth's volume** .
- Core has the **heaviest mineral materials** of the highest density.
- It is composed of nickel and iron **[NiFe]** .
- The **outer core is liquid** while the **inner core is solid** .
- They are separated by **Lehman's Discontinuity.**





Sl. No.	Elements	By Weight(%)
1.	Oxygen	46.60
2.	Silicon	27.72
3.	Aluminium	8.13
4.	Iron	5.00
5.	Calcium	3.63
6.	Sodium	2.83
7.	Potassium	2.59
8.	Magnesium	2.09
9.	Others	1.41



- **Discontinuities** : the places inside the Earth's crust where the seismic waves experience an abrupt change in their velocities and/or direction.

PYQ

Q.1) How does the Juno Mission of NASA help to understand the origin and evolution of the Earth? **(2017, 10m)**

⚡ **Quick revise** Earth's interior & earthquakes Fold ▲

🥚 **Earth = a boiled egg.** Shell = **crust** (0.5–1% of volume, 30–50 km) · White = **mantle** (**83%**, to 2,900 km) · Yolk = **core** (**16%**, 2,900–6,400 km, **NiFe**; outer liquid, inner solid).

Boundaries, top → bottom: Conrad · **Moho** · **R**epetti · **G**utenberg · **L**ehmann → *"Cool Men Really Get Love"*. Conrad (continental/oceanic crust) · Mohorovičić (crust/mantle) · Repetti (upper/lower mantle) · Gutenberg (mantle/core, 2,900 km) · Lehmann (outer/inner core).

Crust	Mantle	Core
Oceanic (SiMa): dense, thin, 5–30 km, basaltic · **Continental (SiAl)**: light, 50–70 km (70–100 km under Himalaya), granitic	Upper 2.9–3.3, lower 3.3–5.7, avg **4.5 g/cm³**. **Lithosphere** = crust + uppermost mantle (10–200 km, brittle). **Asthenosphere** to ~400 km: semi-solid, **source of magma**	Heaviest, densest; nickel + iron

How we know: *Direct* – deepest mines (Mponeng, TauTona **3.6 km**), deepest drill (**Kola, 12 km**), volcanic eruptions. *Indirect* – meteorites, **gravity anomalies** (heavy material = higher gravity), magnetic field shifts, and best of all **earthquake waves**.

⚡ **Quick revise** Earthquake basics — Fold ▲

Mains 2017 (10m): "How does NASA's Juno mission help understand the origin and evolution of the Earth?" – the handout prints the question only, no answer. Notes on Jupiter above (gas giant, storms) are all it offers.

15 · EARTHQUAKES

- **Abrupt release of energy along a fault** causes earthquake waves.



- A fault is a sharp break in the crustal rock layer.
- Rocks along a fault tend to move in opposite directions. But the friction exerted by the overlying rock strata prevents the movement of the rock layer. With time pressure builds up.
- Under intense pressure, the rock layer, at a certain point, overcomes the friction offered by the overlying layer and undergoes an abrupt movement generating shockwaves.
- This causes a **release of energy** , and the **energy waves travel in all directions** .
- The **point where the energy is released** is called the **focus** of an earthquake, alternatively, it is called the **hypocentre.**



- The **point on the surface, nearest to the focus** , is called the **epicentre** .

⚡ **Quick revise** Focus (hypocentre) Fold ▲

Focus (hypocentre) = where energy is released; **epicentre** = surface point nearest the focus. Distribution: mostly at plate boundaries; **two belts – Circum-Pacific, Mid-continental**; **deep-focus** at convergent, **shallow-focus** at divergent boundaries.

Scales: Richter – magnitude from surface-wave amplitude, log scale 0–10 (5 → 6 = ×10 amplitude) · **Mercalli** – intensity (damage) **1–12** · **Moment magnitude** – energy released (fault area, slip, rock rigidity), log, **most widely used now**.

16 · Seismic waves

- **Recorded by Seismographs** installed across the world in Seismic stations.
- **Body Waves:** Generated due to the release of energy at the focus and move in all directions travelling through the body of the earth. **P-waves & S-waves** are the 2 body waves.
- The **velocity** of the Body waves **changes as they travel through materials with different elasticity** (stiffness) (Generally density with few exceptions). The **more elastic the material** is, the **higher the velocity** . Their **direction also changes** as they reflect or refract when coming across materials with different densities.
- When an earthquake or underground nuclear test sends shock waves through the Earth, the cooler areas, which generally are rigid, transmit these waves at a higher velocity than the hotter areas.
- The **body waves interact with the surface rocks** and **generate** new sets of waves called **surface waves.**



- **P-waves** :
  - **Longitudinal or Compressional** Waves
  - **Particles** of the medium **vibrate along the direction of propagation of the wave** .
  - **Fastest, first to arrive** ; **Primary Waves**
  - Their velocity depends on the shear strength or elasticity of the material.
  - Can travel in all mediums: Velocity in **Solids > Liquids > Gases.**



  - **Shadow zone: 103-143 degrees on both sides** .
  - This gives clues about Solid inner core
- **S-waves:**
  - **Transverse or distortional** waves.
  - Analogous to **water ripples** or light waves.
  - Arrive at the surface with **some time lag. Secondary waves** .
  - Travel at varying velocities (proportional to shear strength) through the solid part of the Earth's crust, mantle.
  - **Cannot pass through gases or liquids.**
  - **Particles move perpendicular to the direction of propagation** of the wave.
  - **Shadow zone:** between **103 degrees and 180 degrees** .



- This observation led to the **discovery of a liquid outer core** . Since S waves cannot travel through liquid, they do not pass through the liquid outer core.

- **L-Waves:**
  - **Love waves.**
  - Shear surface waves (transverse like the S-waves)
  - Have the **highest velocity of all surface waves** and can cause **significant horizontal shaking** of the ground.
  - **3rd** to reach the seismograph.
  - Cause **displacement of rocks** , and hence, the collapse of structures occurs.
- **R-Waves:**
  - **Rayleigh Waves.**
  - They are characterized by an **elliptical or rolling motion** that combines both vertical and horizontal particle motion. This motion is similar to the motion of ocean waves.



  - **Slowest surface waves**
  - Responsible for **Tsunamis** due to the vertical shift.
  - 8. **Shadow Zones:** (i) The **Body waves** get recorded in seismographs located at far-off locations. However, there exist some **specific areas where the waves are not reported.** Such a zone is called the ‘shadow zone’. (ii) The study of different events reveals that an altogether different shadow zone exists for each earthquake. (iii) It was observed that seismographs located at any distance within 103° from the epicentre, recorded the arrival of both P and S-waves. (iv) However, the seismographs located beyond 143 ° from the epicentre, record the arrival of P-waves, but not that of S-waves.

(v) Thus, a zone between 103° and 143° from the epicentre was identified as the shadow zone for both types of waves. The entire zone beyond 103 ° does not receive S-waves. (vi) The **shadow zone of the S-wave is much larger than that of the P-wave.** The shadow zone of P-waves appears as a band around the earth between 103 ° and 143 ° away from the epicentre. (vii) The shadow zone of S-waves is not only larger in extent, but it is also a little over 40 per cent of the earth's surface.

- **How does this help us determine Earth's interior ?**
  - Reflection causes waves to rebound whereas refraction makes waves move in different directions.



  - The variations in the direction of waves are inferred with the help of their record on seismographs.
  - Change in densities greatly varies the wave velocity. By observing the changes in velocity, the density of the earth can be estimated.
  - By observing the changes in the direction of the waves (emergence of shadow zones), different layers can be identified.
- **Distribution:**
  - Mostly at the **plate boundaries**
  - 2 major belts: Circum-Pacific belt; Mid continental belt.
  - **Deep focussed** earthquakes at **convergent boundaries** .
  - **Shallow focussed** earthquakes at **diverging boundaries** .
- **How are Earthquakes Measured ?**
  - **Richter's Scale** : Measures earthquake magnitude by **calculating the amplitude of the surface waves** generated during the event.
    - It is a **logarithmic scale** , which means that the amplitude of earthquake waves at Ritcher scale 5 is 10 times less than that at 6.
    - The number indicating magnitude **ranges between 0 to 10** .
  - **Mercalli's Scale** : Measures the intensity of the earthquakes, based on the **actual level of destruction** caused by the event.

- The number indicating intensity **ranges between 1 to 12**

- **Moment Magnitude Scale:**
  - **Most widely used** now.
  - Unlike the Richter scale, which measures the amplitude of seismic waves recorded by seismographs, the moment magnitude scale considers the **total energy released by an earthquake** , including the **area of the fault rupture** , the **average slip along the fault** , and the **rigidity of the rocks** involved.
  - Also, a **logarithmic scale** .



⚡ Quick revise Earthquake basics

Fold ▲

Wave	Type	Motion	Medium	Order
P (primary)	Body, longitudinal	to-and-fro along travel	solid, liquid, gas (faster in solids)	**1st, fastest**
S (secondary)	Body, transverse	up-down, at right angles	**solids only**	2nd
L (Love)	Surface, shear	horizontal shaking; fastest surface wave	–	3rd; collapses buildings
R (Rayleigh)	Surface, rolling ellipse	vertical + horizontal	–	**slowest**; **tsunamis**

🌊 **P = Push-pull, Prompt. S = Sideways, Solids only.** *Love shakes Left-right; Rayleigh Rolls like a surf.* **Shadow zones: P: 103°–143°** (a band) · **S: everything beyond 103°** (bigger, ~40% of the globe). S-waves vanish → **liquid outer core**; P-shadow clue → **solid inner core**.

PYQ 2023 (ans c, both right): P waves are recorded earlier than S; P particles vibrate along the direction of travel, S at right angles.

17 · EARTH MOVEMENTS



Endogenic Forces:

- Caused **due to the internal energy of the Earth** .
- This energy is mostly **generated by radioactivity, rotational and tidal friction and primordial heat** from the origin of the earth. This energy due to geothermal gradients and heat flow from within induces diastrophism and volcanism in the lithosphere.
- Can be of 2 types: Diastrophic and Sudden movements.
- **Diastrophism: bending, folding, warping** (bending and twisting of a large area) and **fracturing** .
  - **Orogenic processes** involving **mountain building** through severe folding and affecting long and narrow belts of the earth's crust. **Tension and Compression** . **Horizontal forces and vertical effects.**
  - **Epeirogenic processes** involving **uplift/subsidence or warping** of large parts of the earth's crust. **Vertical forces and horizontal effects.**
    - Epeirogenic or **continent-forming movements act along the radius of the earth** ; therefore, they are also called **radial movements** . Their direction may be towards (subsidence) or away (uplift) from the centre. The results of such movements may be clearly defined in the relief.
    - Upliftment examples: Colorado Plateau, Brazilian Highlands, Australian Shield, Siberian Platform.
    - Subsidence examples: Siberian Basin, Konkan region.
  - Orogeny is a mountain-building process whereas epeirogeny is a continent-building process.
- **Sudden Movements:**
  - Include **earthquakes and volcanism** involving local relatively minor movements.
  - Due to their sudden-ness, cause a **large amount of destruction** , and are included under natural disasters.
  - **Can also cause upliftment or submergence** .

- The 2004 Indian Ocean earthquake caused Indira Point to submerge. An earthquake in New Zealand (1885) caused an upliftment of up to 3 metres in some areas while some areas in Japan (1891) subsided by 6 metres after an earthquake.

- Through the processes of orogeny, epeirogeny, earthquakes and plate tectonics, there can be faulting and fracturing of the crust.

Exogenic Forces:

- A direct **result of stress induced** in earth materials **due to various forces** that come into existence **due to the sun's heat** .
- **Temperature and precipitation** are the two **important climatic elements** that control various processes by inducing stress in earth materials.
  - Weathering, Erosion, Deposition & Mass Movement.

Denudation

- **All the exogenic geomorphic processes** are covered under a general term, denudation.



⚡ **Quick revise** Earth movements & weathering — Fold ▲

	Endogenic (internal energy: radioactivity, tidal/rotational friction, primordial heat)	**Exogenic** (Sun-driven: temperature + rain)
Slow	**Diastrophism** – *Orogenic* (mountain-building, horizontal force → vertical effect, long narrow belts) and *Epeirogenic* (continent-building, vertical force → horizontal effect, **radial** up/down)	Weathering · Erosion · Deposition · Mass movement = **denudation**
Sudden	Earthquakes, volcanism	–

Mass movement = movement **only under gravity**. **Denudation gives:** soils, ore concentration (iron, manganese, aluminium, copper), landforms.

Types of Weathering:

- **Chemical: Change in chemical composition**

  - **Solution/Hydration:** Rocks formed of **water-soluble minerals.** Soluble rock-forming minerals like **nitrates, sulphates, potassium, etc.** are affected.
  - **Carbonation** : CO2 and water form **Carbonic acid** , which reacts with rocks.
  - **Oxidation:** combination of a **mineral with oxygen to form oxides** . Iron, Manganese, Sulphur rich rocks experience this.
  - **Reduction** : When **oxidized minerals** are **placed in an environment where oxygen is absent** . E.g. Red iron oxides turn greenish or bluish grey.
- **Physical:**
  - Due to gravity, temperature and moisture changes, agents of erosion, and biological activity.
  - **Unloading and Expansion:** Removal of load from above the rocks, and resultant expansion of rocks, cracking and weathering.
  - **Granular Disintegration** (**sedimentary rocks**): Rocks composed of coarse mineral grains commonly fall apart grain by grain or undergo granular disintegration.
  - **Exfoliation (diurnal temperature changes):**
    - **Onion Peeling**
    - With rise in temperature, every mineral expands and pushes against its neighbor and as temperature falls, a corresponding contraction takes place.

    - Because of **diurnal changes in the temperatures** , this internal movement among the mineral grains takes place regularly.
    - This process is **most effective in dry climates and high elevations** where diurnal temperature changes are drastic.



- The surface layers of the rocks tend to expand more than the rock at depth and this leads to the formation of stress within the rock resulting in heaving and fracturing parallel to the surface.
- Exfoliation results in smooth rounded surfaces in rocks.

- **Block separation:**
  - This type of disintegration takes place **in rocks with numerous joints** acquired by mountain-making pressures or by shrinkage due to cooling.
  - This type of disintegration in rocks can be achieved by comparatively weaker forces.
- **Shattering (angular pieces):**
  - A huge rock may undergo disintegration along weak zones to **produce highly angular pieces with sharp corners and edges** through the process of shattering.
  - This occurs when there is a **temperature change** ; when the rock is exposed to the effects of wind, rain and waves also contributes to disintegration.
- **Frost wedging:**
  - During the warm season, the water penetrates the pore spaces or fractures in rocks.
  - During the cold season, the water freezes into ice and its volume expands as a result.
  - This exerts tremendous pressure on rock walls to tear apart even where the rocks are massive.
  - Frost weathering occurs due to the growth of ice within pores and cracks of rocks during **repeated cycles of freezing and thawing (melting)** .
- **Salt weathering (crystallization):**
  - Salts in rocks expand **due to thermal action, hydration, and crystallisation** .

- Many salts like calcium, sodium, magnesium, potassium, and barium tend to expand.
- **High-temperature ranges in deserts favour** such salt expansion.
- Salt crystals in near-surface pores cause the splitting of individual grains within rocks, which eventually fall off.
- This process of falling off of individual grains may result in granular disintegration or granular foliation.

- **Biological:**
  - **Burrowing and wedging** exposes more surface for physical or chemical weathering
  - Physical weathering by **plant roots and human actions**
  - Chemical weathering by:
- **Decaying plant and animal matter** help in the production of **humic, carbonic, and other acids** which enhance decay and solubility of some elements.
- **Removal of minerals by algae and plants** .

MASS MOVEMENTS:

- Movement of material **ONLY under the impact of gravity.**



Failure Type	Type of Material	Type of Motion	Rate of Motion

Failure Type	Type of Material	Type of Motion	Rate of Motion
Rock fall	Rock fragments	Vertical or near-vertical fall (plus bouncing in many cases)	Very fast (>10s m/s)
Rock slide	A large rock body	Motion as a unit along a planar surface (translational sliding)	Typically very slow (mm/y to cm/y), but some can be faster
Rock avalanche	A large rock body that slides and then breaks into small fragments	Flow (at high speeds, the mass of rock fragments is suspended on a cushion of air)	Very fast (>10s m/s)
Creep or solifluction	Soil or other overburden; in some cases, mixed with ice	Flow (although sliding motion may also occur)	Very slow (mm/y to cm/y)
Slump	Thick deposits (m to 10s of m) of unconsolidated sediment	Motion as a unit along a curved surface (rotational sliding)	Slow (cm/y to m/y)
Mudflow	Loose sediment with a significant component of silt and clay	Flow (a mixture of sediment and water moves down a channel)	Moderate to fast (cm/s to m/s)
Debris flow	Sand, gravel, and larger fragments	Flow (similar to a mudflow, but typically faster)	Fast (m/s)

Significance of Denudation:

- **Soil formation and enrichment**
- Helps in the **enrichment and concentrations of certain valuable ores** of iron, manganese, aluminium, copper etc.
- **Landform Creation**

⚡ **Quick revise** Earth movements & weathering Fold ▲

🧂 **Weathering – chemical "S-C-O-R" = "Some Chefs Often Roast"**: **S**olution/hydration · **C**arbonation (CO_2 + water → carbonic acid) · **O**xidation (iron, manganese, sulphur rocks) · **R**eduction (red iron oxide → green/blue-grey). **Physical:** unloading · granular disintegration · **exfoliation** (diurnal temperature, "onion peeling", smooth rounded surfaces, dry/high places) · block separation (jointed rocks) · **shattering** (angular pieces) · **frost wedging** (water freezes, expands) · **salt weathering** (crystallisation, deserts). **Biological:** burrowing, roots, acids from decay.

PYQ

Q1.) Consider the following statements :

- In a seismograph, P waves are recorded earlier than S waves.
- In P waves, the individual particles vibrate to and fro in the direction of Wave propagation whereas. in S waves, the particles vibrate up and down at right angles to the direction of wave propagation.

Which of the statements given above is/are correct? (2023)

- 1 only
- 2 only
- Both 1 & 2
- Neither 1 nor 2

Ans: C

Explanation:

- P or Primary waves are the fastest seismic waves, thats why they are the first to be received. P waves are longitudinal, i.e. direction of propagation of waves and the direction of movement of particles is the same. S waves are transverse i.e. particles move perpendicular to the movement of the wave.

Q2.) Consider the following statements : (2024)

Statement-I: Rainfall is one of the reasons for weathering of roc

Statement-II: Rain water contains carbon dioxide in solution.

Statement-III: Rain water contains atmospheric oxygen.

Which one of the following is correct in respect of the above s

- Both Statement-II and Statement-III are correct and both
- Both Statement-II and Statement-III are correct, bu Statement-I
- Only one of the Statement II and III is correct and that exp
- Neither Statement-II nor Statement-III is correct

Answer : (a) Both Statement-II and Statement-III are cor Statement-I.

⚡ **Quick revise** Earth movements & weathering — Fold ▲

Epeirogenic: uplift – Colorado Plateau, Brazilian Highlands, Australian Shield, Siberian Platform; subsidence – Siberian Basin, **Konkan**. Sudden: **2004 earthquake submerged Indira Point**; NZ 1885 uplift up to 3 m; Japan 1891 subsidence 6 m.

⚡ **Quick revise** PYQ 2024 (ans a) — Fold ▲

PYQ 2024 (ans a): Rain weathers rock because rainwater carries dissolved **CO_2 and O_2** – both statements explain it.

18 · CONTINENTAL DRIFT & SEA FLOOR SPREAD THEORIES

Earlier Theories:

- **Permanency Theory**
- Works of **Antonio Pellegrini (1858)** : **identical fossil plants in North American and European coal deposits** could be explained if the two continents had formerly been connected.
- **Edward Suess (late 19th century)** : large ancient continents had been composed of several of the present-day smaller ones
- **Continental Drift Theory by F. B. Taylor** : arcuate (bow-shaped) mountain belts of Asia and Europe resulted from the creep of the continents toward the Equator .

Continental Drift Theory

- By **Alfred Wegener in 1912** .
- There existed **one big landmass** which he called **Pangaea** which was covered by **one big ocean** called **Panthalassa** .
- **Later** , a sea called **Tethys Sea** divided the Pangaea into two huge landmasses: **Laurentia (Laurasia) to the north and Gondwanaland to the south** of Tethys.
- **Assumptions:**
  - **Free Float of SiAl over SiMa** .
  - **Oceanic Crust was passive** .
  - **Forces** responsible: **Tidal force** (caused the **Westward** Movement) & **Gravity + Polar Fleeing Force** (caused the **Equator-ward Movement**).
    - **Tidal forces** played a **bigger role** .
  - **Disintegration of Pangea** in the **Triassic Period** (251-199.6 mya; Mesozoic Era).



- **Arguments supporting the theory:**
  - **Jigsaw fit of shape:**
    - **South America and Africa** seem to fit in with each other, especially, the bulge of Brazil fits into the Gulf of Guinea.
    - **Greenland** seems to fit in well with **Ellesmere and Baffin islands** .
    - The **east coast of India, Madagascar and Africa** seem to have been joined.
  - **Jigsaw fit of structure** (rocks, minerals and fossil deposits):



- **Rich placer deposits of gold** are found **on the Ghana coast** (West Africa), but the **source** (gold-bearing veins) is **in Brazil** and it is obvious that the gold deposits of Ghana are derived from the Brazil plateau when the two continents lay side by side.
- **Mesosaurus** was a small reptile **adapted to shallow brackish water** .

The **fossils** of these are **found only in South Africa and Iraver formations of Brazil** . The two localities presently are **4,800 km apart** with an ocean in between them.

- **Paleoclimatic Evidence: Tillite sedimentary rock** formed out of deposits of glaciers. The Gondwana system of sediments from India is known to have its counterparts in six different landmasses of the Southern Hemisphere. The glacial Tillite provides unambiguous evidence of paleoclimates and of the drifting of continents.
- **Paleomagnetic evidence (not given by Wegener)** :
  - The **Caledonian and Hercynian mountains of Europe and the Appalachians of the USA** seem to be one continuous series.
  - **Radiometric Dating** has confirmed the presence of similar rocks, 2 billion years old, on both the Brazilian coast as well as West African Coast.

- **Drawbacks:**
  - **No free float of SiAl over SiMa** . It is the lithosphere that moves over the Asthenosphere.
  - **Ocean Floor is not passive** : Hess's Sea Floor Spread Theory.
  - **Tidal and Gravitational Forces are not enough** to cause such massive land masses to move. Convective Currents cause this movement.
  - He talked about the **movement of landmasses** only towards the Equator, and the West direction, when the movement was **seen in all directions** . E.g. Antarctica moved away from the Equator, and Australia moved East. Moreover, India moved in the Northeast direction, initially towards the equator, and later passing it altogether.
  - Did not mention **why the Pangea suddenly broke** only precisely in the Triassic period. **Why did nothing happen before it?**

Seafloor Spread Theory:

- By **Harry Hess in 1960** .



- Detailed research during World Wars revealed that the **ocean floor** is not just a vast plain but it is **full of relief** with mountain ranges, deep trenches, etc.
- The **mid-oceanic ridges** were found to be **most active** in terms of volcanic eruptions.
- Mid-oceanic ridges prove that ocean floors are not static.
- **Rocks on either side of the crest of oceanic ridges** and having **equidistant locations** from the crest were found to have **remarkable similarities** both in terms of their constituents and their age.



- Shallow focus earthquakes at mid-oceanic ridge areas. (Deep focus EQs at the Himalayan belt)
- **Oceanic rocks** are **much younger than continental** , as former subduct.

Convectional Current Theory

- **Arthur Holmes** in the **1930s**
- According to this theory, the **intense heat generated by radioactive substances in the mantle** (100-2900 km below the earth's surface) seeks a path to escape and **gives rise to the formation of convection currents** in the mantle.
- **Rising limbs → divergence; falling limbs → convergence**

Paleomagnetism

- **Certain minerals in rocks lock in a record of the direction and intensity of the magnetic field when they form.**



- **Give details of both age and the location of formation of the rocks.**
- Basaltic rocks formed in the sea floor spread regions contain magnetic material that locks in the information.
- Continental drift could be tested by ascertaining the magnetic characteristics of ancient rocks.
- When paleomagnetists [scientists who study past magnetic fields], took a look at the **ocean floor going out away from oceanic ridges** (either side of the oceanic ridges), they found **magnetic stripes** that were flipped so that **one stripe would be normal polarity and the next reversed** .
- As the new rock is formed near the ridge, older rock, which formed millions of years ago when the magnetic field was reversed, got pushed farther away, resulting in this magnetic striping.
- When Earth's geomagnetic field undergoes a reversal, the change in polarity is recorded in the magma, which contributes to the alternating pattern of magnetic striping in the rocks on the seafloor.

PYQ

Q.1) Which of the following phenomena might have influenced the evolution of organisms ?

- Continental drift
- Glacial cycles

Select the correct answer using the code given below. (2014)

- 1 only
- 2 only

- Both 1 and 2
- Neither 1 nor 2.

Answer: (c) Both 1 and 2

9. About 200 million years ago, all the continents on Earth were actually one huge "supercontinent" surrounded by one enormous ocean. This gigantic continent, called Pangaea , slowly broke apart and spread out to form the continents we know today. All Earth's continents were once combined in one supercontinent, Pangaea.

- One of the most prominent examples of the effect of continental drift on the evolution of organisms is the unique primates found in Australia and South America.
- The final stages of evolution of Genus Homo occur in the last 3 glacial cycles.

Q.2) Which of the following are the evidences of the phenomenon of continental drift?

- The belt of ancient rocks from Brazil coast matches with those from Western Africa.
- The gold deposits of Ghana are derived from the Brazil plateau when the two continents lay side by side.
- The Gondwana system of sediments from India is known to have its counterparts in six different landmasses of the Southern Hemisphere.

Select the correct answer using the code given below:

- I and III only
- I and II only
- I, II and III
- II and III only

Answer: (c) I, II and III

- **Statement I:** Ancient rock belt from Brazil matches with Western Africa **Correct.** Geological studies show a remarkable similarity between the Precambrian rock formations on the coasts of Brazil and Western Africa.
- **Statement II:** Gold deposits of Ghana derived from Brazil plateau **Correct.** Certain mineral deposits such as gold and diamonds found in Ghana and Brazil match in age and type, indicating they were formed in the same geological settings when South America and Africa were joined.
- **Statement III:** Gondwana sediments in India match those in other Southern Hemisphere landmasses **Correct.** The Gondwana sedimentary sequences in India show close similarity with those in Africa, South America, Australia, Antarctica, Madagascar, and New Zealand.

⚡ **Quick revise** Pre-Wegener Fold ▲

Pre-Wegener: Pellegrini 1858 (same fossil plants either side of the Atlantic) · Suess (ancient big continents) · F. B. Taylor (arc mountains from creep toward Equator).

Wegener 1912: one landmass **Pangaea** + ocean **Panthalassa**; **Tethys Sea** split it into **Laurentia (Laurasia)** north and **Gondwanaland** south; forces = **tidal** (westward) + **gravity & polar-fleeing** (equator-ward); break-up in **Triassic (251–199.6 mya)**.

Evidence	Detail
Jigsaw fit	Brazil bulge ↔ Gulf of Guinea; Greenland ↔ Ellesmere/Baffin; India–Madagascar–Africa
Rocks/fossils	**Ghana gold placers** derived from Brazil plateau; **Mesosaurus** fossils only in South Africa & Brazil (4,800 km apart); 2-bn-yr-old rocks on both coasts
Climate	**Tillite**; India's **Gondwana sediments** have counterparts in **six** southern landmasses
Mountains	Caledonian/Hercynian (Europe) continue as the **Appalachians**

Drawbacks: no free float of SiAl on SiMa – the **lithosphere** moves over the asthenosphere · ocean floor isn't passive · tides/gravity too weak (**convection currents** are the engine) · only equator/west movement (but Antarctica moved away, Australia east, India NE) · no reason for Triassic break-up.

> 🧩 **Relay of ideas:** Wegener (1912, continents) → **Holmes** (1930s, **convection currents**; rising limbs diverge, sinking limbs converge) → **Hess** (1960, **sea-floor spreading**: young rocks mirror each other either side of the ridge; shallow quakes at ridges) → **Paleomagnetism** (magnetic **stripes** alternate normal/reversed) → **McKenzie & Parker 1967, Morgan 1968 = plate tectonics.**

19 · PLATE TECTONICS THEORY

- First **suggested** by **McKenzie and Parker in 1967**
- First **outlined by Morgan in 1968** .
- Both ‘convectional current theory’ and ‘seafloor spreading’ paved the way for the Theory of Plate Tectonics.
- Theory: **Lithosphere is broken into multiple plates (7 major, numerous minor)**
- It **floats on the Asthenosphere** under the **impact of Convectional Currents** .
- Lithospheric plates (sometimes called crustal plates, or tectonic plates) vary from minor plates to major plates, continental plates (Arabian plate) to oceanic plates (Pacific plate), and sometimes a combination of both continental and oceanic plates (Indo-Australian plate).
- The **oceanic plates** contain mainly the **Simatic crust** and are **relatively thinner** , while the **continental plates** contain **Sialic material** and are **relatively thicker** .



- **Major Plates:**
  - Antarctica and the surrounding oceanic plate
  - North American plate
  - South American plate
  - Pacific plate
  - India-Australia-New Zealand plate
  - Africa with the eastern Atlantic floor plate
  - Eurasia and the adjacent oceanic plate
- **Minor Plates:** There are many more minor plates other than the above-mentioned (in the image) plates. Most of these minor plates were **formed due to stress created by converging major plates** . For example: the Mediterranean Sea is divided into numerous minor plates due to the compressive force exerted by the Eurasian and African plates.

- **Forces for plate movement: the convectional currents** . Arthur Holmes gave the idea about these in 1930s.
- **Explains Earthquakes, Volcanoes, Fold and Block Mountains**
- **70% earthquakes; 90% of volcanoes (75% active) in the Circum-Pacific Ring of Fire.**
- **Circum-Pacific Ring of Fire:**
  - The **boundary of the Pacific plate** , where it is s **ubducting beneath the Indo-Australian, Eurasian, and American plates** .

- From **New Zealand to Chile.**
- **20% earthquakes** in **Mid-Continental region** .
- The Arctic Ridge has the slowest rate (less than 2.5 cm/yr), and the East Pacific Rise in the South Pacific [about 3,400 km west of Chile], has the fastest rate (more than 15 cm/yr).

Plate Boundaries:

- Major geomorphological features such as fold and block mountains, mid-oceanic ridges, trenches, volcanism, earthquakes etc. are a direct consequence of interaction between various lithospheric plates.
- **Diverging/Constructive Boundaries:**
  - The sites where the **plates move away** from each other are called **spreading sites** .
  - Such edges are sites of earth **crust formation (hence constructive)** and **volcanic landforms** are common along such edges.



  - **Earthquakes (shallow focus)** are common along divergent edges.
  - **Examples:** Mid-oceanic ridges (some of the longest mountain ranges of the world); On land, most prominent: Great African Rift Valley (divergence of African and Somali Plate; Lebanon in North to Mozambique in South.)
  - Leads to the formation of **Volcanoes, Block Mountains** and in case of Oceanic ridges, **volcanic Islands** , like the Chagos in the Indian Ocean.
- **Converging/Destructive Boundaries:**
  - **2 types:**
    - **C-C Convergence:** When it's Continent-Continent Convergence, it's called a **colliding boundary** . E.g., Indian and the Eurasian Plates
    - **O-C Convergence** (aka **Cordilleran convergence** , as leads to formation of fold mountains on the continental plate): When its Ocean-Continent Convergence, it’s called a **subducting boundary.** E.g. Pacific Ocean and South American Plate.
  - The zone of collision may undergo crumpling and folding, and **folded mountains may emerge.**
  - When one of the plates is an oceanic plate, it gets embedded in the softer asthenosphere of the continental plate and as a result, trenches are formed at the zone

of subduction.

- The subducted material gets heated up and is thrown out forming volcanic islands and dynamic equilibrium is achieved.
- Destructive: Subduction of the denser plate under the less dense one.
- Colliding: E.g. Himalayas
- Lead to the **formation of Fold mountains, and in case of O-C convergence, formation of Oceanic Trenches and Volcanoes** too.
- Why are there no volcanoes in colliding boundaries? Since no subduction, no molten material is available to burst out from the volcanoes. That is why the Himalayas do not have any volcanoes, while the Andes and the Alps do!

- **Transforms/Conservative:**
  - Formed when two **plates move past each other** .
  - In this kind of interaction, two plates grind against each other and there is **no creation or destruction of landform but only deformation** of the existing landform. [Crust is neither produced nor destroyed as the plates slide horizontally past each other].
  - No creation or destruction. Only deformation.
  - **Generally perpendicular to oceanic ridges** .
  - The San Andreas Fault along the western coast of the USA is the best example.
- **Indian Plate:**
  - Indo-Australian Plate.
  - East Margin: through Rakin Yoma Mountains (Arakan Yoma) of Myanmar towards the island arc along the Java Trench.
  - West Margin: Kirthar mountains of Pakistan
  - Northern Margin: Himalayas
  - Eurasia plate on its northern and eastern boundary; Arabian plate on its, western boundary; Somalia, Capricorn, and Australia plates to the south
  - Movement of Indian Plate:
    - India was a large island situated off the Australian coast, in a vast ocean.
    - The Tethys Sea separated it from the Asian continent till about 225 million years ago.
    - India is supposed to have started her northward journey about 200 million years ago at the time when Pangaea broke.
    - India collided with Asia about 40-50 million years ago causing rapid uplift of the Himalayas.
    - About 140 million years before the present, the subcontinent was located as south as 50◦ S. latitude. The two major plates were separated by the Tethys Sea and the Tibetan block was closer to the Asiatic landmass.
    - During the movement of the Indian plate towards the Asiatic plate, a major event that occurred was the outpouring of lava and formation of the Deccan Traps. This started somewhere around 65 million years ago and continued for a long period.
    - Note that the subcontinent was still close to the equator. From 40-50 million years ago (disputed by some experts who say it started around 60 million years ago) and thereafter, the event of formation of the Himalayas took place.

- Scientists believe that the process is still continuing and the height of the Himalayas is rising even to this date.

LIMITATIONS OF PLATE TECTONICS THEORY

Does not explain about Hotspots Cannot explain the positions/movements of the ancient plates, also cannot predict where the plate boundaries will get created in the future.

PYQ

Q.1) Consider the following:

- Electromagnetic radiation
- Geothermal energy
- Gravitational force
- Plate Movements
- Rotation of the earth
- Revolution of the earth **Which of the above are responsible for bringing dynamic changes on the surface of the earth? (2013)**
- 1, 2, 3 and 4 only
- 1, 3, 5 and 6 only
- 2, 4, 5 and 6 only
- 1, 2, 3, 4, 5 and 6

Ans. D All the factors are responsible for bringing dynamic changes on the surface of the earth.

- Wind, water, and ice erode and shape the land. Volcanic activity and earthquakes alter the landscape in a dramatic and often violent manner. And on a much longer timescale, the movement of earth's plates slowly reconfigures oceans and continents. Each one of these processes plays a role in the Arctic and Antarctica.
- Electro-magnetic radiation causes temperature change which induces metamorphism of rocks. Gravity besides being a directional force activating all down slope movements of matter also causes stresses on the earth's materials. Plate movement results into continent building. The energy emanating from within the earth is the main force behind endo-genic geomorphic processes. This energy is mostly generated by radioactivity, rotational and tidal friction and primordial heat from the origin of the earth. This energy due to geothermal gradients and heat flow from within induces diastrophism and volcanism in the lithosphere. Due to variations in geothermal gradients and heat flow from within, crustal thickness and strength, the action of endo-genic forces are not uniform and hence the tectonically controlled original crustal surface is uneven.

⚡ **Quick revise** Plate tectonics — Fold ▲

7 major plates: Antarctica+ocean · North American · South American · Pacific · India-Australia-NZ · Africa+E. Atlantic floor · Eurasia+ocean. Minor plates arise from stress where majors converge (e.g. the Mediterranean). Plates can be continental (Arabian), oceanic (Pacific) or both (Indo-Australian).

Ring of Fire (Pacific plate boundary, NZ → Chile): **70% of earthquakes, 90% of volcanoes (75% of active)**; 20% of quakes in mid-continental belt. **Slowest** ridge: **Arctic** (<2.5 cm/yr); **fastest**: **East Pacific Rise** (>15 cm/yr).

💔 **Divorce · Collide · Slide**

Boundary	Plates…	Creates	Example
Divergent (constructive)	move **apart**	new crust, mid-ocean ridges, rift valleys, volcanic islands, shallow quakes	Mid-oceanic ridges, **Great African Rift** (Lebanon → Mozambique), Chagos
Convergent C–C (colliding)	continent meets continent	fold mountains, **no volcanoes** (no subduction)	**India–Eurasia → Himalayas**
Convergent O–C (subducting)	dense ocean plate dives	trench, volcanoes, fold mountains (Cordilleran)	Pacific–South American → **Andes**
Transform (conservative)	slide **past**	only deformation; usually ⊥ ridges	**San Andreas Fault**

Himalayas have no volcanoes; the Andes and Alps do.

Indian plate story: island off Australia → **Tethys** separated it from Asia till ~**225 mya** → started north **~200 mya** (Pangaea breaks) → **~140 mya** at **50°S** → **Deccan Traps ~65 mya** → collision **40–50 mya** (some say 60) → Himalayas still rising. Margins: **E** Arakan Yoma → Java Trench · **W** Kirthar · **N** Himalaya.

Limits of plate tectonics: doesn't explain **hotspots**; can't place ancient plates or predict future boundaries.

⚡ **Quick revise** Plate tectonics — Fold ▲

PYQs: *2014 (ans c)* – continental drift and glacial cycles both influenced evolution. *Drift evidences (no year, ans c)* – Brazil–W. Africa rocks, Ghana gold, Gondwana sediments: **all three** right. *2013 (ans d)* – electromagnetic radiation, geothermal energy, gravity, plate movement, rotation, revolution: **all** cause dynamic surface change.

20 · VOLCANISM

Volcanism: The **process** by which **molten rock material from the asthenosphere finds a way to enter the lithosphere** , where it may or may not reach the surface of the Earth. A **volcano** is a **vent in the earth's crust from which molten rock material (magma), explosive bursts of gases and volcanic ashes erupt** , or A mountain or hill having a crater or vent through which lava, rock fragments, hot vapour, and gas are or have been erupted from the earth's crust.



Types of Magmas:

- **Under Earth: Magma; After eruption: Lava**
- **Acidic Lava:**
- 1. **Light-coloured, of low density** , and have a **high percentage of silica** .
- 2. **Highly Viscous** ; flow slowly; do not spread much.
- 3. Get solidifies in the vent, causing **explosive eruptions** and release of Pyroclastic material.
- **Basic Lava:**
- **1. Dark, high density, less percentage of silica**
- 2. **Highly fluid** ; flow readily; spread to large areas
- 3. **Lava plateaus** ; gently sloping volcanic mountains
- **Fumaroles: openings** in the earth's surface that **emit steam and volcanic gases** , such as sulphur dioxide and carbon dioxide.
- 1. A **fumarole rich in sulphur gases** is called a **solfatara** ; a **fumarole rich in carbon dioxide** is called a **mofette** .
- Volcanoes often release large amount of gases like Carbon, Sulphur and Nitrogen Oxides, Water Vapour, hydrogen sulphide, trace amounts of Chlorine.

Geysers and Hot Springs

Geyser	Hot water spring
Steam or water at high pressure, along its path, gets accumulated in small reservoirs, fissures and fractures. Once the pressure exceeds the threshold limit, the steam bursts out to the surface disrupting the water at the mouth. Hence the name geyser. Usually a carter like structure is created at the mouth. Silicate deposits at mouth gives them their distinct colours Found in very few regions. **Iceland** is famous for its geysers.	Steam or water at high pressure smoothly flows to the top through the vent and condense at the surface giving rise to a spring. Usually a carter like structure is created at the mouth of the spring. Some springs are very colorful because of the presence of **cyanobacteria** of different colors. Found all across the world

- **Almost all the world's geysers** are confined to three major areas: **Iceland, New Zealand and Yellowstone Park of U.S.A.**



Distribution of Volcanoes in the World:

- **Only 10 to 20 per cent** of all volcanic activity is **above sea**
- **Pacific Ring of fire: 75% active volcanoes**
- **Mid-Atlantic ridge** : Iceland (above sea surface presence)
- **Great African Rift** : Mt. Kilimanjaro; Mt. Kenya;
- **Other Volcanoes**
  - **Africa** : Mt. Cameroon (only active volcano in West Africa)
  - **West Indies** : Mt Pelee
  - **Mediterranean** : Spain to Caucasus; Alps: Vesuvius, Stromboli; Asia Minor: Ararat, Elburz
- **No volcanoes in Australia**
- **Only one active lava volcano in South Asia: Barren Island, India.**

- **Mud Volcano in South Asia: Chandragup, Pakistan.** Does not emit any lava or pyroclast, so not a true volcano.
- **Volcanoes in India:**
  - Narcondam island and Barren Island are volcanic islands of India, of them, the latter is active. Erupted last in 2022.
  - Baratang: Only mud volcano of India.
  - **No volcanoes in the Himalayas,** as it is a continent-continent collision boundary.
  - There is a mud volcano “Chandragup” located in Balochistan, Pakistan.

⚡ **Quick revise**

Where: only 10–20% of volcanic activity is above sea · Fold ▲

Where: only 10–20% of volcanic activity is above sea · **Ring of Fire 75% of active** · Mid-Atlantic ridge (**Iceland**) · Great African Rift (**Kilimanjaro, Kenya**) · **Cameroon** (only active volcano in West Africa) · **Pelée** (West Indies) · Mediterranean: **Vesuvius, Stromboli**, Ararat, Elburz · **none in Australia** · South Asia's only active lava volcano: **Barren Island** (India; last erupted **2022**); **Narcondam** extinct; **Baratang** = India's only mud volcano; Chandragup (Pakistan, mud, "not a true volcano"). **No volcanoes in the Himalaya.**

Volcanic Landforms:

- **Extrusive: above earth’s surface**
  - Conical vents and fissure vents (basaltic plateaus)
  - Mid-Oceanic Ridges
  - Pyroclastic Debris,
  - Caldera & Crater Lakes (E.g. Crater Lake, Oregon);

- **Intrusive: Inside Earth’s surface**



- **Appear on Earth's surface after erosion** has removed the upper layers.
- **Massive: Batholiths or Plutons** . Large rock masses formed due to the cooling down and the solidification of hot magma inside the earth. E.g. Mt. Wicklow of Ireland
- **Smaller:**
  - **Horizontal: Sill** (**Horizontal;** E.g. Great Whin Sill, England), **Laccolith** (**dome-shaped** ; E.g. Henry Mountains, Utah, USA), **Lopolith** (**saucer-shaped** ; E.g. Transvaal, South Africa), **Phacolith** (**wavy** ; E.g. Cordon Hill, USA)
  - **Vertical: Dyke** E.g. Alaskan Dyke, USA.

⚡ **Quick revise** Landforms Fold ▲

Landforms

Extrusive (surface)	Intrusive (inside; show after erosion)
Conical & fissure vents, basalt plateaus · mid-ocean ridges · pyroclastic debris · **caldera** & crater lakes (Crater Lake, Oregon)	**Batholith/pluton** (huge; Mt Wicklow) · **Sill** (flat; Great Whin Sill) · **Laccolith** (dome; Henry Mts) · **Lopolith** (saucer; Transvaal) · **Phacolith** (wavy; Cordon Hill) · **Dyke** (vertical; Alaska)

🪟 **Intrusives: Sill** = window-**sill**, lies flat · **Dyke** = wall, stands upright · **Laccolith** = **L**oaf (dome) · **Lopolith** = **Lota** (bowl/saucer) · **Phacolith** = wavy. 🍛 **Eruption styles, calm → crazy:** *"I Have Some Vada-Pav, Please"* = **I**celandic (fissure, basalt) → **H**awaiian (central vent, most lava) → **S**trombolian (short bursts, pasty lava) → **V**ulcanian (dark cauliflower clouds) → **P**elean (most dangerous, "glowing clouds") → **P**linian (ash up to 50,000 ft, deadly pyroclastic flows).

Types of Volcanoes:

- **On the basis of the frequency of eruption:**
  - **Active** : erupt fairly frequently. E.g. Mt. St. Helena, USA; Mt. Merapi, Indonesia.



- **Dormant** : No eruption recently but can erupt. E.g. Mt.

Vesuvius was dormant before it erupted in 79 AD.

- **Extinct** : No eruptions in the recorded history. E.g. Narcondam.

- **On the basis of structure:**
  - **Cinder cones** : Small, steep-sided, conical volcanic landforms built primarily from pyroclastic fragments like cinders. These volcanoes form through explosive eruptions of gas-rich basaltic magma, creating a distinctive and relatively simple structure. Cinder cones often have relatively short lifespans compared to other types of volcanoes, as their eruptions tend to be singular events rather than prolonged periods of volcanic activity.
  - **Composite (Stratovolcanoes):** A special type of Composite volcano is a caldera volcano.
  - **Shield**
- **On the basis of Eruption**
  - **Icelandic** : fissure type; basaltic lava
  - **Hawaiian** : similar to Icelandic. Difference: Hawaiian eruptions have the greatest quantity of lava pouring out of the main vent at the volcano's summit, not alongside fissures.
  - **Strombolian** : short-lived explosive eruptions releasing pasty lava into the air along with bursts of steam and gas.
  - **Vulcanian** : more violent than Strombolian. Along with pasty lava, creates high dark clouds of steam, ash, and gas.



(Cauliflower cloud)

- **Pelean:** Most dangerous and explosive. Release "Glowing clouds" of gas and ash.
- **Plinian** : Most explosive. Very high ash clouds that rise upwards to 50,000 feet (almost 10 miles) high. Very deadly pyroclastic flows.
- Vulcanian, Pelean and Plinian volcanoes release huge amount of Pyroclastic material that acts as Hygroscopic Nucleii, helping in creation of raindrops. The force of the volcano, pushes the air up, causing updraft. A combination of these processes, causes often thunderstorms and rainfall after the volcanoes take place. The rainwater gets mixed with the Pyroclastic mud, and create mudflows called LAHARS.

⚡ **Quick revise** By frequency Fold ▲

By frequency: *Active* (St Helens, Merapi) · *Dormant* (Vesuvius before AD 79) · *Extinct* (Narcondam). **By shape:** cinder cone · composite (strato) · shield. Vulcanian/Pelean/Plinian ash = **hygroscopic nuclei** → rain + ash → **lahars** (mudflows).

Mantle Plumes & Hotspots:

- Region within the Earth's mantle from which heat rises through the process of convection.
- This rising arm = Mantle Plume.
- This does not occur at plate boundaries. It is an **intra-plate event** .



- **Cause Hotspot Volcanism**
  - A volcano above a hotspot **does not erupt forever** . Attached to the tectonic plate below, the volcano moves and is eventually cut off from the hot spot.
  - Without any source of heat, the volcano becomes extinct and cools.
  - According to the latest studies, which are yet to be accepted by geologists, **hotspots may be the first stage in the formation of diverging plate boundaries.**



Azores Island: They are created both due to hotspot activity as well as the action of the Mid-Atlantic ridge. **Create island chains. E.g. Hawaii**



- **Reunion Hotspot:**
  - Believed to be active for over **65 million years** .
  - Reunion Island is over reunion hotspot.
  - When the Indian plate moved over the Reunion hotspot, Deccan traps were created.



⚡ **Quick revise** Mantle plumes & hotspots Fold ▲

Mantle plumes & hotspots: rising heat arm, **intra-plate** (not at boundaries); plate moves over it → **island chains** (Hawaii); volcano cut off from heat goes extinct. **Azores** = hotspot + Mid-Atlantic ridge. **Réunion hotspot:** active **>65 million yrs**; when the Indian plate passed over it → **Deccan Traps**.

PYQ

Q.1) Consider the following : (2024)

- Pyroclastic debris
- Ash and dust
- Nitrogen compounds
- Sulphur compounds

How many of the above are products of volcanic eruptions?

- Only one
- Only two
- Only three
- All four

Answer : (d) All four

⚡ **Quick revise** Magma (underground) Fold ▲

- **Magma** (underground) → **lava** (surface). **Acidic lava:** light, high silica, **viscous**, solidifies in vent → **explosive**, pyroclasts. **Basic lava:** dark, dense, low silica, **runny**, spreads → **lava plateaus**, gentle cones.
- **Fumarole** vents steam/gas: sulphur-rich = **solfatara**; CO_2-rich = **mofette**. **Geysers** are confined to **Iceland, New Zealand, Yellowstone**.
- Volcanic gases: carbon, sulphur, nitrogen oxides, water vapour, H_2S, trace chlorine. **PYQ 2024 (ans d):** pyroclastic debris, ash and dust, nitrogen compounds, sulphur compounds – **all four** are volcanic products.

21 · OROGENY

MOUNTAIN BUILDING

- Process known as **Orogeny** .
- Types:
  - **On basis of Location: Continental, Coastal, Oceanic (Ridge or Hotspots)**
  - **On Basis of formation: Residual, Fold, Fault, Volcanic**

⚡ **Quick revise**

Types: by location – continental, coastal, oceanic; by formation

Fold ▲

Types: by location – continental, coastal, oceanic; by formation – **fold, fault (block), volcanic, residual**.

FOLDS AND FAULTS











FOLD MOUNTAINS

- Fold mountains are formed when sedimentary rock strata are subjected to **compressive forces** .
- They are the **loftiest mountains.**
- The **Young Fold Mountains are found near the active converging boundaries** . E.g. Himalayas (India-Eurasia convergence); Alps & Atlas (Eurasia, Africa); Andes (South America & Pacific), etc. This is the reason why there are seismic and volcanic activities seen here.
- There are old fold mountains too which were formed millions of years ago at the active plate boundaries of that time. E.g. Appalachians (USA), Aravallis (India)

BLOCK MOUNTAINS

- Block mountains are **created when large areas or blocks of earth are broken and displaced vertically.**
- Formed due to **dip-slip faults (vertical displacement)** .
- Block mountains are **also called fault block mountains** since they are formed due to faulting as a result of tensile and compressive forces.
- The **uplifted blocks are termed as horsts** and the **lowered blocks are called graben** .
- The Great African Rift Valley (valley floor is graben and Mitumba & Muchinga mountains are horsts), The Rhine Valley and the Vosges mountain in Europe are examples.
- There are **two basic types** .
  - **Tilted block mountains** have **one steep side contrasted by a gentle slope** on the other side.
  - **Lifted block mountains** have a **flat top and extremely steep slopes**

VOLCANIC MOUNTAINS

- Volcanic mountains are formed **due to volcanic activity** .
- Materials including molten lava, volcanic bombs, cinders, ashes, dust and liquid mud fall around the vent in successive layers, building up a characteristic volcanic cone.
- Volcanic mountains are often called **mountains of accumulation** .
- They are common in the Circum-Pacific belt and include such volcanic peaks as Mt. Fuji (Japan) Mt. Mayon (Philippines), Mt. Merapi (Sumatra), Mt. Etna (Italy) etc.

RESIDUAL MOUNTAINS

- The **mountains left after erosional activities** have eroded the majority portion of the erstwhile fold/fault/volcanic mountains.
- Basically, they **evolve from denudation.**
- E.g. Monadnock mts. In the USA.

⚡ **Quick revise** Mountains, isostasy, rocks — Fold ▲

Mountain	Formed by	Notes / examples
Fold	Compression of sedimentary strata; **loftiest**; young near converging boundaries	Himalaya, Alps, Atlas, Andes (quakes + volcanoes); old: Appalachians, **Aravallis**
Block (fault)	Dip-slip faults, tension/compression; **horst** = raised, **graben** = lowered	Great African Rift (valley floor graben; Mitumba & Muchinga horsts), Rhine valley–Vosges
Volcanic	Accumulated lava/ash cones ("mountains of accumulation")	Fuji, Mayon, Merapi, Etna
Residual	Left by denudation of older mountains	Monadnocks (USA)

ISOSTASY

- **Equilibrium between the Earth's lithosphere and asthenosphere.**
- Balance between buoyant force and gravitational force.
- When weight is added or removed, such as through erosion or mountain building, the crust responds by subsiding or uplifting to achieve isostatic equilibrium.
- The Isostatic re-adjustments **can be due** to both **Tectonic factors (subsidence & Upliftment)** , as well as **Climatic Factors (addition or melting of snow)** .
- Can cause **earthquakes** . Scientists have predicted that **they can also be responsible for the reactivation of volcanoes in Antarctica** , as the ice melts, the lithosphere rebounds, cracks get created, and magma might pour out.

⚡ **Quick revise** Isostasy — Fold ▲

Isostasy: equilibrium between lithosphere and asthenosphere (buoyancy vs gravity); crust **rises/sinks** when load changes (erosion, mountain-building, ice melt); can cause earthquakes and may reactivate Antarctic volcanoes.

IMPORTANT MOUNTAIN RANGES

S. No	Mountain	Location
1.	Mid-Atlantic Ridge	Atlantic Ocean
2.	Andes	South America
3.	Western Cordillera	North America
4.	Great Dividing Range	Australia
5.	Trans-Antarctic Mountains	Antarctica
6.	Ural Mountains	Eurasia
7.	Atlas Mountains	Africa
8.	Appalachian Mountains	North America
9.	Himalayas	Asia
10.	Altai Mountains	Asia (Mongolia)





⚡ **Quick revise** Mountains, isostasy, rocks Fold ▲

📏 **Longest ranges (km):** Mid-Atlantic Ridge **16,000** · Andes **7,000** · W. Cordillera **4,830** · Great Dividing Range & Trans-

> Antarctic **3,500** · Urals & Atlas **2,500** · Appalachians **2,414** · Himalayas **2,400** · Altai **2,000**. *Say it as "16, 7, 48, 35-35, 25-25, 24-24, 20".*

22 · ROCKS & ROCK CYCLE

ROCKS

Igneous Rocks:

Plutonic rocks	Volcanic rocks
Intrusive rocks	Extrusive rocks
Granite	Basalt
Slow cooling allows big-sized crystals (large grains)	Rapid cooling prevents crystallization, as a result such rocks are fine-grained
Less dense and are lighter in colour than basic rocks	Denser and Darker in colour

- **Formed out of magma and lava** and are known as **primary rocks** .
- Having their **origin under conditions of high temperatures** , the igneous rocks are **not fossiliferous.**
- **Granite, gabbro, pegmatite, basalt** , etc. are some of the examples of igneous rocks.
- Based on place and time taken in cooling of the molten matter, igneous rocks can be divided into:
  - **Plutonic Rocks or intrusive rocks**
  - **Lava or Volcanic Rocks or Extrusive rocks**
- If molten material is **cooled slowly at great depths (plutonic/intrusive), mineral grains may be very large.**
- **Sudden cooling (at the surface)** results in **small and smooth grains. (extrusive igneous)**
- Acid igneous rocks, such as granite, are less dense and are lighter in colour than basic rocks.
- **Economic Significance:**
  - Since magma is the **chief source of metal ores** , many of them are associated with igneous rocks. The minerals of great economic value found in igneous rocks are magnetic iron, nickel, copper, lead, zinc, chromite, manganese, gold, diamond and platinum. Amygdales are almond-shaped bubbles formed in basalt due to escape of gases and are filled with minerals like Copper.

- Many igneous rocks like granite are **used as building material** as they come in beautiful shades.
- Source of **fertile soils** like Black Soil.

Sedimentary Rocks:

- These rocks consist of a **number of layers or strata.**

- These rocks **have fossils** of plants and animals.
- These rocks are **generally porous** and allow water to percolate through them.
- Contain **petroleum, natural gas, placer deposits.**
- Formed as a **result of denudation** (weathering and erosion).
- These **deposits through compaction turn into rocks** . This process is called **lithification.**
- **Cover 75 percent of the earth's crust** but **volumetrically occupy only 5 per cent.**
- They are **layered or stratified** of varying thickness. Example: sandstone, shale etc.
- **Till or Tillite: Ice deposited** sedimentary rocks.
- **Loess: Wind deposited** sediments.
- Depending upon the **mode of formation** ,Sedimentary rocks are **classified into**
  - **Mechanically formed** : Sandstone, conglomerate, shale, loess etc.
  - **Organically formed** : Geyserite, chalk, limestone, coal etc.
  - **Chemically formed** : Chert, limestone, halite, potash etc.
- **Economic Significance:**
  - Sedimentary rocks are not as rich in minerals of economic value as the igneous rocks.
  - But **some important minerals** such as hematite iron ore, phosphates, building stones, **coals, petroleum** and material used in cement industry like **limestone** are found. The decay of tiny marine organisms yields petroleum.
  - Petroleum occurs in suitable structures only.
  - Important minerals like bauxite, manganese, tin are derived from other rocks but are found in gravels and sands carried by water. (PLACER DEPOSITS)
  - Sedimentary rocks also yield **some of the richest soils.**

Metamorphic Rocks

- The word metamorphic **means 'change of form'** . Form under the action of pressure, volume and temperature (PVT) changes.
- Metamorphism occurs when rocks are forced down to lower levels by tectonic processes or when molten magma rising through the crust comes in contact with the crustal rocks.
- Metamorphism is a **process by which already consolidated rocks undergo recrystallization and reorganization of materials takes place.**
- In the process of metamorphism in **some rocks grains or minerals get arranged in layers or lines** . Such an arrangement is called **foliation or lineation** .
- Sometimes minerals or materials of different groups are arranged into **alternating thin to thick layers** . Such a structure is called **banding.**
- **Gneissoid, slate, schist, marble, quartzite** etc. are some examples of metamorphic rocks.
- **May or may not have strata and fossils.**



- On the basis of the agency of metamorphism, metamorphic rocks can be of two types:
  - **Thermal Metamorphism:** The change of form or re-crystallisation of minerals of sedimentary and igneous rocks under the **influence of high temperatures** is known as thermal metamorphism. (hot magma, hot gases, vapours and liquids, geothermal heat, etc.)
  - **Dynamic Metamorphism** : This refers to the formation of metamorphic rocks under the **stress of pressure** . Sometimes high pressure is accompanied by high temperatures.

⚡ **Quick revise** Rocks — Fold ▲

	Igneous (primary)	Sedimentary	Metamorphic
Origin	Magma/lava cooling	Denudation → deposits → **lithification**	**PVT** change of existing rock (recrystallise)
Features	No fossils; **intrusive/plutonic** (slow cooling, big grains) vs **extrusive** (fast, small grains)	Layered, **fossils**, porous; **75% of crust area, 5% of volume**	Foliation, lineation, banding; may/may not have strata/fossils

	Igneous (primary)	Sedimentary	Metamorphic
Examples	Granite, gabbro, pegmatite, basalt	Sandstone, shale, limestone, coal, chalk, chert, halite; **tillite** (ice), **loess** (wind)	Gneiss, slate, schist, marble, quartzite
Value	Metal ores (iron, nickel, copper, lead, zinc, chromite, manganese, gold, diamond, platinum); building stone; black soil	**Petroleum, gas, coal**, phosphates, hematite iron ore, limestone; **placers** (bauxite, manganese, tin); rich soils	**Thermal** (heat) and **dynamic** (pressure) types

Sedimentary by mode: mechanical (sandstone, conglomerate, shale, loess) · organic (chalk, limestone, coal, geyserite) · chemical (chert, limestone, halite, potash). *Amygdales* = almond bubbles in basalt filled with minerals (copper).

Rock cycle: igneous → (erosion) sedimentary → (heat/pressure) metamorphic → (subduction, melting) magma → igneous; any rock can go to any other.

Rock Cycle:

- Rock cycle is a **continuous process** through which **old rocks are transformed into new ones** .



- Igneous rocks are **primary rocks** and other rocks form from these rocks.
- **Igneous rocks can be changed into sedimentary or metamorphic rocks.**
- The fragments derived out of igneous and metamorphic rocks form into sedimentary rocks.
- **Sedimentary and igneous rocks themselves can turn into metamorphic rocks**
- The **crustal rocks** (igneous, metamorphic and sedimentary) may be carried down into the mantle (interior of the earth) **through subduction process** and the same melt down

and turn into molten magma, the original source for **igneous rocks.**

SOME IMPORTANT MINERALS

Mineral	Chemical Composition	Hardness/ Softness	Colour	Places Earth's Where
Feldspar	KAlSi3O8 or NaAlSi3O8	6-7	White, pink, grey, or green	Abun igneo metar and sedim rocks
Quartz	SiO2	7	Colourless, white, grey, purple, pink, yellow, or black	Abun igneo metar and sedim rocks
Bauxite	Al2O3•2H2O	1-3	Red, orange, or yellow	Tropi subtr regio
Cinnabar	HgS	2-3	Reddish-brown or black	Mercu depos world
Dolomite	CaMg(CO3)2	3.5-4	White, gray, or pink	Abun sedim rocks
Gypsum	CaSO4•2H2O	2	White, gray, or yellow	Depo basin and regio
Hematite	Fe2O3	5.5-6	Reddish-brown or black	Abun sedim and metar rocks
Magnetite	Fe3O4	5.5-6	Black	Igneo metar rocks

- The numbers in the "Hardness/Softness" column of the table refer to the Mohs Hardness Scale, a qualitative scale for measuring the scratch resistance of minerals. The scale ranges from 1 to 10, with 1 being the softest and 10 being the hardest.

⚡ **Quick revise** Minerals (hardness, Mohs 1–10) Fold ▲

Minerals (hardness, Mohs 1–10): Feldspar 6–7 (glass, ceramics) · **Quartz 7** (jewellery, electronics) · Bauxite 1–3 (aluminium) · Cinnabar 2–3 (mercury) · Dolomite 3.5–4 · Hematite 5.5–6 & Magnetite 5.5–6 (iron).

PYQ

Q.1) Describe the characteristics and types of primary rocks. **(2022, 10m)**

⚡ **Quick revise** Rocks Fold ▲

Mains 2022 (10m) – "Describe the characteristics and types of primary rocks": **Intro** – igneous rocks form from magma/lava, hence "primary". **Body** – non-fossiliferous; plutonic vs volcanic by cooling place/time; acid (granite, light) vs basic; examples; economic value (ores, building stone, black soil). **Close** – all other rocks are derived from them via the rock cycle.

23 · CYCLES OF EROSION

EROSION

- Removal, transportation and deposition of Earth Material. TRANSPORTATION is what differentiates it from weathering.
- **Agents of Erosion: Flowing water; Underground water; Wind; Glaciers; Coastal Waves**
- Cycles of Erosion:
  - Fluvial landforms and Cycle of Erosion
  - Glacial landforms and Cycle of Erosion
  - Marine landforms and Cycle of Erosion
  - Arid landforms and Cycle of Erosion
  - Karst landforms and Cycle of Erosion

⚡ **Quick revise** Erosion = removal + transport + deposition Fold ▲

Erosion = removal + transport + deposition (transport separates it from weathering). Five cycles: fluvial · glacial · marine · arid · karst.

Depositional: Alluvial fans/cones · **natural levees** (breach = flood) · **delta** (arcuate fan; **bird's-foot** weak tides; **cuspate** strong waves) · braided channel · **estuary** (drowned valley; ports/fisheries).

Fluvial Erosion:

- **Terms associated** with Fluvial Erosion:
  - How does water erode?
    - **Hydraulic Action** : the **force of running water** wearing down rocks.
    - **Corrosion** : **chemical action** that leads to weathering.
    - **Attrition** : **river load particles striking, colliding** against each other and breaking down in the process.
    - **Corrasion or abrasion** : solid **river load striking against rocks** and wearing them down.
  - Classification of erosion based on the location:
    - **Downcutting (vertical erosion)** : the erosion of the base of a stream (downcutting leads to **valley deepening**).
    - **Lateral erosion** : the erosion of the walls of a stream (leads to **valley widening**).
    - **Headward erosion** : **erosion at the origin** of a stream channel, which causes the origin to move back away from the direction of the stream flow, and so causes the stream channel to lengthen. It is responsible for **river Piracy** .



- **Evolution of Fluvial Erosion:**
  - **Splash** : A drop of water falls on the surface and disturbs the sediments, creating a tiny crater due to its kinetic energy.
  - **Sheet:** A sheet of water flows over the surface, and carries away the loose sediments of the topsoil.
  - **Rills, Gullies & Valleys:** After the sheet erosion, the areas with weaker rocks, or with rock joints, get more eroded than the rest. This creates small, narrow channels called Rills. Rills grow to form Gullies, which grow to form Valleys.

- **Stages of Erosion:**
  - **Youth stage:** Young rivers close to their source tend to be **fast-flowing, high-energy** environments with **rapid headward erosion** , despite the hardness of the rock over which they may flow. Steep-sided "V-shaped' valleys, waterfalls, and rapids are characteristic features.
  - **Mature stage:** Mature rivers are **lower-energy systems** . Erosion takes place on the outside of bends, creating looping meanders in the soft alluvium of the river plain. Deposition occurs on the inside of the bends and the riverbed.
  - **Old Stage:** At a river's mouth (C), sediment is **deposited** as the velocity of the river slows. As the river becomes shallower more deposition occurs, forming islands and braiding the main channel into multiple, narrower channels. As the sediment is laid down, the actual mouth of the river moves away from the source into the sea or lake, forming a delta.
- **Valley Formation:**
- The extended depression on the ground through which a stream flows throughout its course is called a river valley. At different stages of the erosional cycle, the valley acquires different profiles.
- **At a young stage** , the valley is **deep and narrow** with steep wall-like sides and a convex slope. The erosional action here is characterized by a predominantly vertical downcutting nature.



The profile of the valley here is typically ‘V’ shaped.

- A **deep and steeper valley is also referred to as a gorge** and may result in downcutting erosion because of the recession of a waterfall. A gorge is a valley which is much narrower compared to its depth. Antecedent rivers like the Indus and Brahmaputra form gorges at Nanga Parbat and Namche Barwa respectively.
- It is characterized by cliffs or steep walls, and it is usually formed by more intense erosional forces, such as the rapid erosion of a river through resistant rock.
- A **gorge** is typically considered a **narrower and steeper landform** , whereas a **canyon** is broader with more open valley walls. However, there is no universally strict scientific

boundary between the two, and regional usage often overlaps.

- The Grand Canyon of the Colorado River in Arizona (USA) runs for 483 km and has a depth of 2.88 km.
- A tributary valley lies above the main valley and is separated from it by a steep slope down which the stream may flow as a waterfall or a series of rapids.
- **As the cycle attains maturity, the lateral erosion becomes prominent, and the valley floor flattens out** . The valley profile now becomes typically 'U' shaped with a broad base and a concave slope.



- **Major Erosional Fluvial Landforms:**
  - **Waterfall:** A waterfall is simply the **fall of an enormous volume of water from a great height.** They are mostly seen in the youth stage of the river. Relative resistance of rocks, relative difference in topographic reliefs, fall in the sea level and related rejuvenation, earth movements etc. are responsible for the formation of waterfalls. For example, Jog or Gersoppa Falls on Sharavati has a fall of 260 meters.



  - **Rapids:** Formed due to a sporadic presence of soft and hard rocks along the river valley. The hard rocks don't get eroded as much as soft rocks, causing rivers to jump around, creating rapids. Called Cataracts in larger dimensions.
  - **Potholes: The kettle-like small depressions in the rocky beds of the river valleys** are called potholes which are usually cylindrical in shape.

- **Potholing or pothole-drilling** is the mechanism through which the grinding tools (fragments of rocks, e.g. boulders and angular rock fragments) when caught in the water eddies or swirling water start dancing circularly and grind and drill the rock beds of the valleys like a drilling machine.
- They thus form small holes which are gradually enlarged by the repetition of the said mechanism. The potholes go on increasing in both diameter and depth.





- **River Piracy:** River piracy, also known as **stream capture,** occurs when a river or **stream is diverted from its original course,** often due to natural geological processes such as tectonic shifts, glacial activity, or erosion. Examples:
  - **Saraswati River** : Ancient river piracy is believed to have caused the once-mighty Saraswati River to lose its course and dry up. The river's waters were

likely captured by the Yamuna and Sutlej rivers due to tectonic shifts and changes in the Himalayan landscape.

- **Shyok River and Nubra River** : The Shyok River in Ladakh captured the Nubra River, altering the drainage pattern in the region. This is an example of river piracy due to tectonic activity in the Himalayas.

- **Terraces: Stepped benches along the river course in a floodplain** are called terraces. Terraces represent the level of former valley floors and remnants of former (older) flood plains. Can be paired or unpaired. Created due to rejuvenation.
- **Meanders** : A meander is defined as a pronounced **curve or loop in the course of a river channel.**
  - The **outer bend** of the loop in a meander is characterized by intensive erosion and vertical cliffs and is called the **cliff-slope side** . This side has a concave slope.



  - The **inner side** of the loop is characterized by deposition, a gentle convex slope, and is called the **slip-off side** . The depositions re called the **point-bars** .
- **Ox-Bow Lake:**
  - **Both Erosional and Depositional** landform.
  - Sometimes, because of intensive erosion action, the outer curve of a meander gets accentuated to such an extent that the inner ends of the loop come close enough to get disconnected from the main channel and exist as independent water bodies.
  - These water bodies are **temporary** and get **converted into swamps in due course of time.**
  - In the Indo-Gangetic plains, southwards shifting of Ganga has left many ox-bow lakes to the north of the present course of the Ganga
- **Peneplains: End Product of the erosion cycle.**
  - This refers to an **undulating featureless plain** punctuated with low-lying residual hills of resistant rocks.
  - Gently undulating (wave-like), almost featureless plain that, in principle, would be produced by fluvial erosion that would, in the course of geologic time, reduce the land almost to base level (sea level), leaving so little gradient that essentially no more erosion could occur.

- **Monadnocks:** An **isolated hill** of bedrock standing conspicuously above the general level of the surrounding area. Monadnocks are left as **erosional remnants** because of their **more resistant rock** composition; commonly they consist of quartzite or less jointed massive volcanic rocks.

- **Major Depositional Fluvial Landforms:**
  - **Alluvial fans and cones:** When a stream leaves the mountains and comes down to the plains, its velocity decreases due to a lower gradient. As a result, it sheds a lot of material, which it had been carrying from the mountains, **at the foothills.** This **unconsoliated deposited material** acquires a conical shape and appears as a series of continuous fans. These are called alluvial fans or cones.





- **Natural Levees:** These are **narrow ridges of low height on both sides of a river** , formed due to deposition action of the stream, appearing as natural embankments. These act as a **natural protection against floods** but a breach in a levee causes sudden floods in adjoining areas.
- **Delta** : A delta is a **tract of alluvium at the mouth of a river** where it deposits more material than can be carried away. The river gets divided into distributaries which may further divide and rejoin to form a network of channels. A delta is **formed by** a combination of two processes:
  - The **load-bearing capacity of a river is reduced** as a result of the check to its speed (due to reduced gradient) as it enters a sea or lake
  - Clay particles carried in suspension in the river coagulate in the presence of salt water and are deposited. The **finest particles are carried farthest** to accumulate as bottom-set beds.
  - Depending on the conditions under which they are formed, deltas can be of many types.
  - **Arcuate (fan-shaped); Bird's Foot (elongated** , as in regions with weak tides and waves); **Cuspate (Triangle** ; strong wave action; almost nil distributaries)
- **Braided Channel:** A river channel that splits into multiple smaller, intertwining channels separated by sediment bars or islands, typically formed due to excessive sediment load and variable flow rates
- **Estuaries:** Sometimes the mouth of the river appears to be submerged. This may be due to a drowned valley because of a rise in sea level. Here fresh water and the saline water get mixed.

- When the river starts ‘filling its mouth’ with sediments, mud bars, marshes and plains seem to be developing in it.
- These are **ideal sites for fisheries, ports and industries** because estuaries provide access to deep water, especially if protected from currents and tides. Example: Hudson estuary.

- **Rejuvenation: Change in the stage of erosion** of the river due to tectonic (submergence/emergence) or climatic (sea level rise/ fall) activities.

⚡ **Quick revise** Cycles of erosion — Fold ▲

🏞️ **River life: Young – V, Mature – meander, Old – delta. Youth:** fast, **V-valley, waterfalls, rapids**, headward erosion. **Mature:** lateral erosion, **meanders**, U-profile. **Old:** deposition, braiding, **delta**. *Rejuvenation* (tectonic or sea-level change) restarts the cycle → **terraces**. **Erosion types:** hydraulic action · corrosion (chemical) · attrition (load vs load) · **corrasion/abrasion** (load vs rock). **Downcutting** deepens, **lateral** widens, **headward** lengthens (→ **river piracy**: Saraswati; Shyok captured Nubra).

Erosional fluvial forms: Gorge (narrow; Indus at Nanga Parbat, Brahmaputra at Namche Barwa – antecedent rivers) vs **canyon** (broader; Grand Canyon 483 km long, 2.88 km deep) · **Waterfall** (youth; Jog/Gersoppa on Sharavati **260 m**) · **Rapids/cataracts** · **Potholes** (swirling boulders drill) · **Meanders** (outer = **cliff-slope**, erosion, concave; inner = **slip-off**, deposition, **point bars**) · **Ox-bow lake** (cut-off meander; Ganga shifted south leaving ox-bows north) · **Peneplain** (end product) with **monadnocks**.

Drainage Patterns:

- The **typical shape of a river course as it completes its erosional cycle** is referred to as the drainage pattern of a stream. A drainage pattern reflects the structure of basal rocks, resistance and strength, cracks or joints and tectonic irregularity, if any.



- **Dendritic or Pinnate:**

This is an **irregular tree branch-shaped** pattern. Examples: Indus, Godavari, Mahanadi, Cauvery, Krishna

- **Trellis:** In this type of pattern, the **short tributaries meet the mainstream at right angles** , and differential erosion through soft rocks paves the way for tributaries. Examples: The Seine and its tributaries in the Paris basin (France).
- **Rectangular: The mainstream bends at right angles and the tributaries join at right angles** creating rectangular patterns. E.g. Colorado River in the USA, and Mahanadi River in India.
- **Angular** : The **tributaries join the mainstream at acute angles.** This pattern is common in Himalayan foothill regions.
- **Parallel:** The **streams/tributaries seem to be running parallel to each other** in a uniformly sloping region. Example: rivers of lesser Himalayas.
- **Radial:** The **tributaries from a summit follow the slope downwards and drain down in all directions.** Examples: The Central French Plateau, Mt. Kilimanjaro, Amarkantak and Saurashtra region are good examples of radial drainage in India.
- **Annular: When the upland has an outer soft stratum, the radial streams develop subsequent tributaries** which try to follow a **circular drainage around the summit.** Example: Black Hill streams of South Dakota
- **Centripetal: In a low-lying basin the streams converge from all sides.** Examples: streams of Ladakh, Tibet, and the Bagmati and its tributaries in Nepal.

⚡ **Quick revise** Cycles of erosion Fold ▲

🌳 **Drainage patterns: Dendritic** = tree (Indus, Godavari, Mahanadi, Cauvery, Krishna) · **Trellis** = garden fence, tributaries at right angles (Seine, Paris basin) · **Rectangular** = right-angle bends (Colorado; Mahanadi) · **Angular** = acute angles (Himalayan foothills) · **Parallel** = lesser Himalayas · **Radial** = wheel spokes outward (Amarkantak, Saurashtra, Kilimanjaro) · **Annular** = ring (Black Hills, S. Dakota) · **Centripetal** = all into one basin (Ladakh, Tibet, Bagmati).

> **Handout alert:** Mahanadi is listed under both **dendritic** and **rectangular**. Take it as dendritic unless the question says rectangular.

KARST CYCLE:

Terminologies associated with Groundwater:

- Groundwater: Water that percolates into soil and rock pores.
- Porous Rocks: Rocks which have a lot of pore spaces between their grains. E.g. sandstone
- Permeable/Pervious Rocks: Rocks that allow water to pass through them easily.
  - Most (but not all) porous rocks are permeable. E.g. Clay: highly porous, but the pores are not big enough to allow easy movement of water.
- **Aquifer** : A geological formation that can store and transmit water.
- **Water Table** : The upper surface of the zone of saturation in an aquifer, where the pressure is equal to atmospheric pressure.
- **Confined Aquifer** : An aquifer overlain by an impermeable layer, causing the water to be under pressure.
- **Unconfined Aquifer** : An aquifer without an overlying impermeable layer, where the water table is open to the atmosphere.



- **Artesian Well** : A well drilled into a confined aquifer where the water pressure allows it to rise without pumping.
- **Groundwater Springs** : Natural discharge points where groundwater flows to the surface, typically occurring where the water table intersects the ground.
- **Perched Water Table** : A localized zone of saturation above the main water table, formed by an impermeable layer.



Karst Cycle

- **Named after a province of Yugoslavia (now Slovenia) on the Adriatic Sea coast** where such formations are most noticeable.
- In locations with **thick deposits of Limestone dolomite or gypsum** .
- These rocks should be **dense, highly jointed and thickly bedded**
- **Erosional Features:**

  - **Sinkholes/Swallow holes:** Sinkholes are **funnel-shaped depressions** having an average depth of three to nine meters. These holes are developed by enlargement of the cracks found in such rocks, as a result of **continuous solvent action of the rainwater** . The surface streams that sink disappear underground through swallow

holes. **Larger sinkholes can be called Dolines, Uvalas and Polje (faulting+erosion)** .

- **Lappies:** usually formed when the **limestones are vertically bedded** . Lappies are uneven grooves and ridges that form when the majority of the limestone's surface is removed by the solution process.

  - Deep grooves separate the etched, fluted, and pitted rock pinnacles that make up the weathered limestone surface seen in the karst region.
- **Karst Windows: When several adjoining sinkholes collapse** , they form an open, broad area called a karst window.
- **Karst Arch/Bridge:** When a part of the cavern collapses the portion which keeps standing forms an arch.
- **Karst Cave:** This is an underground cave formed by water action by various methods in a limestone or chalk area.
- **Blind Valley:**
  - A stephead valley, or blind valley is a deep, narrow, f **lat-bottomed valley with an abrupt ending.**
  - Karst valley abruptly terminated by the passage underground of the watercourse which has hitherto resisted the karst processes and remained at the surface.





- **Depositional Features:** Found **in the Karst Caves** and known as **Speleothems** .
  - **Stalactites, Stalagmites and Karst Pillars:** The water containing limestone in solution, seeps through the roof in the form of a continuous chain of drops.
    - A portion of the solution hangs on the roof and on evaporation of water, a small deposit of limestone is left behind contributing to the formation of a **stalactite, growing downwards from the roof.**
    - The remaining portion of the drop falls to the floor.
    - This also evaporates, leaving behind a small deposit of limestone aiding the formation of a **stalagmite, thicker and flatter, rising upwards from the floor.**



    - Sometimes, **stalactite and stalagmite join** to form a complete pillar known as the **column or Karst pillar** .
  - **Flow stones:** Flowstone is one of the most common speleothems. It has been described as **“melted cake icing” and “frozen waterfalls.”** Flowstone is usually composed of calcite or other carbonate minerals, and deposition is in layers or bands. Where composed of calcite, individual flowstone layers **may be very colourful: yellows, reds, and oranges.**
  - **Cave Pearls:** Cave pearls are **concentrically banded concretions** that form in shallow cave pools. Cave pearls vary in shape; they can be spherical, cylindrical, irregular, cubical, or even hexagonal. They can range in size from smaller than a sand grain up to eight inches (20 cm) in diameter.

- **Banded draperies:** When water drops flow down a sloped ceiling before dripping to the floor, calcite can build up in a line. These lines gradually form “draperies” **or “cave bacon.”**
  - This type of speleothem is found in almost every cave in the world and is universally popular because of its close resemblance to its namesake.

  - Iron oxide or organic solutions form the baconlike stripes.

⚡ Quick revise

Karst (limestone, dolomite, gypsum; dense, jointed, thickly bedded)

Fold ▲

Named after a province of Yugoslavia (now Slovenia). **Groundwater words:** porous (pores), permeable (water passes; clay is porous but *not* permeable), aquifer, water table, confined/unconfined, **artesian well** (rises unpumped), spring, perched table.

Erosional: sinkholes/swallow holes (3–9 m; dolines, uvalas, polje) · **lapies** (grooves and ridges) · karst window · karst arch/bridge · caves · blind valley. **Depositional (speleothems): stalactite** hangs from roof, **stalagmite** rises from floor, **column** when joined; flowstone, cave pearls, draperies.

> 🕳️ **Stalactite = Ceiling (holds *tight* to the roof). Stalagmite = Ground (*might* reach the roof).**

PYQ 2025 (ans c): chalk = porous **and** permeable; clay = porous but **not** permeable. So Statement III ("clay not porous at all") is wrong; Statement II is right and explains I.

MARINE CYCLE:

- **Sea waves, aided by winds, currents, tides and storms** carry on the erosional and depositional processes.
- The **erosive work** of the sea depends upon size and strength of waves, slope, height of the shore between low and high tides, shape of the coast, composition of rocks, depth of water, human activity, etc.
- The **wave pressure compresses the air trapped inside rock fissures, joints, faults, etc** . forcing it to expand and rupture the rocks along weak points.
- **Waves also use rock debris as instruments of erosion.** These rock fragments carried by waves themselves get worn down by striking against the coast or one another.



- The **solvent or chemical action of waves** is another mode of erosion, but it is pronounced only in the case of soluble rocks like limestone and chalk.
- **Erosional Features:**
- 1. **Chasms: These are narrow, deep indentations** (a deep recess or notch on the edge or surface of something) carved **due to headward erosion** (downcutting) through vertical planes of weakness in the rocks by wave action.



- 1. With time, further headward erosion is hindered by lateral erosion of the chasm mouth, which itself keeps **widening till a bay is formed.**
- 2. **Wave Cut Platform:** Formed due to the **backward retreat of the sea cliffs.**

- 1. The **bottom of the cliff** suffers the **maximum intensive erosion** by waves and, as a result, a **notch appears** at this position.
- 3. **Sea Cave** : Differential erosion by sea waves through a rock with varying resistance across its structure produces arched caves in rocks called sea caves. **With further erosion, they get converted into sea arches.**



Fig. 79 Gloup

- **4. Stack:** When the **sea caves get extremely eroded** , **all that is left is some rock pillars** standing in the offshore location called stacks. When these stacks further get eroded (often below the Low Tide limit), we get **Stumps.**
- 5. **Blow Holes/ Spouting holes** : The burst of water through a small hole on a sea cave due to the compression of air in the cave by strong waves. They **make a peculiar noise.** With further erosion these holes get bigger, the walls collapse, and we get **Geos/gloup** .
- **Depositional Features:**
- 1. **Beach:** This is the **temporary** covering of rock debris on or along a wave-cut platform
- 2. **Bar:** Currents and tidal currents **deposit rock debris and sand along the coast at a distance from the shoreline** . The resultant landforms which remain submerged are called bars. The enclosed water body so created is called a **lagoon** . They are of various types:
- 1. **Barrier:** It is the overwater counterpart of a bar.



- 2. **Spit and Hook:** A spit is a projected deposition joined at one end to the headland, with the other end free in the sea. The mode of formation is like a bar or barrier. A shorter spit with one end curved towards the land is called a hook.
- 3. **Tombolo:** Sometimes, islands are connected by a bar called tombolo.
- **3. Marine Dunes** : Deposited in coastal locations, but due to the action of Wind and not Waves.

⚡ **Quick revise** Marine (waves) Fold ▲

🌊 **Cliff → Cave → Arch → Stack → Stump** ("**C**ats **A**lways **S**it **S**till"). Wave-cut **platform** forms as the cliff retreats (notch at base). **Chasm** → bay. **Blowhole** (air compressed in cave) → geo/gloup. **Depositional:** beach · **bar** (submerged; encloses a **lagoon**) · barrier · **spit** (one end tied to headland) → **hook** (curved end) · **tombolo** (bar joining an island) · marine dunes (wind).

TYPES OF COASTLINES:

Coastline of emergence:

- These are formed either by an **uplift of the land or by the lowering of the sea level** .
- **Bars, spits, lagoons, salt marshes, beaches, sea cliffs, and arches** are the typical features.

Coastline of submergence:

- A submerged coast is produced **either by subsidence of land or by a rise in sea level. Ria, Fjord, Dalmatian, and drowned lowlands** are its typical features.
- **Ria Coast:** When a region is **dissected by streams** into a system of valleys and divides, submergence produces a **highly irregular shoreline** called Ria coastline. The coast of **south-west Ireland** is a typical example of the ria coastline.

- **Fjord:** Some coastal regions have been **heavily eroded by glacial action** and the **valley glacial troughs (U-shaped valley)** have been excavated below sea level.
- 1. After the glaciers have disappeared, a fjord coastline emerges.

- 2. These coasts have **long and narrow inlets with very steep sides.**

- 3. The fjord **coasts of Norway** are a typical example.
- **Dalmatian:** The Dalmatian coasts result from the **submergence of mountain ridges with alternating crests and troughs** which run parallel to the seacoast. The Dalmatian coast of **Yugoslavia** is a typical example.
- **Drowned Lowland:** A drowned lowland coast is **low and free from indentations** , as it is formed by the submergence of a low-lying area. The **Baltic coast of eastern Germany** is an example of this type of coastline.

Neutral Coastline:

- These are coastlines **formed as a result of new materials being built out into the water.** The word ‘neutral’ implies that there need be **no relative change between the level of the sea and the coastal region of the continent.**
- Whatever is taken away by the wave, gets deposited by the agents of erosion, such that there is no net change.
- Neutral coastlines include the alluvial fan-shaped coastline, delta coastline, volcano coastline, and the coral reef coastline.

Compound Coastline:

- Such coastlines show the forms of two of the previous classes combined, for example, **submergence followed by emergence or vice versa.**
- The coastlines of **Norway and Sweden** are examples of compound coastlines.

Fault Coastline:

- Such coastlines are unusual features and result from the **submergence of a downthrown block along a fault** , such that the uplifted block has its steep side (or the Faultline) standing against the sea forming a fault coastline. E.g. **Konkan Coast of India** .

⚡ **Quick revise** Marine (waves) Fold ▲

Coast type	Cause	Examples
Emergence	land up / sea down	bars, spits, lagoons, cliffs, arches

Coast type	Cause	Examples
Submergence	land sinks / sea up	**Ria** (SW Ireland) · **Fjord** (Norway; drowned U-valley) · **Dalmatian** (ridges ‖ coast, Yugoslavia) · **Drowned lowland** (Baltic, E. Germany)
Neutral	new material built out; no net change	fan, delta, volcano, coral-reef coasts
Compound	submergence + emergence	Norway & Sweden
Fault	down-thrown block, steep fault face to sea	**Konkan coast**

GLACIAL CYCLE:

- A glacier is a **moving mass of ice at speeds averaging a few meters a day** .



- **Types** of Glaciers: **continental glaciers, ice caps, piedmont glaciers, and valley glaciers.**
- A glacier is **charged with rock debris** which are used for erosional activity by moving ice.
- **Erosional Features:**
- 1. **Cirque/Corrie/Cwm: Hollow basin cut into a mountain ridge.** It has a **steep-sided slope** on three sides, an open end on one side, and a slightly concave bottom/base. When the ice melts, tarn lakes may develop at the base of the cirques.

2. **Horn/Pyramidal Peak:** Ridge that acquires a ‘horn’ shape when the glacial activity cuts it from more than two sides

3. **Arete:** It is the **knife-edged boundary between two cirques.**

4. Bergschrund: Crevasse that forms between a moving glacier and the stationary ice or rock of a mountain, typically at the upper end of a glacier, where it separates the flowing ice from the stagnant ice above.



5. **Glacial Trough:**

Original stream-cut valley, further modified by glacial action. It is a **'U' Shaped Valley** . It is at the mature stage of valley formation.

Since glacial mass is heavy and slow-moving, erosional activity is uniform – horizontally as well as vertically. A steep-sided and flat-bottomed valley results, which has a 'U' shaped profile. These U-shaped valleys, when filled with water, form Fjords.

6. **Hanging Valley:** Formed when **smaller tributaries are unable to cut as deeply as bigger ones and remain 'hanging' at higher levels than the main valley** as discordant tributaries. A valley carved out by a small tributary glacier that joins with a valley carved out by a much larger glacier.

7. **Roche Moutonnee:** Roche Moutonnée, a term derived from French **meaning "sheep rock,** " refers to a **glacially eroded bedrock formation** characterized by a **smooth, gently sloping side facing the direction from which the glacier advanced, and a rough, steep side where the glacier plucked** and quarried the rock as it moved. These features provide insights into the direction and nature of past glacial movements.

8. Craig & Tail: Landform consisting of a resistant rock mass (the Craig) with a gently sloping ridge of softer material (the Tail) extending downstream, formed by glacial erosion.





- **Depositional Features:**
- 9. Includes **both depositional and erosional features** .
- 10. **Outwash Plain:** When the **glacier reaches its lowest point and melts, it leaves behind a stratified deposition material** , consisting of rock debris, clay, sand, gravel, etc. This layered surface is called a **till plain or an outwash plain** .
- 11. **Esker: Winding ridge of un-assorted depositions** of rock, gravel, clay, etc. running along a glacier in a till plain. The eskers resemble the features of an embankment and are often used for making roads. They are **analogous to levees.**
- 12. **Kame Terraces: Broken ridges or un-assorted depositions** looking like a hump in a till plain.
- 13. **Drumlin: Inverted boat-shaped deposition in a till plain** . They **help in determining the direction of movement of the glacier** , as the **glacier side is always high and steep** , while the lee side is smooth and tapers gently in the direction of ice movement.



14. **Kettle Holes** : Formed when the **deposited material in a till plain gets depressed locally and forms a basin.**

15. **Moraine:** General term applied to **rock fragments, gravel, sand, etc. carried by a glacier.** Depending on its position, the moraine can be ground moraine, lateral moraine, and end moraine.

⚡ **Quick revise** Glacial — Fold ▲

Glacier = moving ice (a few metres/day); types: continental, ice cap, piedmont, valley.

🏔️ **Cirque = armchair** (steep on 3 sides, open on 1; tarn lake) · **Arête** = knife-edge between two cirques · **Horn** = pyramid peak (cut from >2 sides) · **U-valley** (glacial trough; flooded = **fjord**) ·

> **Hanging valley** (tributary left high) · **Roche moutonnée** = "sheep rock" (smooth side faces ice advance, rough steep side plucked) · **Bergschrund** = crevasse at the head. **Deposits: outwash/till plain** · **esker** (snake ridge, like a levee) · **kame** (hump) · **drumlin** (inverted boat; steep side faces the glacier) · **kettle** (hole) · **moraine** (ground, lateral, end).

ARID CYCLE:

- **Types of Deserts:**

1. Hamada (Rocky) : Consist of large rocks that have been smoothened by exogenic forces.

2. Reg (Stony)/Serir : Covered with extensive sheets of gravel and pebbles. More accessible than sandy deserts (easier mobility).

3. Erg (Sandy)/Koum: Typical pop-culture desert with vast expanses of sand.

4. Cold Deserts: Either at high altitudes or high latitudes.

- **Both Water (flash floods) and wind** are agents of erosion here.
- **Water-related landforms:**
- 1. **Bolsons:** The intermontane basins in dry regions are generally known as bolsons.
- **2. Erosional:**
- 1. A **gully** is a landform created by running water. Gullies resemble **large ditches or small valleys** but are meters to tens of meters in depth and width.
- 2. **Ravine** : A ravine is a landform narrower than a canyon and is often the product of stream-cutting erosion. Ravines are typically classified as **larger in scale than gullies, although smaller than valleys.**
- 3. **Badland Topography:** In arid regions, occasional **rainstorms produce numerous rills and channels which extensively erode weak sedimentary formations.** Lack of any vegetation causes this extreme level of erosion. As the top soil gets eroded, any chance of undertaking agriculture as a source of livelihood also erodes away. This **lack of utility gives these regions their name** . Ravines and gullies are developed by linear fluvial erosion leading to the formation of badland topography. Example: Chambal Ravines
- 4. **Playas:** Small streams flow into bolsons, and this water accumulates in the depressions. These **temporary lakes** are called playas. **After the evaporation of water, salt-covered playas are called Salinas.**
- 5. **Pediments:** Pediment is an erosional landform. A true pediment is a **rock-cut surface at the foot of mountain.**
- **3. Depositional:**
- 1. **Bajada:** Bajadas are moderately sloping depositional plains **in the desert regions.**
- Wind related processes:
- 1. **Deflation** ⇒ removing, lifting, and carrying away dry, unsorted dust particles by winds.

- 2. **Abrasion** ⇒ When wind loaded with sand grains erodes the rock by grinding against its walls.
- 3. **Attrition** ⇒ Attrition refers to wear and tear of the sand particles while they are being transported.
- **Wind Related landforms:**

- **1. Erosional:**
- 1. **Blowouts/ Deflation Hollows:** hollows formed by the removal of particles by wind. They are generally small but may be up to several kilometres in diameter.

2. **Mushroom rocks:** a cap or cap-like structure atop a narrower stem.

3. **Zeugen: A table-shaped** area of rock found in arid and semi-arid areas formed when more resistant rock is reduced at a slower rate than softer rocks around it. Rocks are in horizontal layers.

4. **Yardang: Rocks are in vertical layers** . The hard rocks stay resistant while the soft rocks get eroded.

5. **Wind bridges and windows:** Powerful wind continuously abrades stone lattices, creating holes. Sometimes the holes are gradually widened to reach the other end of the rocks to create the effect of a window—thus forming a wind window.

1. Window bridges, are formed when the holes are further widened to form an arch-like feature

6. **Inselbergs:** A inselberg is an isolated hill, knob, ridge, outcrop, or small mountain that rises abruptly from a gently sloping or virtually level surrounding plain. **Analogous to Monadnocks** in the arid regions.

7. **Pediplain: equivalent of Peneplain** in the arid regions.



2. Depositional:

1. **Ripple Marks:** These are depositional features on a small scale formed by saltation (the transport of hard particles over an uneven surface in a turbulent flow of air or water).

2. **Dunes (in-situ):** Dunes are **heaps or mounds of sand found in deserts.** Generally, their heights vary from a few meters to 20 meters but in some cases, dunes are several hundred meters high and 5 to 6 km long.

1. Formed **both in deserts and coastal locations.**

2. **Longitudinal dunes:** Formed **parallel to the wind movement** . These dunes are commonly found at the heart of trade-wind deserts like the Sahara, Australian, Libyan, South African and Thar deserts.

3. **Transverse dunes** : Dunes deposited **perpendicular (transverse) to the prevailing wind direction.**

4. **Barchans: Crescent-shaped dunes.** The **windward side is convex** whereas the **leeward side is concave** and steep.

5. **Parabolic dunes:** They are **U-shaped** and are **much longer and narrower than barchans** . The **windward side is concave.**

6. **Star dunes:** Have a **high central peak, radically extending three or more arms** . Formed when the **wind blows from multiple directions.**

3. **Loess (ex-situ):** Deposition of very fine particles, sourced from the deserts, in far-off locations, by the action of winds. E.g. Loess Plateau near Gobi desert in China.

⚡ **Quick revise** Arid — Fold ▲

Deserts: Hamada = rocky · **Reg/Serir** = stony gravel · **Erg** = sandy (*H-R-E: Hard → Rubble → Ergs of sand*) · cold deserts (high altitude/latitude).

Water forms: bolson (basin) · gully · ravine · **badlands** (Chambal ravines) · **playa** (temporary lake) → **salina** (salt-covered) · **pediment** (rock-cut slope at mountain foot) · **bajada** (depositional plain).

Wind: deflation · abrasion · attrition. **Landforms:** blowout · **mushroom rock** · **Zeugen** (table-shaped; *horizontal* layers) vs **Yardang** (*vertical* layers) · wind window/bridge · **inselberg** (= monadnock of deserts) · **pediplain** (= peneplain of deserts).

> 🏜️ **Zeugen = Zzz, lies flat (horizontal). Yardang = Y stands up (vertical). Dunes: Barchan** crescent (windward convex, leeward steep concave) · **Longitudinal** ∥ wind (Sahara, Thar) · **Transverse** ⊥ wind · **Parabolic** U-shaped, windward concave · **Star** (wind from many sides). **Loess** = fine desert dust carried far (Loess Plateau, China).

LAKES

- **Body of water of considerable size, localized in a basin, surrounded by land apart from a river or other outlet that feeds/drains it.**
- Larger and deeper than ponds.
- **Most** lakes **have at least one natural outflow** to maintain its average level by allowing the drainage of excess water.
- **Endorheic Lakes: do not have a natural outflow** and **lose water solely by evaporation or underground seepage or both** .
- **Formation of lakes:**
  - Lakes formed by **earth movements: Tectonic lakes** (Titicaca, Caspian Sea), **Rift valley lakes** (Tanganyika, Malawi, Dead Sea)
  - **By glaciation: Tarns, Rock hollow lakes** . Many such lakes in Scandinavia, Canada, Siberia.
  - **By Volcanic Activities: Crater and Caldera lakes** (Lonar lake, Krakatoa lake)
  - **By Erosion: Karst, Playa, Wind deflated** lakes (Great Basin of Utah)
  - **By Deposition: Ox-Bow lakes, Lagoons** (Chilika), **Glacial lakes** (moraine-dammed)
  - **Man-made lakes**

⚡ **Quick revise** Lakes by origin — Fold ▲

Lakes by origin: **tectonic** (Titicaca, Caspian) · **rift valley** (Tanganyika, Malawi, Dead Sea) · **glacial** (tarns; Scandinavia, Canada) · **volcanic** crater/caldera (**Lonar**, Krakatoa) · **erosional** (karst, playa, wind-deflated – Great Basin, Utah) · **depositional** (ox-bow, lagoon e.g. **Chilika**, moraine-dammed) · man-made. **Endorheic** = no outflow (lose water by evaporation/seepage).

PLATEAUS

- **Flat-topped tableland.**

- Plateaus occur in every continent and take up a **third of the Earth's land**
- **Sometimes,** a plateau is so **eroded** that it is **broken up into smaller raised sections called outliers.**
- Many outlier plateaus are composed of very old, dense rock formations. Iron ore and coal often are found in plateau outliers.
- Plateaus are **rich in mineral deposits.** As a result, many of the mining areas in the world are in the plateau areas.
- **Tectonic plateaus** are formed from processes that create mountain ranges – volcanism (Deccan Plateau), crustal shortening (thrusting of one block of crust over another, and folding occurs. Example: Tibet (an intermontane plateau)), and thermal expansion (Ethiopian Highlands).
- Plateaus **can also be erosional.** Erosion of mountains can lead to their creation. E.g. Alleghany plateau in Appalachians.

⚡ **Quick revise** Plateaus Fold ▲

Plateaus: flat tablelands, **a third of land**; mineral-rich (mining; iron ore and coal in outliers). **Tectonic:** volcanic (**Deccan**), crustal shortening (**Tibet**, intermontane), thermal expansion (**Ethiopian**); **erosional:** Allegheny (Appalachians).

24 · PYQ

Q.1) Consider the following statements (2025): Statement I: In the context of effect of water on rocks, chalk is known as a very permeable rock, whereas clay is known as quite an impermeable or least permeable rock. **Statement II:** Chalk is porous and hence can absorb water. **Statement III:** Clay is not at all porous.

Which one of the following is correct in respect of the above statements?

(a) Both Statement II and Statement III are correct and both of them explain Statement I

(b) Both Statement II and Statement III are correct but only one of them explains Statement I

(c) Only one of the Statements II and III is correct and that explains Statement I

(d) Neither Statement II nor Statement III is correct

Answer: (c) Only one of the Statements II and III is correct and that explains Statement I

- Chalk is both porous and permeable → absorbs and transmits water easily.
- Clay is porous but not permeable → holds water but does not allow it to flow through.
- USGS Water Science Glossary: Clay has very high porosity, but low permeability.

Recall sheet

Q	A
CMB found by whom, when?	**Penzias & Wilson, 1965**; ~380,000 yrs after Big Bang
White dwarf vs neutron star vs black hole (mass)	< 1.44 · 1.44–3 · > 3 (Chandrasekhar)
Dark matter / dark energy share	27% / 68%
Moons: Jupiter, Saturn, Uranus, Neptune	101, 274, 28, 16
Kuiper belt distance; Oort start	30–50 AU; 2,000–5,000 AU
Flares classes; CME time	A B C M X; 15–18 h
Earth tilt; sidereal day	23.5°; 23 h 56 m 4.1 s
Time zones champion	France 12
Earth volume: crust / mantle / core	0.5–1% / 83% / 16%
Boundaries top → bottom	Conrad, Moho, Repetti, Gutenberg, Lehmann
P shadow / S shadow	103–143° / beyond 103°
Richter vs Mercalli	Magnitude 0–10 vs intensity 1–12
Drift / spreading / plates	Wegener 1912 / Hess 1960 / McKenzie-Parker 1967
Which plate boundary has no volcanoes?	Continent–continent (Himalaya)
Only active lava volcano in S. Asia	Barren Island (India)
Indian plate collision with Asia	40–50 mya (Deccan Traps ~65 mya)
Fastest / slowest ridge	East Pacific Rise / Arctic
Karst depositional forms	Stalactite, stalagmite, column
Fjord vs ria	Glacial U-valley drowned vs river valley drowned
Barchan windward side	Convex

Traps: "Himalayas have volcanoes" ✘ · "P waves can't pass liquids" ✘ (that's S) · "orogeny = continent-building" ✘ (that's epeirogeny) · "solar flares cause tsunamis/forest fires" ✘ · "Blue Moon = eclipse" ✘ · "Sun overhead on equator on 21 June" ✘ (equinoxes).

Connects to: Himalaya & Deccan story → *Indian Geography*; Tilt and 21 June → *Climatology*; Ridges, trenches, coral reefs → *Oceanography*; Karst/gorge examples (Indus, Brahmaputra) → *Indian drainage*.

GEOGRAPHY · GS FOUNDATION · SUBJECT 2

Climatology

Handout + mind maps: no class slides | Handout pages 48

VERY HIGH · 32 Prelims, 1 Mains

The whole Climatology handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 21 mind-map pages are in Chapter 8.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: VERY HIGH for Prelims. 32 Prelims PYQs** are printed in this part, the most of any chapter: 2011×5 · 2012×3 · 2013 · 2014 · 2015×3 · 2018 · 2019×2 · 2020 · 2021×2 · 2022 · 2023×3 · 2024×6 · 2025 · plus 2 with no year printed. **Mains quoted:** *"Troposphere determines weather processes – how?"* (2022, 15m).
- **Hot clusters:** heat budget & lapse rate (2012, 2023, 2024) · isotherm bending (2024, 2025) · westerlies/Coriolis/jet streams (2011, 2015, 2020, 2024) · clouds & dew (2019, 2022) · cyclone genesis (2015) · climate-type identification (2011–2024: rainforest, savannah, equatorial, Marine West Coast).
- **Prepare:** *Prelims* – cause-and-effect statement pairs (the Statement I/II format keeps coming back: 2023, 2024, 2025). *Mains* – troposphere, jet streams & climate change, tropical vs temperate cyclones.
- *No teacher ink in this PDF; priority = PYQ count in the handout.*

1 · ATMOSPHERE

- **Layer of air surrounding the Earth.**
- **Importance:**
  - It **protects the earth from the harmful radiation** from the sun. The **Ozone Layer absorbs harmful UV radiations.**
  - The **Greenhouse Effect by certains gases in the atmosphere helps keep the Earth warm,** to allow the survival of life.
  - The atmosphere also takes care of extra-terrestrial objects like **meteors that get burnt up while passing through it,** due to air friction.

- The gasses of the present atmosphere are a **product of volcanic eruptions** , **hot springs, chemical breakdowns of solid matter, and redistribution from the biosphere.**
- **Composition of the atmosphere:**
  - The atmosphere is a mixture of many gases. In addition, it contains huge numbers of **solid and liquid particles, collectively called aerosols.**



  - **Nitrogen and oxygen make up nearly 99% of the clean, dry air** . The remaining gases are mostly inert and constitute about 1% of the atmosphere.
  - **Oxygen constitutes 21%** of the total volume of the atmosphere. **All living organisms inhale oxygen** . Besides, oxygen can combine with other elements to form important compounds, such as oxides. Also, combustion is not possible without oxygen.
  - **Nitrogen accounts for 78%** of total atmospheric volume. It is a relatively inert gas and is an important constituent of all organic compounds. The main function of nitrogen is to **control combustion by diluting oxygen** .
  - **Carbon Dioxide: constitutes only about 0.038% (now 0.04%)** of the dry air and is a product of combustion. **Green plants, through photosynthesis, absorb carbon dioxide** from the atmosphere and use it to manufacture food and keep other biophysical processes going. It is an important greenhouse gas, and helps maintain the Heat Budget of Earth.
  - **Carbon dioxide and water vapour are found only up to 90 km from the surface of the earth.**
  - **Water vapour depends on temperature** and therefore it decreases from the equator poleward in response to changes in temperature (and evaporation).
    - **0 to 5 % by volume; More than 90% found up to the height of 5 km.**
  - **Argon: The third important gas which constitutes only about 0.93%.**
  - **Ozone (O3)**
    - **Formed at higher altitudes and transported downwards** .
    - The ozone layer is between **20 km and 25 km** altitude where the greatest concentrations of ozone are found.
    - **Protects from harmful UV radiations.**

- **Other gases: Neon, helium, hydrogen, xenon, krypton, methane.**
- **Particulate Matter: Consists of sand particles (from weathered rocks, and volcanic ash), pollen grains, small organisms, soot, and ocean salts** ; the upper layers of the atmosphere may even have **fragments of meteors** .
  - **Help in the absorbing, reflecting, and scattering of the solar radiation** which adds the varied charming colours of red and orange at sunrise and sunset.
  - **Salt particles become hygroscopic nuclei and thus help in the formation of water drops** , clouds, and various forms of condensation and precipitation.
  - Recent research: Microplastics in the atmosphere can act as nuclei for cloud formation, leading to ice crystal formation at temperatures 5 to 10 degrees Celsius warmer than usual.

Layers Of The Atmosphere:

- **Based on Composition:**



  - **Homosphere** : **Up to 80 km** . The **gases are distributed uniformly** , therefore, there is a homogenous composition.
  - **Heterosphere: Beyond 80 km** . Here the **gases** are **layered according to their molecular masses.**
- **Based on Change in the temperature:**
  - **Troposphere: 8-18 km (13 km avg.)**
    - **It is thicker at the equator (18 km)** due to the strong convection experienced here. It's thinner at the poles (8 km).
    - **Temperature change:** Temperature gradually decreases with height, approx. 1°C for every 165 meters (normal lapse rate).
    - **Why does the temperature reduce?** The prime source of heat for the troposphere is terrestrial radiation. As the **distance from the surface of the earth increases** , the temperature reduces.



- **All weather phenomena like cloud formation, movement of air masses and jet streams, etc** , take place in this layer.
- It is **separated from the stratosphere by Tropopause.**

- **Stratosphere: Upto 50 kilometres** above the earth's surface, beyond the tropopause.
  - **Ozone layer** : Present **in the lower part of the stratosphere** . The absorption of ultraviolet (UV) energy from the sun by this ozone causes the temperature to rise with height.
  - **Temperatures are highest over the summer pole** and lowest over the winter pole in the stratosphere.
  - This **layer is almost free from clouds and associated weather phenomena,** making conditions most **ideal for flying aeroplanes.**
  - Sometimes, rarely, **cirrus clouds are present at lower levels** in this layer.
- **Mesosphere: 50-80 km.**
  - **Up to the Mesopause**
  - The **coldest layer** of the atmosphere, where the **temperature again reduces with the height.**
  - The air is so cold just below the mesopause that **even the scarce water vapour** can be **sublimated into polar-mesospheric noctilucent clouds** .
  - **Most of the meteors burn up in this layer.**
- **Thermosphere: 80-400 km.**
  - Above the mesopause, **temperatures again increase with height.** Due to the **absorption of energetic ultraviolet and X-ray radiation** from the sun.
  - The **International Space Station and satellites orbit here.**
  - The **Kármán line** , located at an **altitude of 100 km** : is widely used to describe the **border between Earth's atmosphere and outer space** .
  - **Ionosphere is a part of this layer,** where electrically charged currents flow in the air in this sphere. **Radio waves are reflected on the earth from this sphere** and due to this radio broadcasting has become possible.

- **Exosphere: Beyond 400 km; High temperature of air particles** , but **less density** , so **temperature reduces with height** .
  - **Above the ionosphere and extends to 400 km** above the earth.
  - The **air** here is **extremely rarefied** .
  - Due to proximity to solar radiations, the **temperature of air particles can go up to 1000 °C.**
  - But due to low density, the air particles are so far apart that they are unable to transfer the temperature.
  - Thus, the **temperature feels cold.**

⚡ **Quick revise** Atmosphere & heat Fold ▲

🏢 **The atmosphere is a 5-floor building** (bottom → top)

Floor	Height	What happens there
5 Exosphere – open terrace	> 400 km	Particles can reach **1,000 °C** yet it *feels* cold (too thin to pass heat)
4 Thermosphere – rooftop antenna	80–400 km	Temperature **rises** again (UV, X-ray). **ISS & satellites**; **Kármán line 100 km**; **ionosphere** bounces radio waves
3 Mesosphere – freezer	50–80 km	**Coldest**; meteors burn; noctilucent clouds
2 Stratosphere – AC floor	to 50 km	**Ozone** (lower part) → temperature **rises** with height; almost cloud-free = **jets fly here**
1 Troposphere – kitchen	8–18 km (avg 13)	**All weather**, clouds, jet streams. 18 km at equator (strong convection), 8 km at poles. **Cools ~1 °C per 165 m**

- **Gases:** N_2 **78%** · O_2 **21%** (≈99%) · Argon **0.93%** · CO_2 **~0.04%** · water vapour 0–5% (**>90% below 5 km**; falls from equator to poles) · CO_2 and water vapour only **up to 90 km** · **ozone layer 20–25 km**. By composition: **homosphere** (<80 km, uniform) and **heterosphere** (>80 km, layered by molecular mass). Aerosols/salt act as **hygroscopic nuclei**; microplastics can seed ice 5–10 °C warmer than usual.
- **Why cooler with height (PYQ 2012, ans C):** air is heated **from the ground up** and thinner air holds less heat. *Not* because of "more moisture" (vapour falls with height). Handout also quotes **6.5 °C per 1,000 m** (≈ the 1 °C/165 m above).
- **Troposphere – PYQ 2024 (ans a):** thicker at the equator **because** strong convection carries heat high. *(Mains 2022, 15m: "troposphere determines weather processes – how?" → clouds, air masses, jet streams all sit here; prime heat source is terrestrial radiation.)*
- **CO_2 warms (PYQ 2012, ans D):** it absorbs **infrared**; transparent to incoming sunlight, opaque to outgoing earth radiation. **Water vapour**

also absorbs infrared (2023 ans C).

- **Ionosphere (PYQ 2011, ans D):** radio works because charged particles reflect waves – *not* ozone, and radio waves are *not* "very long" (1 mm–100 km).
- **Heat budget (PYQ 2024, ans d):** the atmosphere is heated more by **terrestrial (long-wave)** than by incoming (short-wave) radiation, and greenhouse gases absorb long-wave well → only Statement II correct.

Heat & Temperature

- The **interaction of insolation with the atmosphere and the earth's surface creates heat** , which is **measured in terms of temperature.**
- The distribution of temperature **varies both horizontally and vertically.**
- **Factors affecting the distribution of temperature:**
  - **The latitude,** which determines the angle of inclination. The higher the angle of inclination, more the distance sun rays have to travel within the atmosphere, which increases the heat lost by the sun rays. So, less heat is transferred to the Earth. This is why the poles, which get slanting solar radiation, are much cooler than the equator, which gets direct sunlight.
  - **Duration of sunlight:** During the summers, the duration of sunlight is more than during the winters (except at the equator, where day and night are equal). So, summers are warmer.
  - **Sunspot Cycle** (discussed in the Universe part)
  - **Transparency of the atmosphere** : On a cloudy day, the clouds obstruct the sunlight from reaching the earth's surface, which is why cloudy days are cooler. On the other hand, cloudy nights are warmer as the clouds do not allow terrestrial radiation to escape. The same happens when there are lots of aerosols and particulate matter in the atmosphere.
  - **Altitude of the place:** as the altitude increases, the temperature reduces per the Normal Lapse rate.
  - **Orientation:** The places to the North of mountain ranges in the Northern Hemisphere, north of the Tropic of Cancer, and those to the South of the Mountain Ranges, south of the Tropic of Capricorn, are perpetually in the shadow of the mountains (until the mountains are eroded).
  - **Vegetation cover:** Trees undertake transpiration, which has a cooling effect on the environment.
  - **Distance from the sea:** The continental locations have more extreme temperatures than the coastal locations, due to the moderating impact of the ocean/sea through the land and sea breezes.
  - **Ocean Currents:** Warm and cold ocean currents increase or reduce the temperature of the place respectively.
  - **Local factors like local winds, urban area distribution, etc** , can also affect the temperature of a location.

⚡ **Quick revise** What controls temperature Fold ▲

What controls temperature: latitude (slant rays) · duration of daylight · sunspot cycle · **transparency** (cloudy day cooler, cloudy night warmer) · altitude · mountain orientation · vegetation (transpiration cools) · **distance from sea** · ocean currents · local winds/urban.

Ways of heat transfer:

- **Conduction:** Transfer of heat **through a solid material** from **molecule to molecule** .
  - **Mechanism: Direct contact between particles** , with **heat moving from a region of higher temperature to one of lower temperature.**



  - Example: A metal rod heated at one end will gradually transfer heat to the other end.
- **Convection:** Transfer of heat by the **movement of a fluid** (liquid or gas).
  - **Mechanism: Warmer, less dense fluid rises while cooler, denser fluid sinks** , creating a circulation pattern.
  - Example: Boiling water, where hot water rises to the surface and cooler water sinks to the bottom.
- **Advection:** Transfer of heat through the **horizontal movement of air or water** .



- Mechanism: Large-scale horizontal movement of fluid carrying heat from one place to another.
- Example: Warm air moving into a region from another area, such as warm winds bringing heat to a cooler area.

- **Radiation:** Transfer of heat **through electromagnetic waves without the need for a medium** .
  - Mechanism: Heat is emitted as infrared radiation and **can travel through a vacuum** .
  - Example: The heat from the sun warming the Earth.

Horizontal temperature distribution:

- **Distribution of temperature across the latitudes** over the surface of the earth.
- **Shown by "Isotherms"**
- Generally, **from the equator to poleward** , the **temperature** keeps on **declining** .
- In general, the **equatorial region is hot** , and its temperature is high throughout the year.
- The **lowest temperature is at and near the poles.**
- The amount of insolation received varies from latitude to latitude.
- **Regions within the equator and 40° N and S latitudes** receive abundant sunlight and hence more heat will be gained than lost. Hence, they are **energy surplus** regions.
- **Regions beyond 40° N and S latitudes** lose more heat than that gained from sunlight. Hence, they are **energy deficit** regions.
- **The atmosphere (planetary winds) and the oceans (ocean currents) transfer excess heat from the tropics** (energy surplus region) towards the poles (energy deficit regions)
- Most of the **heat transfer takes place across the mid-latitudes (30° to 50°),** hence the high frequency of storms here.

- **Heat Budget:** The **earth receives** a certain amount of **Insolation (incoming short-wave solar radiation)** and **gives back heat** into space **by terrestrial radiation (outgoing long-wave terrestrial radiation).** Through this give and take, or the heat budget, the earth maintains a constant temperature.



⚡ **Quick revise** Atmosphere & heat Fold ▲

🔥 **4 ways heat moves: Con**duction = **Con**tact (spoon in hot chai) · **Con**vection = **Currents** (boiling milk rises/sinks) · **Ad**vection = **A**side, sideways (Loo carrying heat in) · **Rad**iation = **Rays**, no medium (Sun → Earth). 👕 **Albedo:** snow/ice = **white shirt** (reflects, cools) · ocean, forest, city = **black shirt** (absorbs, warms). Hotter body → **shorter** wavelength: Sun = **short-wave**, Earth = **long-wave**.

Albedo

- Albedo is a **measure of the reflectivity of a surface** , indicating how much solar radiation is reflected by that surface compared to how much is absorbed.



- **Surfaces with high albedo** , such as **snow and ice** , reflect a large portion of incoming solar radiation, contributing to cooling.
- Conversely, **dark surfaces like oceans, forests, and urban areas** have **low albedo** and absorb more solar radiation, leading to warming.
- **Short and Long wave radiation:** The **higher the temperature of the body, the shorter the wavelength of the electromagnetic radiation released** by it. Since the sun has a higher temperature than the Earth, the EMRs from the sun to the Earth are called Short wave radiations, and the EMRs released by the Earth are called Long wave radiations.
- **Vertical Distribution of Temperature:** Covered under “Layers of Atmosphere”.= **ISOTHERMS**
- **Isotherms are the lines joining locations with same temperature.**

- Bending of isotherms refers to the way lines of equal temperature (isotherms) on weather maps deviate from their expected straight east-west pattern.



Figure : The distribution of Surface Air Temperature in the Month of January

Figure : The distribution of surface air temperature in the month of July

  - This occurs due to factors like:
    - **Land-sea contrast:** Land heats and cools faster than water, causing isotherms to **bend inland in summer and towards the coast in winter** .
- In Northern Hemisphere (NH):
  - In July, isotherms shift towards equator on land, and towards pole on oceans. In January the isotherms deviate to the pole over the ocean and to the equator over the continent.
- In Southern Hemisphere (SH), this shift is not very evident due to lack of land. But, though at smaller scale, due to opposite seasons in this hemisphere compared to NH, the shift here is opposite. What NH sees in January, is seen here in July, and vice versa.
- To summarize, **in July, isotherms shift to south over oceans, and north over land. In January the isotherms deviate to the north over the ocean and to the south over the continent.**
  - **Ocean currents:** Warm or cold currents **can cause isotherms to bulge or curve depending on their temperature.**
  - **Topography: Mountains can block or channel air masses** , leading to localized temperature variations and bending of isotherms.

⚡ **Quick revise** Energy: 0–40° N/S = Fold ▲

Energy: 0–40° N/S = **surplus**; beyond 40° = **deficit**; winds and currents carry heat poleward, mostly across **30°–50°** (hence storms there).

2 · PYQ

Q.1) A layer in the Earth's atmosphere called Ionosphere facilitates radio communication. Why?

1. The presence of ozone causes the reflection of radio waves to Earth.

2. Radio waves have a very long wavelength.

Which of the statements given above is/are correct? (2011)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: D The ionosphere is defined as the layer of the Earth's atmosphere that is ionized by solar and cosmic radiation.

- It lies 75-1000 km (46-621 miles) above the Earth.
- (The Earth’s radius is 6370 km, so the thickness of the ionosphere is quite tiny compared with the size of Earth.)
- Because of the high energy from the Sun and from cosmic rays, the atoms in this area have been stripped of one or more of their electrons, or “ionized,” and are therefore positively charged. The ionized electrons behave as free particles.
- The Sun's upper atmosphere, the corona, is very hot and produces a constant stream of plasma and UV and X-rays that flow out from the Sun and affect or ionize the Earth's ionosphere.
- Only half the Earth’s ionosphere is being ionized by the Sun at any time.
- The ionosphere has major importance to us because, among other functions, it influences radio propagation to distant places on the Earth, and between satellites and Earth.
- For the very low frequency (VLF) waves that the space weather monitors track, the ionosphere and the ground produce a “waveguide” through which radio signals can bounce and make their way around the curved Earth
- The ionosphere is composed of three main parts, named for obscure historical reasons: the D, E, and F regions.
- The electron density is highest in the upper, or F region.
- The F region exists during both daytime and nighttime.
- During the day it is ionized by solar radiation, during the night by cosmic rays.
- The D region disappears during the night compared to the daytime, and the E region becomes weakened.
- Wavelength of radio waves can be 1 milimeter to 100 km, so they are not “very long wavelength waves.

Q.2) Normally, the temperature decreases with the increase in height from the Earth’s surface, because

1. the atmosphere can be heated upwards only from the Earth’s surface

2. there is more moisture in the upper atmosphere

3. the air is less dense in the upper atmosphere

Select the correct answer using the codes given below : (2012)

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: C The first statement given in this question is correct.

- Temperature falls off with height at a predictable rate because the air near the surface is heated and becomes light, and the air higher up cools to space and becomes heavy.
- The second statement is obviously incorrect. This is because the water vapour, which accounts for around 1% of air, varies greatly in the troposphere and decreases rapidly with altitude.
- As we increase elevation, there is less air above us thus the pressure decreases. As the pressure decreases, air molecules spread out further (i.e. air expands), and the temperature decreases and the atmosphere can be heated upwards only from the Earth's surface.
- Near the Earth's surface, air gets cooler the higher you climb. As you climb a mountain, you can expect the air temperature to decrease by 6.5 degrees C for every 1000 meters you gain. This is called the standard (average) lapse rate.
- The basic answer is that the farther away you get from the earth, the thinner the atmosphere gets. The total heat content of a system is directly related to the amount of matter present, so it is cooler at higher elevations

Q.3) The increasing amount of carbon dioxide in the air is slowly raising the temperature of the atmosphere because it absorbs - (2012)

(a) the water vapour of the air and retains its heat

(b) the ultraviolet part of the solar radiation

(c) all the solar radiations

(d) Infrared part of solar radiation

Answer: D Carbon dioxide is meteorologically a very important gas as it is transparent to the incoming solar radiation but opaque to the outgoing terrestrial radiation.

- It absorbs a part of terrestrial radiation and reflects back some part of it towards the earth’s surface. It is largely responsible for the greenhouse effect.
- Its concentration is greater close to the earth’s surface as it is denser than air.

Q.4) The annual range of temperature in the interior of the continents is high as compared to coastal areas. What is/are the reason/reasons? (2013)

1. Thermal difference between land and water

2. Variation in altitude between continents and oceans

3. Presence of strong winds in the interior

4. Heavy rains in the interior as compared to coasts

Select the correct answer using the codes given below.

(a) 1 only

(b) 1 and 2 only

(c) 2 and 3 only

(d) 1, 2, 3 and 4

Answer: A

- The major factor affecting the distribution of the temperature of Earth is distribution of Land and Oceans. Since there is more land in the Northern Hemisphere and more waters in the Southern hemisphere, there is a big difference between the specific heat of land and water; the loss of heat from the continents is bigger than the oceans. The continents get heated faster and get cooled faster in comparison to the Oceans. This is the reason that the temperatures of the Oceans are moderate while that of continents is extreme. The moderating effect on temperature of the land due to proximity of the seas is called Maritime influence. The increasing effect on temperature of the land at the interior of the continents is called Continental Influence.
- Due to this **moderating effect of the sea** , places near the coast have a low range of temperature and high humidity.The places in the interior of the continent do not experience the moderating effects of the sea. These places have extreme temperatures.
- Statement 4 claims that Heavy rains in the interior as compared to coasts (are responsible for why the annual range of temperature in the interior of the continent is high)
- But the interiors in Central Asia receive very poor rainfall due to the distance from the sea. Annual precipitation is usually low in the interior areas. Hence 4 cannot be the answer

Q.5) Consider the following statements: Statement-I: The temperature contrast between continents and oceans is greater during summer than in winter. **Statement-II:** The specific heat of water is more than that of land surface.

Which one of the following is correct in respect of the above statements? (2023)

(a) Both Statement-I and Statement-II are correct and Statement-II is the correct explanation for Statement-I

(b) Both Statement-I and Statement-II are correct and Statement-II is not the correct explanation for Statement-I

(c) Statement-I is correct but Statement-II is incorrect

(d) Statement-I is incorrect but Statement-II is correct

Answer: D

- Water has a very high specific heat, which means that in order to increase or reduce its temperature by 1 degree celsius, it will have to absorb or release huge amount of heat. Compare t that, the land has low specific heat, and can heat up or cool down fast.

Q.6) With reference to the Earth's atmosphere, which one of the following statements is correct? (2023)

(a) The total amount of insolation received at the equator is roughly about 10 times of that received at the poles.

(b) Infrared rays constitute roughly two-thirds of insolation.

(c) Infrared waves are largely absorbed by water vapour that is concentrated in the lower atmosphere.

(d) Infrared waves are a part of the visible spectrum of electromagnetic waves of solar radiation.

Answer: C

- Water vapour is an important Greenhouse Gas, which absorbs the heat in the form of Infrared Radiations.
- The troposphere is a very significant atmospheric layer that determines weather processes. How ? (2022, 15m)

Q.7) Consider the following statements: (2024) Statement-I: The atmosphere is heated more by incoming solar radiation than by terrestrial radiation.

Statement-II: Carbon dioxide and other greenhouse gases in the atmosphere are good absorbers of long wave radiation.

Which one of the following is correct in respect of the above statements?

(a) Both Statement-I and Statement-II are correct and Statement-II explains Statement-I

(b) Both Statement-I and Statement-II are correct but Statement-II does not explains Statement-I

(c) Statement-I is correct, but Statement-II is incorrect

(d) Statement-I is incorrect, but Statement-II is correct

Answer:

(d) Statement-I is incorrect, but Statement-II is correct

Q.8) Consider the following statements: (2024) Statement-I: Thickness of the troposphere at the equator is much greater as compared to poles. **Statement-II:** At the equator, heat is transported to great heights by strong convectional currents.

Which one of the following is correct in respect of the above statements?

(a) Both Statement-I and Statement-II are correct and Statement-II explains Statement-I

(b) Both Statement-I and Statement-II are correct, but Statement-II does not explains Statement-I

(c) Statement-I is correct, but Statement-II is incorrect

(d) Statement-I is incorrect, but Statement-II is correct.

Answer:

- Both Statement-I and Statement-II are correct and Statement-II explains Statement-I

Q.9) Which of the following is/are correct inference/inferences from isothermal maps in the month of January? (2024)

1. The isotherms deviate to the north over the ocean and to the south over the continent.

2. The presence of cold ocean currents, Gulf Stream and North Atlantic Drift make the North Atlantic Ocean colder and the isotherms bend towards the north.

Select the answer using the code given below:

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer:

(a) 1 only

Q.10) Consider the following statements: Statement I: In January, in the Northern Hemisphere, the isotherms bend equatorward while crossing the landmasses, and poleward while crossing the oceans. **Statement II:** In January, the air over the oceans is warmer than that over the landmasses in the Northern Hemisphere.

Which one of the following is correct in respect of the above statements? (2025)

(a) Both Statement I and Statement II are correct and Statement II explains Statement I

(b) Both Statement I and Statement II are correct but Statement II does not explain Statement I

(c) Statement I is correct but Statement II is not correct

(d) Statement I is not correct but Statement II is correct

Answer:

(a) Both Statement I and Statement II are correct and Statement II explains Statement I Statement I: Correct

- In January (winter in the Northern Hemisphere), isotherms (lines of equal temperature) bend equatorward over landmasses because land cools faster than water.
- Over oceans, where temperatures remain relatively warm, isotherms bend poleward.
- This distortion of isotherms is a well-known climatological feature seen on global isotherm maps for January.

Statement II: Correct

- Due to higher specific heat and slower cooling, oceans retain more heat than land during winter. Hence, in January, air over oceans is warmer than over land.
- This temperature differential explains why maritime climates are milder than continental ones in winter.

Q.11) Consider the following statements:

1. Without the atmosphere, temperature would be well below freezing point everywhere on the Earth’s surface.

2. Heat absorbed and trapped by the atmosphere maintains our planet’s average temperature.

3. Atmosphere’s gases, like carbon dioxide, are particularly good at absorbing and trapping radiation.

Which of the statements given above are correct?

(a) I and III only

(b) I and II only

(c) I, II and III

(d) II and III only

Answer: (d) II and III only Statement 1: It's overgeneralized and incorrect to say temperatures would be "well below freezing everywhere." While nights would be freezing, daytime temperatures could become extremely hot. **Statement II:** Correct The atmosphere traps infrared radiation, regulating Earth's temperature. **Statement III:** Correct Greenhouse gases like CO_2, CH_4, and H_2O vapor absorb and re-radiate heat.

⚡ **Quick revise** Isotherm bending (PYQs 2024, 2025) Fold ▲

Isotherm bending (PYQs 2024, 2025): *January, Northern Hemisphere* – isotherms bend **poleward over oceans** (oceans stay warmer) and **equatorward over continents** (land cools fast). July is the reverse. *Reason:* water has high **specific heat** (2023: Statement I wrong – contrast is greater in **winter**; Statement II right). **Continental interiors have a bigger annual range** because of land–water thermal difference only (2013, ans A).

> **Handout alert:** the isotherm section first says "in July isotherms shift towards the equator on land" and then summarises "in July, north over land, south over oceans". Trust the **summary** (and the Jan rule above).

3 · Temperature Inversion:

- **Inversion of the normal lapse rate** , such that as the altitude increases, the temperature starts to increase too.
- **Conditions for Temperature Inversion:**
  - **Clear, Calm Conditions:** Clear skies and calm winds allow for efficient radiative cooling of the Earth's surface during the night.
  - **Long Nights or Extended Darkness:** Longer nights or periods of extended darkness allow for more significant radiative cooling of the surface air.

- **Absence of Atmospheric Mixing:** A lack of vertical mixing, such as from strong winds or convective activity, prevents the mixing of warmer air aloft with cooler air near the surface, allowing the inversion layer to form and persist.
- **Geographic Features** : Certain geographic features, such as valleys or basins, can enhance the likelihood of temperature inversion by trapping cold air near the surface, especially during calm, clear nights.

- **Most commonly seen at:**
  - **Fronts**
  - **Winter season**
  - **Mountains & Valleys**

Positive and Negative Impacts

Positive:

- **Frost Protection for Plants:** The warmer air above the inversion layer can prevent ground temperatures from dropping too low, protecting crops from frost damage.

Negative:

- **Pollution Dome Effect**
- **Reduced Visibility** due to fog formation, affects transportation.
- Can create conditions where **pests and diseases thrive** , potentially harming crops if not managed properly.

Pollution Dome Effect:

- The pollution dome effect refers to the phenomenon where **pollutants become trapped beneath a layer of warm air during temperature inversion** events.
- During a temperature inversion, a layer of warm air traps cooler air beneath it, along with any pollutants present in the lower atmosphere. This creates a **"dome" or "cap" of stagnant air that prevents the vertical dispersion of pollutants.**



- This usually happens during **cold weather, when the air in contact with the cold ground becomes cool,** resulting in temperature inversion.
- **Temperature Inversion's Role:** Temperature inversion **exacerbates** the pollution dome effect by **preventing vertical mixing of the atmosphere** . Normally, warmer air rises and cooler air sinks, facilitating the dispersal of pollutants. However, during inversion, this vertical movement is suppressed, leading to the accumulation of pollutants near the surface.
- **Consequences:** The pollution dome effect can lead to **increased concentrations of pollutants in affected areas** , resulting in **poor air quality and health hazards** for residents. In severe cases, prolonged exposure to high levels of pollutants trapped within the dome can lead to **respiratory problems, cardiovascular issues, and other health complications** .
- **Mitigation** : Strategies to mitigate the pollution dome effect include **reducing emissions of pollutants** at their sources, implementing **air quality monitoring and forecasting systems** , and promoting **public awareness and behaviour change** to reduce individual contributions to air pollution.
- In 2013, China launched a $100-billion initiative to combat air pollution in Beijing, implementing strict regulations on vehicle emissions, relocating polluting industries, and promoting cleaner energy sources. As a result, Beijing's air quality improved substantially, with PM2.5 levels decreasing by 35% between 2013 and 2017.

⚡ **Quick revise** Temperature inversion Fold ▲

Temperature inversion: temperature *rises* with height. Needs **clear, calm, long nights, no mixing, valleys/basins**; common at **fronts, in winter, in mountains and valleys**. *Good:* frost protection. *Bad:* **pollution dome**, fog, pests. (Beijing's 2013 US$100-bn drive cut PM2.5 by 35% by 2017.)

4 · AIR PRESSURE & PRESSURE BELTS

- Total **weight of a mass of column of air above per unit area at sea level.**



Major Pressure Belts and Wind System

- **Maximum at sea level**
- **High-temperature** results in **Low Pressure and vice versa.**
- When the air gets heated it expands, becomes light and rises vertically. As air rises, the pressure it exerts on the Earth's surface is reduced, causing a low-pressure area. On the other hand, cool air is dense and heavy. As a result, it sinks vertically. It results in additional weight and pressure which causes a high pressure on the ground.
- **Lines joining places with equal pressure** are called **Isobars.**
- Based on temperature distribution, **7 pressure belts** are present on the Earth.
  - **3 of these are thermally induced** , created due to the temperature.
  - **4 are dynamically induced** due to the movement of the winds from the thermally induced pressure belts.
- The Equatorial Low-Pressure belt **(10 degrees north and south) is also known as the ITCZ (Inter-Tropical Convergence Zone)** or the **"Doldrums"** (calms), as this region experiences **monotonous windless weather** , due to the convergence of the trade winds around it.
- **Pressure belts** are **not homogenous** , due to **uneven distribution of land and water.**
- **Pressure belts shift** due to the revolution of Earth **because of Earth's 23.5-degree tilt** , depending upon where the sunlight falls directly.
- In the Northern Hemisphere, January denotes winter, and July denotes summer. In the Southern Hemisphere, the opposite is true. **The pressure belts shift approx 5°- 10° northward (can be as high as 25° over India) when the sun shines vertically overhead on the Tropic of Cancer (21 June), and approx 5°- 10° southward when the sun shines vertically overhead on the Tropic of Capricorn** (22 December) from their initial position.
- **Seasonal climate fluctuations are caused by the movement of the pressure belts** , especially between the latitudes of 30° and 40° (Mediterranean Climate) in both hemispheres. During winter, Westerlies prevail, resulting in rain. During the summer, dry Trade Winds blow offshore, preventing rain from falling in certain areas.

- The pressure belts in both hemispheres stay balanced when the sun shines vertically over the Equator on March 21st and September 23rd (the Equinoxes).

WINDS: FACTORS AFFECTING THEIR MOVEMENT

- **Pressure Gradient (Direction and Speed)** : It denotes the pressure difference between two locations. The winds always move from the High to the Low-pressure region. This determines the direction of the winds.
  - The steep Pressure Gradient is denoted on the maps via closely spaced isobars. The steeper the gradient is, the higher the speed of movement of the winds.
  - So overall, this force determines **both the speed and the direction of the wind movement** i.e. it helps determine the overall velocity of the Winds.
- **Coriolis Force (Only Direction)**
  - This force is generated by virtue of the movement of the Earth and **affects only the objects or entities moving on the surface of the Earth** , like winds and ocean currents. That is why it is an apparent and not a real force.
  - It affects **only the direction** of movement of the winds.
  - The **magnitude of deflection depends upon the speed of the wind** . Higher speed means higher deflection.
  - Deflects the winds to the **Right in the Northern Hemisphere.**
  - Deflects the winds to the **Left in the Southern Hemisphere.**
  - **0 at the equator, maximum at the poles.**
- **Frictional Force (Direction and Speed):**
  - It is the force experienced by the winds in the **direction opposite to its movement.**
  - It can affect **both the direction and the speed** of wind movement.
  - It is effective only up to a few thousand meters above the Earth's surface.
  - The oceans provide much less frictional force than land, as the latter is covered with vegetation, infrastructure, landforms, etc. That is why the wind speeds are higher over water bodies than on the land.

⚡ **Quick revise** Three forces on wind Fold ▲

Three forces on wind: Pressure gradient (speed + direction; close isobars = fast) · **Coriolis** (direction only; deflects **right in N, left in S**; **0 at equator, max at poles**; bigger for faster winds – PYQ 2024, ans c) · **Friction** (both; only a few thousand metres up; less over sea so winds are faster there). Winds are **named for where they come from**.

TYPES OF WINDS

- **Prevailing/Primary/Planetary/Permanent winds:** These winds get **affected by all the abovementioned forces** and affect the Earth's surface throughout the year. E.g. Trade winds, westerlies, polar easterlies.

- **Geostrophic Winds:** They move at tropopause-ic level, and therefore are **not affected by the frictional force** . As a result, they travel parallel to the isobars. E.g. Jet streams.
- **Seasonal Winds:** They are experienced only during a particular season. E.g. Monsoonal winds.
- **Diurnal winds: change direction between day and night.** E.g. in coastal regions, the sea breeze blows during the day and the land breeze blows during the night. Another example is the mountain valley breeze.
- **Local Winds:** Winds that **affect only a specific location.** May be all year round, or during a particular season.

NOMENCLATURE OF WINDS

- Winds are **named after the direction from which they seem to be coming** . E.g. Polar easterlies are named so as they come from the eastern direction. They have an east-to-west movement.

PLANETARY WINDS AND TRI-CELLULAR MERIDIONAL CIRCULATION:

- Planetary winds are **large-scale wind patterns that circulate the Earth** , driven by differences in temperature and pressure between regions.



- They are present **throughout the Year** .
- These winds play a **crucial role in redistributing heat and moisture across the planet** , influencing weather patterns and climate. There are several key types of planetary winds:
  - **Trade Winds** : Trade winds are persistent **easterly winds** that **blow from the subtropical high-pressure belts towards the equator in both hemispheres.** They are named "trade winds" because they historically facilitated trade routes across the world's oceans. Trade winds play a significant role in the movement of ocean currents and are important for sailors and aviation routes.
  - **Westerlies** : Westerlies are prevailing winds that blow **from the subtropical high-pressure belts towards the sub-polar low-pressure region in both hemispheres.** They are characterized by their **westerly direction** (from west to east) and play a crucial role in the movement of weather systems across mid-latitudes. The westerlies are particularly strong and well-defined in the Southern Hemisphere and flow at

extremely high speeds. This has given them the names like roaring fourties, furious fifties, and shrieking sixties.

- **Polar Easterlies** : Polar easterlies are cold winds that blow from the polar **high-pressure belts towards the mid-latitudes in both hemispheres** . They are characterized by their **easterly direction** (from east to west) and are influenced by the temperature gradient between the polar regions and the mid-latitudes. Polar easterlies are weaker and less consistent than the trade winds and westerlies.

- **Tricellular Meridional Circulation:** Tricellular meridional circulation refers to the **three-cell circulation pattern** in each hemisphere of the Earth's atmosphere, characterized by the movement of air from the equator towards the poles and back again.
  - **Hadley Cell** : The Hadley cell is the largest and most prominent cell in the tricellular circulation, extending **from the equator to about 30 degrees latitude** . Warm, moist air rises at the equator, forming low-pressure zones and generating trade winds. As the air cools and descends around 30 degrees latitude, it creates high-pressure belts and generates the trade winds.
  - **Ferrel Cell** : The Ferrel cell is a **mid-latitude cell** that lies between the Hadley and polar cells. It operates in the **opposite direction to the Hadley** cell, with air rising at around 60 degrees latitude and descending at around 30 degrees latitude. The Ferrel cell is driven by the interaction between the Hadley and polar cells.
  - **Polar Cell** : The polar cell is the **smallest and weakest** in the tricellular circulation, extending **from around 60 degrees latitude to the poles.** Cold, dense air sinks at the poles, creating high-pressure zones and generating the polar easterlies. As the air warms and rises around 60 degrees latitude, it creates low-pressure belts and generates westerly winds.
  - **Importance** : Tricellular meridional circulation plays a crucial role in **regulating global climate and weather patterns** by transporting heat, moisture, and energy around the planet. It helps to redistribute heat from the equator towards the poles and vice versa, influencing temperature gradients, atmospheric pressure systems, and wind patterns.

⚡ **Quick revise** Pressure, winds & circulation Fold ▲

🪜 **Pressure ladder – "Lazy Hens Lay Happily":** Equator **L**ow (ITCZ / **Doldrums**, 10°N–S) → **H**igh at ~30° → **L**ow at ~60° → **H**igh at poles. **7 belts = 3 thermal + 4 dynamic.** Belts **shift 5–10° (up to 25° over India)** with the overhead Sun; balanced at the equinoxes (21 Mar, 23 Sep). At 30–40° this shift gives the **Mediterranean** pattern: winter westerlies = rain; summer dry trades.

Types: planetary · **geostrophic** (above friction, parallel to isobars, e.g. jet streams) · seasonal (monsoon) · diurnal (sea/land breeze, mountain-valley) · local.

⚡ **Quick revise** Pressure, winds & circulation Fold ▲

Planetary wind	From → to	Notes
Trade winds	subtropical high → equator, easterly	helped trade routes; push ocean currents
Westerlies	subtropical high → sub-polar low, W→E	strongest in S. Hemisphere (little land): **Roaring 40s, Furious 50s, Shrieking 60s**
Polar easterlies	polar high → mid-latitudes	cold, weaker

> 🌀 **Three cells: Hadley** (equator–30°, biggest, trades) · **Ferrel** (30–60°, middle, runs opposite) · **Polar** (60°–pole, smallest, polar easterlies). *Hadley Hot, Ferrel Fill-in, Polar Puny.*

5 · PYQ

Q.1) Westerlies in the southern hemisphere are stronger and persistent than in the northern hemisphere. Why?

1. The Southern hemisphere has less landmass as compared to the Northern hemisphere.

2. Coriolis force is higher in Southern hemisphere as compared to the Northern hemisphere

Which of the statements given above is/are correct? (2011)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: A

- The Westerlies are strongest in the Southern hemisphere and at times when the pressure is lower over the poles, while they are weakest in the Southern hemisphere and when pressures are higher over the poles.
- The Westerlies are particularly strong in areas where land is absent because land mass obstructs the movement of westerlies.
- Westerlies wind of the Southern hemisphere are known by different names like at 40 degrees South latitude they are called the Roaring forties, at 50 degrees latitude, they are called Furious fifties.
- The Westerlies play an important role in carrying the warm, equatorial waters and winds to the western coasts of continents, especially in the southern hemisphere because of its vast oceanic expanse.

- Wind never proceeds or moves in a straight line manner, during the movement of the wind deflection is caused due to Coriolis Force.
- Coriolis Force is highest at the Poles and zero at Equator.

Q.2) The jet aircraft fly very easily and smoothly in the lower stratosphere. What could be the appropriate explanation?

1. There are no clouds or water vapour in the lower stratosphere.

2. There are no vertical winds in the lower stratosphere.

Which of the statements given above is/are correct in this context? (2011)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: C

- Commercial jet aircraft fly in the lower stratosphere to avoid the turbulence.
- The Stratosphere is very dry; air there contains very little water vapour. Because of this, few clouds are found in this layer.
- Vertical wind shear is a change in wind speed or direction with change in altitude.
- Wind shear has significant effects on control of an aircraft, and it has been a sole or contributing cause of many aircraft accidents.
- There is no vertical wind shear in the lower stratosphere.
- That's plane fly in that region to avoid turbulence or an accident

Q.3) With reference to "Coriolis force", which of the following statements is/are correct? (2024)

1. It increases with increase in wind velocity.

2. It is maximum at the poles and is absent at the equator.

Select the answer using the code given below:

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: (c) Both 1 and 2

⚡ **Quick revise** PYQ 2011 (ans A) Fold ▲

PYQ 2011 (ans A): Southern westerlies are stronger **because the southern hemisphere has less land** – *not* because Coriolis is higher there. **PYQ 2015 (ans B):** westerlies blow between **30° and 60°** (not

"30°N to 60°S") and bring NW India's winter rain (western disturbances).

6 · GEOSTROPHIC WINDS:

- Geostrophic winds are large-scale **winds that blow parallel to isobars** (lines of constant pressure) in the atmosphere. They are the result of a **balance between the pressure gradient force and the Coriolis force** , acting perpendicular to the pressure gradient.
- **Formation** : Geostrophic winds form when the **pressure gradient force** , which drives air from areas of high pressure to areas of low pressure, is **balanced by the Coriolis force** , which deflects moving air to the right in the Northern Hemisphere and the left in the Southern Hemisphere due to the Earth's rotation.

7 · JET STREAMS

- Jet streams are **narrow bands of fast-flowing, high-altitude winds** found in the upper atmosphere, typically at altitudes of 8 to 15 kilometres. They are characterized by their high wind speeds and meandering, ribbon-like appearance.
- **Formation** : Jet streams are primarily formed by the **interaction between the temperature gradients of different air masses.** The largest and most well-known jet streams are the polar jet stream and the subtropical jet stream, which are located near the boundaries of the polar and tropical air masses, respectively.
- **Wavy Jet Streams** are said to have " **Rossby Wave** " characteristics.
  - Coriolis force and reduction in temperature contrast, can lead to the Rossby wave characteristic. **Due to climate change, the Jet streams are weakening and becoming more meandering.**
- **Types** : They can be permanent or seasonal.
  - **Permanent:** Polar Jet Streams, Sub-tropical Jet streams
  - **Temporary** : Tropical Easterly Jet, Somali Jet Stream.
- **Polar Jet Stream** : The polar jet stream is located at **high latitudes, typically between 50 and 60 degrees north and south of the equator.** It forms along the boundary between cold polar air to the north and warmer subtropical air to the south. The polar jet stream is **strongest in winter and weaker in summer** , and it plays a crucial role in steering weather systems and influencing storm tracks.
- **Subtropical Jet Stream** : The subtropical jet stream is located at **lower latitudes** , typically **between 20 and 30 degrees north and south of the equator** . It forms along the boundary between warm tropical air to the south and cooler mid-latitude air to the north. The subtropical jet stream is **weaker and less consistent than the polar jet stream but still influences weather patterns and atmospheric circulation.**
- **Impact on Weather & Economy** :

- Jet streams play a significant role in shaping weather patterns by steering the movement of weather systems and influencing the development of storms and precipitation.
- They **move air masses** from one place to another, leading to the development of fronts and extratropical cyclones.
- They can also **affect aviation routes** , with aircraft often taking advantage of the strong tailwinds associated with jet streams to reduce flight times and fuel consumption.
- Jet Stream may **transport ozone-depleting substances** higher up in the stratosphere. This vertical air circulation causes a rapid rate of mixing of air between the Troposphere & Stratosphere.
- The **Monsoon** of South Asia is largely controlled & affected by Jet Streams.
- In India, they bring **Western disturbances** which are extra-tropical cyclones of the Mediterranean, dragged into India by the subtropical westerly jet streams.

- **Jet Streams and Climate Change:**
  - One potential consequence of climate change on jet streams is their potential **weakening in strength.** As the Arctic region experiences accelerated warming compared to other parts of the globe, the temperature contrast between the Arctic and lower latitudes diminishes. Since the movement of jet streams is driven by these temperature variations, a less pronounced gradient can lead to jet streams that are weaker and exhibit more meandering paths. This phenomenon can result in prolonged periods of certain weather conditions, such as heatwaves, coldwaves, droughts, or heavy rainfall.
  - Furthermore, the altered temperature differentials may **induce shifts in the positioning of jet streams.** As they migrate closer to the poles or display more erratic trajectories, they can exert influence on the trajectories of storms and weather systems. Consequently, this can bring about alterations in precipitation distribution, storm intensity, and the frequency of extreme weather occurrences.

Heat Dome Effect:

- A heat dome refers to a weather phenomenon **characterized by the trapping of hot air beneath a dome of high pressure in the atmosphere.** This leads to prolonged periods of extremely hot temperatures and stifling conditions, particularly during the summer months.



- **Formation:** Heat domes typically form when a **large, stagnant area of high pressure settles over a region** , causing the **air beneath it** to **sink and compress** . As the air descends, it warms up due to compression, **creating a dome of hot air near the surface.**
- **Consequences:** The Heat Dome effect can result in **dangerously high temperatures, heatwaves, and heat-related illnesses** such as heat exhaustion and heatstroke. Prolonged exposure to extreme heat can also **impact agriculture, infrastructure, and energy demand** .
- **Regional Variability:** Most common in regions with **continental climates** . Also **since Jet Streams have a role in creating** these Heat Domes, they are **mostly found in Middle or higher latitudes.**
- **Climate Change:** Climate change is **expected to increase the frequency, intensity, and duration** of heat domes in many regions, exacerbating the risk of heat-related impacts and challenging efforts to adapt to a warming climate.

DANA (Depresion Aislada en Niveles Altos)

Isolated Depression at high levels/Cut-off Low **Polar Vortex**



- **Cold air trapped in its place due to Polar Jet Stream** .
- Can be **in both the poles** , but **impact on human life is mainly due to the North pole vortex** .
- Due to global warming & Arctic Amplification and resultant **meandering of the Jet streams, this parcel of cold air can intrude in the non-polar, lower latitudes, causing extremely cold conditions.**



Stratospheric Cooling

Stratospheric cooling refers to the **gradual decrease in temperature in the stratosphere** , primarily due to **human-induced changes** like greenhouse gas emissions and ozone depletion.

1. Greenhouse Gas Effect (Counterintuitive Cooling)

- Greenhouse gases (especially CO_2) **trap heat in the troposphere** , warming the lower atmosphere.
- However, in the **stratosphere** , CO_2 acts differently, it **radiates heat away into** space.
  - CO_2 molecules absorb infrared (IR) radiation and then **re-emit it in all directions** .
  - In the **dense troposphere** , that emitted energy often gets **reabsorbed by other molecules** , keeping heat trapped (warming effect).
  - But in the **thin stratosphere** , there are **fewer molecules** to absorb that radiation.
  - Because the stratosphere is less dense and lies higher in the atmosphere, much of the IR radiation emitted by CO_2 simply **escapes into space** .
- As more CO_2 accumulates, the **stratosphere loses more heat** , causing it to **cool down** .
- **2. Ozone Layer Depletion**
- Ozone in the stratosphere **absorbs UV radiation** , releasing heat and maintaining stratospheric temperature.
- Due to CFCs and other ozone-depleting substances, the **ozone layer thins** , especially over the poles.
- With **less ozone** , there is **less UV absorption** , leading to **less heating** and hence cooling. Anthropogenic greenhouse gases began affecting the stratosphere by ~1885, five decades before the detection of CO_2's impact on tropospheric warming. **IMPACT:**
- **Slows Ozone Layer Recovery**
  - Cooling reduces chemical reaction rates needed to **rebuild ozone** , delaying ozone layer recovery.
  - **Polar Stratospheric Clouds (PSCs)** form more readily in cold conditions, which **enhance ozone depletion** , especially over Antarctica.
- **Strengthens the Polar Vortex**
  - A cooler stratosphere **intensifies the polar vortex** , a band of strong westerly winds circling the poles.
  - This can trap cold air over polar regions or, when destabilized, lead to **cold outbreaks** in mid-latitudes (e.g., North America, Europe).
- **Disrupts Jet Streams and Weather Patterns**
  - Changes in the thermal structure of the stratosphere can shift **jet streams** , impacting **storm tracks** , **monsoons** , and **mid-latitude cyclones** .
- **Extreme Weather Events**
  - This vertical temperature gradient between the stratosphere and troposphere (due to global warming) can **enhance convection** and influence **weather extremes** .
- **Affects Aviation and Satellite Orbits**
  - **Temperature and density changes** in the upper atmosphere can alter aircraft flight performance and affect **satellite drag** , impacting orbital lifetimes.
- **Key Indicator of Climate Change**
  - The simultaneous **tropospheric warming** and **stratospheric cooling** is a **distinct fingerprint of anthropogenic climate change** , not caused by solar activity or natural variability.

⚡ **Quick revise** Jet streams Fold ▲

- **Jet streams:** narrow fast bands at **8–15 km**. **Permanent:** Polar (50–60°, strongest in winter) and Subtropical (20–30°). **Temporary:** Tropical Easterly, Somali. Occur in **both** hemispheres (PYQ 2020: "Northern only" is wrong). Wavy = **Rossby waves**. They steer storms, carry **western disturbances** (Mediterranean extra-tropical cyclones, dragged in by the subtropical westerly jet), control the **South Asian monsoon**, mix troposphere and stratosphere (ozone-depleting substances).
- **Climate change:** Arctic warms faster → smaller temperature contrast → **weaker, more meandering jets** → long heatwaves, coldwaves, droughts or floods.
- **Heat dome:** stagnant high pressure; sinking air compresses and warms; mid/high latitudes, continental climates. **Polar vortex:** cold air held by the polar jet; meanders let it spill south. **DANA:** isolated depression at high levels (cut-off low).
- **Stratospheric cooling:** CO_2 in thin stratosphere **radiates heat to space**; ozone loss = less UV heating. Effects: slower ozone recovery, more **PSCs**, stronger polar vortex, disrupted jets. "Warm troposphere + cooling stratosphere" is the **fingerprint of human-caused change** (GHGs affected it by ~1885).

8 · LOCAL WINDS

- They affect **only a certain specific part of the world** . These winds **can be both warm and cold** . Here is a list of the major local winds of the world:

Wind Name	Location	Main Charac
Chinook	Eastern slopes of Rockies (USA, Canada)	Warm, dry, c snow, "snow
Foehn	Leeward side of Alps (Central Europe)	Strong, dry, rapid temper
Khamsin	Egypt, eastern Mediterranean	Dry, dust-lac days
Sirocco	Sahara to Mediterranean (Italy, Balkans, North Africa)	Hot, dry w moist/rainy i "Ghibli" ("Khamsin" (E
Harmattan	West Africa (Sahara to Gulf of Guinea)	Dry, dust-la lowers humi doctor/Docto
Loo	North India, Pakistan (Indo-Gangetic Plain)	Furious, ho (May-June), deserts
Bora	Adriatic Sea coasts (Italy, Slovenia, Croatia)	Extremely c onset, hurric
Mistral	Rhône Valley and Provence, France	Cold, stron accelerates s
Santa Ana	Southern California (USA)	Hot, dry, gus worsens wild
Blizzard	Tundra/Polar regions, North America, Antarctica	Severe snov <0.25 mile vi
Purgas/ Purga	Russia (Siberia, steppes)	Bitterly co northeasterly visibility
Pamperos	Pampas of Argentina, Uruguay, Brazil	Speedy, viole stormy wea drops

⚡ **Quick revise** Local winds (name → hook) Fold ▲

Wind	Where	Hook
Chinook	E. Rockies (USA, Canada)	warm dry, "**snow eater**"
Foehn	Alps (leeward)	warm dry descending
Sirocco (Ghibli, Chili, Khamsin)	Sahara → Mediterranean	hot, dry; rainy in S. Europe
Harmattan	W. Africa (Sahara → Guinea)	dry, dusty NE trade; "**Doctor**" wind
Loo	N. India, Pakistan, May–June	furious, hot, heatstroke
Bora	Adriatic (Italy, Slovenia, Croatia)	extremely cold dry NE, katabatic
Mistral	Rhône valley (France)	cold NW
Santa Ana	S. California	hot dry; worsens wildfires
Khamsin	Egypt	dusty spring wind, up to 50 days
Purga	Siberia, Russian steppes	icy NE blizzard wind
Pamperos	Pampas (Argentina, Uruguay, Brazil)	violent cold SW

🧣 **Hot vs cold:** Chinook, Foehn, Sirocco, Harmattan, Loo, Santa Ana = **hot/warm** · Bora, Mistral, Purga, Pamperos = **cold**. *Mistral & Bora = "Brrr" winds.*

9 · MONSOONAL WINDS:

- These are **seasonal winds** that are **experienced in the tropical regions of the world** . The most peculiar part about these winds is the **reversal they experience with the switch in the apparent positioning of the sun.**
- This portion will be covered in more detail in the Indian Physical Geography section, as India is highly dependent on these monsoonal winds for its climate, vegetation, economy, as well as society.

⚡ **Quick revise** Monsoon (PYQ 2014, ans C) Fold ▲

Monsoon (PYQ 2014, ans C): seasonal **reversal of winds** is typical of the **monsoon** climate.

10 · HUMIDITY & CONDENSATION

- **Humidity** refers to the **amount of water vapor present in the air.** It is a measure of the moisture content of the atmosphere and plays a significant role in determining weather and climate conditions.
- There are several different **types** of humidity:
  - **Absolute Humidity** : Absolute humidity is the **actual amount of water vapor present in a unit volume of air** , typically expressed in grams per cubic meter (g/m^3). It provides a direct measure of the moisture content of the air.
  - **Relative Humidity** : Relative humidity is the **ratio of the amount of water vapor present in the air to the maximum amount of water vapor that the air can hold at a given temperature** , **expressed as a percentage** . It indicates how close the air is to saturation and is commonly used to describe how "humid" or "muggy" the air feels.
  - **Specific Humidity** : Specific humidity is the **ratio of the mass of water vapor to the total mass of air** (including dry air and water vapor) in a sample of air. It is expressed in grams of water vapor per kilogram of air and provides a measure of the actual moisture content of the air without regard to temperature.
- **Condensation** is the process by which **water vapor in the air transforms into liquid water** . There are several types of condensation, including:
  - **Dew** : Dew forms when the **surface temperature** of an object, such as grass or car windows, **falls below the dew point temperature** of the surrounding air. **Water vapor** in the air condenses onto the cooler surface, **forming small droplets of liquid water** .
    - **Dew Point:** Dew point is the **temperature at which air becomes saturated** with water vapor and condensation begins to form. It is a measure of how much moisture is present in the air and is often used as an indicator of comfort level, with lower dew points indicating drier air and higher dew points indicating more humid air.
  - **Fog** : Fog is a **type of condensation** that occurs when moist air cools and reaches its dew point, causing water vapor to condense into tiny water droplets suspended in the air near the ground. Fog reduces visibility and is commonly observed in low-lying areas, near bodies of water, or during temperature inversions. **Fogs are near-surface clouds.**
  - **Clouds** : Clouds are **large masses of water droplets or ice crystals suspended in the atmosphere.** They form when moist air rises, cools, and reaches its dew point, causing water vapor to condense onto tiny particles called condensation nuclei. Clouds play a crucial role in the Earth's climate system and weather patterns.
  - **Frost** : Frost **forms when water vapor in the air undergoes deposition, transitioning directly from a gas to a solid** without passing through the liquid phase. Frost forms on surfaces such as grass, leaves, or windows when the surface temperature falls below the frost point temperature of the surrounding air.
- **Smog** :
  - **Smog** = **smoke + fog (smoky fog)** caused by the burning of large amounts of coal, vehicular emission, and industrial fumes (Primary pollutants).
  - Smog contains soot particulates like smoke, sulphur dioxide, nitrogen dioxide, and other components.

- At least two distinct types of smog are recognized: **sulphurous smog and photochemical smog.**

- **Haze:** An atmospheric phenomenon where **dust, smoke, and other dry particles obscure the clarity** of the sky.

ATMOSPHERIC RIVERS

- Atmospheric Rivers are **long, narrow corridors of concentrated moisture (water vapour) in the atmosphere.** Sometimes they can carry more water than the Amazon River. They are **responsible for transporting vast amounts of water vapor from tropical regions to mid-latitudes** (under the impact of Westerlies), often leading to heavy rainfall and flooding.
- **Creation of Atmospheric Rivers:**
  - **Evaporation:** Warm ocean waters, particularly in tropical regions, cause large-scale evaporation, adding significant moisture to the atmosphere.
  - **Transport by Wind:** Strong winds in the atmosphere, especially within the jet streams, transport this moisture in narrow bands. These winds can extend over thousands of kilometers.
  - **Convergence and Condensation:** When these moisture-laden air currents encounter mountains or cooler air masses, the moisture condenses into clouds and precipitation, often leading to intense rainfall.

TYPES OF CLOUDS

- Clouds are categorized into different types based on their altitude, appearance, and the processes that form them. Based on altitude, clouds are generally classified into three main categories:



  - **High-Level Clouds (Cirrus Clouds)** : These clouds typically form at altitudes above **6,000 meters.** They are **composed of ice crystals** and appear **thin, wispy, and feathery.** Examples include cirrus, cirrostratus, and cirrocumulus clouds.

- **Mid-Level Clouds (Alto Clouds)** : Mid-level clouds typically form between **2,000 to 6,000 meters** above the ground. They are **composed of water droplets and sometimes ice crystals** . Examples include altocumulus and altostratus clouds.
- **Low-Level Clouds** : These clouds form at altitudes **below 2,000 meters** . They are **composed of water droplets** and are often associated with weather phenomena such as rain or drizzle. Examples include stratus, stratocumulus, and nimbostratus clouds.

- Additionally, **some clouds form vertically through the troposphere** , extending through multiple altitude levels:
  - **Cumulonimbus Clouds** : These towering clouds extend **vertically through the troposphere** and are associated with **thunderstorms, heavy rain, lightning, and sometimes hail and tornadoes.**

⚡ **Quick revise** Moisture, clouds, precipitation, storms Fold ▲

☁️ **Clouds by height: High (>6,000 m) = Cirrus** – wispy ice "chicken feathers" · **Mid (2,000–6,000 m) = Alto-** · **Low (<2,000 m) = Stratus, stratocumulus, nimbostratus** · **Cumulonimbus** = vertical tower (thunder, hail, tornado). **Clouds & climate (PYQ 2022, ans D):** *high thin clouds trap* heat; *low thick clouds reflect* – so both statements (high cool, low warm) are wrong. **PYQ 2019 (ans D):** cirrus thinning and sulphate aerosols in the stratosphere are **geo-engineering to reduce global warming**.

Polar Stratospheric Clouds

Polar stratospheric clouds (PSCs) **form in the stratosphere at altitudes of 15-25 kilometers in polar regions** during the **winter months** , where temperatures drop below -78°C . Typically, clouds do not form in the stratosphere because it is usually too dry and stable; however, the extreme cold in polar regions during winter allows PSCs to form. Often called **nacreous or mother-of-pearl clouds** due to their iridescent colors, PSCs **appear bright and colorful during twilight.** PSCs **play a critical role in ozone depletion** by providing surfaces for chemical reactions that release ozone-depleting chlorine and bromine, significantly contributing to the formation of the ozone hole over Antarctica.

Cloud Seeding

Cloud seeding is a type of **weather modification** technique where certain substances are introduced into clouds to stimulate rainfall or snowfall. **Process**

- Substances like **Silver Iodide (AgI), Potassium Iodide, Sodium Chloride (common salt), or Dry Ice (solid CO_2)** are dispersed into clouds.
- These act as **Hygroscopic Nuclei** .
- Water vapor condenses/freezes around them, forming larger droplets or ice crystals.
- These eventually grow heavy enough to fall as rain or snow.

Methods of Delivery

- **Aircrafts** : release particles directly into target clouds.
- **Ground-based generators** : shoot particles upward using rockets or flares.

Applications

- **Enhancing rainfall** in drought-prone regions.
- **Increasing snowpack** for water supply in mountains.
- **Dispersing fog** at airports for visibility.
- **Hail suppression** in agriculture by reducing hailstone size.
- **Pollution Control Limitations & Concerns**
- Works only if **there is moisture in the air** .
- Effectiveness is variable and debated.
- Environmental concerns over **silver iodide accumulation** .
- Expensive and logistically complex.

⚡ **Quick revise** Polar stratospheric clouds Fold ▲

- **Polar stratospheric clouds:** 15–25 km, polar winter, below **−78 °C**; "nacreous / mother-of-pearl"; surfaces for **ozone destruction** (Antarctic ozone hole).
- **Cloud seeding:** silver iodide, potassium iodide, salt, dry ice → **hygroscopic nuclei**; by aircraft or ground generators; for rain,

snowpack, fog clearing, hail suppression; works only if moisture exists.

TYPES OF PRECIPITATION

- Precipitation **occurs when water droplets in clouds coalesce and grow large enough to overcome air resistance and fall to the ground** . Raindrops form through the process of collision and coalescence or the freezing of supercooled water droplets at high altitudes.



- **Convective (Convectional)**

Precipitation:

- Occurs when the air near the Earth's surface is heated, becomes buoyant, and rises rapidly, forming cumulus clouds that grow into **towering cumulonimbus clouds.**
- Intense and localized, often resulting in **heavy showers, thunderstorms, and short-lived periods of rainfall.**
- Latitudes: Primarily found in tropical regions **near the equator** (0° to 30° latitude).
- Precipitation: **Rainfall** is the most common type of precipitation associated with convective rainfall. It is these clouds that produce thunderstorms and can also produce **hail** in some cases.

- **Orographic Precipitation** :
  - Forms when **moist air is forced to rise over elevated terrain such as mountains or hills.**
  - **As the air rises** , it **cools and condenses** , leading to the formation of clouds and precipitation on the windward side of the mountain.
  - The **leeward side** of the mountain may experience a **rain shadow effect** , with drier conditions and less precipitation.
  - Latitudes: Orographic rainfall can occur at various latitudes but is **most common in regions with mountainous terrain.**
  - Precipitation: **Rainfall** is the typical outcome of orographic lifting. However, at higher elevations and colder temperatures, precipitation can fall as **snow** . In some cases it can also fall as **hail** .
- **Frontal (Cyclonic) Precipitation:**

- Occurs along the leading edge of a **frontal boundary between two contrasting air masses** , such as a cold front or warm front.
- **Warm air is lifted over denser, cooler air, leading to the formation of clouds and precipitation.**
- Can result in prolonged periods of rainfall over a large area in case of warm front, and short periods of intense rainfall in case of cold fronts.
- **Often associated with mid-latitude cyclones and frontal systems.**
- Latitudes: Frontal rainfall is common in mid-latitude regions (30° to 60° latitude), where contrasting air masses meet.
- Precipitation: **Rainfall** is the primary type of precipitation associated with frontal systems. However, in colder regions or during transitional seasons, precipitation can fall as **snow or sleet** .

- **Cyclonic Precipitation:**
  - This is **convectional/frontal rainfall on a large scale.**
  - Associated with the circulation of air around a low-pressure system, such as a tropical cyclone or extratropical cyclone.
  - **Intense and widespread,** with heavy rainfall occurring over large areas for an extended period.
  - Can result in flooding, landslides, and other severe weather impacts.
  - **Precipitation:**
    - **Tropical cyclones** typically produce **heavy rainfall** , sometimes accompanied by **thunderstorms and hail** .
    - In colder regions, **extratropical cyclones** can result in **snowfall.**

⚡ **Quick revise** Moisture, clouds, precipitation, storms Fold ▲

Precipitation	How	Where
Convectional	heated air rises → cumulonimbus; short, intense, thunder	tropics 0–30°
Orographic	air forced up mountains; **rain shadow** on lee side	mountains
Frontal	warm air over cooler (warm front = long steady; cold front = short intense)	mid-latitudes 30–60°
Cyclonic	convection/frontal on a large scale around a low	cyclones

THUNDERSTORMS:

- Thunderstorms are atmospheric disturbances characterized by the presence of **thunder and lightning,** often accompanied by **heavy rain, strong winds, and sometimes hail.**
- **Formation:**
  - Thunderstorms typically **form in warm, moist air masses** where there is instability and upward vertical motion, of air. This **upward motion, i.e. convection (updraft)** leads to the **development of towering cumulonimbus clouds.**



- **Stages:** Thunderstorms go through three main stages: the cumulus stage, the mature stage, and the dissipating stage.
  - During the **cumulus stage** , **warm, moist air rises** and **forms cumulus clouds** .
  - In the **mature stage** , updrafts and downdrafts within the **cumulonimbus cloud** lead to the development of **lightning, thunder, heavy rain, and sometimes hail** .
  - Finally, in the **dissipating stage,** the **storm weakens as downdrafts dominate** and the supply of warm, moist air diminishes.
- The cumulo nimbus clouds have a large amount of rain drops and ice crystals these particles constantly collide with each other and get charged the lighter positive charges accumulated at the top of clouds while heavier negative charges accumulate at the cloud bottom due to the accumulation of charges at cloud bottom a potential difference (voltage) appears to get created between the ground and the cloud bottom. Initially the air between the cloud and the ground act as an insulator, preventing the passage of any current. However as the potential difference between the cloud and the grounds keeps on increasing the current starts to move through the air exciting the nitrogen and oxygen molecules thus creating a bluish/ purpulish glow called lightning full stop the sound associated with this lightning discharge is called as thunder. Apart from ground and clouds the lightning discharge can also occur:
  - Within a cloud (top and bottom)



  - Between the clouds
- The taller objects on the ground (building/trees etc) have a high probability of experiencing a lightning event, as they are closer to the cloud bottom. To prevent lightning from destroying the electrical system of a building and endangering the life of the people inside, electric conductors are installed at the top of buildings.

- **Clouds in Thunderstorms:**
  - The **main cloud** type associated with thunderstorms is the **cumulonimbus** cloud, which is tall and vertically developed. These clouds can reach heights of up to 10 miles (16 kilometers) or more and are characterized by their anvil-shaped tops. **Other cloud** types, such as **cumulus clouds and stratocumulus clouds, may also be present** during the early stages of thunderstorm development.
- **Hazards:**
  - Thunderstorms can pose various hazards, including **lightning strikes** , which can cause fires and electrical damage, **heavy rainfall** leading to flash flooding, **strong winds** capable of causing property damage and power outages, and **hail** , which can damage crops, vehicles, and buildings.

⚡ **Quick revise** Thunderstorm Fold ▲

Thunderstorm: 3 stages – **cumulus → mature → dissipating**. Ice/water collisions give **lighter + charge at the top, heavier – charge at the bottom** → lightning (air excited N_2/O_2), thunder; lightning conductors on tall buildings. **Tornado:** rotating funnel from a thunderstorm; >300 km/h; **Tornado Alley** (Texas → Nebraska) where warm moist Gulf air meets cold dry Canadian air and warm dry desert air → **supercells**. **Waterspouts:** over water; fair-weather vs tornadic.

TORNADO

- A tornado is a **violently rotating column of air** that **extends from a thunderstorm to the ground** .
- It's formed when warm, moist air rises rapidly and meets cooler air, creating a strong updraft. This updraft causes the air to spin, and as it spins faster, it becomes more focused and descends towards the ground. If the rotating air column touches the ground, a tornado is formed.
- **Formation:**
  - **Thunderstorm Development:** Tornadoes form within powerful thunderstorms, often associated with strong updrafts of warm, moist air.
  - **Wind Shear:** As the warm air rises, it encounters cooler air at higher altitudes. This creates wind shear, a difference in wind speed and direction at different levels in the atmosphere.
  - **Rotation:** The **wind shear causes the rising air column to begin rotating** . This rotation can be further **intensified by the Coriolis force** .
  - **Condensation Funnel:** As the air column rotates faster and becomes more focused, it cools and condenses, forming a visible funnel cloud. This funnel can be various colors, including white, gray, or even black depending on the amount of debris it picks up.
- **Tornado Characteristics:**
  - **Size:** Tornadoes can **vary greatly** in size, ranging from a few feet wide to over a mile.

- **Windspeed:** Wind speeds in tornadoes **can reach** incredible speeds, **exceeding 300 kmph in the strongest ones.**
- **Duration:** Most tornadoes are **short-lived** , lasting only minutes to tens of minutes. However, some can persist for longer periods.
- **Damage:** Tornadoes can cause significant damage due to their **high wind speeds** , lifting and throwing debris, and even causing structural damage to buildings.

- Tornadoes can be incredibly destructive, with wind speeds reaching over 300 miles per hour. They are **most common in the central United States** , particularly in an area known as **Tornado Alley,** which stretches from Texas to Oklahoma, Kansas, and Nebraska. However, they can occur in many parts of the world, including Europe, Asia, Africa, and Australia.
- **Why the majority in "Tornado Alley"?**
  - The majority of tornadoes occur in Tornado Alley in the USA due to the **unique convergence of different air masses.**
  - **Warm, moist air** from the **Gulf of Mexico** meets **cold, dry air** from **Canada (Rockies)** and **warm, dry air** from the **southwestern deserts** .
  - This **convergence creates instability,** which, combined with strong wind shear (changes in wind speed and direction with height), leads to the **formation of powerful thunderstorms called supercells** . These supercells are the primary drivers for the high frequency of tornadoes in Tornado Alley.
  - The **geography and flat terrain of the region also help** funnel and intensify these weather patterns, making conditions ideal for tornado formation.

WATER SPOUTS

- A swirling column of air and water droplets that **forms over a body of water** , typically resembling a tornado.

- It occurs when **warm, moist air rises rapidly and meets cooler air above** , causing the air to spin and create a vortex.
- Usually **less intense than tornadoes** and are most commonly seen over warm ocean or lake waters.
- There are two **types: fair-weather** waterspouts, which form in calm conditions and don't usually cause much damage, and **tornadic waterspouts** , which are more intense and can be dangerous.

⚡ **Quick revise** Humidity Fold ▲

- **Humidity:** absolute (g/m³) · relative (% of what air can hold) · specific (g per kg). **Dew point** = saturation temperature. **Dew** forms when a surface falls below dew point; **no dew on cloudy nights** because clouds send the Earth's radiation back (PYQ 2019, ans B). **Frost** = deposition (vapour → ice). **Fog** = near-surface cloud. **Smog** = smoke + fog (sulphurous and photochemical). **Haze** = dry particles.
- **Atmospheric rivers:** narrow moisture corridors (can carry more water than the Amazon) pushed by westerlies/jet streams to mid-latitudes → heavy rain, floods.

Water vapour (PYQ 2024, ans a): amount decreases with altitude; it is *not* maximum at the poles.

11 · PYQ

Q.1) The seasonal reversal of winds is the typical characteristic of - (2014)

(a) Equatorial climate

(b) Mediterranean climate

(c) Monsoon climate

(d) All of the above climates

Answer: C

- Monsoon is traditionally defined as a seasonal reversing wind accompanied by corresponding changes in precipitation but is now used to describe seasonal changes in atmospheric circulation and precipitation associated with the asymmetric heating of land and sea.
- A monsoon is a seasonal reversal in wind patterns over a region. The word “monsoon” comes from the Arabic word mausim, meaning “season.” The seasonal wind shift is usually accompanied by a dramatic change in precipitation.
- Winds that blow for approximately six months from the northeast and six months from the southwest. The most prominent monsoons occur in South Asia, Africa, Australia, and the Pacific coast of Central America.

Q.2) Consider the following statements

1. The winds that blow between 30 degrees N and 60 degrees S latitudes throughout the year are known as westerlies.

2. The moist air masses that cause winter rains in the North-Western region of India are part of westerlies.

Which of the statements given above is/are correct? (2015)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: B

- The westerlies, anti-trades, or prevailing westerlies, are prevailing winds from the west toward the east in the middle latitudes between 30 and 60 degrees latitude.
- They originate from the high-pressure areas in the horse latitudes and trend towards the poles and steer extratropical cyclones in this general manner.
- Tropical cyclones which cross the subtropical ridge axis into the westerlies recurve due to the increased westerly flow.
- Westerly disturbances originate in the Mediterranean sea, this sea is surrounded on all sides by high mountains and plateau, as the result high pressure is developed there in the month of September- October, while low pressure still developed over the Mediterranean water body.
- This disturbance is dragged to India or brought to India by the Jet stream (westerly winds (flowing west to east)).
- This is the reason that J&K and NW part of India receives highest rainfall from westerly disturbances.

Q.3) In the context of which of the following do some scientists suggest the use of cirrus cloud thinning technique and the injection of sulphate aerosols into the stratosphere? (2019)

(a) Creating the artificial rains in some regions

(b) Reducing the frequency and intensity of tropical cyclones

(c) Reducing the adverse effects of solar wind on the Earth

(d) Reducing global warming

Answer: D

- Cirrus clouds do not reflect a lot of solar radiation back into space, but because they form at high altitudes and cold temperatures, they trap long-wave radiation and have a climate impact similar to greenhouse gases.
- Thus Thinning cirrus clouds would reduce the absorption of infrared radiation and proposed a form of climate engineering to reduce global warming.
- Another proposed Geo-engineering technique involves injecting reflective sulphate aerosol particles into Earth’s lower stratosphere to cast a small proportion of the inbound sunlight back into space and cool the planet off.
- Thus both are climate engineering/ Geo-engineering techniques to reduce global warming.

Q.4) Why are dewdrops not formed on a cloudy night? (2019)

(a) Clouds absorb the radiation released from the Earth’s surface.

(b) Clouds reflect back the Earth’s radiation.

(c) The Earth’s surface would have low temperature on cloudy nights.

(d) Clouds deflect the blowing wind to ground level.

Answer: B

- Less dew is formed on cloudy nights because radiation during cloudy nights occurs at a very slow pace.
- The radiation occurs very slowly because the clouds are thick during cloudy nights and trap the heat and water vapour inside the clouds.
- All this indicates why the level of dew is reduced during nights that are heavy with clouds.
- Dew is a natural form of water, formed as water vapour condenses. Dew, like the glistening drop on this grassy field in Anaconda, Montana, forms as water near the surface of the ground is cooled to its dew point, the temperature at which water vapour condenses. When the sky is cloudy, the trees and plants do not get cool in the night and hence there is less dew formation because on a cloudy night, the clouds send the heat (earth's radiation) back to the ground so the ground never gets cold enough for the dew to be formed.

Q.5) Consider the following statements:

1. Jet streams occur in the Northern Hemisphere only.

2. Only some cyclones develop an eye.

3. The temperature inside the eye of a cyclone is nearly 10°C lesser than that of the surroundings.

Which of the statements given above is/are correct? (2020)

(a) 1 only

(b) 2 and 3 only

(c) 2 only

(d) 1 and 3 only

Answer: C

- Jet streams occur in both the Northern and Southern Hemispheres. The actual appearance of jet streams result from the complex interaction between many variables - such as the location of high and low pressure systems, warm and cold air, and seasonal changes. Statement 1 is incorrect.
- The warmest part of Erin's eye was almost 21 degrees (Fahrenheit) warmer than the surrounding air, a dramatic difference from the air around it. Above 7.5 miles high, the eye's temperature dropped quickly to the same temperature as the air outside the eye. Statement 3 is incorrect.

Q.6) Consider the following statements:

1. High clouds primarily reflect solar radiation and cool the surface of the Earth.

2. Low clouds have a high absorption of infrared radiation emanating from the Earth's surface and thus cause warming effect.

Which of the statements given above is/are correct? (2022)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: D

- High clouds are often thin and do not reflect very much. They let lots of the Sun's warmth in. They radiate less energy into space than the lower, warmer clouds.

Therefore, high clouds work to "trap" more energy than the low clouds.

- Low clouds are often quite thick and reflect lots of sunlight back to space. Low clouds are excellent reflectors. But, they don't stop the longwave energy from escaping to space. Therefore, low clouds help to cool the Earth.

Q.7) With reference to “water vapour”, which of the following statements is/are correct? (2024)

1. It is a gas, the amount of which decreases with altitude.

2. Its percentage is maximum at the poles.

Select the answer using the code given below:

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer :

(a) 1 only

12 · AIR MASSES

- When the air remains over a homogenous area for a sufficiently longer time, it acquires the characteristics of the area.
- This **air parcel with distinctive homogeneous characteristics in terms of temperature and humidity** is called an air mass.
- Due to these homogenous conditions, they **lack any vertical movement of winds within themselves.**
- The homogenous regions can be the vast ocean surface or vast plains and plateaus.
- Air masses form an integral part of the global planetary wind system. Therefore, they are associated with one or other wind belts.

- They **extend from the surface to the lower stratosphere** and are across thousands of kilometers. This is why the **Jet streams can carry them** from their source location to other places.
- **Source Region:**
  - They are **often the places where the wind movement is absent** so that the air can stay at the same place for a longer period of time, resulting in the uniform transfer of the land characteristics to the air.
  - That is why the **source regions are High-Pressure regions that experience atmospheric stability.**
  - The main source regions are the **high-pressure belts in the subtropics (giving rise to tropical air masses) and around the poles (the source for polar air masses):**
    - Warm tropical and subtropical oceans
    - The subtropical hot deserts
    - The relatively cold high-latitude oceans
    - The very cold snow-covered continents in high latitudes
    - Permanently ice-covered continents in the Arctic and Antarctica
  - **Tropical air masses are warm, while polar air masses are cold.**
  - **Continental Air masses are dry while maritime ones are with moisture.**
  - **When an air mass moves away from a source region, the upper level maintains the physical characteristics for a longer period.**
  - This is possible because air masses are stable with stagnant air which does not facilitate convection.
  - Conduction and radiation in such stagnant air are not effective.
- **Major Air Masses:**
  - Maritime tropical (mT)
  - Continental tropical (cT)
  - Maritime polar (mP)
  - Continental polar (cP)
  - Continental Arctic (cA)
- **Climatic Impact of Air Masses:**
  - **Tropical (T):** Warm temperatures.
    - **Maritime Tropical (mT):** Warm, moist air from tropical oceans.
- Brings **heavy rainfall and thunderstorms to coastal regions and mid-latitudes** , and contributes to **tropical cyclones** and hurricanes.
- **Southeast Asia:** The maritime tropical air masses from the South China Sea cause significant rainfall during the wet season, contributing to the region's lush tropical climate.
- **Amazon Basin:** The mT air mass from the Atlantic Ocean brings abundant rainfall, supporting the Amazon rainforest ecosystem.
    - **Continental Tropical (cT):** Hot, dry air from desert regions.
- **Impact:** Causes **heatwaves and dry conditions in continental interiors** , influences summer climates in regions like the southwestern USA.

- **Sahara Desert:** The hot, dry cT air mass dominates, leading to extremely high temperatures and arid conditions.
- **Central Australia:** The interior experiences hot, dry conditions due to the influence of cT air masses, resulting in arid and semi-arid climates.
  - **Polar (P):** Cold temperatures, can vary in humidity.
    - **Maritime Polar (mP):** Cool, moist air from high-latitude oceans.
- **Brings cool, damp, and cloudy weather with precipitation to coastal and inland regions** , particularly in the winter.
- **Western Europe:** The mP air mass from the North Atlantic Ocean brings cool, moist conditions, resulting in frequent rain and cloudy weather, especially in the UK and Ireland.
- **Japan:** The Sea of Japan coast experiences cold, snowy winters due to mP air masses from the Sea of Okhotsk.
    - **Continental Polar (cP):** Cold, dry air from high-latitude land areas.
- Causes **cold, dry conditions in winter** , can lead to clear, cold weather and **influence winter storm tracks in mid-latitudes** .
- **Russia (Siberia):** cP air masses from the interior lead to extremely cold, dry winters with clear skies and very low temperatures.
- **Canada (Prairies):** The influence of cP air masses results in cold, dry winters with clear skies, contributing to the region's continental climate.
  - **Arctic (A):** Extremely cold, very dry.
    - **Continental Arctic (cA):** Very cold, dry air from Arctic regions.
- Brings **frigid temperatures and clear skies** , influences polar vortex events and can push cold air far into mid-latitudes, causing severe winter weather.
- **Northern Canada and Greenland:** cA air masses bring extremely cold, dry conditions, influencing the Arctic climate and contributing to polar vortex events that can push cold air southward.
- **Scandinavia:** Periodic invasions of cA air masses can cause severe cold spells and harsh winter weather.
  - **Equatorial (E):** Very warm, very humid.
    - **Maritime Equatorial (mE):** Extremely warm and moist air from equatorial oceans.
- Contributes to **heavy rainfall and thunderstorms in equatorial regions** , influences monsoon climates and enhances tropical cyclone development.
- **Indonesia and Malaysia:** The mE air mass brings high humidity and significant rainfall, supporting the region's tropical rainforest climate.
- **West Africa (Gulf of Guinea):** The mE air mass contributes to heavy rainfall and the humid tropical climate of the coastal regions.
  - Overall Impacts on Global Climate:
    - **Temperature Regulation:** Air masses redistribute heat globally, balancing temperature differences between the equator and poles.
    - **Precipitation Patterns:** Maritime air masses (mT, mP) bring moisture and precipitation, impacting rainfall patterns and water availability.

- **Weather Systems and Storms:** Interactions between different air masses lead to the formation of weather fronts, cyclones, and storms, influencing weather patterns and extremes.
- **Seasonal Variations:** The movement of air masses with seasons contributes to seasonal climate variations, such as monsoons and winter storms.

FRONTS

- These are the **3-dimensional boundary zones** , **between 2 air masses** of **contrasting characteristics** .
- They are typically **found in the mid-latitudinal regions** , and are uncommon in tropics and the polar regions.
- The process of formation of a front is known as **Frontogenesis** (the war/face-off begins), and **the dissipation of a front** is known as **Frontolysis** (one of the air masses wins, and the war is now over).
- The temperate cyclone owe their genesis to the process of frontogenesis.
- **Characteristics of Fronts:**
  - The **more the temperature contrast between the 2 air masses, the lesser the thickness** of the front formed.
  - With a sudden change in temperature through the front, there is a **change in pressure also.**
  - Sudden **wind shifts** too.
  - The zone experiences precipitation due to the **upliftment (gradual or forced) of the warm air.**
  - The **cold air masses and fronts moves faster** than the warm front as the compact and denser cold air can push the boundaries with more force.
- **Types of Fronts:**
  - **Stationary Front:**
    - Forms when two air masses meet but **neither advances significantly.**
    - **Clouds:** A mix of stratocumulus, nimbostratus, and altostratus.
    - Clouds and light precipitation can persist along a stationary front for an extended period.



- **Warm Front:**
  - Forms when a **warm air mass advancesinto the region of a cold air mass.**
  - Clouds: Stratus, altostratus, cirrostratus.
  - These clouds often produce **steady rain or drizzle** and can lead to overcast skies and fog.
- **Cold Front:**
  - **Forms when a cold air mass advances and displaces a warm air mass.**
  - Clouds: Cumulus, towering cumulonimbus.
  - These clouds often produce **thunderstorms, heavy rain, hail, and tornadoes.**
- **Occluded Front:**
  - Forms when a **fast-moving cold front overtakes a slower-moving warm front.**
  - Clouds: A mix of cumulus, stratocumulus, and nimbostratus.
  - Can produce a **variety of weather conditions** , including thunderstorms, rain, and cooler temperatures as warm air is lifted off the surface.
  - **Warm Occlusion:**

- Forms when a fast-moving cold front overtakes a slower-moving warm front, and **the air ahead of the warm front is cooler than the air behind the cold front.**
- Warm air is lifted off the surface, creating a mix of clouds, including cumulus, stratocumulus, and nimbostratus.
- Typically associated with moderate to heavy precipitation, including rain and sometimes thunderstorms.
- The warm sector ahead of the occluded front may experience warmer temperatures compared to the cold sector behind it.



- **Cold Occlusion:**

- Forms when a fast-moving cold front overtakes a slower-moving warm front, **but the air behind the cold front is colder than the air ahead of the warm front.**
- Cold air wedges underneath the warm air, lifting it off the surface.
- Clouds: A mix of stratocumulus, nimbostratus, and altostratus.
- Typically associated with widespread precipitation, including rain or snow, and cooler temperatures. A mix of cold & warm front weathers.
- The resulting weather can be like that of a warm occlusion but with colder temperatures and a greater likelihood of snow in the colder air mass.

⚡ Quick revise Air mass — Fold ▲

- **Air mass:** big air body that took on the character of its source (**high-pressure, calm regions**). Types: **mT, cT, mP, cP, cA** (+ equatorial). *Tropical = warm, polar = cold, continental = dry, maritime = moist.* Examples: mT → Amazon, SE Asia rain; cT → Sahara, central Australia; mP → UK, Ireland, Japan Sea coast snow; cP → Siberia, Canadian prairies; cA → N. Canada, Scandinavia cold spells.
- **Fronts:** boundary of two air masses; **frontogenesis** (formation) / **frontolysis** (death); mid-latitudes; bigger contrast = thinner front; **cold fronts move faster**.

⚡ Quick revise Air masses, fronts, cyclones — Fold ▲

> ⚔️ **Cold front = Crash** (cumulonimbus, thunder, hail) · **Warm front = Wet blanket** (stratus/cirrostratus, steady drizzle, fog) · **Stationary** = stalemate (light rain for days) · **Occluded** = cold overtakes warm (warm or cold occlusion).

13 · CYCLONES

- Cyclones are large-scale atmospheric systems characterized by **low-pressure centers and circulating winds that rotate counterclockwise in the Northern Hemisphere and clockwise in the Southern Hemisphere.**
- There are different types of cyclones, including tropical cyclones, extratropical cyclones, and subtropical cyclones.
- Tropical cyclones, also known as hurricanes or typhoons depending on the region, form over tropical or subtropical waters and are characterized by intense thunderstorms and strong winds.
- Extratropical cyclones form outside the tropics and are associated with frontal systems and changes in temperature gradients.

TROPICAL CYCLONES:

- **Location:**
  - Tropical cyclones primarily occur in **tropical and subtropical regions** over warm ocean waters, typically between **10° and 30° latitude** in both hemispheres.
  - They are known as **Cyclones in the Indian Ocean, Hurricanes in the Atlantic, Typhoons in the Western Pacific and South China Sea, and Willy-willies in Western Australia.**
- **Necessary Conditions:**
  - **Warm Ocean Waters:** Sea surface temperatures of at least 26.5°C (80°F) are necessary to fuel tropical cyclone development.
  - **Moisture:** High levels of atmospheric moisture are essential for sustaining convective activity and cloud formation.
  - **Low Vertical Wind Shear:** Minimal change in wind speed and direction with height allows for vertical development of the storm.
- **Formation:**
  - **A disturbance** , **such as a low pressure centre** like monsoon trough, **serves as the initial seed** for tropical cyclone development.
  - **If the atmospheric and oceanic conditions are favourable, the disturbance may undergo cyclogenesis** , the process by which a tropical cyclone forms. Warm, moist air rises rapidly, creating an area of low pressure at the surface. As the air converges and rises, it cools and condenses, releasing latent heat and further fuelling the convective activity.



- As the disturbance intensifies and becomes better organized, it may develop into a tropical depression, characterized by a closed circulation and maximum sustained winds of up to 38 miles per hour (62 kilometres per hour).
- **If the conditions persist, the depression continues to strengthen into a cyclone.**
- Once formed, tropical cyclones can track across the ocean and intensify, fuelled by warm ocean waters and favourable atmospheric conditions. They may reach their peak intensity at some point before eventually weakening as they move over cooler waters or make landfall.

- **Structure:**
  - **Eye:** A well-defined center of low pressure characterized by calm winds and clear skies.
  - **Eyewall:** A ring of intense thunderstorms surrounding the eye, where the strongest winds and heaviest rainfall occur.
  - **Rainbands:** Bands of clouds and thunderstorms extending outward from the center, bringing additional rainfall and gusty winds.
- **Types of Clouds Associated:**
  - **Cumulonimbus Clouds:** Towering clouds with strong updrafts and potential for severe weather, including heavy rainfall, lightning, and hail.
  - **Cirrus Clouds:** High-altitude clouds that may indicate the presence of upper-level outflow associated with the storm.
- **Impact:**
  - **Strong Winds:** Tropical cyclones can produce sustained winds of 64 knots (74 mph) or greater, capable of causing widespread damage to structures, vegetation, and infrastructure.
  - **Heavy Rainfall:** Intense rainfall can lead to flooding, landslides, and mudslides, especially in areas with poor drainage or in mountainous terrain.
  - **Storm Surge:** The onshore flow of water driven by strong winds can result in storm surges, causing coastal flooding and inundation of low-lying areas.
  - **Tornadoes:** Embedded within rainbands, tornadoes can form and cause additional localized damage.
  - **Economic and Humanitarian Impacts:** Tropical cyclones can result in loss of life, displacement of populations, disruption of essential services, and significant economic losses.

⚡ **Quick revise** Tropical vs temperate cyclones Fold ▲

Names: **Cyclone** (Indian Ocean) · **Hurricane** (Atlantic) · **Typhoon** (W. Pacific, S. China Sea) · **Willy-willy** (W. Australia).

TEMPERATE CYCLONES

- **Location:**
  - Temperate cyclones, also known as extratropical cyclones, occur in the mid-latitudes, typically between **30° and 60° latitude** in both hemispheres.



re contrast between polar and tropical air masses, such as North

  - They are most common **in regions with a strong temperature contrast between polar and tropical air masses** , such as North America, Europe, and Asia.
- **Necessary Conditions:**
  - **Temperature Contrast:** A significant temperature difference between polar and tropical air masses is essential for the development of temperate cyclones.
  - **Baroclinic Instability:** The interaction between warm and cold air masses along a frontal boundary creates instability, leading to the formation of cyclones.
- **Formation:**
  - **Temperate cyclones form along frontal boundaries,** where warm and cold air masses meet and create a zone of instability.
  - The process **begins with** the **development of a stationary front,** where warm air rises over denser, cooler air.
  - As the temperature gradient steepens and instability increases, a low-pressure system forms and begins to intensify.
- **Structure:**
  - **Warm and Cold Fronts:**



Temperate cyclones are associated with both warm and cold fronts, which mark the leading edges of advancing air masses.

- **Cyclonic Rotation:** In the **Northern Hemisphere** , **temperate cyclones rotate counterclockwise** around their centers, while in the **Southern Hemisphere** , they rotate **clockwise.**
- **Occluded Front:** As the cyclone matures, the cold front catches up to the warm front, resulting in an occluded front and the eventual dissipation of the cyclone.

- **Types of Clouds Associated:**
  - **Cumulonimbus Clouds:** Towering clouds associated with thunderstorms and heavy rainfall, especially along the cold front.
  - **Nimbostratus Clouds:** Layered clouds that produce steady rain or snow, often associated with the warm sector of the cyclone.
  - **Altostratus Clouds:** Mid-level clouds that may precede the warm front and bring overcast conditions and light precipitation.
- **Impact:**
  - **Precipitation:** Temperate cyclones can produce a variety of precipitation types, including rain, snow, sleet, and freezing rain, depending on temperature and atmospheric conditions.
  - **Strong Winds:** Gusty winds associated with the cyclone can cause damage to structures, trees, and power lines.
  - **Flooding:** Heavy rainfall from temperate cyclones can lead to flash flooding, river flooding, and landslides.
  - **Storm Surge:** Along coastal areas, the onshore flow of water driven by strong winds can result in storm surges and coastal flooding.

IMPACT OF CLIMATE CHANGE

Tropical Cyclones:

- Rising ocean temperatures due to climate change provide more energy for tropical cyclones, leading to more intense storms with higher wind speeds and heavier rainfall. There has been an increase in the proportion of severe (Category 4 & 5) cyclones, especially in regions like the Arabian Sea and the North Atlantic.

- Climate models show a decreased overall frequency of tropical cyclones globally, with some regions like the North Atlantic bucking the trend by becoming slightly more active.
- The cyclones are encroaching even higher latitudes (poleward shift), and their forward motion has slowed down, which can prolong rainfall and increase flood risks.
- Climate change leads to an increase in rapid intensification events (storms that quickly strengthen), making forecasts and disaster responses more difficult.
- Sea level rise exacerbates storm surges from cyclones, increasing coastal flooding hazards.

Temperate Cyclones:

- Temperate cyclones (also called extratropical cyclones or mid-latitude storms) may experience shifts in their tracks due to climate change, often moving poleward (lack of availability of cold air masses at lower latitudes), which could alter weather patterns in affected regions.
- Warming impacts the temperature gradients and jet stream patterns that drive temperate cyclones, potentially modifying their intensity, frequency, and seasonal behavior.
- Increased atmospheric moisture from climate change can lead to more intense precipitation events associated with these cyclones, enhancing the risks of flooding, landslides, and transport disruption. There is evidence of excess rainfall and unseasonal weather in areas affected by temperate cyclones, which can impact crops and infrastructure.

TROPICAL v/s TEMPERATE CYCLONES

	Tropical Cyclone	Tempera
Origin	Thermal Origin	Dynamic
Latitude	10 - 30 degree N and S of equator	35 - 65 More hemisph tempera
Frontal system	Absent	Cyclone
Formation	Only on warm seas	Can forn
Season	Seas	Irregular more in
Size	100 – 500 kms in diameter. Varies with the strength of the cyclone	Typical diamete region.
Rainfall	Heavy but does not last beyond a few hours.	Rainfall i days, so
Wind Velocity and destruction	100 – 250 kmph, and 200 – 1200 kmph in upper troposphere. Greater destruction due to winds, storm surges and torrential rains	Compar – 150 k winds b flooding
Lifetime	Doesn't last for more than a week	Last for
Influence on India	Both coasts affected. But east coast is the hot spot	'Westerr

⚡ **Quick revise** Tropical vs temperate cyclones Fold ▲

	Tropical	Temperate (extra-tropical)
Origin	**Thermal**	**Dynamic** (frontal)
Latitude	10–30°	35–65° (more pronounced in N. Hemisphere – bigger temperature contrast)

	Tropical	Temperate (extra-tropical)
Forms over	warm seas only	land **and** sea
Size	100–500 km	300–2,000 km
Rain	heavy, but not beyond a few hours	slow, for many days, even weeks
Wind	100–250 km/h; more damage from wind, surge, torrential rain	30–150 km/h; more damage from flooding
Life	not more than a week	2–3 weeks
Fronts	absent	created by fronts
India	both coasts hit, **east coast hot-spot**	**Western Disturbances**

Needs: sea-surface ≥ 26.5 °C (the PYQ note says >27 °C), moisture, **low wind shear**, a pre-existing disturbance, Coriolis. *Depression ≤ 62 km/h → cyclone ≥ 64 knots (74 mph).* Structure: **eye** (calm, clear), **eyewall** (worst winds/rain), **rainbands**. Impacts: wind, rain, **storm surge**, tornadoes.

PYQ 2015 (ans B): no cyclones in the S. Atlantic / SE Pacific because the **ITCZ seldom occurs there** (also strong wind shear, cooler SST, no spin to start). **PYQ 2020 (ans C):** *only some* cyclones develop an eye; the eye is **not** 10 °C cooler.

Climate change: warmer seas → **more intense** storms, more Cat 4–5 (Arabian Sea, N. Atlantic), fewer overall, **poleward shift**, slower movement, **rapid intensification**, sea-level rise worsens surge. Temperate cyclones also shift poleward and bring heavier rain.

14 · PYQ

Q.1) In the South Atlantic and South Eastern Pacific regions in tropical latitudes, cyclones do not originate. What is the reason? (2015)

(a) Sea Surface temperature are low

(b) Inter-Tropical Convergence Zone seldom occurs

(c) Coriolis force is too weak

(d) The absence of land in those regions

Answer: B The most proximate reasons for the lack of cyclone formation in the South Atlantic are sea surface temperatures that tend to run a shade cooler than ideal for tropical cyclone formation even in the southern summer, climatologically high values of vertical wind shear across that basin throughout the year, and a lack of pre-existing centres of rotation (vorticity) in that area.

- Tropical cyclones are the zones of low pressure systems around which there is closed air circulation in counterclockwise direction(Northern hemisphere) and circular direction(Southern hemisphere).
- Important condition for the origin of tropical cyclones:
  - Sea Surface temperature should be greater than 27 degree C.
  - Pre existing weak tropical disturbances
  - Continuous supply of energy.
  - Coriolis force.
  - Weak wind shear
- Strong Wind shear present at South Atlantic removes the heat and moisture from the place where it is required to drive the heat engine of cyclone i.e near the centre.
- Without an Inter-tropical Convergence Zone (ITCZ) to provide synoptic vorticity and convergence (i.e large scale spin and thunderstorm activity) as well as having strong wind shear, it becomes very difficult to nearly impossible to have genesis of tropical cyclones.

15 · Koppen's Classification

Table 12.2 : Climatic Types According to Koeppen

Group	*Type*	*Letter Code*	*Characteristics*
A-Tropical Humid Climate	Tropical wet	Af	No dry season
	Tropical monsoon	Am	Monsoonal, short dry season
	Tropical wet and dry	Aw	Winter dry season
B-Dry Climate	Subtropical steppe	BSh	Low-latitude semi arid or dry
	Subtropical desert	BWh	Low-latitude arid or dry
	Mid-latitude steppe	BSk	Mid-latitude semi arid or dry
	Mid-latitude desert	BWk	Mid-latitude arid or dry
C-Warm temperate (Mid-latitude) Climates	Humid subtropical	Cfa	No dry season, warm summer
	Mediterranean	Cs	Dry hot summer
	Marine west coast	Cfb	No dry season, warm and cool summer
D-Cold Snow-forest Climates	Humid continental	Df	No dry season, severe winter
	Subarctic	Dw	Winter dry and very severe
E-Cold Climates	Tundra	ET	No true summer
	Polar ice cap	EF	Perennial ice
H-Highland	Highland	H	Highland with snow cover

⚡ **Quick revise** Climatic regions (Köppen) Fold ▲

🔤 **Köppen letters: A**–tropical humid · **B**–dry · **C**–warm temperate · **D**–cold snow-forest · **E**–polar · **H**–highland. Second letters: **f** no dry season · **w** winter dry · **s** dry summer · **m** monsoon. *"All Big Cities Deserve Extra Halwa."*

Handout alert: the Köppen group table labels **Df = humid continental and Dw = subarctic**, but the regional table labels **Taiga = Df and Manchurian = Dw**. Remember the **names and places**, not the D-

codes. Numbers in the regional table (e.g. "150–1000 cm" for Af) are printed as-is and look scrambled – don't memorise them.

16 · CLIMATIC REGIONS OF THE WORLD

Climate	Koppen Name	Temp.	Rainfall	Location
Hot, Wet Equatorial	Af	24–27 degree. Low Annual And Diurnal Range	150–1000 cm. Avg. annual.Daily Rainfall (4 pm rains).Convectional. Norainlessmonth. Equinox Maxima; Solstice minima. High Relative Humidity (~80%)	10 degree north & South of Equator. E.g.Amazon Basin,Congo Basin,Indonesia, PNG.
Savannah (Sudan Type)	Aw	High diurnal range (in summers, up to 38 in	Hot rainy summer, dry cold winters. 100–150	10–30 degrees continent interiorsE.g.

MY NOTES

		day, 10 in night). About 10 degrees annual range Highest temperatures in April (NH) or October (SH).	cm average annual	South Sudan, Venezuela, Brazil, North Australia, etc.
Tropical Monsoon	Am	3 distinct seasons. Summer, Winter, Rainy. High annual range of temperature	Lot of rain in short period of time. 200-250 cm	10-30 degree; East Continental Margins E.g. India, Vietnam, parts of Australia, Central America.
Hot Desert	BWh	Very high temperature. High diurnal and annual range of temperature. No cold season, but frost may occur on winter	Small amount of convective rain. Not more than 25-50 cm	West margins of Tropical landmass. Sub-tropical HP belt, & Impact of Cold Currents. E.g. Sahara,

		nights.		Kalahari, Mojave, Atacama deserts.

MY NOTES

Cold Deserts	BWk	Wide range of annual temperature (more than hot deserts). Annual avg <18 degrees C. Severe winters. Flooding during early summers due to ice thawing.	Lesser than hot deserts. 0- 30 cm, due to occasional depressions. (Gurbantunggut desert: 0 rain).	Sub-Tropical Continental interiors E.g. Gobi desert, Siberia, Patagonia
Steppe Climate	BS	Warm summers (18-20 degree), Cold winters (can be sub-zero). High annual range. However, not that extreme in the Southern Hemisphere (narrow land; impact of oceans)	25-75 cm; decreases with continentality. Summer maxima. Effect of Temperate cyclones too. More rain in Southern Hemisphere due to maritime influence	45-60 degree mid continental regions. Westerly influence E.g. Prairies (impact of Chinook), Steppes, Downs, etc.
Mediterranean	Cs	Long hot summers. (>22 degree C). Cool Winters (0 degree)	Winter rainfall (onshore Westerlies).	30-45 degree; West margin; Medi. shoreline;

		degree). Impact of many local winds (Sirocco, Mistral, Bora)	Summer drought(off shore trade) Shift of pressure	shoreline, Central California; central Chile, SW South Af, SW Aust.

MY NOTES

			belts. 50-75 cm	
Wet West Marine (British Type)	Cf	5-15 degree C; Warmer in Europe (NAD & Norwegian current); cold in rest regions. Low annual range of temperature (cool throughout the year)	Onshore westerlies throughout year. 125cm. Wettest Temperate climate. More rain on western margins.	45-60 degree West margins. North West Europe, South West S.America, parts of New Zealand, Canada
China Type/ Gulf type/ Natal type	Cw (also known as Cfa, but Cw also accepted due to dry winters)	Warm Moist summer; Cool dry winter. Heavy maritime influence. **China:** high annual range **Gulf:** narrow range **Natal**: small annual range	**China type:** SE monsoons in summers; minor snowfall in winters; Typhoons (summer maxima) **Gulf type:** No distinct dry period (summer maxima); Hurrican	Temperate East margins (30-45 degrees). **China type:** Central & North China; South Japan **Gulf type:** Gulf of Mexico **Natal type:** Southern hemisphere (Uruguay-Paraguay; parts of Brazil & Argentina; East Australia;

			Hurricanes & tornadoes. **Natal type:** even rain by	Australia, parts of South Africa.

MY NOTES

			onshore winds throughout the year.	
Taiga/ Siberian Climate	Df	Exceptionally cold winters (- 30 to -40 deg. C); cool brief summers. V. High temp range	30cm. Mostly convectional and frontal. Summer Maxima	50/60-70 degree latitude. Northern Canada and Eurasia. Continuous strip
Manchurian/ Laurentian Climate	Dw	Cold dry winters; wet warm summers	Up to 100 cm (More in N.Am than Asia due to Great lakes); Onshore maritime in summer; During winter off-shore winds.	Temperate East Margin. Only in Northern Hemisphere Manchuria, Japan, Koreas; NE North America
Polar	Sub-Polar (ET); Frosted (EF)	ET: Summer temp. Less than 10 degrees Frost can	Mostly Snowfall; In summers Rain & sleet; Winter	Greenland, Northern Canada & Alaska; Antarctica

		occur thorough ut the year.	maxima due to Extra-Tropical Cyclones ;	

Name of Shifting Cultivation/	Location/Country
Humah/ Ladang	Java and Indonesia
Tamrai	Thailand
Chena	Sri Lanka
Taungya	Myanmar
Caingin	Philippines
Chetemini	Uganda, Zambia and Zimbabwe
Roka	Brazil
Konuko	Venezuela
Milya	Mexico and Central America
Echalin	Guadeloupe
Milpa	Yucatan and Guatemala
Comile	Mexico
Logan	Western Africa
Fang	Equatorial African Countries
Masole	Congo (Zaire river Valley)
Tavi	Madagascar
Ray	Vietnam

Famous Grasslands in the world

List of all Important Grassland around the world:

Name of the Grassland	Location
Prairie Grassland	North America (USA Canada)
Pampas Grassland	South American (Argentina, Uruguay Brazil)
Downs Grassland	Australia
savanna Grassland	Central Africa
Campos Grassland	Brazil
Pustaz Grassland	Hungary and it's surrounding regions
Veldt Grassland	Southern Africa (South Africa, Lesotho, Zimbabwe Botswana)
Steppe Grassland	European Russia, Siberia and Asian Russia
Manchurian Grassland	Mongolia, China
Canterbury grassland	New Zealand

PYQs

rainforest is removed, it does not regenerate quickly as compared to a s forest. This is because – (2011)

nforest is deficient in nutrients
f the trees in a rainforest have poor viability
st species are slow-growing
s invade the fertile soil of rainforest

⚡ Quick revise Climatic regions (Köppen)

Fold ▲

Climate (code)	Where	Look-for hooks (as printed)
Equatorial (Af)	10°N–S: Amazon, Congo, Indonesia, PNG	24–27 °C, tiny annual range; convectional **4 pm rain**, no dry month; **evergreen selvas**, hardwoods (mahogany, rosewood, ebony), almost no undergrowth, **leached soil**; hunters-gatherers, slash-and-burn
Savannah (Aw)	10–30°, continental interiors: Sudan, Venezuela, Brazil, N. Australia	hot rainy summer, dry winter; tall grass (elephant grass), baobab; **Masai** pastoralism; "Big Game country"
Tropical monsoon (Am)	10–30°, E. continental margins: India, Vietnam	3 seasons; 200–250 cm; teak, sal; intensive paddy

Climate (code)	Where	Look-for hooks (as printed)
Hot desert (BWh)	W. margins of tropical land, **subtropical high**, cold currents: Sahara, Kalahari, Mojave, Atacama	xerophytes (waxy/thorny leaves, deep roots, succulent stems), date palms at oases; Bedouin, Tuareg, Bushmen
Cold desert (BWk)	continental interiors: Gobi, Patagonia	wide annual range, cold winters; bunchgrasses
Steppe (BS)	45–60°, mid-continental: Prairies, Steppes, Downs	warm summers, cold winters; **Chernozem**; wheat/maize, ranching
Mediterranean (Cs)	30–45°, west margins: California, Chile, SW S. Africa, SW Australia	hot dry summer, **winter rain** (westerlies); orchards (citrus, olives), vineyards; cork oak, eucalyptus
Marine West Coast (Cfb, "British type")	45–60°, west margins: NW Europe, SW S. America, NZ	mild, rain all year, small annual range; temperate deciduous forest; dairy
China type (Cw / Cfa; Gulf & Natal types)	30–45°, east margins: SE China, Japan	warm moist summer, cool dry winter; typhoons; most productive mid-latitude farming
Taiga / Siberian (Df)	50/60–70°: N. Canada, Eurasia	−30 to −40 °C winters; **evergreen conifers** (softwood, pure stands); fur farms, lumbering
Manchurian / Laurentian (Dw)	E. margins, N. Hemisphere only: Manchuria, Japan, NE America	cold dry winters, wet warm summers; cool temperate forest
Polar (ET, EF)	Greenland, N. Canada, Alaska, Antarctica	summer < 10 °C; mosses, lichens; Eskimos, Lapps, Yakuts

17 · PYQs

Q.1) If a tropical rainforest is removed, it does not regenerate quickly as compared to a tropical deciduous forest. This is because - (2011)

(a) the soil of rainforest is deficient in nutrients

(b) propagules of the trees in a rainforest have poor viability

(c) the rain forest species are slow-growing

(d) exotic species invade the fertile soil of rainforest

Answer: A Due to heavy rainfall and weathering, most of the nutrients have been washed out of the soil.

- Only a thin layer of decaying organic matter is found, unlike in temperate deciduous forests.That's why the soil of most of the tropical rainforest is poor in nutrients.
- The high temperature and moisture of tropical rainforests cause dead organic matter in the soil to decompose more quickly than in other climates, thus releasing and losing its nutrients rapidly. The high volume of rain in tropical rainforests washes nutrients out of the soil more quickly than in other climates.
- Why are rainforest soils low in nutrients? One reason the rain forest soil is so poor is that most of the nutrients are stored in the plants themselves. In any forest, dead organic matter falls to the ground, providing valuable nutrients for new growth. In cooler or drier climates, the nutrients build up in the soil.

Q.2) What could be the main reason/reasons for the formation of the African and Eurasian desert belt?

1. It is located in the sub-tropical high-pressure cells.

2. It is under the influence of warm ocean currents.

Which of the statements given above is/are correct in this context? (2011)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: A

- Hot deserts are located between 20°30° latitude in both the hemisphere on the western coast of the continents.
- They are also called Tropical Deserts.
- They are different from the midlatitude deserts and cold deserts, hot deserts are found in High-pressure belts along the Tropics in the trade winds region.
- Distribution Of Hot Deserts:
  - Asia Thar desert and the Arabian Desert;
  - Africa The Sahara and the Kalahari;
  - N. America Lower California and Arizona states of the USA;
  - S. AmericaAtacama desert;
  - Australia The Great Australian desert.
  - The Sahara desert of Africa is the biggest hot desert in the world.
  - The Hot desert regions are located in the subtropical high-pressure belts, which are not favourable regions for rainfall.
  - There are three main causes of deserts:
    - Most of the world's deserts are made by the equatorial wind belts.
    - At the equator, warm air is always rising.
    - This creates wind belts that blow north and south away from the equator.

- When air descends at the tropics of cancer and Capricorn it is too dry for clouds to form.
- No clouds mean no rain producing Deserts. E.g. the Sahara desert
- Rainshadow also produces Deserts.
- Any moisture in the air will fall as air passes high mountain ranges, so land beyond the mountains receives little or no rain.
- The Gobi desert is produced by Rainshadow.
- Cold Currents also produce deserts.
- The southwestern coast of Africa and South America are swept by cold currents upwelling from the ocean floor. These currents cool the air that passes over them, causing the water in the air to fall as rain before it reaches the land. The Namib and Atacama desert have formed because of these cold currents.
- The Eurasian Belt is related to cold currents NOT warm current, so statement 2 is incorrect.

Q.3) Which one of the following is the characteristic climate of the Tropical Savannah Region? (2012)

(a) Rainfall throughout the year

(b) Rainfall in winter only

(c) An extremely short dry season

(d) A definite dry and wet season

Answer: D The tropical savanna climate is also called the tropical wet and dry climate. It is the climate experienced in savanna or tropical grassland regions of the world.

- These places are located near the equator, and they lie between the Southern and the Northern Tropics.
- The climate dominates many parts of the African continent, the northern region of South America, and parts of Asia such as India.
- The tropical savanna climate has alternating dry and wet seasons, hence its name.
- It shares some similar characteristics with the tropical monsoon climate, but it receives less annual rainfall as compared to the tropical monsoon climate.
- Tropical savanna climates are relatively hot since they lie within the tropical latitudes.
- Precipitation in the tropical savanna climate is mainly in the form of rainfall. On average, this climatic zone receives between 800 and 1,600 millimetres of rain annually.

Q.4) "Each day is more or less the same, the morning is dear and bright with a sea breeze; as the Sun climbs high in the sky, heat mounts up, dark clouds form, then rain comes with thunder and lightning. But rain is soon over".

Which of the following regions is described in the above passage? (2015)

(a) Savannah

(b) Equatorial

(c) Monsoon

(d) Mediterranean

Answer: B Regions with this climate experience high temperatures all year round.

- The average monthly temperatures are about 26 – 28 degrees Celsius.
- The annual temperature range (the difference between the average temperature of the hottest and coldest months) is very small.
- The annual temperature range may be as low as 3 degrees Celsius.
- The diurnal or daily temperature range (the difference between the highest temperature in the day and the lowest temperature at night) is usually greater. Humidity is usually very high.
- Another major characteristic of this climate is high rainfall.
- These regions usually experience 2000 mm of rainfall or more in a year.
- Rainfall is high for most of the year.
- Many equatorial regions are affected by the ITCZ.
- As the ITCZ passes over these areas it brings heavy rainfall and thunderstorms.
- In some areas, the ITCZ causes two periods of very heavy rainfall every year.
- One occurs when the ITCZ crosses these areas on its way north and another occurs when it crosses these areas again on its way south.
- The climograph below shows the rainfall and temperature pattern of an area which experiences an equatorial climate.

Q.5) Which of the following leaf modifications occur(s) in the desert areas to inhibit water loss?

1. Hard and waxy leaves

2. Tiny leaves

3. Thorns instead of leaves

Select the correct answer using the code given below: (2018)

(a) 2 and 3 only

(b) 2 only

(c) 3 only

(d) 1, 2 and 3

Answer: D Leaf modifications of Desert Plant to inhibit water loss for adaptations and survival of Desert Plant-

- Wax coatings on leaves prevent water loss through transpiration, which in the hot desert can cause loss of water from both the surface and the inside of the leaves. Leaves are also smaller on desert plants, further reducing the possibility for water loss.
- Deciduous plants in desert ecosystems have adapted through the activity of their leaves. Leaves on these plants are typically smaller and coated with wax to prevent evapo-transpiration.
- Plants are equipped with fleshy leaves that contain much of their water supply. Because of their moist inner bodies, these plants are called succulents. They typically feel

spongy and when cut open are filled with a pulpy flesh, protected by a waxy outer layer.

- Many plants in the desert conserve water by not having any leaves at all. Cacti are the most prolific of this plant type. Many cacti have spines instead of leaves, which conduct photosynthesis and catch dew when the climate is right. These small structures also reflect light, further reducing water loss. During heavy rains, cacti will grow temporary root systems and absorb water. They will then shed the roots when the ground has dried.

Q.6) The vegetation of savannah consists of grassland with scattered small trees. The forest development in such areas is generally kept in check by one or more or a combination of some conditions.

Which of the following are such conditions?

1. Burrowing animals and termites

2. Fire

3. Grazing herbivores

4. Seasonal rainfall

5. Soil properties

Select the correct answer using the code given below (2021)

(a) 1 and2

(b) 4 and 5

(c) 2, 3 and 4

(d) 1, 3 and 5

Answer: C

- Savanna vegetation includes scrub , grasses and occasional trees, which grow near water holes , seasonal rivers or aquifers . Plants and animals have to adapt to the long dry periods. Many plants are xerophytic - for example, the acacia tree with its small, waxy leaves and thorns.
- The crops grown in the savanna are mainly annuals, the staple food crops being maize, millets and sorghum, but in many regions cassava is planted as a 'hedge' against crop failure. Cash crops include groundnuts, cotton and, at higher altitudes in East Mrica, coffee.
- So, the Vegetation is deliberately reduced in size by keeping in mind of the various factors which can play an important role in increasing the growth of the disasters.

Q.7) “Leaf litter decomposes faster than in any other biome and as a result, the soil surface is often almost bare. Apart from trees, the vegetation is largely composed of plant forms that reach up into the canopy vicariously, by climbing the trees or growing as epiphytes, rooted on the upper branches of trees.” This is the most likely description of (2021)

(a) Coniferous forest

(b) Dry deciduous forest

(c) Mangrove forest

(d) Tropical rain forest

Answer: D

- Option d is correct. In tropical rainforest biome, hot, moist atmosphere and all the dead plant material create the perfect conditions in which bacteria and other microorganisms can thrive. This leads to rapid decomposition of leaf litter material. The tropical rainforest is known for its dense canopies of vegetation that form three different layers. The top layer or canopy contains giant trees that grow to heights of 75 m (about 250 ft) or more. Thick, woody vines are also found in the canopy. They climb trees in the canopy to reach for sunlight.

Q.8) Consider the following statements: Statement-I : The soil in tropical rain forests is rich in nutrients. **Statement-II** : The high temperature and moisture of tropical rain forests cause dead organic matter in the soil to decompose quickly.

Which one of the following is correct in respect of the above statements? (2023)

(a) Both Statement-I and Statement-II are correct and Statement-II is the correct explanation for Statement-I

(b) Both Statement-I and Statement-II are correct and Statement-II is not the correct explanation for Statement-I

(c) Statement-I is correct but Statement-II is incorrect

(d) Statement-I is incorrect but Statement-II is correct

Answer: D

- The soil here is not rich in nutrients. It is in fact leached, as the fast growing vegetation constantly keep on absorbing the nutrients being created in the soil.

Q.9) Consider the following description: (2024)

1. Annual and daily range of temperatures is low.

2. Precipitation occurs throughout the year.

3. Precipitation varies between 50 cm – 250 cm.

What is this type of climate?

(a) Equatorial climate

(b) China type climate

(c) Humid subtropical climate

(d) Marine West coast climate

Answer: (d) Marine West coast climate

Q.10) Consider the following statements: Statement I: The amount of dust particles in the atmosphere is more in subtropical and temperate areas than in equatorial and polar regions. **Statement II:** Subtropical and temperate areas have less dry winds.

Which one of the following is correct in respect of the above statements?

(a) Both Statement I and Statement II are correct and Statement II explains Statement I

(b) Both Statement I and Statement II are correct but Statement II does not explain Statement I

(c) Statement I is correct but Statement II is not correct

(d) Statement I is not correct but Statement II is correct

Answer: (c) Statement I is correct but Statement II is not correct Statement I: Correct

- Dust concentration is higher in subtropical and temperate regions, particularly in arid and semi-arid zones like the Sahara Desert, Arabian Peninsula, Central Asia, and parts of China.
- These areas have scarce vegetation, high wind activity, and dry conditions, ideal for dust uplift.
- In contrast, equatorial regions are humid and forested (e.g., Amazon, Congo), limiting dust production. Polar regions are ice-covered, also limiting dust.

Statement II: Incorrect

- Subtropical regions, especially deserts (e.g., Sahara, Thar), experience strong dry winds like:
  - Trade winds
  - Harmattan (West Africa)
  - Santa Ana (California)
- These dry winds are key agents for transporting dust particles across continents and oceans.
- Therefore, the statement that these areas have “less dry winds” is factually incorrect.

⚡ **Quick revise** Climate PYQs in one glance Fold ▲

Climate PYQs in one glance | Year | Asked | Answer / rule | |---|---|---| | 2011 | Rainforest regenerates slowly | **Soil poor in nutrients** (leached, nutrients stored in plants; rapid decay) | | 2011 | African–Eurasian desert belt | **Subtropical high-pressure cells** (not warm currents). Rain-shadow (Gobi) and **cold currents** (Namib, Atacama) are other causes | | 2012 | Tropical savannah | **Definite dry and wet season** | | 2015 | "Clear morning, heat builds, thunder, short rain" | **Equatorial** (ITCZ; two rainfall peaks) | | 2018 | Desert leaf modifications | **Hard waxy, tiny, thorns – all three** | | 2021 | What keeps savannah forest in check | **Fire, grazing herbivores, seasonal rainfall** (2, 3, 4) | | 2021 | Fast leaf-litter decay, bare soil, epiphytes | **Tropical rain forest** | | 2023 | Rainforest soil rich in nutrients? | Statement I **wrong**; II right (fast decay, nutrients absorbed) | | 2024 | Low annual/daily range, rain all year, 50–250 cm | **Marine West**

Coast | | no year | Dust in subtropical/temperate areas | I correct (deserts, wind); II wrong – these have **strong dry winds** |

Recall sheet

Q	A
Troposphere height; lapse	8–18 km (13 avg); 1 °C per 165 m (≈6.5 °C/km)
Ozone layer height	20–25 km
Coldest layer; layer for jets	Mesosphere; lower stratosphere
Kármán line	100 km
Energy surplus/deficit boundary	40° N/S
Coriolis: where zero / max; direction	Equator / poles; right in N, left in S
Roaring / Furious / Shrieking	40s / 50s / 60s
Polar jet vs subtropical jet	50–60° (winter-strong) / 20–30°
Warm cold-front cloud types	Warm: stratus, cirrostratus; Cold: cumulonimbus
Cyclone SST	≥ 26.5 °C
Where does Chinook blow; its nickname	E. Rockies; snow-eater
PSC altitude & temperature	15–25 km; below −78 °C
Cloud seeding agents	Silver iodide, potassium iodide, salt, dry ice
Which hemisphere has stronger westerlies, why	Southern – less land
Desert belts' main cause	Subtropical high-pressure cells (plus rain shadow, cold currents)

Traps: "Jet streams only in the Northern Hemisphere" ✘ · "Temperature contrast of land-sea is greater in summer" ✘ (winter) · "Westerlies 30°N–60°S" ✘ · "Ozone causes radio reflection" ✘ · "More moisture aloft makes it colder" ✘ · "Gulf Stream is a cold current" ✘ (statement 2 in PYQ 2024 is wrong) · "Eye is 10 °C cooler" ✘ · "Rainforest soil is rich" ✘.

Connects to: Pressure belts shift with the Sun → *Geomorphology* (23.5° tilt, 21 June) · Western disturbances, monsoon, Loo → *Indian Geography* (climate) · Ocean currents shape isotherms and deserts → *Oceanography* · Climatic regions → *Economic Geography* (agriculture).

GEOGRAPHY · GS FOUNDATION · SUBJECT 3

Oceanography

Handout + mind maps: no class slides | Handout pages 29

MEDIUM · 6 Prelims, 1 Mains

The whole Oceanography handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 18 mind-map pages are in Chapter 9.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: MEDIUM-HIGH. 6 Prelims PYQs** printed: **2013×2** (freshwater share; fishing grounds) · **2015×2** (equatorial counter-current; tides) · **2021×2** (freshwater; warm/cold sides of oceans). **Mains quoted:** *"Account for variations in oceanic salinity and discuss its multi-dimensional effects"* (2017, 15m).
- **Pattern:** the exam asks *why* – why the counter-current flows east, why tides occur, why fishing grounds sit where warm and cold currents meet.
- **Current-affairs heavy handout:** AMOC weakening, coral bleaching, dead zones, marine heatwaves, Southern Ocean carbon sink – good Mains material.
- *No teacher ink in this PDF; priority = PYQ count in the handout.*

1 · OCEANS AND THEIR RELIEF

- The hydrosphere covers **nearly 71% of the total surface area** of the earth.
- Divided into Oceans, Inland Seas, Small enclosed seas, Bays, etc. based on size and location.
- **5 major oceans** : **Pacific Ocean, Atlantic Ocean, Indian Ocean, Arctic Ocean and Southern Ocean**
- The **ocean floors are characterized by various relief zones** – Continental Shelf, Continental Slope, Deep Sea Plain, and Oceanic deep or trench.
- Ocean relief is **created largely due to tectonic, volcanic, erosional, and depositional processes** and their interactions.
- The **average depth of the ocean is 3,800 m against the 840 m average height of the lithosphere.**

Water on Earth



Oceans	:	97.3	Saline Water
Ice-caps	:	02.0	
Ground water	:	0.68	
Fresh water lakes	:	0.009	
Inland seas and salt lakes	:	0.009	Fresh Water
Atmosphere	:	0.0019	
Rivers	:	0.0001	
		100.00	

Oceanic Relief Features

⚡ **Quick revise** Oceans and their relief — Fold ▲

🏠 **Ocean floor = a house, from the beach inwards:**

Part	House picture	Facts
Continental shelf	front **porch**	gentle **1–2°**; to ~**200 m** (up to 500 m); continental crust + sediment; **oil & gas**, fisheries; North Sea, Gulf of Mexico; **Great Barrier Reef** (>2,300 km)
Continental slope	steep **stairs**	**3–6°** (up to 25°); ~200–3,000 m; **submarine canyons** (Monterey, Hudson); Blake Plateau
Continental rise	**landing**	gentle sediment apron at slope base; Amazon rise
Abyssal (deep-sea) plain	living-room **floor**	**> 2,000 m** (2,000–6,000+); flat, fine sediment; **polymetallic nodules** – Clarion–Clipperton Zone (Hawaii–Mexico)
Trench	**cellar**	**> 6,000 m**; convergent boundaries; subduction, quakes; **Mariana Trench**

Continental Shelf:

- The continental shelf is the **submerged extension of a continent** , sloping gently (1-2 degrees) from the shoreline to the continental slope.
- The **width of the continental shelf varies widely** , ranging from a few kilometers to hundreds of kilometers.



- The depth of the continental shelf is **relatively shallow** , typically extending to depths of about 200 meters (656 feet) but can reach depths of up to 500 meters (1,640 feet) in some areas.
- The continental shelf is **composed of continental crust and sedimentary deposits** , including sand, silt, and clay.
- It **may feature submerged ridges, valleys, and canyons** formed by geological processes such as erosion, sedimentation, and tectonic activity.
- **Economic Significance:**
  - The continental shelf is often rich in **hydrocarbon resources** such as oil and natural gas, which accumulate in sedimentary layers over millions of years. Offshore drilling and exploration are conducted on continental shelves to extract these resources.
  - The continental shelf supports **diverse marine ecosystems** , including coral reefs, kelp forests, and commercially important fish species.

- It provides habitat and feeding grounds for marine organisms and supports **commercial fisheries** .
- This biodiversity can also attract **tourism** .

- **Examples:**
  - **The North Sea** : Located off the coasts of Norway, Denmark, Germany, the Netherlands, and the United Kingdom, the North Sea has a wide continental shelf and is known for its oil and gas reserves.
  - **The Gulf of Mexico** : Bordered by the southern United States and Mexico, the Gulf of Mexico has an extensive continental shelf and is a major oil-producing region.
  - **The Great Barrier Reef** : Off the coast of Australia, the Great Barrier Reef is the world's largest coral reef system, extending over 2,300 kilometers along the continental shelf of Queensland.

Continental Slope:

- The continental slope is **the steeply sloping transition zone between the continental shelf and the deep ocean floor.**
- The continental slope begins at the outer edge of the continental shelf and extends to the continental rise or the abyssal plain.
- The gradient of the continental slope is much **steeper** than that of the continental shelf, typically **ranging from 3 to 6 degrees but can be as steep as 25 degrees in some areas.**
- The depth of the continental slope varies widely, ranging from depths of about 200 meters (656 feet) to 3,000 meters (9,843 feet) or more.
- The continental slope is **characterized by underwater cliffs, canyons, and steep slopes formed by erosion, sedimentation, and tectonic activity.**
- Submarine canyons, such as the Monterey Canyon off the coast of California, are common features of continental slopes.

- Sediments eroded from the continental shelf are transported down the continental slope by underwater currents and gravity, contributing to the formation of deep-sea fans, Continental Rise, and turbidity currents.
- It supports **deep-sea ecosystems** adapted to high pressure, low light, and cold temperatures.
- **Resource Exploration** :
  - The continental slope is of interest for resource exploration, including oil and gas deposits, mineral resources, and deep-sea mining opportunities. But we are **yet to develop proper technologies** for resource extraction.
  - Deep-sea research and exploration expeditions study the geological, biological, and environmental characteristics of the continental slope.
- Examples:
  - The Blake Plateau: Located off the southeastern coast of the United States, the Blake Plateau features a prominent continental slope and is known for its rich biodiversity and geological formations.
  - The Antarctic Slope Front: Surrounding the Antarctic continent, the Antarctic Slope Front marks the boundary between the continental shelf and the deep ocean and plays a key role in ocean circulation and climate dynamics.

Deep Sea Plains

- Deep-sea plains, also known as **abyssal plains** , are **vast, flat expanses of the ocean floor located at depths greater than 2,000 meters** (6,560 feet).
- Deep-sea plains are **found in the abyssal zones of the ocean** , typically beyond the continental slope and rise, covering large areas of the ocean basin.
- Deep-sea plains are characterized by depths ranging from 2,000 meters (6,560 feet) to 6,000 meters (19,685 feet) or more, with relatively gentle slopes and low relief.
- Deep-sea plains have **relatively flat topography** compared to the rugged terrain of the continental slope and rise.
- They are **composed of fine-grained sediments** , including clay, silt, and sand, deposited over millions of years by marine currents and processes.
- **Biological Activity:**
  - Deep-sea plains support **diverse ecosystems adapted to extreme conditions, including high pressure, low light, and cold temperatures.**
  - **Benthic organisms** such as deep-sea corals, sponges, and echinoderms inhabit the sediments and substrates of deep-sea plains.
- **Mineral Resources:**
  - Deep-sea plains are of interest for their mineral resources, including **polymetallic nodules, manganese crusts, and hydrothermal vent deposits** .
  - These resources contain valuable metals and minerals used in various industries, including technology, manufacturing, and construction.
  - However, their **extraction is a challenge.** We are yet to develop technologies for that.
- **Geological Processes** :

- Deep-sea plains are shaped by geological processes such as **sedimentation, erosion, and tectonic activity.**
- Sediments accumulate slowly over time, forming flat layers that may preserve geological and environmental records.
- There are also volcanic features like sea mounds, guyots, trenches, etc.

- Example: The Clarion-Clipperton Zone: Located in the Pacific Ocean between Hawaii and Mexico, the Clarion-Clipperton Zone is one of the largest deep-sea plains in the world and is known for its abundance of polymetallic nodules.
- **Trenches:**
  - Trenches are **long, narrow, and deep depressions in the ocean floor** , **formed at convergent plate boundaries** where one tectonic plate is forced beneath another.
  - They are the **deepest parts of the ocean** , with depths exceeding 6,000 meters (19,685 feet) in some cases.
- **Submarine Canyons** :
  - Submarine canyons are **deep, steep-sided valleys** cut into the continental slope and extending across the continental shelf and sometimes into the abyssal plain.
  - They are **formed by erosional processes such as turbidity currents, underwater landslides, and river discharge.**
  - Submarine canyons serve as **conduits for sediment transport from the continental shelf to the deep ocean** , and they provide habitat for diverse marine life.
  - E.g. Hudson Canyon, off the coast of New York.
- **Continental Rise:**
  - The continental rise is a **gently sloping sedimentary deposit located at the base of the continental slope** , where it meets the abyssal plain.
  - It consists of **accumulated sediments transported from the continental shelf and slope** by gravity, turbidity currents, and sediment gravity flows.
  - The continental rise marks the **transition between the continental margin and the deep ocean basin.**
  - E.g. Amazon Continental Rise off the coast of northeastern South America
  - Trenches are **associated with intense geological activity** , including subduction zones, earthquakes, and volcanic eruptions.
  - E.g. Mariana Trench in the Western Pacific Ocean, east of the Mariana Islands
- **Seamounts:**
  - Seamounts are **underwater mountains rising from the ocean floor** , typically with heights of 1,000 meters (3,280 feet) or more above the surrounding seabed.
  - They **can be either active or extinct volcanoes** , formed by hotspot activity or tectonic processes.
  - Seamounts provide **habitat for diverse marine life** , including deep-sea corals, sponges, and fish, and they serve as important fishing grounds and navigation hazards.
  - E.g. Davidson Seamount Off the coast of central California, USA
- **Guyots:**
  - Guyots, also known as table mounts, are **flat-topped seamounts** that have been **eroded by wave action and submerged beneath the ocean surface** .

- They are characterized by a **flat, plateau-like summit with steep slopes** and are typically found in ocean basins far from tectonic plate boundaries.
- Guyots **provide valuable insights into past sea level changes and geological processes** , and they support unique ecosystems adapted to their deep-sea environments.
- E.g. Emperor Seamounts in the Northwestern Pacific Ocean, near the Hawaiian Islands

- **Hydrothermal vents:**
  - These are **openings in the sea floor** from where **mineral-rich water comes out** .
  - It is probably here that Earth's **life first started** .
  - If black smoke comes out of them, they are known as **Black Smokers.**

⚡ **Quick revise** Seamounts Fold ▲

- **Seamounts:** ≥ **1,000 m** underwater volcanoes (Davidson). **Guyots (table mounts):** flat-topped seamounts, wave-eroded then submerged (Emperor Seamounts). **Hydrothermal vents:** mineral-rich water; **black smokers**; life may have begun here.
- **Terms (with the handout's examples): Strait** – Palk · **Isthmus** – Kra · **Archipelago** – Japanese · **Lake** – Baikal, Wular · **Bay** – Bengal · **Cape** – Comorin (Kanyakumari; southernmost tip of the Peninsula), Good Hope · **Lagoon** – **Chilika** (brackish) · **Gulf** – Kutch · **Peninsula** – Indian.

General Terminologies Associated With Oceans:

- **Strait:** A narrow waterway connecting two larger bodies of water. Example: Palk Strait, separating India and Sri Lanka.
- **Isthmus:** A narrow strip of land connecting two larger landmasses, with water on either side. Example: Isthmus of Kra, connecting the Malay Peninsula with mainland Asia.
- **Archipelago** : A group or chain of islands clustered together in a body of water. Example:Japanese Archipelago, in Western Pacific Ocean.
- **Lakes** : Bodies of water surrounded by land, larger and deeper than ponds. Example: Lake Baikal in Russia, Wular Lake in India.
- **Bay:** A recessed, coastal body of water that is partly surrounded by land. Example: Bay of Bengal.
- **Cape** : A pointed or curved piece of land extending into a body of water. Example: Cape Comorin (Kanyakumari), the southernmost tip of the Indian Peninsula; Cape of Good Hope, Africa.
- **Lagoon** : A shallow body of water separated from a larger body of water by a barrier island or reef. Example: Chilika Lake, a brackish water lagoon in Odisha, India.
- **Island** : A landmass surrounded by water. Example: Lakshadweep Islands, located in the Arabian Sea off the western coast of India.
- **Gulf:** A inlet with constricted entrance compared to a bay. Example: Gulf of Kutch, along the western coast of Gujarat, India.

- **Peninsula** : A piece of land surrounded by water on three sides, connected to a larger landmass. Example: Indian Peninsula, surrounded by Arabian Sea, Bay of Bengal and Indian Ocean.

Properties Of The Oceanic Water: Temperature

- Temperature is a vital property of ocean water that influences a wide range of physical, chemical, and biological processes.
- **Factors Affecting Temperature Distribution** :
  - **Solar Radiation (Latitude)** : Solar radiation is the primary driver of ocean temperature distribution. Near the equator, where solar radiation is most intense, ocean temperatures tend to be warmer. At higher latitudes, solar radiation is less direct, resulting in cooler surface temperatures.
  - **Ocean Currents:** Ocean currents redistribute heat horizontally across the ocean's surface. Warm ocean currents, such as the Gulf Stream and the Kuroshio Current, transport heat from the equator toward higher latitudes, influencing regional climates and ocean temperatures.
  - **Atmospheric Circulation** : Atmospheric circulation patterns, such as the Hadley, Ferrel, and Polar cells, affect ocean temperature distribution by influencing wind patterns, evaporation rates, and heat exchange between the atmosphere and the ocean.
  - **Continentality:** The water body surrounded by land on its majority of shores, will be warmer than a body surrounded by land on less of its sides. This is because just like oceans have a moderating impact on the land’s temperature, similarly land has the opposite i.e. **“extreming” effect** on the sea’s temperature.
  - **Ocean Depth** : Temperature distribution in the ocean varies vertically with depth. Surface waters are generally warmer due to solar heating and mixing by wind and waves, while deeper waters tend to be colder.
  - **Seasonal Variation** : Seasonal changes in solar radiation and atmospheric conditions lead to seasonal variations in ocean temperature. For example, ocean temperatures may be warmer in summer and cooler in winter due to changes in solar insolation and atmospheric circulation.
- **Horizontal Distribution of Temperature:**
  - The horizontal distribution of ocean temperature is influenced by factors such as latitude, solar radiation, ocean currents, and atmospheric circulation patterns.
  - **Near the equator, ocean temperatures are generally warm, with surface temperatures often exceeding 25°C (77°F) due to direct exposure to intense solar radiation.**
  - **As you move away from the equator toward higher latitudes, ocean temperatures tend to decrease.** In polar regions, surface temperatures can drop below freezing due to the influence of cold polar air masses.
  - Ocean currents play a crucial role in redistributing heat horizontally across the ocean surface. Warm ocean currents, such as the Gulf Stream in the North Atlantic and the Kuroshio Current in the North Pacific, transport heat from the equator toward higher latitudes, influencing regional climates and ecosystems.
- **Vertical Distribution of Temperature:**

- The vertical distribution of ocean temperature **varies with depth** , forming distinct layers known as water masses.



Figure : Thermocline

- **The uppermost layer of the ocean, called the surface mixed layer up to 200-400m, is typically warmer** due to direct solar heating and mixing by wind and waves. Surface temperatures can vary widely depending on factors such as location, season, and ocean currents.
- **Below the surface** mixed layer lies the **thermocline, a transition zone characterized by a rapid decrease in temperature with depth.** The thermocline separates the warm, well-mixed surface layer from the colder, deeper layers of the ocean.
- **In the deep ocean, temperatures are relatively uniform and cold,** typically ranging from 2°C to 4°C (35.6°F to 39.2°F). This deep layer, known as the abyssal zone, extends from the base of the thermocline to the ocean floor and is largely unaffected by solar heating or surface processes. Here sporadically certain locations are heated due to volcanic activities, but that is an exception and not a rule.

- **Significance of Temperature:**
  - Ocean temperature influences the **distribution and behaviour of marine organisms,** including fish, plankton, and marine mammals.
  - **Temperature gradients** in the ocean **drive ocean currents** , which play a crucial role in regulating Earth's climate and transporting heat around the globe.
  - Changes in ocean temperature **can impact weather patterns, sea level rise, and the frequency and intensity of extreme events** such as hurricanes and droughts.
- Mean annual temperature **: 26.7° C in equatorial areas; 12.7° C in 45° N and S latitudes; freezing point at the poles.**
- The **Indian Ocean is the warmest.**
- The **Northern Hemisphere is warmer** than the Southern Hemisphere, due to more presence of land.

Properties Of The Ocean Water: Salinity

- Salinity, the **concentration of dissolved salts in ocean water** , is a crucial property that influences various oceanographic processes.

Salts	Percentage
1. Sodium chloride	77.8%
2. Magnesium chloride	10.9%
3. Magnesium sulphate	4.7%
4. Calcium sulphate	3.6%
5. Potassium sulphate	2.5%

- **Factors Affecting Salinity Distribution:**
  - **Evaporation and Precipitation** : Regions with high evaporation rates and low precipitation, such as subtropical oceans, experience higher salinity due to water loss and salt accumulation. Conversely, areas with high precipitation rates, such as the equator and high latitudes, may have lower salinity due to dilution by freshwater inputs.
  - **Riverine Input: Rivers transport freshwater** and dissolved minerals from land to the ocean, contributing to variations in salinity along coastlines and estuaries. Regions near river mouths tend to have lower salinity due to freshwater influx, while areas further offshore may exhibit higher salinity.
  - **Sea Ice Formation and Melting: During sea ice formation, salt is excluded from the ice crystals** , increasing the salinity of surrounding seawater. **Conversely, when sea ice melts, freshwater is added to the ocean, reducing salinity** . These processes influence salinity distribution in polar regions and areas affected by seasonal ice cover.
  - **Ocean Currents** : Ocean currents transport water masses with different salinities across the ocean's surface and depths. **Warm, saline currents, such as the Gulf Stream, contribute to higher salinity in regions they influence** , while cold, freshwater currents, such as the Labrador Current, can lower salinity in adjacent areas.
  - **Atmospheric Circulation** : Atmospheric circulation patterns, such as trade winds and westerlies, **influence surface ocean currents and evaporation** rates, affecting regional salinity patterns.
- **Horizontal Distribution of Salinity:**
  - **Latitudinal Variations:**



        - Salinity levels tend to be **higher in subtropical regions and lower in equatorial and polar regions.** Subtropical regions, around 20-30 degrees latitude, experience high evaporation rates and limited precipitation, leading to elevated salinity levels.
        - **Equatorial regions receive high levels of precipitation but also experience significant evaporation, resulting in relatively lower salinity.**
        - Polar regions have lower salinity due to freshwater input from melting ice, precipitation, and runoff from land.
    - **Coastal Influence: Salinity near coastlines can vary significantly** due to factors such as river discharge, freshwater input, and local weather patterns.
        - **Riverine input** from streams and rivers can reduce salinity near coastlines, especially during periods of high runoff.
        - **Estuaries,** where rivers meet the ocean, often have lower salinity due to mixing with freshwater.
    - **Inland Seas:**
        - **The salinity of the inland Seas and lakes is very high because of the regular supply of salt by 'the rivers falling into them.**
        - Their water becomes progressively more saline due to **evaporation** .
        - For instance, the salinity of the **Great Salt Lake, (Utah, USA), the Dead Sea, and the Lake Van in Turkey are 220, 240, and 330 respectively** .
        - The oceans and salt lakes are becoming saltier as time goes on because the rivers dump more salt into them, while freshwater is lost due to evaporation.
- **Vertical Distribution of Salinity:**
    - **Surface Layer:** The surface layer of the ocean, known as the mixed layer, generally has salinity variations latitudinally, as stated above.
    - **Halocline:** Below the surface layer lies the thermocline, a transition zone characterized by **a rapid change in temperature with depth, creating a layer of Halocline.** Salinity levels may also change within this zone but generally remain relatively stable compared to temperature changes.
    - **Deep Ocean:** Salinity in the deep ocean is relatively uniform compared to surface and intermediate layers. It is approx. **34.7 ppt** .
- **Significance of Salinity** :
    - Salinity **influences seawater density, which drives ocean circulation patterns, mixing processes, and vertical stratification** .
    - Changes in salinity can affect ocean density gradients, leading to alterations in thermohaline circulation, nutrient distribution, and marine habitats.
    - Salinity variations play a **crucial role in shaping marine ecosystems, species distributions, and biogeochemical cycles** , influencing the productivity and biodiversity of marine environments.

⚡ **Quick revise** Temperature, salinity, density Fold ▲

🧂 **Salt rule: "Sun + No Rain = Salty."** Subtropics (hot, dry) are saltiest; equator and poles are diluted. **Enclosed seas keep**

> **getting saltier** (rivers add salt, evaporation removes water): *220 – 240 – 330 = Great Salt Lake → Dead Sea → Lake Van.*
> **Continentality twist:** a sea ringed by land is *warmer* – land "extremes" the sea just as sea moderates the land.

Properties Of Ocean Water: Density

- Density is a critical property of ocean water that influences various physical processes, including ocean circulation, stratification, and mixing.



- **Factors affecting** Density Distribution:
  - **Temperature** : **Cold water is denser than warm water** , so temperature variations play a significant role in determining ocean density.
    - Cold polar waters tend to be denser than warmer equatorial waters, leading to density gradients across latitudes.
    - Vertical variations in temperature also contribute to density differences within the ocean.
  - **Salinity** : Salinity affects seawater density, with **higher salinity generally increasing water density.**
    - Evaporation and precipitation influence salinity levels, leading to variations in density in regions with different water balances.
  - **Pressure:** Pressure **increases with depth in the ocean** , **compressing water molecules and increasing density.**
    - Deep ocean waters are denser than surface waters due to the higher pressures experienced at depth.
- The **impact of temperature on density is higher than the impact of Salinity on it.**
- **Horizontal Distribution:**
  - **Latitudinal Variations: Density varies horizontally across latitudes due to temperature and salinity gradients.**
    - **Subtropical regions often have higher density** due to higher salinity and lower temperatures associated with subtropical gyres.
    - **Equatorial regions** may **have lower** density due to warmer temperatures and lower salinity associated with precipitation.

- **Coastal Influence: Coastal areas may exhibit density variations due to factors such as river runoff, freshwater input, and mixing with terrestrial sources.**
  - Estuaries and coastal upwelling zones often experience density gradients due to interactions between freshwater and seawater.

- **Vertical Distribution** : Vertical variation of density is **not constant** and is different for different latitudinal areas. However, a general trend is that it increases with depth due to an increase in pressure.
- **Significance:**
  - Density variations **drive vertical and horizontal ocean circulation** , influencing heat transport, nutrient distribution, and marine habitats.
  - Ocean density gradients contribute to the formation of ocean currents, **upwelling and downwelling zones** , and the global conveyor belt circulation system.

⚡ **Quick revise** Temperature, salinity, density — Fold ▲

	Temperature	Salinity	Density
Key numbers	**26.7 °C** equatorial mean; **12.7 °C** at 45° N/S; deep ocean **2–4 °C**; mixed layer to 200–400 m; **thermocline** = rapid fall	deep ocean ≈ **34.7 ppt**; **Great Salt Lake 220 · Dead Sea 240 · Lake Van 330**	cold > warm; high salinity > low; deep > surface
Pattern	warmest at equator; **Indian Ocean warmest**; **N. Hemisphere warmer** (more land)	**highest at ~20–30° (subtropics)**: high evaporation, little rain; lower at equator & poles	**temperature matters more than salinity**
Factors	latitude, currents (Gulf Stream, Kuroshio warm), atmospheric circulation, continentality, depth, season	evaporation/precipitation, river inflow, **sea-ice** (freezing leaves salt behind), currents (Gulf Stream saline; **Labrador** cold fresher), winds	temperature, salinity, pressure

OCEAN ACIDIFICATION

- Reduction in the pH of Oceanic water. Pre-industrial oceanic pH was 8.2, 8.11 in 1985 to 8.04 in 2024 . (30% increase in acidity in the past 200 yrs).
- Seen more in arctic region compared to tropics, as colder water have a higher capacity to absorb CO2.
- Causes:
  - Dissolution of Carbon Dioxide in Water.
  - Volcanoes
  - Agricultural runoff
- Impact: Coral bleaching, weakening of shells & exoskeleton of shellfish, changes in the growth of some species of phytoplankton.

⚡ **Quick revise** Basics: hydrosphere — Fold ▲

Basics: hydrosphere ≈ **71%** of Earth's surface · 5 oceans: **Pacific, Atlantic, Indian, Arctic, Southern** · average ocean depth **3,800 m** vs average land height **840 m**.

⚡ **Quick revise** Ocean acidification Fold ▲

Ocean acidification: pH **8.2** (pre-industrial) → **8.11** (1985) → **8.04** (2024); ~**30% more acidic** in 200 yrs. *More* in the Arctic than tropics (cold water absorbs more CO_2). Causes: CO_2 dissolving, volcanoes, agricultural runoff. Effects: coral bleaching, weaker shells, changes in phytoplankton.

2 · PYQ

Q.1) On the planet earth, most of the freshwater exists as ice caps and glaciers. Out of the remaining freshwater, the largest proportion (2013)

(a) is found in the atmosphere as moisture and clouds

(b) is found in freshwater lakes and rivers\

(c) exists as groundwater

(d) exists as soil moisture

Ans: C According to the U.S. Geological Survey, over 68 percent of the freshwater on Earth is found in ice caps and glaciers, and just over 30 per cent is found in groundwater.

- Only about 0.3 percent of our freshwater is found in the surface water of lakes, rivers, and swamps.
- Over 68 percent of the freshwater on Earth is found in icecaps and glaciers, and just over 30 percent is found in groundwater. Only about 0.3 percent of our freshwater is found in the surface water of lakes, rivers, and swamps.
- US and Canadian researchers recently calculated the total amount of the world's groundwater and estimated that it is equivalent to a lake 180 metres deep covering the entire Earth. This makes groundwater the largest active freshwater resource on the planet.

Q.2) With reference to the water on the planet Earth, consider the following statements: (2021)

1. The amount of water in the rivers and lakes is more than the amount of groundwater.

2. The amount of water in polar ice caps and glaciers is more than the amount of groundwater.

Which of the statements given above is/are correct?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Ans.- B

- Groundwater represents about 30% of world's fresh water. From the other 70%, nearly 69% is captured in the ice caps and mountain snow/glaciers and merely 1% is found in river and lakes.
- Of the total freshwater, 69% resides in glaciers and 30% underground. Polar ice caps and glaciers has more water than groundwater.
- 3% of the earth's water is fresh. 2.5% of the earth's fresh water is unavailable: locked up in glaciers, polar ice caps, atmosphere, and soil; highly polluted; or lies too far under the earth's surface to be extracted at an affordable cost. 0.5% of the earth's water is available fresh water.

Q.3) Account for variations in oceanic salinity and discuss its multi-dimensional effects. (2017, 15m)

⚡ **Quick revise** Mains 2017 (15m) skeleton – salinity Fold ▲

Mains 2017 (15m) skeleton – salinity. *Intro:* salinity = dissolved salts (avg ~35 ppt, deep 34.7). *Body (variation):* latitude (subtropical high), evaporation vs rain, rivers, sea ice, currents, inland seas (220/240/330). *Effects:* controls **density → thermohaline circulation**, stratification (halocline), nutrient distribution, marine habitats and productivity. *Close:* freshwater from melting ice is now altering salinity and circulation.

Freshwater PYQs. *2013 (ans C):* after ice caps/glaciers, the largest share of freshwater is **groundwater**. *2021 (ans B):* ice caps and glaciers hold **more** than groundwater (≈69% vs 30%); rivers and lakes ≈ **1%** (handout also gives 0.3% for surface water). Only ~3% of Earth's water is fresh.

3 · OCEAN MOVEMENTS: OCEAN CURRENTS

- Ocean currents are continuous **flows of large volumes of water in a definite direction.**
- Unlike waves, ocean currents are **continuous** in nature, **flowing all year round** .
- Just like the prevailing winds, the ocean currents are also responsible for the **transfer of heat from the tropics to the poles.**

Types of currents:

- **Surface Currents** :

- Surface currents are **driven primarily by wind patterns** and occur in the upper layer of the ocean (typically the top 400 meters).
- They **affect only 10% of the ocean water** .
- They flow horizontally and can extend over large distances, influencing weather patterns, climate, and marine ecosystems.
- Examples include the Gulf Stream in the North Atlantic, the Kuroshio Current in the North Pacific, and the Antarctic Circumpolar Current in the Southern Ocean.
- **Open Ocean Currents**
  - **Equatorial Currents** : Equatorial currents are primarily **driven by the trade winds** , which blow from east to west across the equatorial region.

- These winds generate surface currents that flow in the same direction as the wind, known as the equatorial drift. This is **supported further by** the West to east **movement of the Earth** , which, due to inertia makes the equatorial water move west.
- **The Coriolis effect** , caused by the Earth's rotation, deflects the equatorial drift to the right in the Northern Hemisphere and the left in the Southern Hemisphere, resulting in the formation of the North Equatorial Current and South Equatorial Current.
- These currents **converge near the equator and form the eastward-flowing Equatorial Counter Current.**
  - **Circumpolar Currents:**
- Ocean currents that **flow around the polar regions** of the Earth.
- **Continuous Flow** : Circumpolar boundary currents form a continuous loop **around the poles** , unlike other ocean currents that may be interrupted by landmasses.
- Isolation of Polar Waters: These currents **help to isolate the polar waters** from the warmer waters of the mid-latitudes, playing a crucial role in the Earth's climate system.
- E.g.: Antarctic Circumpolar Current, West wind drift
  - **Boundary currents** : **Flowing at the oceanic boundaries** , cutting across the latitudes. Depending upon the latitudes from where they start, they can be warm or cold currents.
- **Warm Currents or the West Boundary Currents:** Western boundary currents are **strong, narrow, and fast-moving currents** that flow along the western edges of ocean basins.
  - They are characterized by **warm water and high speeds** .
  - They are **warm as they flow from the tropics towards the poles** .
  - In the Northern Hemisphere, the trade winds blow from northeast to southwest, pushing surface water towards the western boundary of ocean basins. The Coriolis effect deflects this water to the right, resulting in the formation of strong western boundary currents.
  - Similarly, in the Southern Hemisphere, the trade winds blow from southeast to northwest, causing the formation of western boundary currents.
- **Eastern boundary currents are slower, broader, and cooler currents that flow along the eastern edges of ocean basins.**
  - They are driven by the **upwelling of deep, nutrient-rich water and the return flow of equatorial currents.**
  - They are **cold as they flow from the poles towards the tropics** .

- They are **created due to the impact of the Westerlies** , causing the water to accumulate at the Eastern boundaries of the oceans in the upper latitudes.
- **Deep Water Currents** :
  - Deep ocean currents, also known as **thermohaline currents or abyssal currents** , occur below the surface layer of the ocean and are driven by density differences.
  - These currents **result from variations in temperature and salinity** , with cold, dense water sinking and warmer, less dense water rising.
  - The **down-welling happens at the polar locations** , and the **upwelling at the tropical regions of cold currents.**
  - Deep ocean currents play a crucial role in global thermohaline circulation, which helps redistribute heat and nutrients around the world's oceans.



- **Factors affecting Ocean Currents:**
  - **Latitude** : It determines whether the current will be warm or cold.
  - **Wind** : Surface currents are primarily driven by wind patterns, especially in the upper layer of the ocean. Prevailing winds, such as the trade winds and westerlies, play a significant role in generating ocean currents.
  - **Coriolis Effect** : The Coriolis effect, caused by the Earth's rotation, deflects moving objects to the right in the Northern Hemisphere and to the left in the Southern Hemisphere. This effect influences the direction of ocean currents, causing them to curve as they flow.
  - **Gravity** : Gravity influences the movement of ocean water, causing it to flow downhill along the ocean's surface. Gravity-driven currents, such as tidal currents and density-driven currents, play a role in shaping ocean circulation patterns.
  - **Temperature** : Temperature differences between water masses contribute to the formation of density gradients, which drive thermohaline circulation. Cold, dense water sinks, while warmer, less dense water rises, creating vertical movement and influencing ocean currents.
  - **Salinity** : Higher salinity water tends to be denser and sinks, while lower salinity water is less dense and rises. This density-driven circulation, known as thermohaline circulation, plays a significant role in global ocean circulation.
  - **Continental Margins** : Ocean currents are influenced by the shape and topography of the ocean floor, particularly around continental margins. Submarine features such

as underwater ridges, seamounts, and canyons can deflect or channel ocean currents.

- **Gyres:** Gyres are **large-scale circular ocean currents** that **dominate the surface waters** of the world's oceans. These circular motion patterns are **driven primarily by** a combination of factors, including **global wind patterns, the Coriolis effect, and the configuration of coastlines and ocean basins.**



  - **Gyres in the Northern Hemisphere rotate clockwise, while gyres in the Southern Hemisphere rotate counterclockwise** . This rotation direction is influenced by the Coriolis effect.
  - **Important Gyres:**
    - **North Atlantic Gyre:**
- The North Atlantic Gyre is a **clockwise** circulation pattern in the North Atlantic Ocean.
- It consists of several currents, including the **Gulf Stream, North Atlantic Current, Canary Current, and North Equatorial Current.**
- The Gulf Stream, a warm and swift Atlantic Ocean current, is one of the most prominent features of this gyre.
- **Sargasso Sea** , the only sea without a coastline lies **in the middle of this Gyre** . It is **named after the Sargassum weed** , which grows here, sustaining a huge biodiversity.
    - **South Atlantic Gyre:**
- The South Atlantic Gyre is a **counterclockwise** circulation pattern in the South Atlantic Ocean.
- It includes currents such as the **Brazil Current, Benguela Current, and South Equatorial Current.**
    - **North Pacific Gyre:**
- The North Pacific Gyre is a **clockwise** circulation pattern in the North Pacific Ocean.
- It features currents like the **Kuroshio Current, North Pacific Current, and California Current.**
    - **South Pacific Gyre:**
- The South Pacific Gyre is a **counterclockwise** circulation pattern in the South Pacific Ocean.
- It includes currents such as the **East Australian Current, Peru Current, and South Equatorial Current.**

- **Indian Ocean Gyre:**

- The Indian Ocean Gyre is a **anti-clockwise** circulation pattern in the Indian Ocean.
- It consists of currents like the **Agulhas Current, West Australian Current, and South Equatorial Current.**

Southern Ocean Anomaly

The **Southern Ocean Carbon Anomaly** refers to the **unexpectedly high absorption of anthropogenic CO_2 by the Southern Ocean** , making it a stronger carbon sink than predicted by climate models. The anomaly is primarily driven by **surface stratification** , where **freshwater from Antarctic glacier melt and increased rainfall** forms a **light, stable surface layer** over denser saline water. This **"freshwater lid" inhibits vertical mixing** , trapping **carbon-rich deep waters 100–200 m below the surface** and preventing CO_2 from escaping into the atmosphere despite strong westerly winds. The Southern Ocean currently absorbs **about 40% of the carbon dioxide taken up by the global oceans** , temporarily slowing global warming. However, this is considered a **"fleeting reprieve"** , as weakening of the freshwater layer could trigger rapid CO_2 release and accelerate climate change in the future.

⚡ Quick revise Ocean currents

Fold ▲

> 🌀 **West = Warm & Wild; East = cold & Easygoing. Western boundary currents** (Gulf Stream, Kuroshio, Brazil, East Australian, Agulhas): **warm, narrow, fast, strong**. **Eastern boundary currents** (Canary, Benguela, California, Peru, West Australian): **cold, broad, slow**, upwelling nutrient-rich water. **Gyres: N. Hemisphere clockwise, S. Hemisphere anticlockwise** (Coriolis). *Indian Ocean gyre* is anticlockwise.

- **Surface currents** (top ~400 m, only **~10%** of ocean water; wind-driven) vs **deep (thermohaline) currents** (density-driven; **down-welling at poles, up-welling in tropics**).
- **Equatorial system:** trade winds push water **east → west** (also Earth's eastward spin leaves water behind) → deflected by Coriolis into **North and South Equatorial Currents** → water **piles up in the west** → flows back east as the **Equatorial Counter Current** (3–10° N; between the two). **PYQ 2015 (ans B):** the counter-current flows east because of the **convergence of the two equatorial currents** (pile-up + gravity), not Earth's rotation directly. *(Don't confuse with the Equatorial Undercurrent, deeper at ~200 m.)*
- **Circumpolar:** Antarctic Circumpolar Current, West Wind Drift – isolate polar water.
- **Gyre members:** *N. Atlantic* – Gulf Stream, N. Atlantic Current, Canary, N. Equatorial; **Sargasso Sea** in its middle = **the only sea without a coastline**. *S. Atlantic* – Brazil, Benguela, S. Equatorial. *N. Pacific* –

Kuroshio, N. Pacific, California. *S. Pacific* – East Australian, Peru, S. Equatorial. *Indian* – Agulhas, West Australian, S. Equatorial.

- **PYQ 2021 (ans C, both right):** in the tropics the **west** of an ocean is warmer (trade winds pile warm water west); in temperate zones the **east** is warmer (westerlies carry warm water to the east side).
- **Factors:** latitude, wind, **Coriolis**, gravity, temperature, salinity, continental margins.

Climate change on currents: weaker thermal and salinity gradients → thermohaline slowdown, less upwelling nutrients and less oxygen in downwelling zones, shifted winds, ice melting from below, biodiversity decline.

Marine heatwaves: prolonged unusually warm sea; episodes **doubled since 1982**; warming rate nearly doubled since 2005 (Copernicus); causes – climate change, high pressure, shifting currents, El Niño, feedback loops.

WATER MASSES

- **Water masses** are **large, homogenous bodies of ocean water** that have distinct **temperature** , **salinity** , and **density** characteristics formed in specific source regions. Once formed, they **retain these properties** for long distances and time periods as they move through the ocean.
- Water masses primarily **define** the **physical and chemical environment** for marine life. Their temperature, salinity, and nutrient content **determine the types of organisms that can thrive in a particular region.**
- Water masses, once formed, **move and circulate** through the ocean due to a combination of **gravity, density differences, wind stress, and Earth's rotation** . This movement happens in **three dimensions** — vertically and horizontally — and forms the backbone of the **global ocean circulation system** .



⚡ **Quick revise** Water masses — Fold ▲

Water masses = big bodies with distinct temperature/salinity/density from a source region → set habitat (warm clear = coral; cold nutrient-rich = kelp; deep = specialised life).

Southern Ocean carbon anomaly: absorbs ~**40%** of the CO_2 taken up by oceans because glacial-melt freshwater forms a light **lid** that traps carbon-rich deep water below; a "fleeting reprieve" – if the lid weakens CO_2 could rush out.

IMPACT OF WATER MASSES

- **Defining the Physical and Chemical Environment:** Water masses primarily determine the "habitat" for marine life by influencing characteristics like temperature, salinity, and nutrients.
- **Supporting Specific Ecosystems:** Different water masses create unique environments that favor specific types of ecosystems:
  - **Coral Reefs:** Warm, nutrient-poor water masses with high clarity often support vibrant coral reef ecosystems.
  - **Kelp Forests:** Cold, nutrient-rich water masses can support the growth of kelp forests, which provide a habitat for diverse marine life.
  - **Deep-sea ecosystems:** Deep-sea water masses with constant cold temperatures and high pressure create a unique environment for specialized deep-sea organisms.
- **Impact on Marine Productivity:** The **temperature, salinity, and nutrient content of a water mass directly influence the overall productivity of an area** . Nutrient-rich, cooler water masses can support larger populations of phytoplankton (the base of the marine food chain), leading to higher overall productivity.

Atlantic Meridional Overturning Circulation: (AOC)

- The AMOC is a crucial component of global ocean circulation, responsible for redistributing heat around the Atlantic Ocean.



- It involves the northward transport of warm surface waters and the southward flow of cooler, denser waters at depth.
- **Warm, salty surface waters from the tropics are transported northward by currents like the Gulf Stream.**
- **These waters cool and sink in the North Atlantic, forming the North Atlantic Deep Water (NADW) current.**

- NADW flows southward along the ocean floor, completing the overturning circulation.
- The AMOC plays a **vital role in regulating climate patterns, and influencing weather, and precipitation in regions bordering the Atlantic.**
- Changes in the AMOC can lead to shifts in regional climate patterns, affecting temperature, rainfall, and storm frequency.
- **Climate change is affecting the AMOC through** various mechanisms, including **increased freshwater input from melting ice caps and increased precipitation.**
- These changes can disrupt the formation of NADW and weaken the overturning circulation, by disrupting the temperature and salinity properties of the polar water leading to regional climate disruptions.
- AMOC is weaker than at any point in the past millennium, primarily due to global warming
- Weakening of the AMOC could lead to altered weather patterns, more frequent extreme events (such as heatwaves, coldwaves and storms), and changes in ocean currents and ecosystems.
    - **Reduced heat transport** from the tropics to higher latitudes could lead to cooler temperatures in regions typically influenced by warm ocean currents, such as Western Europe.
    - The weakening of the AMOC may also reduce the ocean's capacity to absorb carbon dioxide, exacerbating global warming.
    - **Increased frequency and intensity of hurricanes** along the east coast of North America due to changes in ocean temperature gradients and atmospheric dynamics. Human-induced climate change has elevated Atlantic hurricane wind speeds by 18 miles per hour (29 kilometers per hour) over the past six years.
    - **Changes in the strength and position of major ocean currents** , such as the Gulf Stream, could affect the distribution and abundance of marine life in the North Atlantic.
    - **Declines in fish populations** due to changes in ocean currents and temperature regimes could have cascading effects on marine ecosystems and fisheries.
    - It could also contribute to **sea level rise** and impact global climate stability.
- **Impact of Climate Change on Ocean Currents:**
    - **Disruption of the Thermohaline circulations** . Can even cause their collapse, due to a reduction in the thermal gradient between tropical and polar waters, and a reduction of salinity difference as a result of the entry of fresh waters into the oceans due to the melting of glaciers.
    - Disruption in thermohaline currents can affect upwelling and downwelling, thereby **reducing nutrients in upwelling zones, and reducing Oxygen availability in downwelling zones.**
    - **Can change how winds move** , thereby **altering the movement of both equatorial and boundary currents** .
    - Warmer waters beneath the Arctic and Antarctic ice caps **can affect the melting of ice from below.**
    - Changes in ocean currents can **disrupt marine ecosystems** , affecting species distribution, migration patterns, and breeding grounds. This can lead to a **decline in biodiversity and changes in the availability of marine resources.**

- Coastal communities may experience changes in weather patterns, increased frequency of extreme weather events, and impacts on fisheries, tourism, and shipping routes. Altered ocean currents can also affect regional climates, agriculture, and water resources.

⚡ **Quick revise**

AMOC (Atlantic Meridional Overturning Circulation) Fold ▲

AMOC (Atlantic Meridional Overturning Circulation): warm salty surface water flows north (Gulf Stream) → cools, **sinks** in the North Atlantic (**NADW**) → flows south at depth. **Weaker than at any point in the past millennium** (global warming; melting-ice freshwater disrupts sinking). *If it weakens:* cooler W. Europe, more extreme events, stronger east-coast US hurricanes, less CO_2 uptake, sea-level rise, fisheries shifts.

Ekman Currents:

- Ekman Current: Ekman currents are **ocean currents caused by wind blowing over the surface of the sea.** Because of Earth's rotation (the Coriolis effect), water near the surface moves at an angle to the direction of the wind—usually about 45° to the right in the Northern Hemisphere and to the left in the Southern Hemisphere.
- **As you go deeper, each layer of water moves slightly differently, forming a spiral pattern called the Ekman spiral** (in the image).
- Significance:
  - **Nutrient Upwelling** : Ekman currents cause **upwelling** , particularly near coastlines, where deep, cold, nutrient-rich waters rise to the surface, **enhancing marine productivity and supporting rich fisheries.**
  - **Heat Distribution** : Ekman currents help in the **redistribution of heat** in the ocean, affecting local and global climate patterns by moving warm and cold water masses.
  - **Wind Stress and Transport** : **Ekman transport** , the net movement of surface water, occurs at an angle to the wind, influencing **coastal circulation** and the formation of important features like **gyres** .
  - **Climate and Weather** : Ekman currents and upwelling regions have significant effects on **regional climates** and contribute to weather phenomena such as **El Niño** and **La Niña** .
  - **CO_2 Absorption** : The upwelling associated with Ekman currents aids in the **exchange of carbon dioxide** between the atmosphere and ocean, impacting the **carbon cycle** .

⚡ **Quick revise** Ekman current Fold ▲

Ekman current: wind-driven; water moves **~45° to the right** (N) / left (S) of the wind; layers form the **Ekman spiral**; causes **coastal upwelling** (nutrients, fisheries), heat redistribution, El Niño/La Niña links.

Marine Heatwaves

- Prolonged **periods of unusually warm ocean temperatures,** that **can disrupt marine ecosystems.** They can lead to significant impacts on marine life, coral bleaching, and changes in ocean currents.
- According to a report by EU Monitor Copernicus, ocean warming rate has nearly doubled since 2005.
- Their **episodes have doubled since 1982.**
- **Causes:**
  - **Climate Change** : Global warming leads to increased sea surface temperatures, making marine heatwaves more frequent and intense.
  - **Atmospheric Conditions** : High-pressure systems can reduce cloud cover and wind, allowing more sunlight to warm the ocean surface.
  - **Ocean Currents** : Changes in ocean circulation, like the slowing of cold currents or the intensification of warm currents, can concentrate heat in certain areas.
  - **El Niño Events** : These natural climate cycles can cause significant warming of sea surface temperatures in the Pacific Ocean, triggering marine heatwaves.
  - **Feedback Loops** : Warm water can reduce upwelling of cooler, nutrient-rich waters, leading to further warming and prolonged heatwaves.

Ocean Movements: Tides

- **Rise (High tide/ Flood) and Fall (Low Tide/ Ebb) of Seawater**



- **Due to the gravitational pull of the Moon and Sun.**
- **On average, every place experiences a high and a low tide twice a day** (semi-diurnal).
- **Period: 12 hr 26 minutes.**

- A particular tide centre takes 24 hours 52 minutes to come under the moon but by that time there is another tide at the opposite side of the referred tide centre due to centrifugal force of the Earth.
- So, a place experiences the same tide after 12 hours 26 minutes.
- **Diurnal tides occur in specific regions** due to the unique geography and resonance effects, resulting in **one dominant high tide and one dominant low tide each day.** (Gulf of Mexico, Gulf of Thailand);
- **Semi-diurnal tides** , on the other hand, are **more widespread** .
- **Spring & Neap Tides:** There is a lot of **temporal and spatial variation in the tide-producing forces** due to different positions of the sun and the moon.
  - **Spring Tides** : Very high tides are caused when the sun, the moon, and the earth are almost in the same line.
    - This is the **syzygy position.**
    - **Height 20% more** than normal tides.
    - **Highest high and lowest low** tides created.
    - Occur twice every month (during new moon & full moon)
  - **Neap Tides** :
    - The sun, the earth, and the moon come in the **position of quadrature** (form right angle); tide-producing forces of the **sun and the moon** work in **opposite directions** , with the result, of lower high tides and higher low tides are created.
- **Factors affecting tides:**
  - **Relative distances of Sun, Moon, and Earth:**
    - When the Sun, Moon, and Earth are aligned (during full and new moons), their combined gravitational pull creates higher high tides (spring tides) and lower low tides (neap tides).
    - When the Sun and Moon are at right angles to each other (during quarter moons), their gravitational forces partially cancel out, resulting in lower high tides and higher low tides.
  - **Shape of Shoreline:**
    - **Funnel-shaped bays and estuaries can amplify tidal ranges** , leading to higher high tides and lower low tides compared to open coastlines.
    - Irregular coastlines with headlands and bays can cause tidal currents to intensify and create areas of convergence and divergence, affecting local tidal patterns.
  - **Local winds:**
    - Wind can influence tidal currents by **creating surface water movement and altering the distribution of water masses.**
    - Strong onshore winds can push water toward the shore, causing higher high tides, while offshore winds can result in lower high tides.
    - Wind-driven currents can interact with tidal currents, enhancing or opposing their direction and intensity, especially in coastal areas with complex bathymetry.
  - **Pressure systems:**
    - Atmospheric pressure variations associated with weather systems, such as high and low-pressure systems, can affect sea level and tidal height.

        - Low-pressure systems can cause sea level to rise, leading to higher high tides, while high-pressure systems can cause sea level to fall, resulting in lower high tides.
        - These pressure-induced changes in sea level can influence tidal ranges and coastal flooding events, particularly during storm surges.
- **Impact of tides:**
    - **Sustenance of life (nutrient circulation):**
        - Tidal movements play a crucial role in the circulation of nutrients within coastal ecosystems, including estuaries, salt marshes, and intertidal zones.
        - Tidal currents **transport nutrients such as nitrogen, phosphorus, and organic matter from the open ocean to coastal areas** , where they support the growth of phytoplankton and other primary producers.
        - As tides ebb and flow, they mix and distribute nutrients throughout coastal waters, enhancing productivity and biodiversity.
        - Nutrient-rich tidal flats and marshes serve as important feeding and breeding grounds for a variety of marine organisms, including fish, birds, and invertebrates.
    - **Temperature moderation:**
        - Tidal movements help regulate water temperatures in coastal areas by **mixing surface waters with cooler, deeper waters** .
        - During incoming tides (flood tide), cooler water from deeper layers is brought to the surface, lowering surface temperatures.
        - Conversely, during outgoing tides (ebb tide), warmer surface waters are mixed with cooler, deeper waters, moderating temperatures.
        - This temperature moderation effect is particularly important in coastal regions with extreme temperature fluctuations, helping to create more stable and hospitable environments for marine life.
    - **Removal of pollutants:**
        - Tidal movements contribute to the **removal and dispersion of pollutants in coastal waters through the process of flushing.**
        - During flood tides, incoming currents carry pollutants away from coastal areas and disperse them into the open ocean.
        - As tides ebb, outgoing currents transport pollutants back out to sea, diluting their concentration and reducing their impact on coastal ecosystems.
        - Tidal flushing helps mitigate the accumulation of pollutants in estuaries, harbors, and coastal zones, improving water quality and reducing the risk of contamination to marine organisms and human populations.
- **Tidal Bore** : Tidal bores are **rare phenomena** where incoming tides form **a wave that travels upstream against the river flow** , typically occurring in narrow estuaries or rivers with a shallow and gradually sloping bed even if it has deltas.
    - India: most prominent in Hooghly River



- Most prominent in the world:

Qiantang River, China.

- **Only during high tides.**
- **Absent during Neap Tides**

- **Impact of Climate Change on Tides:**
  - Higher sea levels will create **higher high tides** . Due to this, larger areas will experience the effects of tides.
  - In some regions, the rising sea level **may amplify the existing tidal range** , increasing the difference between high and low tides. This can exacerbate coastal flooding during storm surges and extreme weather events.
  - Many marine species depend on predictable tidal patterns for nesting and breeding. Mangroves too stay in such tidal regions. Alterations in tidal ranges and flooding patterns **can disrupt** these **critical habitats.**

⚡ **Quick revise** Tides, waves, tsunamis Fold ▲

> 🌗 **Spring tide = SPrung Straight line (syzygy)** – new & full moon, Sun-Moon-Earth aligned, highest highs and lowest lows, **~20% higher**, twice a month. **Neap = Nearly right-angle (quadrature)** – quarter moons, forces oppose, **lower highs, higher lows**. *Spring has nothing to do with the season.*

- **Cause (PYQ 2015, ans D):** gravity of **Moon (main)** and **Sun** *plus* **centrifugal force** of Earth → bulges on the near and far sides.
- Most places: **two highs + two lows** a day (**semi-diurnal**); a place sees the same tide every **12 h 26 min** (Moon takes **24 h 52 min** to return). **Diurnal** tides: Gulf of Mexico, Gulf of Thailand.
- **Tide height depends on:** Sun-Moon-Earth distances, **funnel-shaped bays/estuaries** (amplify), local winds, **pressure** (low pressure raises sea level).
- **Use:** nutrient circulation, temperature moderation, **flushing pollution**. **Tidal bore:** wave moving up a narrow river on a high tide, absent at

neaps; **India – Hooghly**; world's biggest – **Qiantang (China)**.

- **Climate change:** higher sea level → larger areas tidal; may amplify ranges; threatens mangrove habitats.

Tsunami: very long **seismic sea waves**; causes – earthquakes at **subduction zones** (size ∝ vertical shift), volcanoes, landslides, meteorite impact, nuclear tests. **500–1,000 km/h in deep water, < 1 m high** (invisible), slows and **grows near the coast**; many waves; warning – sea level change, odd ocean behaviour.

Rogue waves: freak giants from **constructive interference** or **currents/topography** focusing energy (Agulhas Current). **Seiche:** standing wave sloshing in lake/bay (wind, pressure, quakes, resonance).

Ocean Movements: Waves

- Waves are the **energy, not the water** as such, which moves across the ocean surface. **Water particles only travel in a small circle** as a wave passes.



- Wind **provides energy to the waves** .
- Waves **travel because the wind pushes the water body in its course while gravity pulls the crests** of the waves downward.
- The **falling water pushes** the **former troughs upward** , and the wave moves to a new position.
- The **actual motion of the water** beneath the waves is **circular.**
- **Impact of Climate Change:**
  - **Global warming can lead to stronger and more frequent storms** , particularly hurricanes and typhoons, which **generate larger and more powerful waves.** This increased wave energy can contribute to coastal erosion and damage infrastructure.
  - **Higher sea levels** mean that **waves** , including those from storm surges, **can travel further inland** , increasing the potential for coastal flooding and erosion.
  - **Changes in wind patterns and ocean currents** due to global warming **can alter wave patterns.** This can impact coastal erosion rates, sediment transport, and the stability of coastal structures.

- **Increased wave energy and frequency can damage coral reefs** , which are already stressed by warmer water temperatures and ocean acidification. The physical destruction of reefs reduces their ability to act as natural barriers against waves.

- Tsunamis: **Extremely long waves** ,
  - **Caused by a large and sudden displacement of the ocean** due to earthquakes, volcanic eruptions, etc.
  - These are **also called seismic sea waves** .
  - When they reach the coast, they **can cause dangerous coastal flooding and powerful currents** that can last for several hours or days.
  - **Causes:**
    - Just like winds are the source of energy in the case of normal surface waves, for Tsunamis, this energy comes from some other sources.
    - **Earthquake –** Most tsunamis are generated by earthquakes that occur along the subduction boundaries of plates along the ocean trenches. The size of the Tsunami is related to the size of the earthquake and the vertical shift.
    - **Volcanic eruption**
    - **Landslides**
    - **Meteorite Impacts –** Though no historical example of meteorite impact has caused tsunamis, the apparent impact of a meteorite about 5 million years ago produced tsunamis leaving deposits along the Gulf Coast of Mexico and the United States.
    - Can also be caused due to **Anthropogenic causes** like Nuclear Explosions. E.g.- Nuclear Testing by the US generated Tsunami in 1940 and the 1950s in Marshall Island.
  - **Characteristics:**
    - **Long Wavelengths** : Tsunamis have long wavelengths, often hundreds of kilometers, which allow them to travel rapidly across ocean basins.
    - **High Speeds** : Tsunamis travel at high speeds in deep water, typically around 500 to 1000 kilometers per hour (300 to 600 miles per hour).
    - **Low Wave Heights** : In the deep ocean, tsunamis may have low wave heights, often less than one meter, making them difficult to detect.
    - **Amplification near Coast** : As tsunamis approach shallow coastal areas, their speed decreases while their wave height increases, leading to amplification and potential devastation upon landfall.
    - **Destructive Power** : Tsunamis can cause catastrophic flooding, inundation of coastal areas, destruction of infrastructure, and loss of life.
    - **Multiple Waves** : A single tsunami event may consist of multiple waves, with successive waves arriving minutes to hours apart.
    - **Warning Signs** : Before the arrival of a tsunami, warning signs may include rapid sea level changes, unusual ocean behavior, and the sound of approaching waves.

⚡ **Quick revise** Waves carry Fold ▲

Waves carry **energy, not water** (particles move in circles); wind

provides energy.

Rogue Waves

Creation:

Rogue waves , also called **freak waves** , **monster waves** , or **killer waves** , are **unusually large, spontaneous, and unpredictable ocean surface waves** that are **far larger** than the surrounding waves. They can occur even in **relatively calm seas** and pose significant danger to ships and offshore structures. **Creation:**

1. Constructive Interference

- When several wave systems **coincide in phase** , their energies **combine** to form a single **giant wave** .
- This is a **random and rare** alignment, but can create extremely high waves.

2. Focusing by Currents and Topography

- **Strong ocean currents** (e.g., Agulhas Current) flowing opposite to wave direction can **compress wave fronts** , increasing wave height. **Seafloor features** like ridges or continental shelves can also focus energy.



Seiches

Creation:

A **seiche** is a **standing wave** oscillation of water in an **enclosed or semi-enclosed water body** , such as a **lake, bay, reservoir, or even a swimming pool** , caused by disturbances like **strong winds, atmospheric pressure changes** , **earthquakes** , etc. **Creation:**



- **Wind Action:** Strong, persistent winds push water toward one end of a lake or basin. When the wind stops suddenly, the water sloshes back, creating oscillations.
- **Atmospheric Pressure Changes:** Rapid changes in air pressure can depress water levels on one side and cause it to oscillate when the pressure equalizes.
- **Seismic Activity:** Underwater earthquakes or landslides can displace water, initiating back-and-forth wave motion.
- **Resonance Effect:** If the disturbance (like wind or pressure) matches the natural rhythm of the lake, the waves grow stronger and keep going longer—like pushing a swing at just the right time.
- **Basin Geometry:** The size, shape, and depth of the lake decide how big the waves will be and how long they will last.

CRYOSPHERE

Earth’s ice in all its forms



- **Significance:**
  - **69% Earth's freshwater** stored in Cryosphere.
  - During the warm season, these ice reservoirs feed the rivers of the world.
  - **Albedo** , which prevents heating of the earth via insolation.
  - These ice sheets/permafrost **have trapped within themselves many greenhouse gasses** when they were freezing. E.g. Methane, CO2
  - Their seasonal melting and freezing **affects the oceanic circulations** .
  - Their melting due to Global warming will affect the global oceanic salinity, further affecting the oceanic circulations.
  - In hilly regions, like the Himalayas, Alps, Andes, etc, the melting of the cryosphere will cause increased GLOF events.
- **Impacts of melting of Cryosphere:**
  - Affects thermohaline circulations.
  - Impact on prevailing winds, jet streams
  - Affect freshwater availability (increases in short-term, reduces in long term)
  - Albedo reduces, earth's heat budget will get affected
  - Release of GHG, thus global warming
  - Release of heavy elements.
  - Increase GLOF events
  - Release of pathogens and micro-organisms
  - Isostatic rebounce and volcanic activity
  - Impact on climate, flora, fauna & economy.

How Arctic and Antarctic melting affect us?

- **Arctic:**
  - **Climatic Impacts**

        - **Weakening of Polar vortex** , and more frequent chilly conditions in North America and Europe
        - Due to the southward movement of cold air masses, **more frequent Temperate storms** , as these cold air masses meet the warmer air masses at lower latitudes.
    - **Impact on Human Activities**
        - Opening up of **new navigation routes.** However, recent studies on the Northwest passenger of Canada have shown that global warming is causing thick ice blocks to flow into this channel from Greenland. This can pose a danger to shipping vessels, reducing the navigability of the shipping passages, at least in the short term.
        - **Easing resource exploration** in these regions.
        - **Change in fish and wildlife patterns** , that may affect the local communities and their traditional livelihoods.
        - **Increased GLOF events** : In Greenland, hundreds of lakes have burst since 2008.
        - According to a latest study, toxic Mercury trapped in the Arctic is now leaking into the environment.
- (How this mercury reaches the Arctic? Through atmospheric and oceanic circulation)
- **Antarctica:**
    - **Climate**
        - **Jet stream weakening** can cause changes in weather in South America, southern Africa, and Australia.
        - The weakening of the polar vortex is causing winter season heatwaves here. Some parts of Antarctica, in mid-July had temperatures as high as -25 to -30 degrees Celcius compared to -50 and -60 degrees Celcius.
    - **On Humans**
        - Will **affect research activities** in Antarctica.
        - **Affect the wildlife** in the region (Penguins, seals, microorganisms, etc)

Impact of Climate Change on Climate & Oceans

- **Climate**
    - **Urban Areas:** Quarter of global urban population to face 0.5°C temperature increase by 2040. The temperature rise will disproportionately impact vulnerable populations, leading to increased health risks, reduced labor productivity, and greater energy demands for cooling.
    - Increase instances of extreme weather events.
- **Biodiversity**
    - 2023 report by the Intergovernmental Panel on Climate Change had warned that if the carbon dioxide emission and global warming continued unabated, the tropical rainforest community may altogether collapse much before the end of this century.
    - According to a BHU study, while increased CO_2 can initially enhance crop growth through the "CO_2 fertilization effect," surpassing certain temperature thresholds due to climate change leads to significant reductions in crop yields.
- **Oceans**
    - Increased acidification

- Increased temperature due to global warming: a r ise in s ea surface temperature from 0.06°C per decade in the 1980s to 0.27°C per decade now.
- Changes in oceanic circulations, and thus, nutrient circulation.

⚡ **Quick revise** Cryosphere = Earth's ice Fold ▲

- **Cryosphere = Earth's ice**: **69%** of freshwater; albedo; stores methane/CO_2; feeds rivers; affects circulation; **GLOFs** in Himalaya, Alps, Andes.
- **Melting effects:** thermohaline change, winds/jet streams, freshwater (short-term ↑, long-term ↓), lower albedo, GHG release, heavy elements and **pathogens** released, **isostatic rebound** and volcanic activity.
- **Arctic:** weaker polar vortex → cold spells in N. America/Europe; more temperate storms; new shipping routes (but thick ice drifting into the Northwest Passage); resource access; **hundreds of Greenland lakes burst since 2008**; trapped **mercury** leaking. **Antarctica:** jet-stream weakening; winter heatwaves (mid-July −25 to −30 °C instead of −50 to −60 °C); effects on wildlife and research.
- **Oceans:** acidification; **SST rise 0.06 °C/decade (1980s) → 0.27 °C/decade now**; circulation and nutrient changes.

4 · PYQ

Q.1) What explains the eastward flow of the equatorial counter-current? (2015)

(a) The Earth’s rotation on its axis

(b) Convergence of the two equatorial currents

(c) Difference in salinity of the water

(d) Occurrence of the belt of calm near the equator

Ans: B The Equatorial Counter Current is an eastward flowing, wind-driven current which extends to depths of 100-150m in the Atlantic, Indian, and Pacific Oceans.

- As the earth rotates from west to east, water near the equator tends to move from east to west (due to the prevailing winds). This leads to the formation of Equatorial currents, both North and South of the equator.
- Now, as the North Equatorial Current and South Equatorial Current take the water from east to the western part of the ocean basins, there is a piling up water in the western parts of ocean basins (or eastern boundaries of land masses near the equator). This piled up water then flows from west to east, under the effect of gravity, resulting in Counter Equatorial Current (CEC).
- So, while the rotation of the earth is the primary causative factor for Equatorial Currents, it is piling up of water and subsequent backflow (convergence of both

equatorial currents) under gravity which causes the eastward flow of CEC.

- More often called the North Equatorial Countercurrent (NECC), this current flows west-to-east at about 3-10°N in the Atlantic, Indian Ocean and Pacific basins, between the North Equatorial Current (NEC) and the South Equatorial Current (SEC).
- The NECC is not to be confused with the Equatorial Undercurrent (EUC) that flows eastward along the equator at depths around 200m in the western Pacific rising to 100m in the eastern Pacific.

Q.2) Tides occur in oceans and seas due to which among the following? (2015)

1. Gravitational Force of the Sun

2. Gravitational Force of the Moon

3. Centrifugal force of the Earth

Select the correct answer using the code given below

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2, and 3

Ans: D Tides are the rise and fall of sea levels caused by the combined effects of the gravitational forces exerted by the Moon and the Sun, and the rotation of the Earth.

- The periodical rise and fall of the sea level is known as tide. The Moon's gravitational pull is the major cause of a tide. Sun’s gravitational pull and centrifugal force due to the gravitational pull of the earth to a lesser extent so causes types.
- Tides are very long waves that move across the oceans. They are caused by the gravitational forces exerted on the earth by the moon, and to a lesser extent, the sun. Gravity pulls the ocean towards the moon and high tide occurs.
- High tides and low tides are caused by the moon. The moon's gravitational pull generates something called the tidal force. The tidal force causes Earth—and its water—to bulge out on the side closest to the moon and the side farthest from the moon.

Q.3) Consider the following statements:

1. In the tropical zone, the western sections of the oceans are warmer than the eastern sections owing to the influence of trade winds.

2. In the temperate Zone, westerlies make the eastern sections of oceans warmer than the western sections

Which of the statements given above is/are correct? (2021)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 and 2

Ans.- C

- Statement 1 is correct. Warmer water is transported westward in the ocean by the Northeast trade winds in the Northern hemisphere of the tropical zone (as the figure given below shows). So, in tropical zones, the western section of ocean is warmer than eastern sections due to trade winds.



- Statement 2 is correct. Similarly, the Westerlies play an important role in carrying the warm, equatorial waters and winds to the western coasts of continents (that is eastern section of the Oceans in temperate zone). Thus, in temperate zones, westerlies make the eastern section of the ocean warmer than the western sections.

CORAL REEFS

- Corals are **marine invertebrates** , which **exist in colonies composed of numerous individual coral polyps** , which are tiny, soft-bodied organisms.
- Corals **secrete calcium carbonate to form hard skeletons** , which accumulate over time **to create** large, underwater structures known as **coral reefs.**
- These reefs provide habitat and shelter for a diverse range of marine life and are often referred to as the **"rainforests of the sea"** due to their high biodiversity.
- Corals rely on **symbiotic algae called zooxanthellae** for energy through photosynthesis, and they thrive in warm, clear, shallow waters with abundant sunlight.
- **Necessary conditions:**
  - **Warm Water Temperatures:** Coral reefs thrive in warm tropical and subtropical waters with temperatures typically ranging **between 20°C to 29°C** . These temperatures support the growth and reproduction of coral polyps.
  - **Clear, Sunlit Waters:** Corals require clear, sunlit waters to facilitate **photosynthesis by their symbiotic algae (zooxanthellae)** , which provides them with essential nutrients and energy. Adequate sunlight penetration is crucial for the growth and survival of corals.
  - **Shallow Depths:** Most coral reefs develop in shallow water environments, typically **less than 50 meters** deep. Shallow depths allow corals to access sunlight for photosynthesis and promote the growth of reef-building corals.

- **Stable Salinity Levels:** Corals prefer stable salinity levels within the range of **32 to 40 parts per thousand** (ppt). Fluctuations in salinity can stress corals and affect their ability to maintain osmotic balance.
- **Low Nutrient Levels:** Coral reefs **thrive in oligotrophic (nutrient-poor) waters** . Excessive nutrient runoff from land-based sources can lead to eutrophication, which promotes the growth of algae that can outcompete corals for space and resources.
- **Moderate Wave Action:** Moderate wave action helps to oxygenate the water and **remove sediment** , promoting coral growth and reef development. However, excessive wave energy from storms or strong currents can damage coral reefs.

- **Other type of Corals:**
  - **Hard Corals:**
    - Also known as stony corals or reef-building corals.
    - Secrete calcium carbonate skeletons, which contribute to the formation of coral reefs.
    - Examples include branching corals like Acropora and massive corals like Porites.
  - **Soft Corals:**
    - Lack a hard skeleton and have flexible structures.
    - Often referred to as gorgonians or octocorals.
    - Include sea fans, sea whips, and sea pens.
  - **Cold-water Corals:**
    - Found in deep, cold waters, including polar regions and temperate seas.
    - Adapted to low light levels and cooler temperatures compared to tropical corals.
    - Examples include species of Lophelia, Paragorgia, and Primnoa.
  - **Black Corals:**
    - Typically found in deeper waters, ranging from 30 meters to over 300 meters deep.
    - Have a **dark appearance due to the presence of a protein** called **melanin.**
    - Often prized for their use in jewelry and ornamental pieces.
    - Found in India in Gulf of Mannar region, along with other corals.
- **Types of Coral Reefs:**
  - **Fringing Reefs:**
    - Found **close to the shoreline, directly adjacent to land.**
    - Develop along the edges of continental coastlines or islands.
    - Characterized by their shallow depths and proximity to land, with the reef crest often exposed during low tide.
    - They might create small lagoons, known as **boat channels** .
  - **Barrier Reefs:**



    - **Parallel to the shoreline but separated from land by a lagoon.**
    - Form farther offshore compared to fringing reefs and are separated from the mainland by a deep lagoon.
    - Barrier reefs can stretch for long distances and are often associated with large oceanic islands or continental landmasses.
  - **Atolls:**
    - **Circular or horseshoe-shaped reefs** surrounding a central lagoon.
    - Formed from the gradual subsidence of volcanic islands or seamounts over time, leaving behind a ring-shaped coral reef.
    - Atolls are often found in the open ocean, far from continental landmasses, and may contain several small islands or islets within the lagoon.
- **Coral Bleaching:**
  - Coral bleaching occurs when **corals expel the symbiotic algae (zooxanthellae) living within their tissues** , leading to a loss of color and vital nutrients.
  - Bleached corals appear white or pale due to the loss of pigmented algae.
  - **This may be temporary or permanent.**
  - Bleaching weakens corals and can lead to their death if the stress is prolonged.
  - **Factors Responsible for this:**
    - **Elevated Water Temperatures** : High water temperatures, particularly during periods of ocean warming associated with climate change, can trigger coral bleaching.
    - **Ocean Acidification:** Increased carbon dioxide levels in the atmosphere lead to ocean acidification, which can weaken coral skeletons and make them more susceptible to bleaching.
    - **Pollution:** Pollution from agricultural runoff, sewage, and coastal development can increase stress on corals and contribute to bleaching events.
    - **UV Radiation:** Excessive exposure to ultraviolet (UV) radiation, often exacerbated by clear skies and calm seas, can also induce coral bleaching.
    - **Tectonic Activity:** Sudden exposure of the corals above the sea level due to land upliftment. Or drowning of corals to non-photic zones due to subsidence.
    - **Overfishing:** Fish help keep coral reefs clean. If they are removed, sediments and non-useful algae start accumulating over the corals and suffocate them.

- **Diseases:** Pathogenic diseases may also at times force the coral polyps to expel the zooxanthellae.
- **Irresponsible Tourism:** This may cause pollution, sedimentation, and in certain cases diseases in the corals.



- **Massive coral bleaching event is ongoing since Jan 2023.**

IMPACT OF CLIMATE CHANGE

ON CORAL REEFS

- **Coral Bleaching:** Rising sea temperatures due to global warming are the primary culprit. Bleached corals are weakened and more susceptible to diseases.
  - Example: The Great

Barrier Reef in Australia has experienced large-scale bleaching events in recent years, jeopardizing its vast coral ecosystems. Currently, 4th mass bleaching event going on since Feb 2023.

- **Ocean Acidification:** Increased absorption of carbon dioxide from the atmosphere by the oceans leads to acidification. This makes it harder for corals to build their calcium carbonate skeletons, hindering their growth and survival.
  - Example: Coral reefs in Florida, USA, have shown signs of reduced growth and weakened skeletons due to ocean acidification.
- **Increased Storm Intensity:** Warmer ocean temperatures contribute to stronger storms and hurricanes. These storms can physically damage coral reefs through strong waves and generate large amounts of sediment that can smother corals.
  - Example: Coral reefs in the Caribbean have suffered extensive damage from recent hurricanes, hindering their recovery from other stressors like bleaching.
- **Corals in India:**
  - Indian reefs have also **experienced several bleaching events** in recent decades, with significant damage reported in the Gulf of Mannar and the Andamans.
  - In March 2024, coral bleaching events have been reported from Lakshadweep.
  - These events threaten the vibrant coral ecosystems and the associated biodiversity.

- Pollution from coastal development, destructive fishing practices, and sedimentation from rivers can exacerbate the impacts of global warming on Indian coral reefs.

⚡ **Quick revise** Coral reefs Fold ▲

Corals are fussy beach tourists – "Warm, Clear, Shallow, Salt-stable, Light-snack, Mild-waves": 20–29 °C · clear sunlit water (for **zooxanthellae** algae) · **< 50 m** deep · salinity **32–40 ppt** · **low nutrients** (oligotrophic – runoff feeds algae that choke them) · moderate waves. **Reef types: Fringing = Friend** hugs the shore (boat channels) · **Barrier = Bodyguard** stands off, lagoon between · **Atoll = Wedding ring** around a lagoon (sinking volcanic island/seamount).

- **Kinds:** hard (stony, reef-builders; *Acropora, Porites*) · soft (octocorals, sea fans) · cold-water (deep, polar/temperate) · **black corals** (30–300+ m; found in **Gulf of Mannar**). Reefs = "**rainforests of the sea**"; corals secrete **calcium carbonate**.
- **Bleaching** = corals expel zooxanthellae (white/pale; may recover or die). Causes: **heat**, acidification, pollution, UV, **tectonic** exposure/drowning, **overfishing** (fish keep reefs clean), disease, tourism. **4th mass bleaching ongoing since early 2023** (handout says Jan 2023 once and Feb 2023 once).
- **Climate impacts:** bleaching, acidification (harder to build skeletons), stronger storms. **India:** bleaching in **Gulf of Mannar and Andamans**; **Lakshadweep, March 2024**.

OCEANIC DEPOSITS

- **Terrigenous Deposits:**
  - Derived from **land-based sources** , such as rivers, glaciers, and wind-blown dust.
  - **Found primarily on continental margins** , including continental shelves and slopes.
  - Examples include sand, silt, and clay, as well as larger particles like gravel and boulders.
- **Biogenous Deposits** :
  - **Composed of** remains of **marine organisms** , such as shells, skeletons, and coral fragments.
  - Accumulate on the seafloor over long periods.
  - **Found in both shallow and deep ocean environments** , including continental shelves, slopes, and abyssal plains.
  - Biogenous deposits **include** :
    - **Calcareous Ooze** : Composed primarily of calcium carbonate shells and skeletons of foraminifera, coccolithophores, and other calcareous organisms.

Found in warmer, shallower waters.

- **Siliceous Ooze** : Made up of silica-based remains of diatoms, radiolarians, and sponge spicules. Found in colder, deeper waters.

- **Hydrogenous Deposits:**
  - Formed through **chemical precipitation from seawater** .
  - **Found in various oceanic environments** , including continental shelves, slopes, and deep-sea hydrothermal vent areas.
  - Examples include:
    - **Manganese Nodules** : Formed of manganese, iron, and other metals precipitated from seawater onto the seafloor.
    - **Phosphorite Deposits** : Consist of phosphate minerals precipitated from marine sediments.
- **Cosmogenous Deposits:**
  - **Derived from extraterrestrial sources** , such as **meteorites and interplanetary dust.**
  - Found **distributed throughout the oceans** , with higher concentrations in deep-sea environments.
  - Examples include microscopic meteorite particles and cosmic dust grains.

⚡ **Quick revise** Deposits, dead zones, resources, law Fold ▲

🔠 **Four deposit types – "Land, Life, Liquid, Cosmos":** **Terrigenous** (from land: rivers, glaciers, wind; on margins) · **Biogenous** (organisms) – *calcareous ooze* ($CaCO_3$; **warmer, shallower**), *siliceous ooze* (diatoms, radiolarians; **colder, deeper**) · **Hydrogenous** (chemical: **manganese nodules**, phosphorite) · **Cosmogenous** (meteorite dust).

DEAD ZONES

- Dead zones, **also known as hypoxic zones** , are areas in oceans and large lakes where oxygen levels become so low that most marine life cannot survive.
- **Causes:** The primary culprit behind dead zones is **nutrient pollution** , particularly excess nitrogen and phosphorus. This pollution often comes from:
  - **Agricultural runoff:** Fertilizers used in farming can be washed away by rain, carrying these nutrients into rivers and ultimately into coastal waters.
  - **Sewage discharge:** Untreated or inadequately treated sewage can also contribute excess nutrients.
  - **Industrial waste:** Certain industries may release nitrogen and phosphorus into waterways.
  - **Wind Deposits:** A massive algal bloom off the coast of Madagascar was triggered by dust blown from drought-stricken regions of southern Africa. The dust contained nutrients like iron and phosphorus that acted like fertilizers.

Formation:

- When these nutrients reach coastal waters, they trigger a rapid growth of algae. This phenomenon is called an **algal bloom** . As the algae die and decompose, they sink to the bottom, where they are consumed by bacteria. This bacterial decomposition process consumes a large amount of oxygen dissolved in the water.

Consequences:

The spread of dead zones poses a serious threat to marine ecosystems, with cascading effects: **Loss of Marine Life:** When oxygen levels become depleted, fish, shrimp, and other marine organisms suffocate and die. This disrupts the food chain and can lead to ecosystem collapse. Dead zones can become "ecological graveyards" devoid of most marine life. **Economic Damage:** Dead zones can have a significant economic impact on fisheries, tourism, and other industries that rely on healthy marine ecosystems. As fish populations decline, fishing communities suffer economic losses, and tourism to affected areas can decrease. **Public Health Risks:** Decomposition of organic matter in dead zones can release harmful toxins and bacteria. This can contaminate seafood and pose a risk to human health if affected fish are consumed.

Examples:

- **Northern Gulf of Mexico:** One of the world's largest dead zones occurs annually in the Gulf of Mexico, primarily due to agricultural runoff from the Mississippi River basin.
- **Baltic Sea:** This semi-enclosed sea suffers from chronic oxygen depletion due to nutrient pollution from surrounding countries.
- **Black Sea:** The Black Sea has a naturally occurring dead zone in its deep waters due to limited circulation and high organic matter content.

How to Combat?

- **Reducing Nutrient Pollution:** Implementing stricter regulations on agricultural practices, improving wastewater treatment, and promoting sustainable farming methods are essential to curb nutrient runoff.
- **Monitoring and Management:** Regular monitoring of water quality and implementing measures to manage nutrient loads in vulnerable areas can help prevent dead zone formation.
- **Public Awareness:** Raising public awareness about the causes and impacts of dead zones is essential for promoting behavioral changes and supporting policies that address this environmental threat.

⚡ **Quick revise** Dead zones (hypoxic) Fold ▲

Dead zones (hypoxic): nutrient pollution (N, P: farm runoff, sewage, industry, even wind-blown dust – Madagascar bloom) → **algal bloom → dies → bacteria use up oxygen**. Effects: marine life suffocates, fisheries/tourism lose, toxins. **Examples:** northern **Gulf of Mexico** (Mississippi runoff), **Baltic Sea**, **Black Sea** (natural, poor circulation). **Fix:** cut nutrient pollution, treat sewage, monitor, awareness.

OCEAN RESOURCES:

- **Biotic Resources:**
  - **Edible Resources:**
    - **Fish:** Tuna and salmon are abundant in the Pacific Ocean, while cod fisheries are prominent in the North Atlantic. Sardines are found in the Mediterranean Sea and off the coast of Peru.
    - **Shellfish:** Shrimp and crabs are harvested in the Gulf of Mexico and along the coast of Alaska. Lobsters are abundant in the waters of Maine and the North Atlantic, while oysters are farmed in the Chesapeake Bay and along the Gulf Coast.
    - **Seaweed:** Japan is a major producer of seaweed, particularly nori, used in sushi. Other countries like China, Korea, and Indonesia also cultivate seaweed for culinary and industrial purposes.
  - **Non-Edible Resources:**
    - **Marine Pharmaceuticals:** Marine organisms from coral reefs in the Caribbean and the Great Barrier Reef in Australia have been studied for their potential pharmaceutical properties.
    - **Coral Reefs:** The Great Barrier Reef in Australia, the Coral Triangle in Southeast Asia, and the Caribbean Sea are renowned for their rich coral reef ecosystems.
    - **Pearls:** Pearl farming occurs in regions like the South Pacific, particularly in countries like Japan, China, and Indonesia, where pearl oysters are cultivated for their pearls.
    - **Seal and Whale Products:** Seal fur and whale blubber have historically been harvested in regions such as the Arctic and Antarctic, though regulations have limited commercial harvesting in recent years.
- **Non-Biotic Resources:**
  - **Energy Resources:**
    - **Oil and Gas:** Offshore oil and gas reserves are found in regions such as the North Sea, the Gulf of Mexico, and the Persian Gulf.
    - **Renewable Energy:** Tidal energy, Ocean Thermal Energy Conversion, Wave energy; Offshore wind farms are located in the North Sea, the Baltic Sea, and along the coastlines of countries like the United Kingdom, Germany, and Denmark. Wave energy converters are being tested in regions with strong wave action, such as the Pacific Northwest and the Atlantic coast of the United States.
  - **Mineral Resources:**
    - **Dissolves salts** , especially NaCl.
    - **Manganese Nodules (Poly-Metallic Nodules):** Found on the abyssal plains of the Pacific Ocean, particularly in the Clarion-Clipperton Zone. Recent discovery states that these nodules, also known as Polymetallic nodules, are an important source of “dark oxygen”, sustaining the life in the ocean bottom regions.
    - **Phosphorite Deposits:** Located off the coast of countries like Morocco, Peru, and the United States, where upwelling currents bring nutrient-rich waters to the surface.
- **Intangible Resources**

- **Regulating Services (Environmental Balance)**
  - **Climate regulation** : Oceans absorb **~90% of the excess heat** trapped by greenhouse gases.
  - **Carbon sink** : Absorb ~25–30% of anthropogenic CO_2; “blue carbon” ecosystems like mangroves and seagrass sequester it long-term. In fact, recent research has shown that the ocean's surface absorbs significantly more carbon dioxide than previously believed, especially the upper layers.
  - **Heat & circulation regulation** : Ocean currents redistribute heat, influencing monsoons and global climate.
  - **Weather & water cycle regulation** : Evaporation drives rainfall, storms, and global hydrological balance.
  - **Coastal protection** : Coral reefs, mangroves, wetlands reduce storm surges, erosion, and flooding.
- **Cultural & Recreational Services**
  - **Tourism & recreation** : Beaches, diving, surfing, cruises fuel coastal economies.
  - **Cultural & spiritual values** : Sacred seascapes, heritage of coastal communities.
  - **Education & research** : Source of marine knowledge, climate science, biodiversity studies.
  - **Aesthetic & inspiration** : Oceans inspire literature, art, and human imagination.
- **Supporting Services (Underlying Systems)**
  - **Oxygen production** : Phytoplankton generate **50–60% of Earth’s oxygen** .
  - **Biodiversity hotspots** : Coral reefs, estuaries, deep-sea vents harbor immense species diversity.
  - **Nutrient cycling** : Recycle nitrogen, phosphorus, and carbon essential for global food webs.
  - **Habitat provision** : Support breeding, nursery, and migratory grounds for marine life.

⚡ **Quick revise** Resources Fold ▲

Resources: *Biotic* – fish (tuna, salmon – Pacific; cod – N. Atlantic; sardines – Mediterranean, Peru), shellfish, seaweed (Japan nori), marine pharma, pearls. *Abiotic* – offshore oil and gas (North Sea, Gulf of Mexico, Persian Gulf), tidal/OTEC/wave/offshore wind, **dissolved salts (NaCl)**, **manganese nodules** (a source of "dark oxygen"), phosphorite (Morocco, Peru, USA). *Services:* oceans absorb **~90%** of excess heat and **25–30%** of human CO_2 (blue carbon: mangroves, seagrass); **phytoplankton make 50–60% of Earth's oxygen**.

PYQ 2013 (ans C) – fishing grounds: best where **warm and cold currents meet** (plankton mixing; e.g. **Newfoundland banks, Sea of Japan**). Other helpers: coastal upwelling, indented coasts, nearby forests for boats, rugged land that pushes people to fishing, temperate water below 20 °C.

LAW OF SEA



- The Law of the Sea refers to a **body of international law that governs the use of the world's oceans and seas.** It encompasses rules and principles regarding maritime boundaries, navigation, exploitation of marine resources, conservation, and protection of the marine environment.
- **United Nations Convention on the Law of the Sea (UNCLOS):** Adopted in **1982** , UNCLOS is the primary international legal framework governing the rights and responsibilities of states in maritime zones. It establishes rules on maritime boundaries, navigation, marine resources, and the protection of the marine environment.
- **Components:**
  - **Territorial Sea:** A belt of water adjacent to a coastal state's land territory, usually extending up to **12 nautical miles (22.2 kilometres**) from the baseline.
  - **Contiguous Zone** : A zone adjacent to the territorial sea, extending up to **24 nautical miles (44.4 kilometres)** from the baseline, where a coastal state may exercise limited control to prevent and punish violations of its customs, fiscal, immigration, or sanitary laws and regulations within its territory or territorial sea.
  - **Exclusive Economic Zone (EEZ):** An area extending up to **200 nautical miles (370.4 kilometres)** from a coastal state's baseline, within which the state has sovereign rights over the exploration and exploitation of natural resources.
  - **High Seas:** Areas of the ocean beyond the jurisdiction of any state, where all states enjoy freedoms such as navigation, overflight, fishing, and scientific research.

⚡ **Quick revise** Deposits, dead zones, resources, law Fold ▲

> ⚓ **UNCLOS 1982 – "12 – 24 – 200": Territorial sea 12 nm** (22.2 km) · **Contiguous zone 24 nm** (44.4 km; customs, fiscal, immigration, sanitary laws) · **EEZ 200 nm** (370.4 km; sovereign rights over resources) · **High seas** beyond (freedom of navigation, overflight, fishing, research). *Think: a dozen → a day → a double century.*

PYQ

Q.1) The most important fishing grounds of the world are found in the regions where: (2013)

(a) Warm and cold atmospheric currents meet

(b) Rivers drain out large amounts of fresh water into the sea

(c) Warm and cold oceanic currents meet

(d) Continental shelf is undulating

Ans.- C Reasons that determining Fishing grounds

- The location of major world fishing grounds is governed by many factors:
  - Abundance of plankton
- coastal waters are beneficial to fish in more than one way—
- There is plentiful food supply from the erosion of continents, which provide mineral and organic matter.
- There is enough sunlight penetration, which is critical to the photosynthesis process required by plants.
- Areas where warm and cold currents meet tend to have high biological productivity, because plankton growth is encouraged by the mixing of cold currents and warm currents.
- Some of the world’s most productive fishing grounds are located here, where warm and cold currents converge.
- Examples of these phenomena are the Newfoundland 'banks' and the Sea of Japan.
  - In areas of coastal upwelling high rates of primary production takes place. Conventional mixing provides plenty of oxygen to sustain life in coastal zones as well as assuring a favourable dispersion of nutrients.
  - Broken coastline with numerous indentations provides excellent berthing facility.
  - Accessible forest resources, which favour the manufacture of fishing boats, jetties, etc.
  - **The character of land:** Rugged land, which hinders the development of other activities like agriculture and forces the people to go for fishing.
  - **Climatic condition:** Temperate climate with a temperature of less than 20°C is the best region for marine life.

Recall sheet

Q	A
Ocean average depth	3,800 m
Shelf / slope gradient	1–2° / 3–6°
Trench depth	> 6,000 m (Mariana)
Equatorial vs 45° mean temperature	26.7 / 12.7 °C

Q	A
Saltiest water bodies listed	Lake Van 330 > Dead Sea 240 > Great Salt Lake 220
Ocean pH now	8.04 (2024)
Warm boundary currents lie on which side?	Western edges
Only sea with no coastline	Sargasso Sea
Why counter-current flows east	Convergence/pile-up of the two equatorial currents
Tide repeats after	12 h 26 min
Spring tide happens at	New & full moon (syzygy)
Tidal bore in India	Hooghly
Tsunami speed in deep ocean	500–1,000 km/h
Coral temperature / depth / salinity	20–29 °C / < 50 m / 32–40 ppt
UNCLOS zones	12 · 24 · 200 nm
Share of Earth's oxygen from phytoplankton	50–60%

Traps: "Spring tide = in spring" ✘ · "Counter-current is due to Earth's rotation" ✘ · "Salinity is highest at the equator" ✘ · "Density is driven more by salinity" ✘ (temperature) · "Tsunamis are tall in deep ocean" ✘ · "Rivers/lakes hold most freshwater after ice" ✘ (groundwater) · "Atolls form on continents" ✘ (sinking volcanic islands).

Connects to: Trenches, guyots, hotspots → *Geomorphology*; Coriolis, trade winds, westerlies, cyclones, El Niño-type warming → *Climatology/Indian climate*; Ocean resources, ports, fisheries → *Economic Geography*; Palk Strait, Chilika, Gulf of Kutch, Cape Comorin → *Indian Geography*.

GEOGRAPHY · GS FOUNDATION · SUBJECT 4

Indian Geography

Handout + mind maps: no class slides | Handout pages 78

VERY HIGH · 40 Prelims

The whole Indian Geography handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 29 mind-map pages are in Chapter 10.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: VERY HIGH (largest PYQ bank of the six).** The handout prints **40 Prelims PYQs** here: **physiography 15** (2012, 2014×2, 2015, 2016, 2017×2, 2018×2, 2020, 2022, 2023; three carry no year) · **drainage 12** (2011, 2013, 2014, 2015, 2016×2, 2017, 2021×2, 2022, 2023; one undated) · **climate 3** (2011, 2017, one undated) · **forests & soils 10** (2011, 2013, 2015×3, 2016, 2021, 2023×2, one undated). *Counted from the handout text.*
- **Pattern:** *"Which of these pairs/rivers/peaks/states?"* Map-and-list questions: tributaries of Godavari/Brahmaputra, which hills belong to which range, which state has what, which forest has which tree. **Facts you must be able to place on a map.**
- **Mains quoted:** none quoted in the material. *Hooks:* Western Disturbances, cloudbursts, urban flooding, ELLA, river-linking, mangroves, social forestry – all written as current-affairs notes.
- *No teacher ink in this PDF; priority = PYQ count in the handout.* The handout also has **image-only pages** (peaks, glaciers, passes, waterfalls, lakes) – I read them and kept them below as compact tables.

1 · LOCATION OF INDIA

- With an **area of 32,87,263 km2** , India is the **seventh largest country** in the world.
- India accounts for about **2.4 percent** of the **total surface area of the world** .
- **North-south extent** of the mainland, from Indira Col in Kashmir to Kanyakumari is **3,214 km** .
- **East-west width** from the Rann of Kutch to Arunachal Pradesh is **2,933 km** .
- The **southernmost point of the country is the Pygmalion Point, or Indira Point** is located at 6° 45' N latitude.

- **Entirely Northern Hemisphere.**
- **8° 4′N (6° 45′ for islands) and 37°6′N latitudes; and 68° 7′E and 97° 25′E longitudes.**
- **Standard Meridian 82° 30′ E passes through 5 states** (U C My Old Aunt) **UP, CG, MP, Odisha, Andhra Pradesh.**
- **East-West time difference is nearly 2 hrs** . (A difference of 1° longitude will make a difference of 4 minutes in time. ~30 x 4 = ~120 minutes or ~2 hours)
- **Chai Bagan time** : Introduced by the British to increase productivity in NE India. 1 hour ahead of IST.
- The Tropic of Cancer passes through the middle of the country dividing it into two latitudinal halves. The area to the north of the Tropic of Cancer is nearly twice the area that lies to the south of it.
- **Tropic of Cancer states: Gujarat, Rajasthan, M.P., CG, Jharkhand, West Bengal, Tripura, Mizoram.**

⚡ **Quick revise** Location and neighbours Fold ▲

🚂 **Standard Meridian 82°30′E passes through 5 states – "U C My Old Aunt":** **U**ttar Pradesh · **C**hhattisgarh · **M**adhya Pradesh · **O**disha · **A**ndhra Pradesh. *(Handout mnemonic.)*

Fact	Number
Area	**32,87,263 km²** · **7th** largest · **2.4%** of world area
North–south (Indira Col → Kanyakumari)	**3,214 km** · East–west (Rann of Kutch → Arunachal) **2,933 km**
Latitudes / longitudes	**8°4′N** (6°45′ for islands) to **37°6′N** · **68°7′E** to **97°25′E** · wholly **Northern Hemisphere**
Southernmost point	**Indira Point / Pygmalion Point, 6°45′N**
Time	1° longitude = **4 min** · east–west gap ≈ **2 hrs** · **Chai Bagan time** = 1 hr ahead of IST (British, NE India)
Tropic of Cancer	splits India in two; **north ≈ 2× south**; **8 states:** Gujarat, Rajasthan, MP, CG, Jharkhand, West Bengal, Tripura, Mizoram
Borders	land **15,106.7 km** through **17 states** · coastline **11,098 km**, **13** states/UTs · **longest** = Bangladesh · **shortest** = Afghanistan (Wakhan)
Landlocked / no border states	Telangana, MP, CG, Jharkhand, Delhi, Chandigarh, Haryana have **no international border or coastline** – all others are "frontline"

2 · INDIA AND ITS NEIGHBOURS

- India has 15106.7 Km of **land border** running through **17 States.**

- India has a **coastline of 11,098 km** , touching 13 States and Union Territories (UTs).



State / UT	New Length	Old length
1. Gujarat	2340.62	1217.7 ↑
2. Tamil Nadu	1068.69	906.9 ↑
3. Andhra Pradesh	1053.07	973.7 ↑
4. Maharashtra	877.97	652.6 ↑
5. West Bengal	727.02	157.5 ↑
6. Kerala	600.15	569.7 ↑
7. Odisha	574.71	464.4 ↑
8. Karnataka	343.3	280 ↑
9. Goa	193.95	101 ↑



State/ UT	New Length	Old Length
10. Andaman & Nicobar	3083.5	1962 ↑
11. Lakshdweep	144.8	132 ↑
12. Daman & Diu Dadra & Nagar Haveli	54.38	42.5 ↑
13. Puducherry (includes all 4 parts)	42.65	47.6 ↓

- **Barring Telangana, Madhya Pradesh, Chhattisgarh, Jharkhand, Delhi, Chandigarh, and Haryana** , **all other States** in the country **have one or more international borders or a coastline** and can be regarded as frontline States from the point of view of border management.
- India's **neighbours** include (clockwise): **Pakistan, Afghanistan, China, Nepal, Bhutan, Bangladesh, Myanmar, Sri Lanka, Maldives, and Indonesia.**
- India's **longest border is with Bangladesh while the shortest border is with Afghanistan (Wakhan Border)** .

- States/UTs sharing a border with each land neighbor:
  - **Pakistan** : Shares borders with Jammu and Kashmir, Ladakh, Punjab, Rajasthan and Gujarat.
  - **Afghanistan:** Ladakh
  - **China** : Shares borders with Ladakh, Himachal Pradesh, Uttarakhand, Sikkim, and Arunachal Pradesh.
  - **Nepal** : Shares borders with Uttar Pradesh, Uttarakhand, Bihar, Sikkim, and West Bengal.
  - **Bhutan** : Shares a border with Assam, Sikkim, Arunachal Pradesh, and West Bengal.
  - **Bangladesh** : Shares borders with West Bengal, Assam, Meghalaya, Mizoram, and Tripura.
  - **Myanmar** : Shares borders with Arunachal Pradesh, Nagaland, Manipur, and Mizoram.

⚡ **Quick revise** Neighbours clockwise Fold ▲

Neighbours clockwise: Pakistan, Afghanistan, China, Nepal, Bhutan, Bangladesh, Myanmar, Sri Lanka, Maldives, Indonesia.

Neighbour	Bordering Indian states/UTs
Pakistan	J&K, Ladakh, Punjab, Rajasthan, Gujarat
Afghanistan	Ladakh
China	Ladakh, HP, Uttarakhand, Sikkim, Arunachal
Nepal	UP, Uttarakhand, Bihar, Sikkim, West Bengal
Bhutan	Assam, Sikkim, Arunachal, West Bengal
Bangladesh	West Bengal, Assam, Meghalaya, Mizoram, Tripura
Myanmar	Arunachal, Nagaland, Manipur, Mizoram

PYQs on location. *2015 (ans D):* easternmost/westernmost states = **Arunachal Pradesh and Gujarat**. *2018 (ans A):* the city on the longitude closest to Delhi = **Bengaluru** (both ≈ 77°E). *2017 (ans B):* Kohima → Kottayam by road needs **7 states** (Nagaland, Assam, WB, Odisha, AP, TN, Kerala).

3 · PHYSIOGRAPHY OF INDIA

India has 6 physiographic units:

⚡ **Quick revise** Six physical divisions (the "map" of India)

Fold ▲

> 🏔️→🏝️ **Walk top to bottom: "Mountains, Plains, Desert, Plateau, Coasts, Islands"** – *"My Parents Do Play Cricket Indoors"*. These are the handout's six units.

4 · The Northern and Northeastern Mountains

- The Northern Plains
- The Indian Desert
- Peninsular Plateau
- Coastal Plains
- The Islands

THE NORTHERN AND NORTH-EASTERN MOUNTAINS:

- **Origin:** Convergence of Indian and Eurasian Plates. The Sediments of the Tethys sea got folded, forming the Himalayas.
- They include the **Trans-Himalayas (Karakoram, Ladakh & Zaskar ranges), Himadri, Himachal, and Shivaliks, and the Purvanchals** .
- Created due to the **convergence of the Indian and Eurasian Plates.** (Closing of the Tethys Sea).
- **Multiple episodes of upliftment** have given rise to **parallel ranges** .
- **About the Mountains:**
  - **2500 km spread** ; **Width varies** , with 500 km width in Kashmir and 200 km width in Arunachal.
  - The main Himalayas extend from **Nanga Parbat peak (Indus Gorge) in Ladakh to Namche Barwa peak (Brahmaputra Gorge)** in Arunachal Pradesh (syntaxial bends). Post that, the Purvanchals are present.
  - **Syntaxial Bends** are knee joint bends.

are knee

13 Indian
Territories
mu and
Ladakh,
Himachal
Arunachal
Manipur,
Mizoram,
Sikkim,
and West

to South;
North.
raphy)
ography.
s:
er Himalayas, Central Himalayas.
6100 m;

as:
esser Himalayas
3500–4500 m
haula Dhar (southernmost Himachal range), Kumaon & Mussoorie, ranges.
esent **between Pir Panjal & Zaskar**): These formations consist of , silt, sand, and gravel, deposited by ancient rivers and lakes during ne epoch. They are famous for the **growth of Zafran** (the best quality world), **walnuts, and almonds.**



- Spread across **13 Indian States/Union Territories** (namely Jammu and Kashmir, Ladakh, Uttarakhand, Himachal Pradesh, Arunachal Pradesh, Manipur, **Meghalaya,** Mizoram, Nagaland, Sikkim, Tripura, Assam and West Bengal)
- Steep gradient to South; Gentler in North.

(hogback topography)

- Ridge valley topography.
- **Great Himalayas:**
  - **Himadri,** inner Himalayas, Central Himalayas.
  - **Avg. height: 6100 m** ;
  - Tallest peaks
- **Middle Himalayas** :
  - **Himachals, Lesser Himalayas**
  - Avg. height: 3500-4500 m
  - **Pir Panjal, Dhaula Dhar (southernmost Himachal range), Kumaon & Mussoorie, Mahabharata ranges.**
  - **Karewas** (present **between Pir Panjal & Zaskar**): These formations consist of layers of clay, silt, sand, and gravel, deposited by ancient rivers and lakes during the Pleistocene epoch. They are famous for the **growth of Zafran** (the best quality saffron in the world), **walnuts, and almonds.**
  - **Bugyals** : They are **high-altitude grasslands in the Himachals** , known for their lush greenery, colourful flowers, and grazing pastures.

- The **transhumance activity takes place here** , which is the seasonal movement of herders and their livestock between different elevations in search of grazing pastures.
- During the summer months, herders from lower valleys bring their livestock, such as sheep, goats, and cattle, to graze on the lush grasslands of bugyals situated at higher altitudes. These bugyals provide abundant fodder and grazing opportunities during the warmer months when lower elevations may experience heat and scarcity of forage. In the winter, as bugyals become snow-covered and inaccessible, herders return with their livestock to lower elevations where they can find shelter and fodder.
- Important **tribes** involved in this: **Bakarwals, Gaddis, Bhotiyas and Drokpas.**
  - **Outer Himalayas: Shivaliks**
    - Almost unbroken from Himachal to Arunachal except a **gap of 80-90km occupied by Teesta valley.**
    - Jammu Hills (Jammu), Dafla, Miri, Abor (Arunachal); Dhang and Dudhwa in Uttarakhand; Churia Ghat in Nepal
    - **Duars:**
- Location: **Foothills of the Eastern Himalayas (Himachals) in Northeast India** (Assam, Arunachal Pradesh etc.)
- **Floodplains** in foothills (North of Brahmaputra valley) with dense vegetation and wildlife habitats.
- Soil: Alluvial soil, fertile and good for agriculture.
  - **Duns** : (Sometimes Doons)
- Location: **Valleys between the Shivalik Hills and the Himachals in Wastern Himalayas** (Himachal Pradesh, Uttarakhand, etc.)
- Longitudinal valleys with rivers and fertile lands.
- Examples: Dehradun, Patli Dun, Kota Dun, etc.
  - **Chos:**

- **Location: Foothills of the Shivalik Hills in the north-western part of Punjab** , India.
- **Seasonal streams or rivers** found primarily in the foothills of the Shivalik (Siwalik) range. Sandy riverbeds are characteristic.
- **Alternate names: Choes/Raos.**
- Soil: These regions are **less fertile compared to Duars** , may require irrigation for agriculture.
- **Trans Himalayas:**
  - Refers to the **northernmost part of the Himalayan range** , located beyond the main Himalayan range.
  - Comprises **rugged and high-altitude terrain** with stark landscapes, characterized by barren plateaus, deep valleys, and towering peaks.
  - They have high altitudes with a cold, dry climate. **Vegetation is sparse, and permanent human habitation is limited.**
  - Along with the Himalayas, they act as a **climatic barrier** , influencing weather patterns in Asia. They are also a **source of freshwater** through glaciers and rivers, and a region of cultural and spiritual importance for many communities.
  - **Karakoram** :
    - Located in the **UT of Ladakh,** extending into Afghanistan, Pakistan and China.
    - Home to **some of the world's highest peaks** , including K2 (Mount Godwin-Austen) and Broad Peak.

Ladakh (highest Indian Plateau):

    - Situated in the **eastern part of Ladakh Union territory** , adjacent to the Karakoram Range.
    - Known for its **high-altitude deserts,** deep gorges, and Buddhist monasteries.
  - **Zaskar** :
    - Lies to the **south of the Indus River in Ladakh** , running parallel to the Great Himalayas.
    - Across Ladakh, Himachal, and Uttarakhand.
    - Features rugged terrain with deep canyons, glaciers, and remote valleys **inhabited by the Zanskar people.**
  - **Kailash (entirely outside India):**
    - Located in the **western part of the Tibetan Plateau** , straddling the border between China and the Tibet Autonomous Region.
    - **Houses Mount Kailash** , a **sacred peak** revered by Hindus, Buddhists, Jains, and Bonos.
  - **Known for glaciers** (maximum in Karakoram).
- **Purvanchals:**
  - **Crescent shape** , almost **perpendicular** to the Himalayas, and **convex to West**
  - In the North to South direction, they are present from **Arunachal Pradesh to Mizoram.**
  - Its parts include:
    - **Mishmi & Patkai Bum:** Arunachal (Indo-Myanmar Boundary)

    - **Naga Hills** , **Kohima Hills**
    - **Manipur hills** (Loktak Lake)
    - **Barail hills** separate Manipur & Naga hills. Separate the Brahmaputra & Barak Basins.
    - **Mizo Hills** (earlier called Lushai Hills)
  - They have very rough terrain, and are full of dense forests, making the region challenging from a developmental perspective.

Regional Division of Himalayas:

- **Kashmir/Punjab/Himachal Himalayas (Westernmost Section):**
  - Location: Stretches **from the Indus River in the west to the Sutlej River in the east** .
  - Mountain Ranges: Himadri, Himachal and Shivaliks. Also, Trans-Himalayas.
  - Peaks: Nanga Parbat (8,126 m), K2 (8,611 m - part of the peak lies in Pakistan), Dhaulagiri I (8,167 m)
  - Valleys: Kashmir Valley, Kullu Valley, Kangra Valley
  - Rivers: Indus River, Jhelum River, Chenab River, Ravi River, Beas River, Sutlej River
  - Significance: This section is known for its scenic beauty, and diverse landscapes with valleys, meadows, and glaciers. It's also a region of cultural and historical significance, with Kashmir Valley being a major tourist destination.
- **Kumaon Himalayas (Central Section):**
  - Location: Lies **between the Sutlej River in the west and the Kali River** (eastern border with Nepal) in the east.
  - Mountain Ranges: Himadri, Kumaon Himalayas (Himachal), Shivaliks
  - Peaks: Nanda Devi (7,816 m - highest mountain entirely within India), Nanda Kot (7,818 m), Trishul (7,120 m)
  - Valleys: Valley of Flowers National Park
  - Rivers: Sutlej River, Yamuna River, Ganges River (originates from glaciers in this region)
  - Significance: This section is known for its high peaks, including Nanda Devi, and for housing pilgrimage sites like the source of the Ganges River. The Valley of Flowers National Park is a UNESCO World Heritage Site known for its vibrant alpine meadows.
- **Nepal Himalayas (Eastern Section):**
  - Location: Encompasses the central part of the Himalayas, stretching **from the Kali River in the west to the Teesta River** (eastern border with Bhutan) in the east.
  - Mountain Ranges: Himadri, Himachal and Shivaliks
  - Peaks: Mount Everest (8,848 m - highest peak in the world), Kanchenjunga (8,586 m - third highest peak in the world), Lhotse (8,516 m), Annapurna I (8,091 m)
  - Valleys: Kathmandu Valley
  - Rivers: Karnali River, Gandaki River, Kosi River

- Significance: This section boasts the highest peaks in the Himalayas, including Everest. It's a major center for mountaineering expeditions and adventure tourism.

Nepal is a landlocked country, and the Himalayas play a vital role in its geography and culture.

- **Assam Himalayas (Easternmost Section):**
  - Location: Lies **east of the Teesta River and extends into Bhutan, till Arunachal Pradesh and Purvanchals** in India.
  - Mountain Ranges: Himadri, Himachal and Purvanchals.
  - Peaks: Namche Barwa (7,756 m)
  - Valleys: Manas National Park
  - Rivers: Teesta River, Brahmaputra River
  - Significance: This section is known for its lush green valleys, diverse ecosystems, and abundant rainfall. Manas National Park is a UNESCO World Heritage Site known for its rich biodiversity.



	Peak	Characteristics
1.	Nanga Parbat	In Himadris, in Ladakh; Himalayas make a Syntaxial bend here. (>8000 m high)
2.	Godwin Austin (K2)	In Karakoram Range; 2nd highest peak of the world.
3.	Broad Peak, Gasherbrum 1, Gasherbrum 2.	All three are part of Gasherbrum group of peaks; all are higher than 8000 m.
4.	Nunkun	Nun in J&K; Kun in Ladakh; Part of Himadris
5.	Reo Purgyil	Highest Peak of H.P. Part of Zanskar Range
6.	Satopath	In Uttarakhand. Satopath Lake at its Foot
7.	Kamet	In Greater Himalayas Highest peak of Uttarakhand & of Zaskar Range
8.	Trishul	Group of 3 peaks (look like a Trishul / Trident)
9.	Nanda Devi	In Greater Himalayas. It is a UNESCO World Heritage Site. Also Nanda Devi National Park
10.	Karoh Peak	Part of Shivaliks. Highest peak of Haryana
11.	Amsot Peak	Part of Shivaliks. Highest peak of U.P.
12.	Someshwar Peak	Part of Shivaliks. Highest peak of Bihar
13.	Kunchenjunga	Highest Himalayan Peak in India. Part of Greater Himalayas. In Sikkim. Presence of Biosphere Reserve, National Park, India's First UNESCO mixed Heritage site
14	Sandakphu	Highest Peak of W.B. Part of Himachals
15	Tiger Hill	In Darjeeling. Known for Tea Cultivation Part of Himachal Range.
16	Kangto	Part of Greater Himalayas. Highest peak of Arunachal Pradesh.

	Peak	Characteristics
17	Dapha Bum	Highest Peak of Purvanchals. Located in Mishmi hills.
18.	Saramati	Highest peak of Nagaland.
19.	Mt. Esii / Iso	Highest peak of Manipur
20.	Phwangpui (Blue Mountain)	Highest peak of of Mizoram
21.	Betlingchhip (Thaidawr Tlang)	Highest peak of Tripura
22.	Nokrek	Highest peak of Garo Range. Also a National Park & a Biosphere Reserve.
23.	Shillong Peak	In Khasi hills. Highest peak of Meghalaya
24.	Tumjang / Sielkal / Thingtuibung	In Barail Range. Highest peak of Assam.
25.	Parasnath	Highest peak of Jharkhand. Important Jain pilgrimage site. Part of Chotanagpur Plateau
26.	Guru Shikhar	Highest peak of Aravallis, and of Rajasthan
27.	Gorakhnath	Highest peak of Gujarat. Also an imp. pilgrimage site for Hindus. Located in Girnar hills.
28	Dhupgarh	Highest peak of Satpuras and of M.P.
29.	Sadbhawna Shikhar (Goodwill Peak)	Highest peak of Vindhyas.
30.	Kalsubai	In Western Ghats. Highest peak of Maharashtra
31.	Mahabaleshwar	Part of Western Ghats. Origin region of Krishna river
32.	Sonsogor	In Western Ghats. Highest peak of Goa. Near trijunction of Maharashtra, Goa & Karnataka

Peak	Characteristics
33. Mullayanagari	Part of Western Ghats. Highest peak of Karnataka.
34. Kudremukh	Important peak for mineral resources. Also a Tiger Reserve & at National Park. In Karnataka. Source of Kaveri
35. Brahmagiri	
36. Doddabetta.	Highest peak of Tamil Nadu and of Nilgiris.
37. Anaimudi	Highest peak of Western ghats. Also highest peak of Kerala.
38. Agastyamalai	Part of Western Ghats. At Tamil Nadu - Kerala border.
39. Saddle Peak	Highest peak of Andaman & Nicobar
40. Mt. Harriet	Now known as Mt. Manipur
41. Bailadila	Important peak of Chattisgarh. Part of Deccan Plateau
42 Gaurlata	Highest peak of Chattisgarh. Part of Central Highlands.
43. Doliguta	Highest peak of Telangana. Part of Deccan Plateau
44. Mahendragiri	Imp. peak of Eastern Ghats.
45.	Highest peak of Odisha. Part of Eastern Ghats.
46. Arma Konda/ Jhindagada	Highest peak of Eastern Ghats & of Andhra Pradesh


IMPORTANT GLACIERS OF INDIA
Hunza
BATURA
Shimsal
Shigar
Yarkant
SKARMI
KHURDOPIN
BIAFO
BALTORO
Shigar
SIACHEN
Hispar
HISPAR
Nubra
Shyok
RIMO
CHONG KUMDAN
Shyok
Basha
CHOGO
Basna
LUNGMA
SHAFAT
DRANG DRUNG
Stod
Rupal
RUPAL
SONAPANI
Kutle Nala
NANGA PARBAT
BARASHIGRI
Chandra
GANGOTRI
Ganga
Feeds Indus system
Suru
CHORABARI
Mandakini
MILAM
Gori Ganga
PINDARI
KAMET
Pindari
Dhauliganga
YAMUNOTRI & BANDARPUNCH
Yamuna
ZEMU
Teesta
– Part of Tarim Basin
– Part of Indus Basin
– Part of Ganga Basin
– Part of Brahmaputra Basin


IMPORTANT PASSES OF INDIA
AGHIL
KARAKORAM
ZOJI LA
KHARDUNGLA
BURZIL
MIGLA
CHANG
BANIHAL
UMLING LA
SHIPKI LA
BARALACHA LA
SHINKUN LA
MANA
NITI
ROHTANG
LIPULEKH
YONGYAP
DIPHU
BUMLA
SELA
BOMDILA
NATHULA
JELEP LA
HALDIGHATI
(BURHANPUR) ASIRGHAT
CHAUKAN/ CHANKAN
PANGSAU
THALGHAT
BHORGHAT
CHORLA GHAT
PALGHAT
SHENKOTTA

	Mountain Pass	State/UT	Characteristic
1	Agzil	Ladakh	In Karakoram Range Historically used for trade. Military importance
2.	Karakoram	Ladakh	In Karakoram pass. Was part of ancient silk route
3.	Khardung la	Ladakh	connects Leh & Siachen
4	Mig la	Ladakh	world's highest motorable pass.
5.	Chang la	Ladakh	connects Pangong Tso with rest of Ladakh
6.	Umling La	Ladakh	2nd highest motorable road in the world.
7.	Zoji La	Ladakh & JnK	connects Ladakh and Kashmir valley. Often obstructed by snow
8.	Burzil La	JnK	Currently in PoK. Used for trading activities
9.	Banihal	JnK	Known for Jawahar Tunnel (NH 44)
10.	Shinkun La	Himachal Pradesh	Known for NPD Route (Nimmu - Padum - Darcha) All weather connectivity to Ladakh.
11.	Baralacha La	Himachal Pradesh	Known for Leh - Manali Highway
12.	Rohtang	Himachal Pradesh	Important for Leh connectivity. Atal Tunnel is under Rohtang Pass
13.	Shipki La	Himachal Pradesh	At India - China border. Sutlej enters India from here.

Mountain Pass	State/UT	Characteristic
14. Mana	Uttarakhand	Highest motorable pass of Uttarakhand
15. Niti	Uttarakhand	Northernmost pass of Uttarakhand
16. Lipulekh	Uttarakhand	Nepal claims it, but belongs to India. Used for Mansarovar Yatra.
17. Nathu La	Sikkim	Important for Indo-China trade. Used for mansarovar yatra.
18. Jelep la	Sikkim	Connects India & Bhutan
19. Bum la	Arunachal Pradesh	Used by Dalai Lama to enter India in 1959.
20. Sela	Arunachal Pradesh	Sela Tunnel is located here. Important for connecting Tawang with rest of Arunachal Pradesh.
21 Bomdila	Arunachal Pradesh	Helpful for Tawang connectivity. Used by India for the 1962 Indo-China war.
22. Yongyap	Arunachal Pradesh	Near the entry point of Dichang in India. British built the Sitwell Road to connect Myanmar & China through here.
23. Diphu	Arunachal Pradesh	At trijunction of India-China-Myanmar
24. Chaukan	Arunachal Pradesh	Important for Indo-Myanmar connectivity.
25 Pangsau	Arunachal Pradesh	Sitwell road also passes through here.

Mountain Pass	State/UT	Characteristic
26. Haldighati	Rajasthan	In Aravallis. Historically important for Battle between Maharana Pratap & Mughals.
27. Asirghat / Burhanpur Darra	Madhya Pradesh	In Satpuras. Known for Asirgarh fort. Key to Deccan. Connects Northern & Southern India.
28. Thalghat	Maharashtra	Known for Mumbai – Nashik Highway. In Western ghats
29. Bhorghat	Maharashtra	Known for Mumbai-Pune Highway. In Western Ghats
30. Chorla Ghat	Goa	Near Goa-Maharashtra-Karnataka trijunction. In Western Ghats.
31. Palghat / Palakkad	Kerala – Tamil Nadu	Widest gap in Western ghats. At Kerala – Tamil Nadu Border. Gateway of Monsoon in Tamil Nadu. Between Nilgiris in North & Annamalai in South.
32. Shenkotta	Kerala – Tamil Nadu	At Kerala – Tamil Nadu border. In Western Ghats.

NORTHERN PLAINS

Great depression to the South of the Himalayas was filled with alluvial deposits fr Indus, Brahmaputra, and Ganga, leading to the creation of these plains.



Expanse: **3200 km West to East.**

The Northern Plains are one of the most densely populated regions in the wo supporting a significant portion of India's agriculture.

The rivers provide water for irrigation, transportation, and hydropower.

The region is rich in history and culture, with major cities like Delhi, Varanasi, and Pa located here.

The plains are generally **flat with a slight slope from north to south.**

The climate varies from semi-arid in the west to humid subtropical in the east.

The region faces challenges like soil erosion, water pollution, and depletion groundwater resources.

Terminologies associated with these Plains:

- **Bhabher**: Alluvial cone deposits at the foothill region, where the rivers go underground, after the steep change of gradient, from the mountains to the



⚡ Quick revise 2.1 Northern & North-eastern mountains

Fold ▲

Origin: **Indian–Eurasian plates converge; Tethys sediments fold.** Multiple upliftment episodes → **parallel ranges**; **steep south, gentle north (hogback)**; ridge-valley topography. **2,500 km** long; width **500 km (Kashmir)** → **200 km (Arunachal)**. Main Himalaya runs from **Nanga Parbat (Indus gorge)** to **Namche Barwa (Brahmaputra gorge)** = the two **syntaxial (knee-joint) bends**; then the **Purvanchals**. Spread over **13** states/UTs.

Belt	Also called	Height	Notes
Trans-Himalaya	Karakoram, Ladakh, Zaskar	highest, cold dry	**maximum glaciers in Karakoram**; K2 (Godwin-Austen); **Ladakh = highest Indian plateau**; Zaskar south of the Indus; **Kailash entirely outside India**
Great Himalaya	**Himadri**, inner	avg **6,100 m**	tallest peaks

Belt	Also called	Height	Notes
Middle Himalaya	**Himachal**, Lesser	**3,500–4,500 m**	Pir Panjal, **Dhauladhar** (southernmost HP range), Kumaon, Mussoorie, Mahabharat; **Karewas** (clay-silt-sand-gravel of Pleistocene lakes, between Pir Panjal and Zaskar) → **saffron (zafran), walnuts, almonds**; **Bugyals** (high meadows)
Outer Himalaya	**Shiwaliks**	lowest	nearly unbroken, **gap of 80–90 km at Teesta valley**

🛌 **Foothill sleep-over, top to bottom:** mountain → **Bhabar** (stones – *rivers vanish underground*) → **Terai** (swamp – *rivers reappear*) → plains. **Duns** (Dehradun, Patli, Kota) = valleys between Shiwaliks and Himachals (west); **Duars** = floodplain foothills in NE; **Chos/Choes/Raos** = seasonal Shiwalik streams of Punjab (sandy beds).

- **Transhumance** (seasonal herding to bugyals): **Bakarwals, Gaddis, Bhotiyas, Drokpas.**
- **Purvanchals:** crescent, convex to west, perpendicular to Himalaya; **Mishmi & Patkai Bum, Naga, Kohima, Manipur hills (Loktak), Barail** (separates **Manipur & Naga hills**; divides **Brahmaputra and Barak basins**), **Mizo (Lushai)**. Rough, forested = hard to develop.

⚡ Quick revise 2.1 Northern & North-eastern mountains

Fold ▲

Regional Himalaya	Limits	Peaks / valleys (as printed)
Punjab/Kashmir/Himachal	Indus → Sutlej	Nanga Parbat **8,126 m** · K2 **8,611 m** · Dhaulagiri I **8,167 m** · Kashmir, Kullu, Kangra valleys
Kumaon (central)	Sutlej → Kali	**Nanda Devi 7,816 m** (highest peak *entirely in India*) · Nanda Kot **7,818 m** · Trishul **7,120 m** · Valley of Flowers (UNESCO) · Ganga source
Nepal (eastern)	Kali → Teesta	**Everest 8,848 m** · **Kanchenjunga 8,586 m** (3rd highest) · Lhotse 8,516 m · Annapurna I 8,091 m
Assam (easternmost)	Teesta → Arunachal/Purvanchals	**Namche Barwa 7,756 m** · Manas NP (UNESCO)

⚠️ **Handout alert:** Nanda Devi **7,816 m** vs Nanda Kot **7,818 m** (and the 2022 PYQ note says Nanda Devi **7,817 m**). Learn **~7,816** and, in the exam, rely on *"highest entirely in India"* rather than the digit.

⚡ Quick revise

2.2 Peaks, passes, glaciers (image pages – read for you) Fold ▲

🗺️ **"Highest peak of ..." = the favourite question. Remember by region walking west → east → south:**

Region	Peak → what the handout says
Trans/Himadri	**Nanga Parbat** – Ladakh Himadri, >8,000 m, syntaxial bend · **K2/Godwin-Austen** – Karakoram, 2nd highest in world · **Broad Peak, Gasherbrum I & II** – Gasherbrum group, all >8,000 m · **Nun** (J&K) and **Kun** (Ladakh) – Himadri
HP	**Reo Purgyil** – highest of HP, Zaskar Range
Uttarakhand	**Satopanth** (Satopanth Lake at foot) · **Kamet** – highest of Uttarakhand and of Zaskar · **Trishul** – group of 3 peaks like a trident · **Nanda Devi** – Greater Himalaya, UNESCO World Heritage (with its NP)
Shiwalik peaks	**Karoh** – highest of Haryana · **Amsot** – highest of UP · **Someshwar** – highest of Bihar
East Himalaya	**Kanchenjunga** – highest Himalayan peak *in India*, Sikkim, biosphere reserve + NP, **India's first UNESCO mixed heritage site** · **Sandakphu** – highest of **West Bengal**, Darjeeling, tea · **Tiger Hill** – Himachal range · **Kangto** – highest of **Arunachal**
Purvanchal / NE	**Dapha Bum** – highest of Purvanchals (Mishmi) · **Saramati** – Nagaland · **Mt Esii/Iso** – Manipur · **Phawngpui (Blue Mountain)** – Mizoram · **Betlingchhip (Thaidawr Tlang)** – Tripura · **Nokrek** – Garo, NP + biosphere reserve · **Shillong Peak** – Khasi, highest of **Meghalaya** · **Tumjang/Sielkal/Thingtuibung** – Barail, highest of **Assam**
Peninsula (north→south)	**Parasnath** – highest of **Jharkhand**, Jain pilgrimage, Chotanagpur · **Guru Shikhar** – highest of Aravallis **and Rajasthan** · **Gorakhnath** – highest of **Gujarat**, Girnar hills, pilgrimage · **Dhupgarh** – highest of Satpuras **and MP** · **Sadbhawna Shikhar (Goodwill Peak)** – highest of **Vindhyas** · **Kalsubai** – highest of **Maharashtra** (W. Ghats) · **Mahabaleshwar** – origin region of Krishna · **Sonsogor** – highest of **Goa** (near Maharashtra-Goa-Karnataka trijunction)
Western Ghats south	**Mullayanagiri** – highest of **Karnataka** · **Kudremukh** – minerals, tiger reserve, NP · **Brahmagiri** – source of Kaveri · **Doddabetta** – highest of **Tamil Nadu and Nilgiris** · **Anaimudi – highest of Western Ghats and Kerala** · **Agasthyamalai** – TN–Kerala border
Others	**Saddle Peak** – highest of **Andaman & Nicobar** · **Mt Harriet** – now **Mt Manipur** · **Bailadila** – Chhattisgarh, Deccan · **Gaurlata** – highest of **Chhattisgarh**, Central Highlands · **Doligutta** – highest of **Telangana** · **Mahendragiri** – important Eastern Ghats peak · *(No. 45, highest of Odisha, Eastern Ghats – name left blank in the handout)* · **Arma Konda / Jindhagada** – highest of **Eastern Ghats and Andhra Pradesh**

🔱 **Memory pegs: Trishul = trident** (3 peaks) · **Kamet** = *the "Kamet-ition winner" of Uttarakhand* · **Doddabetta = "Dodo-big-bat" of the Nilgiris** · **Anaimudi = "Anna" is the tallest in the**

Ghats' family (Kerala) · **Kalsubai** = Maharashtra's "kal-subah" (tomorrow morning) climb.

Glaciers (map page; colour = basin). *Tarim:* **Yarkant**, Skarmi · *Indus system (red):* Hispar, Chogo, Basha, Lungma, Rupal (Nanga Parbat), Hunza/Batura/Khurdopin/Biafo/Baltoro → **Shigar**; **Siachen, Rimo → Nubra/Shyok**; Chong Kumdan, Drang Drung → **Shyok/Stod**; Shafat; **Sonapani → Kutle Nala**; **Barashigri → Chandra**; Suru (Nanga Parbat side). *Ganga basin (orange):* **Gangotri → Ganga**, Chorabari → **Mandakini**, Milam → **Gori Ganga**, Pindari → **Pindari**, Kamet, Dhauliganga; **Yamunotri & Bandarpunch → Yamuna**. *Brahmaputra basin (blue):* **Zemu → Teesta**.



Handout map: glaciers and the basin each feeds (colour = basin).



Handout map: where the important lakes sit.

🚪 **Passes = the doors of the Himalaya.** North → east → south (State/UT, hook):

Pass	State/UT	Hook
Aghil	Ladakh	Karakoram; historic trade; military
Karakoram	Ladakh	ancient **silk route**
Khardung La	Ladakh	**connects Leh & Siachen**
Mig La	Ladakh	printed as **"world's highest motorable pass"**
Chang La	Ladakh	connects **Pangong Tso** to rest of Ladakh
Umling La	Ladakh	printed as 2nd highest motorable road
Zoji La	Ladakh –J&K	connects **Ladakh & Kashmir valley**; snow-blocked

Pass	State/UT	Hook
Burzil La	J&K (now PoK)	trade
Banihal	J&K	**Jawahar Tunnel (NH44)**
Shinku La	HP	**NPD route (Nimmu–Padum–Darcha)**, all-weather to Ladakh
Baralacha La	HP	**Leh–Manali highway**
Rohtang	HP	**Atal Tunnel under it**; Leh connectivity
Shipki La	HP	India–China border; **Sutlej enters India**
Mana	Uttarakhand	highest motorable pass of Uttarakhand
Niti	Uttarakhand	northernmost pass of Uttarakhand
Lipulekh	Uttarakhand	Nepal claims it, handout says belongs to India; Mansarovar Yatra
Nathu La	Sikkim	Indo-China trade; Mansarovar Yatra
Jelep La	Sikkim	connects India & Bhutan
Bum La	Arunachal	Dalai Lama entered India **1959**
Sela	Arunachal	**Sela Tunnel**; connects Tawang
Bomdila	Arunachal	Tawang connectivity; used in **1962** war
Yongyap	Arunachal	near entry of Dihang; British **Stilwell Road**
Diphu	Arunachal	tri-junction India–China–Myanmar
Chaukan, Pangsau	Arunachal	Indo-Myanmar; Stilwell Road passes Pangsau
Haldighati	Rajasthan	Aravallis; Maharana Pratap vs Mughals
Asirgarh / Burhanpur	MP	Satpuras; Asirgarh fort; **key to the Deccan** (north–south link)
Thalghat	Maharashtra	**Mumbai–Nashik** highway (Western Ghats)
Bhorghat	Maharashtra	**Mumbai–Pune** highway
Chorla Ghat	Goa	near Goa–Maharashtra–Karnataka junction
Palghat (Palakkad)	Kerala–TN	**widest gap in Western Ghats**; **gateway of monsoon to Tamil Nadu**; between Nilgiris (north) and Annamalai (south)
Shenkotta	Kerala–TN	Western Ghats gap

5 · NORTHERN PLAINS

- **Great depression to the South of the Himalayas was filled with alluvial deposits** from **Indus, Brahmaputra, and Ganga** , leading to the creation of these plains.
- Expanse: **3200 km West to East.**
- The Northern Plains are one of the most densely populated regions in the world, supporting a significant portion of India's agriculture.
- The rivers provide water for irrigation, transportation, and hydropower.
- The region is rich in history and culture, with major cities like Delhi, Varanasi, and Patna located here.
- The plains are generally **flat with a slight slope from north to south.**
- The climate varies from semi-arid in the west to humid subtropical in the east.
- The region faces challenges like soil erosion, water pollution, and depletion of groundwater resources.
- **Terminologies associated with these Plains** :
  - **Bhabher** : Alluvial cone deposits at the foothill region, where the rivers go underground, after the steep change of gradient, from the mountains to the plains.



  - **Terai** : Swampy region after the Bhabher region, where the rivers re-emerge.
  - **Bhangar** : Coarser, older alluvium found in the floodplain region farther away from the river channel. It is less fertile than the newer alluvium. It consists of certain congregations of grains, known as Kanker particles.
  - **Khadar** : The newer alluvium, found in the nearby floodplain regions of the rivers. They are renewed almost every year. That is why, they are extremely fertile.
  - **Reh/Kallar** : Barren saline drier areas found in UP, Haryana. They are extending beyond their natural zone due to bad agricultural practices, like excessive irrigation.
  - **Bhur:** an elevated piece of land along the Ganga. Esp in Ganga Yamuna doab.
  - **Bagar:** Semi-arid plain starting from the foothills of Aravallis, in Rajasthan.
  - **Rohi:** patches of fertile land in the Bagar
  - **Doab** : Land between 2 rivers. Examples: Bari, Bist, Rechna, Chaj
  - **Betlands** : Khadar plains of Punjab
- Includes Rajasthan plains, Punjab Haryana plains, Ganga plains, Brahmaputra plains.
- **Rajasthan Plains:**

- Location: **Westernmost part** of the Northern Plains, including the Rajasthan bagar, which extends into the Thar Desert.
- Characteristics: **Semi-arid climate, saline lakes** due to high evaporation

- **Punjab Haryana Plains:**
  - Location: **West-central part, bordering the Indus River in the west and the Yamuna River in the east** .
  - **Bari and Bist doab** .
  - Characteristics: **Fertile** land, ideal for agriculture (wheat, rice, cotton), network of canals for irrigation.
- **Ganga Plains:**
  - Location: The **most extensive** part, stretching from Yamuna River eastwards to Bangladesh.
  - Parts: **Rohilkhand, Awadh, Mithila, Magadha and Bengal plains** .
  - Characteristics: **Highly fertile** land due to Ganga-Yamuna Doab, densely populated, diverse agriculture (wheat, rice, sugarcane, pulses), home to the Ganges River Delta.
- **Brahmaputra Plains:**
  - Location: **Easternmost part** , bordering Bangladesh.
  - Characteristics: Riverine islands (Point bars), also called **chars** , prone to flooding, lush green due to high rainfall, Brahmaputra River Delta Sundarbans. These plains, in the foothills, are also known for known for tea plantations.

⚡ Quick revise 2.3 Northern Plains

Fold ▲

- **Great depression south of the Himalaya filled by Indus, Ganga, Brahmaputra alluvium. 3,200 km** long; one of the most densely populated regions; semi-arid in the west → humid subtropical in the east.

> 🧱🆕 **Old Bhangar = "Bhang-ar" (bhang, old & coarse, kankar). New Khadar = "Khad" (manure) – renewed every year, super fertile.**

Term	Meaning
Bhabar	alluvial cones at foothills; rivers vanish
Terai	swampy; rivers re-emerge
Bhangar	older coarse alluvium, away from channel, **kankar** nodules, less fertile
Khadar	newer alluvium near rivers, renewed yearly, very fertile
Reh/Kallar	barren saline patches (UP, Haryana), spreading from over-irrigation
Bhur	elevated land along the Ganga (Ganga–Yamuna doab)
Bagar	semi-arid plain from Aravalli foothills, Rajasthan · **Rohi** = fertile patches in it
Doab	between two rivers – **Bari, Bist, Rechna, Chaj**

Term	Meaning
Betlands	Khadar plains of Punjab

Four sections: *Rajasthan plain* (bagar, saline lakes) · *Punjab–Haryana* (Bari and Bist doab, canals) · *Ganga plain* (Yamuna eastward; **Rohilkhand, Awadh, Mithila, Magadha, Bengal**; delta) · *Brahmaputra plain* (**chars/point bars**, floods, tea, Sundarbans delta).

6 · THE INDIAN DESERT: THAR

- The Thar Desert, also known as the Great Indian Desert, is a vast arid region straddling the northwestern part of the Indian subcontinent.
- Created due to a combination of the Sub-Tropical High Pressure belt, and this location being in the rainshadow zone of the Aravallis.
- It is often considered an **extension of the Rajasthan Plains** .
- Covers an area of approximately **200,000 square kilometers** .
- About **85% of the desert lies in India,** with the remaining 15% in Pakistan.
- In India, it spreads across parts of **Rajasthan, Gujarat, Punjab, and Haryana** .
- Bordered by the irrigated Indus River plain to the west.
- The Punjab Plain flanks it to the north and northeast.
- The Aravalli Range acts as a barrier to the southeast.
- The Rann of Kutch lies to the south.
- Locally, it is called **Marusthali.**
- The desert landscape is dominated by:
  - **Sand dunes** : These are constantly shifting due to wind action, with some reaching heights of over 150 meters (500 ft).
  - **Sandy plains** : These plains separate the dune fields. Here the Indira Gandhi Canal Project has provided the irrigational facilities, providing people with a source of employment.
  - **Low barren hills** : Locally known as **bhakars** , these rise abruptly from the surrounding plains.
  - The landscape is punctuated by intermittent rivers, seasonal streams (known as "Rivulets"), and saline lakes (such as the Sambhar Lake, Didwana, etc.).
- The Thar Desert acts as a **natural border between India and Pakistan.**
- It's a source of salt, limestone, and gypsum.

⚡ **Quick revise**

Sub-tropical high pressure + Aravalli rain-shadow — Fold ▲

Sub-tropical high pressure + Aravalli rain-shadow. ~2 lakh km²; 85% India, 15% Pakistan (Rajasthan, Gujarat, Punjab, Haryana). Local name **Marusthali**. Features: **dunes (some >150 m)**, **sandy plains** (Indira Gandhi Canal), low barren hills **bhakars**, seasonal "rivulets", **saline**

lakes – Sambhar, Didwana. Source of **salt, limestone, gypsum**. Indus plain on west, Punjab plain north-east, Aravallis south-east, Rann of Kutch south.

7 · PENINSULAR PLATEAU OF INDIA

- It is the **oldest and the largest** physiographic feature of India.
- It is in the shape of an **irregular triangle** .
- An ancient landmass formed by the breakup of Gondwanaland.
- Due to its age, it is **rich in many metallic minerals** .
- The **Deccan Traps** are a part of the Peninsular Plateau.
- The **Meghalaya Plateau part of this** , which was separated from it during the Indian plate's movement North into the Eurasian plate.
- Separation by **Rajmahal-Garo Gap/Malda Gap/Ganga Suture Zone.**
- Divided into the **Central Highlands and the Deccan Plateau** , separated by the **Narmada Rift.**

Parts of the Peninsular Plateau:

- **Mewar/Marwar Uplands:**
  - Location: Northwestern part of the Plateau, in Rajasthan, near Aravallis.
  - Characteristics: Arid and semi-arid climate with scarce rainfall, supports scrub vegetation and hardy desert plants. The Thar Desert borders its western flank.
- **Malwa Plateau:**
  - Location: From Aravallis in NE, to Bundelkhand Plateau in East and Vindhyas in South. Contain Chambal Badlands.
  - Characteristics: Known for its fertile black soil (regur), ideal for cotton cultivation. The network of rivers like Narmada and Chambal provides water resources for agriculture.vegetation.
- **Bundelkhand Uplands:**
  - Location: Between the Vindhya Range and the Ganga Plains, East of Malwa Plateau, covering parts of Madhya Pradesh and Uttar Pradesh.
  - Characteristics: Rugged terrain with a mix of hills, valleys, and plateaus. The climate supports mixed deciduous forests with trees like sal, teak, and kheria.
- **Baghelkhand:**
  - Location: Northern part of the Deccan Plateau, in Madhya Pradesh and Chhattisgarh.
  - Characteristics: Hilly terrain with dense forests of sal, teak, and bamboo. The region is rich in mineral resources, particularly coal.
- **Eastern Highlands:**

- Chotanagpur Plateau: (Comprises Hazaribagh Plateau, Ranchi Plateau, Rajmahal Hills)
- Location: Eastern India, covering parts of Jharkhand, Chhattisgarh, Odisha, and West Bengal.
- Characteristics: Rolling hills with a network of rivers. Dense forests with sal trees dominate the landscape. The region is known for its mineral wealth, including coal and mica.

- **North-Eastern Tip: Meghalaya** (Garo, Khasi, Jaintia, Mikir and Rengma Hills):
  - Location: Northeastern tip of the plateau, bordering Bangladesh, in Meghalaya and Assam.
  - Characteristics: Home to Mawsynram, the wettest place on Earth. The region receives heavy rainfall, supporting lush green forests with diverse flora and fauna. The region is known for its unique tribal cultures.
- **Deccan Plateau:**
  - Location: Largest and central part of the Peninsular Plateau, covering Chattisgarh, Maharashtra, Karnataka, Telangana, and Andhra Pradesh.
  - It's called Regur in Maharashtra and Malnads in Karnataka.
  - Include the Deccan Traps.
  - Characteristics: A vast, relatively flat tableland with an average elevation of 600 meters. The landscape varies from forests and grasslands to rocky outcrops. The Deccan Traps, lava formations from volcanic eruptions millions of years ago, are a distinctive feature.
  - **Chhattisgarh Plateau & Plains:**
    - Location: Eastern part of the Deccan Plateau, in Chhattisgarh. Includes the **Dandkarnya and Bastar Plateaus.**
    - Also has fertile **plains formed by the Mahanadi River and its tributaries.** The fertile soil supports rice cultivation, making it a vital agricultural region.
- **Important Hills Ranges:**


MOUNTAIN RANGES OF INDIA
KARAKORAM
LADAKH
ZASKAR
HIMADRI
HIMACHAL
SIWALIK
ARAVALLIS
VINDHYAS
KAIMUR
RAJMAHAL
MAIKALA
MAHADEO
RAJPIPLA
GWALIGARH
SATMALA
AJANTA
GARHJAT
BALAGHAT
HARISH-CHANDRA
SAHYADRIS
EASTERN GHATS
ERAMALLA
NALLAMALLA
PALKONDA
VELIKONDA
NAGARI
JAVADI
SHEVROY
PANCHAIMALAI
SIRUMALAI
BABA BUDAN
NILGIRI
ANNAMALAI
PALANI
CARDAMOM
HIMACHAL
① PIR PANJAL
② DHAULADHAR
③ NAG TIBBA
④ MUSSOORIE HILLS
⑤ KUMAON
SIWALIKS
④ DAFLA
⑤ MIRI
⑥ ABOR
PURVANCHALS
ⓓ MISHMI
ⓔ PATKAI BUM
ⓕ NAGA HILLS
ⓖ EAST & WEST MANIPUR HILLS
ⓗ MIZO (LUSHAI) HILLS
ⓘ JAMPUI HILLS
ⓙ BARAIL HILLS
MEGHALAYAN PLATEAU
I GARO
II KHASI
III JAINTIA
IV MIKIR
V RENGMA

Mountain Ranges of India — Highest Peak

1. Karakoram → Ladakh ; Pakistan, India, China ; K2
2. Ladakh → Ladakh ; India, China
3. Zanskar → Ladakh, Himachal Pradesh, Uttarakhand ; India & China ; Mt Kamet
4. Himalayas ; Mt Everest
   - ↳ Nanga Parbat to Namche Barwa
   - ↳ J&K, Ladakh, HP, Uttarakhand, Sikkim, West Bengal, Arunachal Pradesh
   - ↳ Himadri (Greater Himalayas)
   - ↳ Himachal (Lesser Himalayas) : Dissected by rivers
   - ↳ Shivaliks (Outer Himalayas) : Starts from Potwar Plateau
5. Purvanchals : Mt Dapha
   - ↳ Mishmi, Patkai Bum → Arunachal
   - ↳ Naga Hills → Nagaland
   - ↳ Manipur Hills → Manipur
   - ↳ Barail Hills → Assam, Nagaland, Manipur
   - ↳ Mizo Hills → aka Lushai hills, Mizoram
   - ↳ Jampui Hills → Tripura

   Purvanchals extend into Myanmar as Arakan Yoma
6. Aravallis : Gujarat, Rajasthan, Haryana, Delhi ; Guru Shikhar
7. Vindhyas : Gujarat, M.P, Chattisgarh, UP, Bihar
   - ↳ Kaimur is an extension
   - ↳ Sadbhavna Shikhar
8. Satpuras : Maharashtra, M.P., Chattisgarh ; Dhupgarh
   - ↳ Parts : Rajpipla, Gwaligarh, Mahadeo, Maikal

   Vindhyas & Satpuras meet at Amarkantak Plateau
9. Western Ghats (Sahyadris) → Anaimudi
   - ↳ Gujarat, Maharashtra, Goa, Karnataka, Kerala, Tamil Nadu
   - ↳ Extensions → Satmala, Ajanta, Balaghat, Harishchandra, Baba Budan, Nilgiri, Annamalai, Palani, Cardamom.

- **Aravalli** (relict of world's oldest mountains; Its northeast edge is Delhi Ridge; highes pt is Guru Shikhar). Across Gujarat, Rajasthan, Haryana, and Delhi.
- **Vindhyas** (Jobat in Gujarat to Sasaram in Bihar; eastwards extend as Bharner & Kaimur). Across Gujarat, MP, UP, Chhattisgarh, Bihar.
- **Satpuras** (higher than Vindhyas; Its sub ranges include Rajpipla, Mahadev, and Maikala Ranges; Dhupgarh is its highest peak). Across Gujarat, Maharashtra, MP, and Chhattisgarh.
- **Western Ghats:**
  - Location: Along the western coast of India, running parallel to the Arabian Sea.
  - Also known as **Sahyadris.**
  - Its **parts** include **Satmala, Harishchandra, Balaghat, Ajanta, Baba Budan, Nilgiri Annamalai, Cardamom, Palani hills.**
  - Important gaps include **Thalghat** (aka **Kasara** ghat; Mumbai-Nasik Highway) **Bhorghat** (Mumbai-Pune highway), **Chorla Ghat** (Goa), **Palghat** (Palakkad) **Shenkotta gaps** (Gateway of monsoon to Tamil Nadu).
  - **Tallest peak Anaimudi**
  - States: Gujarat, Maharashtra, Goa, Karnataka, Kerala, Tamil Nadu.
  - Characteristics: The mountain range acts as a barrier, causing heavy rainfall o the windward side due to the southwest monsoon. The region is known for its lus green forests, rich biodiversity, and numerous waterfalls. It's a source of severa major rivers flowing eastward, and many minor west flowing rivers. **So it is both climatic and a drainage divide.**
- **Eastern Ghats:**
  - Location: **Along the eastern coast** of India, running parallel to the Bay of Bengal.

- **Aravalli** (relict of world's oldest mountains; Its northeast edge is Delhi Ridge; highest pt is Guru Shikhar). Across Gujarat, Rajasthan, Haryana, and Delhi.
- **Vindhyas** (Jobat in Gujarat to Sasaram in Bihar; eastwards extend as Bharner & Kaimur). Across Gujarat, MP, UP, Chhattisgarh, Bihar.
- **Satpuras** (higher than Vindhyas; Its sub ranges include Rajpipla, Mahadev, and Maikala Ranges; Dhupgarh is its highest peak). Across Gujarat, Maharashtra, MP, and Chhattisgarh.
- **Western Ghats:**
  - Location: Along the western coast of India, running parallel to the Arabian Sea.
  - Also known as **Sahyadris.**
  - Its **parts** include **Satmala, Harishchandra, Balaghat, Ajanta, Baba Budan, Nilgiri, Annamalai, Cardamom, Palani hills.**
  - Important gaps include **Thalghat** (aka **Kasara** ghat; Mumbai-Nasik Highway), **Bhorghat** (Mumbai-Pune highway), **Chorla Ghat** (Goa), **Palghat** (Palakkad) **& Shenkotta gaps** (Gateway of monsoon to Tamil Nadu).

- **Tallest peak Anaimudi**
- States: Gujarat, Maharashtra, Goa, Karnataka, Kerala, Tamil Nadu.
- Characteristics: The mountain range acts as a barrier, causing heavy rainfall on the windward side due to the southwest monsoon. The region is known for its lush green forests, rich biodiversity, and numerous waterfalls. It's a source of several major rivers flowing eastward, and many minor west flowing rivers. **So it is both a climatic and a drainage divide.**

- **Eastern Ghats:**
  - Location: **Along the eastern coast** of India, running parallel to the Bay of Bengal.
  - **Highly dissected** due to the presence of multiple rivers.
  - Its parts include: **Garjhat hills** , **Mahendragiri, Eramalla, Nallamalla, Sesachalam, Velikonda, Palkonda, Javadi, Shevroy, Kalrayan, Biligiri Rangan.**
  - The highest peak is **Jindhagada** (Araku Valley, Andhra Pradesh).
  - States: Odisha, Telangana, Andhra Pradesh, Karnataka, and Tamil Nadu.
  - Characteristics: **Less continuous and lower in elevation** compared to the Western Ghats. The rainfall is lower, resulting in drier ecosystems with deciduous forests and scrub vegetation. The Eastern Ghats play a crucial role in draining many major rivers of the Indian Peninsula.
- **Eastern and Western Ghats meet at the Nilgiri Ranges** .

⚡ **Quick revise** Oldest and largest — Fold ▲

Oldest and largest unit; irregular triangle; broken from **Gondwanaland**; rich in **metallic minerals**; includes **Deccan Traps**. **Meghalaya plateau** was cut off by the **Rajmahal–Garo (Malda) gap / Ganga Suture**. Two halves – **Central Highlands** and **Deccan Plateau** – split by the **Narmada rift**.

Part	Where / what
Mewar/Marwar	NW, Rajasthan, near Aravallis; scrub
Malwa	Aravallis (NE) to Bundelkhand (E) to Vindhyas (S); **Chambal badlands**; black soil, cotton
Bundelkhand	between Vindhyas and Ganga plain (MP, UP); sal, teak, kheria
Baghelkhand	north Deccan, MP/CG; forests; **coal**
Chotanagpur	**Hazaribagh, Ranchi, Rajmahal Hills**; coal, mica; sal
Meghalaya	Garo, Khasi, Jaintia, Mikir, Rengma; **Mawsynram wettest on Earth**
Deccan	largest, central; avg **600 m**; Deccan Traps; **Chhattisgarh plateau & plains** (Dandakaranya, Bastar; Mahanadi rice bowl)

Hill ranges.

Range	Facts (as printed)
Aravalli	relict of world's **oldest** mountains · NE edge = **Delhi Ridge** · highest **Guru Shikhar** · Gujarat, Rajasthan, Haryana, Delhi
Vindhyas	**Jobat (Gujarat) → Sasaram (Bihar)**; east as **Bharner & Kaimur** · Gujarat, MP, UP, CG, Bihar
Satpuras	**higher than Vindhyas** · Rajpipla, Mahadev, Maikala · **Dhupgarh** highest · Gujarat, Maharashtra, MP, CG
Western Ghats (Sahyadris)	Satmala, Harishchandra, Balaghat, Ajanta, Baba Budan, Nilgiri, Annamalai, Cardamom, Palani · **climatic AND drainage divide** · **6 states** (Gujarat, Maharashtra, Goa, Karnataka, Kerala, TN)
Eastern Ghats	Garjhat, Mahendragiri, Eramalla, Nallamalla, **Seshachalam**, Velikonda, Palkonda, Javadi, Shevroy, Kalrayan, Biligiri Rangan · **highly dissected** by rivers · highest **Jindhagada (Araku)** · Odisha, Telangana, AP, Karnataka, TN
Meeting point	**Eastern and Western Ghats meet at the Nilgiris**

🔪 **Ghats compare: Western = Wall** (continuous, high, wet windward, divide). **Eastern = Entry-gate pieces** (broken by rivers, low, drier, scrub). *Western Ghats gaps, north → south:* **Thal – Bhor – Chorla – Pal – Shenkotta** (Thalghat, Bhorghat, Chorla, Palghat, Shenkotta).

8 · COASTAL PLAINS

Western Coastal Plains:

- It is a **mixed shoreline** , including shorelines of emergence, submergence, and also stable shoreline.
- **Kachchh Peninsula** : **shoreline of emergence** due to tectonic and depositional reasons, that has led to the creation of the **salty marshes** i.e. the Rann of Kachchh; It has the Banas and the Luni rivers, which undergo inland drainage. **Banni Grasslands** , one of the largest grasslands of india are here.
- **Kathiawar Plains** : **South of Kuchchh, includes Kathiawad (Saurashtra)** region. It has the **Deccan Trap lava black soil** ; It is also known for the **Gir Range** which is the only existing habitat of the Asiatic Lions. **Rivers Narmada and Tapi form estuaries** here.

region. It has the **Deccan** is also known for the **Gir** ly existing habitat of the **Narmada and Tapi form**

Daman to Goa; It is a **nce**, and that is why the wider, and it consists of eatures of Marine Erosion

oa to Mangalore; It is also **nce**.

stable part (emerging due estern Coast; It goes from umari; Due to Marine **es, lagoons, and backwaters called Kayals**.

arekha river to

deposits of major

mergence, due to ong rivers that flow

Northern Circars) & **Coromandel**

a; includes the Chilika from the south of t Lake; Important include the Kolleru e a lagoon, and is tion due to coastal e Krishna Godavari



ends from the Pilucat lake to Kanyakumari.

- **Konkan Plain** : From **Daman to Goa** ; It is a **shoreline of submergence** , and that is why the continental shelf here is wider, and it consists of many sea cliffs, where features of Marine Erosion are very evident.
- **Karnataka Plain** : From **Goa to Mangalore** ; It is also a **shoreline of submergence** .
- **Malabar Coast** : It is the **stable part (emerging due to deposition)** of the Western Coast; It goes from Mangalore to Kanyakumari; Due to Marine deposits, it **has many lakes, lagoons, and backwaters called Kayals** .

Eastern Coastal Plains:

- Extend **from Subarnarekha river to Kanyakumari** .
- They include **alluvial deposits of major rivers** .
- It is a **shoreline of emergence** , due to **deposition** by multiple long rivers that flow into the Bay of Bengal.
- **2 major parts** include **Northern Circars** (Mahanadi to Krishna) & **Coromandel Coast** (South of Krishna)
- **Regionally divided** into:
  - **Utkal Plain** : Odisha; includes the biggest lake of India Chilika
  - **Andhra Plain** : located from the south of Utkal, till the Pulicat Lake; Important geographic features include the Kolleru Lake, which was once a lagoon, and is now

in an inland location due to coastal deposition; the fertile Krishna Godavari delta.

- **Tamil Nadu Plain** : Extends from the Pilucat lake to Kanyakumari.

- **Eastern and Western** coastal plains **meet at Kanyakumari** .

⚡ **Quick revise** 2.6 Coastal plains and islands Fold ▲

Coast	Facts
Western plain	mixed shoreline (emergence, submergence, stable)
Kachchh	emergence → **Rann of Kachchh** salt marsh; **Banas and Luni** inland drainage; **Banni grassland**
Kathiawar	Deccan Trap black soil; **Gir – only Asiatic lions**; **Narmada, Tapi** estuaries
Konkan (Daman–Goa)	**submergence**; wider shelf; sea cliffs
Karnataka plain (Goa–Mangalore)	submergence
Malabar (Mangalore–Kanyakumari)	**stable/deposition**; **kayals** (lakes, lagoons, backwaters)
Eastern plain	**Subarnarekha → Kanyakumari**; **emergence** by deposition; **Northern Circars** (Mahanadi–Krishna) + **Coromandel** (south of Krishna)
Utkal / Andhra / Tamil Nadu	**Chilika** (biggest lake, Utkal) · **Kolleru** (once lagoon, now inland) + **Krishna–Godavari delta** · plain ends at **Pulicat → Kanyakumari**

East and west coasts **meet at Kanyakumari.**

9 · ISLANDS OF INDIA

Andaman & Nicobar Islands

- An **archipelago of over 500 islands,** stretching from 6° to 13° North latitude and 92° to 93° East longitude in the **Bay of Bengal.**
- Emerged due to the **collision of tectonic** plates millions of years ago. They are an **extension of** the Burmese mountain range **Arakan Yoma** , including **a few volcanic islands** as well.
- The highest peak is the **Saddle Peak** , located in the Andaman Islands.
- The **Andaman and the Nicobar Islands** are **separated** from each other **by** the **10-degree channel. Great Nicobar** separated from **Sumatra** via **6-degree channel.**
- **Barren Island** in the Andaman Islands is the **only active lava volcano in South Asia** , while Narcondam Island is an extinct volcano. (Indian plate under Burmese plate)
- **Baratang Island:** India's only active mud volcano.
- Havelock Island, also known as **Swaraj Dweep** , is the **largest island** of this group.



Lakshadweep Islands:

- It is a **coral archipelago** of **36 islands and coral reefs** , situated about 400 km off the coast of Kerala in the **Arabian Sea.**
- Emerged as **coral atolls on top of submerged volcanic features (Lakshdweep-Chagos Ridge)** millions of years ago. The islands are low-lying and sandy, with no rivers or permanent freshwater sources.
- Landforms: Flat and sandy islands with surrounding coral reefs.
- **Andrott Island** is the **largest** one.
- **Maliku Atoll** (Minicoy Island), the Southernmost of this group, is **separated from the rest of the islands of Lakshadweep** by the **9-degree channel.**

⚡ **Quick revise** 2.6 Coastal plains and islands Fold ▲

Islands	Facts
Andaman & Nicobar	**500+** islands, 6°–13°N, Bay of Bengal · extension of **Arakan Yoma** · **Saddle Peak** highest · **10° Channel** separates Andaman & Nicobar · **6° Channel** Great Nicobar–Sumatra · **Barren Island = only active lava volcano in South Asia**; **Narcondam = extinct**; **Baratang = only active mud volcano** · **Havelock (Swaraj Dweep)** largest of the group
Lakshadweep	**36** coral islands/reefs, **~400 km** off Kerala · coral atolls on submerged volcanic **Lakshadweep–Chagos Ridge** · no rivers/permanent freshwater · **Andrott** largest · **Minicoy (Maliku)** southernmost, split by **9° Channel**

🔟 🔢 **Channels = their own number: 10° Channel** (Andaman|Nicobar) · **9° Channel** (Lakshadweep|Minicoy) · **6°**

> **Channel** (Great Nicobar|Sumatra). *"10 – 9 – 6, count down from the north."*

10 · PYQ

Q1.) When you travel to the Himalayas, you will see the following :

- Deep gorges
- U-turn river courses
- Parallel mountain ranges
- Steep gradients causing land-sliding **Which of the above can be said to be the evidence for the Himalayas being young fold mountains? (2012)**
- 1. 1 and 2 only
- 2. 1, 2 and 4 only
- 3. 3 and 4 only
- 4. 1, 2, 3 and 4

Answer: D The Himalayas are known as the young fold mountains because they were formed only a few million years ago.

- They were formed because of the folding of the earth's crust due to tectonic activities.
- Gorge is a deep cleft resulting from weathering and the erosive activity of a river over geologic timescales.
- These deep gorges are irregular and don't have steep slopes which shows that these are formed recently.
- It shows that the Himalayas are young folded mountains.
- Similarly, other features like U-turn river courses, Parallel mountain ranges, Steep gradients causing landslides etc can be seen generally in young folded mountains only.
- These all features constitute that the Himalayas are young folded mountains.
- The old folded mountain is lower than a young folded mountain.
- The Himalayas have the highest mountain ranges in the world which show that the Himalayas are young folded mountains.

Q2.) Consider these pairs -

- Cardamom Hills Coromandel Coast
- Kaimur Hills Konkan Coast
- Mahadeo Hills Central India
- Mikir Hills North-East India

Which of the above pairs are correctly matched? (2014)

1. 1 and 2 only

2. 2 and 3 only

3. 3 and 4 only

4. 2 and 4 only

Answer: C Kaimur Range is the eastern portion of the Vindhya Range, about 483 kilometres (300 mi) long, extending from around Katangi in Jabalpur district of Madhya Pradesh to around Sasaram in Rohtas district of Bihar.

- It passes through the Rewa and Mirzapur divisions.
- The Cardamom Hills or Yela Mala are a mountain range of southern India and part of the southern Western Ghats located in southeast Kerala and southwest Tamil Nadu in India.
- Their name comes from the cardamom spice grown in much of the hills' cool elevation, which also supports pepper and coffee.
- The Western Ghats and Periyar Sub-Cluster including the Cardamom Hills, are UNESCO World Heritage Sites.

Q3.) Which one of the following pairs of islands are separated from each other by the 'Ten Degree Channel'? (2014)

(a) Andaman and Nicobar

(b) Nicobar and Sumatra

(c) Maldives and Lakshadweep

(d) Sumatra and Java

Answer: A This is a map based question, please refer to map.

Q4.) Which of the following pairs of States of India indicates the easternmost and Westernmost State? (2015)

(a) Assam and Rajasthan

(b) Arunachal Pradesh and Rajasthan

(c) Assam and Gujarat

(d) Arunachal Pradesh and Gujarat

Answer: D

- Go through the Map

Q5.) Consider the following pairs: Famous place Region

- Bodhgaya - Baghelkhand
- Khajuraho - Bundelkhand
- Shirdi: - Vidarbha
- Nasik (Nashik) - Malwa
- Tirupati - Rayalaseema

Which of the pairs given above are correctly matched? (2016)

1. 1, 2 and 4

2. 2, 3, 4 and 5

3. 2 and 5 only

4. 1, 3, 4 and 5

Answer: C

- Malwa region includes districts of Western Madhya Pradesh and parts of South-Eastern Rajasthan. Baghelkhand covers the North-Eastern regions of Madhya Pradesh and a small area of Western Uttar Pradesh. Vidarbha is the Eastern region of the Indian State of Maharashtra, comprising Nagpur division and Amravati division.

Q6.) If you travel by road from Kohima to Kottayam, what is the minimum number of States within India through which you can travel, including the origin and the destination? (2017)

- 6
- 7
- 8
- 9

Answer: B

- States – Kerala, Tamil Nadu, Andhra Pradesh, Odisha, West Bengal, Assam, Nagaland

Q7.) Consider the following statements:

1. In India, the Himalayas are spread over five States only.

2. The Western Ghats are spread over five States only.

3. Pulicat Lake is spread over two States only.

Which of the statements given above is/are correct? (2017)

(a) 1 and 2 only

(b) 3 only

(c) 2 and 3 only

(d) 1 and 3 only

Answer: B

- In India, the Himalayas are spread over more than 5 states.
- The Western Ghats are spread over 6 states, including Gujarat.
- Pulicat Lake is spread over Andhra Pradesh and Tamil Nadu.
- The Indian Himalayan Region (IHR) is the section of the Himalayas within India, spanning the states of Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Sikkim, Arunachal Pradesh, as well as the hill regions of two states - Assam and West Bengal.
- Western Ghats are a chain of mountains running parallel to India's western coast, approximately 30-50 km inland.
- The Western Ghats traverse the States of Kerala, Tamil Nadu, Karnataka, Goa, Maharashtra and Gujarat.
- Pulicat Lake formerly PralayaKaveri is the second largest brackish water lake or lagoon in India, after Chilika Lake.
- It straddles the border of Andhra Pradesh and Tamil Nadu states with over 40% of it in Andhra Pradesh and 60% in Tamil Nadu situated on the Coromandel Coast in South India.
- The lake encompasses the Pulicat Lake Bird Sanctuary. The barrier island of Sriharikota separates the lake from the Bay of Bengal and is home to the SatishDhawan Space Centre.

Q8.) Consider the following statements:

- The Barren Island volcano is an active volcano located in Indian territory.
- Barren Island lies about 140 km east of Great Nicobar.
- The last time the Barren Island volcano erupted was in 1991 and it has remained inactive since then.

Which of the statements given above is/are correct? (2018)

(a) 1 and 3 only

(b) 2 and 3

(c) 3 only

(d) 1 only

Answer: D

- According to the researchers at Goa-based National Institute of Oceanography (NIO), the Barren Island volcano, India’s only live volcano in the Andaman and Nicobar islands.
- The only live volcano in the Andaman and Nicobar islands is erupting once again. The Barren Island volcano, located 140-km north-east of Port Blair in Great Nicobar Island,

dormant for more than 150 years, started erupting in 1991 and has since then shown intermittent activity.

- Last eruption was as recent as 2022.

Q9.) Among the following cities, which one lies on a longitude closest to that of Delhi? (2018)

(a) Bengaluru

(b) Hyderabad

(c) Nagpur

(d) Pune

Answer: A

- Both Delhi and Bangalore fall on the 77° E longitude.

Q10.) Siachen Glacier is situated to the: (2020)

(a) East of Aksai Chin

(b) East of Leh

(c) North of Gilgit

(d) North of Nubra Valley

Answer: D

- Siachen Glacier, one of the world's longest mountain glaciers, lying in the Karakoram Range system of Kashmir near the India–Pakistan border, extending for 44 mi (70 km) from north-northwest to south-southeast.
- The Siachen Glacier is a glacier located in the eastern Karakoram Range in the Himalayas.

Q11.) Consider the following pairs: Peak Mountains

- Namcha Barwa — Garhwal Himalaya
- Nanda Devi — Kumaon Himalaya
- Nokrek — Sikkim Himalaya

Which of the pairs given above is/are correctly matched? (2022)

(a) 1 and 2

(b) 2 only

(c) 1 and 3

(d) 3 only

Answer: B

- Namcha Barwa is in an isolated part of southeastern Tibet. It stands inside the Great Bend of the Yarlung Tsangpo River.
- On other hand the Garhwal Himalayas are mountain ranges located in the Indian state of Uttarakhand

- Kumaun Himalayas rise to 7,817 metres at Nanda Devi, the range's highest peak.
- The Nokrek Biosphere Reserve is located in the northeast of India on the Tura Range, which forms part of the Meghalaya Plateau

Q12.) Which one of the following is the best example of repeated falls in sea level, giving rise to present-day extensive marshland?

(a) Bhitarkanika Mangroves

(b) Marakkanam Salt Pans

(c) Naupada Swamp

(d) Rann of Kutch

Answer: D

- Rann of Kutch is a perfect example of shoreline of emergence.

Q13.) Consider the following statements:

- Amarkantak Hills are at the confluence of Vindhya and the Sahyadri Ranges.
- Biligirirangan Hills constitute the easternmost part of Satpura Range.
- Seshachalam Hills constitute the southernmost part of Western Ghats.

How many of the statements given above are correct? (2023)

(a) Only one

(b) Only two

(c) All three

(d) None

Answer: D

- Amarkantak is at the confluence of Vindhyas and Satpuras.
- Biligiriranjan hills are in South East Karnataka.
- Seshachalam Hills are a part of the Eastern Ghats.



Q14.) Which of the following geographical features or phenomena is/are associated with the Peninsular Block of India?

1. Submergence of parts of the western coast due to tectonic activity

2. Presence of residual mountain ranges such as the Veliconda Hills and Mahendragiri Hills

3. Deep, V-shaped river valleys formed by fast-flowing rivers

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Answer: B The Peninsular Block is an ancient stable cratonic landmass, but parts of the western coast experienced submergence due to tectonic movements. Eastern Ghats are denuded mountains. Most rivers of Peninsular India are mature rivers flowing over old hard rocks.

Q15.) Which of the following with reference to Indian States is/are not correct?

1. Uttar Pradesh shares its boundary with the highest number of other Indian States.

2. Rajasthan shares the longest international border among all Indian States.

3. Sikkim is the only State that shares its boundary with just one other Indian State.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Answer: A West Bengal has longest international border. Meghalaya also shares border with only 1 state.

⚡ **Quick revise** Physiography PYQs Fold ▲

Year	Asked	Answer / trap
2012	Evidence for young fold mountains: deep gorges, U-turn courses, parallel ranges, steep gradients	**D – all four**
2014	Pairs: Cardamom–Coromandel? Kaimur–Konkan? Mahadeo–Central India? Mikir–NE?	**C – Mahadeo and Mikir only**; Kaimur = eastern Vindhya
2014	**Ten Degree Channel** separates	**Andaman and Nicobar (A)**
2015	Easternmost & westernmost states	**Arunachal & Gujarat (D)**
2016	Place–region: Khajuraho–Bundelkhand ✔, Tirupati–Rayalaseema ✔; Bodhgaya–Baghelkhand ✘, Shirdi–Vidarbha ✘, Nashik–Malwa ✘	**C (2 and 5)**
2017	Kohima → Kottayam: minimum states	**7 (B)**
2017	Himalaya in 5 states only? W. Ghats in 5 states only? Pulicat in 2 states only?	**Only 3 is right (B)** – Himalaya >5 states; W. Ghats **6**; **Pulicat = AP + TN** (2nd largest brackish lagoon after Chilika; Sriharikota separates it)
2018	Barren Island: active ✔; ~140 km east of Great Nicobar ✘ (handout: 140 km NE of Port Blair); last erupted 1991, inactive since ✘	**D (1 only)**; erupted again, last as recent as **2022**
2018	City on longitude nearest Delhi	**Bengaluru (A)**
2020	**Siachen Glacier** is…	**North of Nubra Valley (D)**
2022	Namcha Barwa–Garhwal ✘ · Nanda Devi–Kumaon ✔ · Nokrek–Sikkim ✘ (Nokrek = Meghalaya/Garo)	**B (2 only)**
–	Repeated falls in sea level → marshland	**Rann of Kutch (D)** – shoreline of emergence
2023	Amarkantak = Vindhya + Sahyadri? Biligiri = easternmost	**None (D)** – Amarkantak = **Vindhya + Satpura**; Biligiri

Year	Asked	Answer / trap
	Satpura? Seshachalam = southernmost W. Ghats?	Rangan in SE Karnataka; Seshachalam **Eastern Ghats**
–	Peninsular Block: western coast submergence ✔; residual hills (Velikonda, Mahendragiri) ✔; deep V-valleys ✘	**B** – peninsular rivers are mature
–	UP has most neighbouring states · Rajasthan longest int'l border · Sikkim only one-neighbour state	**A as printed**

⚠️ **Handout alert:** the last PYQ's printed answer is **A (statement 1 not correct)**, but its own explanation says **West Bengal** has the longest international border and **Meghalaya** also has just one neighbouring state – which would make statements 2 and 3 wrong. Source disagrees with itself; **check against your class notes / atlas.**

11 · INDIAN DRAINAGE AND WATER RESOURCES

Factors Governing Drainage Of A Location

- **Topography:**
  - **Slope and Relief:** Steeper slopes and higher relief lead to faster drainage. Water flows rapidly downhill, creating well-defined channels and rivers. (Example: The Himalayas with their steep slopes have fast-flowing rivers like the Indus and Brahmaputra)
- **Nature of Rocks:**
  - **Permeability** : Permeable rocks like sandstone allow water to seep through, reducing surface runoff and contributing to groundwater recharge. (Example: The Deccan Plateau with its basalt rocks has low permeability, leading to seasonal streams and some areas with poor drainage). Impermeable rocks like granite don't allow water to pass through, resulting in higher surface runoff and potentially flash floods. (Example: The Aravalli Range with its granitic rocks has a limited capacity for water infiltration, leading to seasonal rivers)
- **Amount of Water:**
  - **Rainfall** : Areas with high rainfall will naturally have more water to drain. Heavy precipitation can overwhelm drainage systems, leading to flooding. (Example: The Western Ghats receive heavy rainfall, leading to numerous perennial rivers)
  - **Snowfall:** Melting snow in mountainous regions contributes significantly to river flow, impacting seasonal drainage patterns. (Example: The Himalayan rivers like the Ganga and Yamuna experience increased flow during the spring thaw)
- **Periodicity of Flow:**

- **Perennial Rivers:** These rivers flow continuously throughout the year, fed by constant sources like glaciers, springs, or consistent rainfall. (Example: The major rivers of India like the Ganga, Brahmaputra, and Indus are perennial)
- **Seasonal Rivers:** These rivers flow only during specific seasons, typically during the rainy season. (Example: Many rivers in the Deccan Plateau are seasonal, flowing only during the monsoon months)

- **Geological Time:**
  - **Long-term processes like tectonic activity, erosion, and deposition** can shape the landscape and influence drainage patterns. (Example: The uplift of the Himalayas millions of years ago created the current drainage pattern of major rivers flowing southwards)
- **Anthropogenic Activities:**
  - **Deforestation** : Cutting down trees reduces water infiltration and increases surface runoff, leading to flash floods and soil erosion.
  - **Urbanization** : Paving over land with concrete reduces natural drainage and increases the burden on storm drains. (Example: Rapid urbanization in Indian cities like Delhi and Mumbai has placed stress on drainage systems during heavy rainfall)
  - **Dams and Reservoirs** : These structures can regulate water flow, reduce flooding, and provide irrigation water, but can also disrupt natural drainage patterns and ecosystems downstream. (Example: The Bhakra Nangal Dam on the Sutlej River in Punjab helps control floods and provides water for irrigation)

⚡ **Quick revise** Drainage and water resources Fold ▲

Factors: slope/relief · rock permeability (basalt low; granite impermeable) · rainfall + snowfall · perennial vs seasonal · geological time · humans (deforestation, urbanisation, dams).

Terminologies Related To Drainage:

- A **catchment region** (also called a drainage basin or watershed) is the **entire land area that drains water to a single point** , like a river mouth or lake.



- A **drainage divide** (also called a water divide or watershed divide) is the elevated landform that **separates different catchment regions** , determining which direction water flows.

- (Drainage patterns already covered in Geomorphology).
- **Drainage Types:**
  - **Sequent/Concordant Drainage:** A concordant drainage system exhibits a **harmonious** relationship **with the underlying geological structure** . The streams and rivers flow in a way that reflects the topography and rock formations.
    - **Consequent Drainage** : Streams flow directly down the **steepest slopes** , creating a dendritic (branch-like) pattern. This is common in mountainous areas with young landscapes. (Example: Streams flowing down the slopes of the Himalayas)
    - **Subsequent Drainage:** These are usually the **tributaries of consequent** streams, created at the locations of softer rocks. (Example: Drainage patterns in the Appalachian Mountains of the US).
  - **Insequent/Discordant:** A discordant drainage system exhibits a dissonance with the underlying geological structure. The streams and rivers flow in a way that **does not reflect the current topography** or rock formations.
    - **Antecedent Drainage:** An **established drainage system persists even after the underlying rock formations have been uplifted or eroded.** The river cuts through the new landscape, creating waterfalls or gorges. (Example: The antecedent drainage of the Indus and Brahmaputra River through the Himalayas)
    - **Superimposed Drainage:** This occurs when a **drainage system developed on a different landscape** (e.g., older, eroded mountains) is superimposed on a new, uplifted landscape. The rivers maintain their original course, creating a mismatch with the current topography. (Example: Subarnarekha River).
  - **Concordant drainage systems suggest a long-term evolution** where the rivers have carved their paths in harmony with the landscape. It is found in regions with tectonic stability.
  - **Discordant drainage systems often indicate geological changes** like uplift, erosion, or past landscapes that have been superimposed upon them.



⚡ **Quick revise** Drainage and water resources Fold ▲

Type	Meaning	Example
Consequent	flows down the original slope	Himalayan streams
Subsequent	tributary on softer rock	Appalachian
Antecedent	river **older than the uplift**, cuts gorges	**Indus, Brahmaputra** (through

Type	Meaning	Example
		Himalaya); Ghaghara
Superimposed	pattern from older landscape laid onto new rocks	**Subarnarekha**
Concordant = follows structure (stable regions) · Discordant = does not (uplift/erosion past)		

River Systems In India

- **77% of drainage is towards the Bay of Bengal; 23% is towards the Arabian Sea.**
- **Delhi Ridge-Aravallis-Vindhyas-Satpuras-Western Ghats form the Drainage Divide of India.**
- **Himalayan rivers are younger** (40-60mn yrs) **than peninsular** (200mn yrs).
- Inland drainage is very commonly seen in the Kachchh region, due to a shoreline of emergence.
- **River basins are divided based on their size** into: Major River basins: >20,000 sq. km; Medium: 2000-20000 sq. km; Minor: <2000 sq. km.



⚡ **Quick revise** Big numbers Fold ▲

Big numbers: 77% drains to Bay of Bengal, **23%** to Arabian Sea · **Divide:** Delhi Ridge – Aravallis – Vindhyas – Satpuras – Western Ghats · Himalayan rivers **40–60 mn yrs**, peninsular **200 mn yrs** · Basins: **major >20,000 km²; medium 2,000–20,000; minor <2,000** · Himalayan basin areas **Indus 11.78, Ganga 8.61, Brahmaputra 5.8 lakh km²**; **Godavari 3.12 lakh km²** is the largest peninsular basin.

Himalayan River System:

- **Origin:** The Shiwalik river got divided into Indus and Brahmaputra rivers.

- **Youthful perennial** rivers, that cause **flooding and experience shifting of river channels (Avulsions)** in their mid to lower courses (Kosi) due to the deposition of the sediments brought about by them from their upper courses.

Important River Systems:

- **Indus** : origin at Bokhar Chu in Kailash Mts. In India, it **moves through Ladakh only.**
- In India, **riparian states/UTs** include: **Ladakh, Jammu & Kashmir, Himachal Pradesh, and Punjab.**
- Makes a **gorge near Nanga Parbat.**
- **Important Left tributaries include:** Panjnad (IJKRBS), which are all left bank tributaries.
  - **Jhelum** : Origin at Verinag; Meanders; Wular lake; joins Chenab in Pak
  - **Chenab** : origin near Baralacha la pass, Lahaul-Spiti; Joins Sutlej in Pak
  - **Ravi:** West of Rohtang, Kullu Hills; Joins Chenab in Pak
  - **Beas:** Beas Kund near Rohtang; Gorges at Kati & Largi in Dhauladhar; entirely in India; Meets Sutlej in Punjab
  - **Sutlej:** Rakshasthal near Mansarovar, Kailash Range; enters India through Shipki La; Joins Indus in Pak as **Panjnad** .
  - Others: Zanskar, Suru
- **Important Right bank tributary:** Shyok River which originates from the Rimo Glacier, one of the tongues of the Siachen Glacier, and joins the Indus River at Keris, in India (POK). Another one is Gilgit river.

- **Ganga System:**
  - Origin: **Gangotri (Bhagirath) & Satopanth (Alaknanda)** glaciers.
  - It is known for the **Panch Prayags** , located in Uttarakhand (see the image).
  - **National Waterway 1** is on the Ganga River (from Haldia in West Bengal to Prayagraj in U.P.)
  - Known as **Padma in Bangladesh**
  - In India **moves through: Uttarakhand, UP, Bihar, Jharkhand, and West Bengal. Riparian states also include H.P., Delhi, Haryana, Rajasthan, M.P., and Chhattisgarh.**

MNEMONIC: Vasudev Ne Karna ko Rasta Dikhaya MNEMONIC FOR RIVERS: Dhul Nandakini ke Pair Mein Bikhri

  - Its important left bank tributaries include:
    - **Ramganga:** Origin at Chamoli District; Meets Ganga at Kannauj;
    - **Gomti:** Originates at Pilibhit; meets Ganga at Ghazipur
    - **Ghaghara:** Originates in Tibet; It is an antecedent river; joins Ganga in Bihar; Major Tributaries: Sarda; Saryu; Rapti, Choti Gandak
    - **Gandak:** Originates near Tibet-Nepal border; Kali Gandak, Mayangadi, and Trishuli are its major tributaries; joins the Ganga near Patna.
    - **Burhi Gandak:** Originates in Bihar (West Champaram). It flows almost parallel to Gandak. Makes one of India's biggest oxbow lake i.e. Kanwar Tal.
    - **Kosi River:** also called Sapt Kosi due to its 7 tributaries; important tributaries being Arun, Tumar, and Sun Kosi; It has a shifting course, giving it the title of Sorrow of Bihar; Originates in Nepal; Joins Ganga near Kursela.
    - **Mahananda:** Originates Near Tiger Hill. Last major right bank tributary to join Ganga in India. MNEMONIC: Rama Ghoomte hue Ghaghara Ganda Kar Mat.
  - **Important right bank tributaries:**
    - **Yamuna: Yamunotri Glacier** ; joins the Ganga at Prayagraj; Important tributaries include Chambal, Sind, Ken, Betwa.

- **Tons:** also called Tamsa and joins Ganga in UP.
- **Son** : Originates as a part of radial drainage (Narmada, Son, Johila) from the Amarkantak plateau in M.P. and joins the Ganga River near Patna.
- **Phalgu:** Passes through Gayaji.

Important Distributary:

- **Hooghly River** : starts near Giria, West Bengal, and flows into the Bay of Bengal near Sagar Island. Makes an Estuary.
  - **Damodar** : It is a tributary of the Hooghly River, which is a distributary of the Ganga. It starts in Chotanagpur Plateau and joins Hooghly near Kolkata.
  - Others: Mayurakshi, Ajay

- **Brahmaputra system:**

- Origin **: Chemayungdung & Angsi glaciers** , Kailash Range, very near to the origin point of Sutlej River.
- It enters India after taking a U-turn from **Namcha Barwa.**
- **Enters India as Dihang (Siang/Siyom), into which Lohit and Dibang meet.**
- Becomes Brahmputra, from **Sadiya to Dhubri (Assam)** , before entering Bangladesh.
- Has multiple names throughout its course: **Tsangpo - Dihang - Brahmaputra - Jamuna- Padma - Meghana**
- States it moves through: Arunachal Pradesh, Assam
- Riparian states: Above + Manipur, Nagaland, Meghalaya, Sikkim, West Bengal
- **Major Right Bank Tributaries: Subansiri, Kameng, Manas, Sankosh, Tista.**
- **Major Left Bank Tributaries: Burhi Dihing and Dhansari (South), Dibang or Sikang and Lohit.**
- **Forms Majuli Island** where Subansiri joins.
- Experiences a **shifting river course and causes flooding in Assam** , especially during the monsoon season.



4. Barak River:

- Origin in Manipur Hills
- Splits into Surma and Kushiyara rivers at Indo-Bangladesh Border
- Called Meghana in Bangladesh after it merges with Padma.
- Separated from the Brahmaputra by the Barail Hills.
- States: Manipur & Assam. Other riparian states: Meghalaya, Nagaland, Mizoram, Tripura.

⚡ **Quick revise** Drainage and water resources — Fold ▲

> 🏞️ **HIMALAYAN rivers = teenagers** (young, V-valleys, perennial, snow+rain, meanders, **big deltas**). **PENINSULAR rivers = grandparents** (mature, shallow valleys, **seasonal** – except **Kaveri**, straight courses, small basins, **Narmada & Tapi make estuaries**, others deltas).

⚡ **Quick revise** Indus: origin — Fold ▲

Indus: origin **Bokhar Chu, Kailash**; in India only **Ladakh** (riparian: Ladakh, J&K, HP, Punjab); gorge near **Nanga Parbat**. **Panjnad = I-J-K-R-B-S** (Indus left tributaries – *as printed*).

River	Origin / hook
Jhelum	**Verinag**; meanders; **Wular Lake**; joins Chenab in Pakistan
Chenab	near **Baralacha La** (Lahaul-Spiti); joins Sutlej in Pakistan

River	Origin / hook
Ravi	west of **Rohtang**, Kullu hills; joins Chenab in Pakistan
Beas	**Beas Kund** near Rohtang; gorges at **Kati & Largi** (Dhauladhar); **entirely in India**; meets Sutlej in Punjab
Sutlej	**Rakshasthal near Mansarovar**; enters India at **Shipki La**; joins Indus in Pak as **Panjnad**
Others	Zanskar, Suru · **Right bank: Shyok** (from **Rimo glacier**, joins Indus at Keris), **Gilgit**

PYQ 2021 (ans D): of Chenab, Jhelum, Ravi, Sutlej – **Sutlej** is the one that joins the Indus directly (handout: the others join into the Sutlej system first).

⚡ **Quick revise** 3.2 Ganga system Fold ▲

Origin: Gangotri (Bhagirathi) + Satopanth (Alaknanda). Panch Prayag: *Vishnu, Nand, Karna, Rudra, Dev.* **NW-1: Haldia → Prayagraj. Padma** in Bangladesh. Flows through Uttarakhand, UP, Bihar, Jharkhand, WB.

Panch Prayag – handout mnemonic "Vasudev Ne Karna ko Rasta Dikhaya" = **V**ishnuprayag, **N**andprayag, **K**arnaprayag, **R**udraprayag, **D**evprayag (in order up→down). **Feeder rivers – "Dhul Nandakini ke Pair Mein Bikhri"** = **Dhauli** (Ganga) · **Nandakini** · **Pindar** · **M**andakini · **Bhagirathi** *(decoded from the figure)*.

Handout figure: the five confluences and the streams that make the Ganga (Dhauli, Nandakini, Pindar, Mandakini, Bhagirathi).

⬅️ **LEFT-bank tributaries – "Rama Ghoomte hue Ghaghara Ganda Kar Mat"** = **Ram**ganga · **Gomti** · **Ghaghara** · **Gandak** · **Kosi** · **Mahananda** (+ **Burhi Gandak** between Gandak and Kosi).

Left bank	Facts
Ramganga	Chamoli; meets Ganga at **Kannauj**
Gomti	**Pilibhit**; meets Ganga at **Ghazipur**
Ghaghara	Tibet; **antecedent**; joins in Bihar; tribs **Sarda, Saryu, Rapti, Choti Gandak**
Gandak	Tibet–Nepal border; **Kali Gandak, Mayangadi, Trishuli**; joins near **Patna**
Burhi Gandak	West Champaran; **Kanwar Tal** (India's biggest oxbow)
Kosi	**Sapt Kosi** (7 tribs: Arun, Tumar, Sun Kosi); **Sorrow of Bihar** (shifting course); Nepal; joins near **Kursela**
Mahananda	near Tiger Hill; **last major right-bank**…*(handout wording)* tributary to join Ganga in India
Right bank	**Yamuna** (Yamunotri; joins at **Prayagraj**; Chambal, Sind, Ken, Betwa) · **Tons/Tamsa** · **Son** (Amarkantak **radial drainage** with Narmada, Johila; joins near Patna) · **Phalgu** (Gayaji)

Left bank	Facts
Distributary	**Hooghly** (Giria → Sagar Island, estuary) · **Damodar** (Chotanagpur → Hooghly near Kolkata) · Mayurakshi, Ajay

⚠️ **Handout alert:** Mahananda is listed under *left-bank* tributaries but described as "last major **right bank** tributary". Treat it as printed (left list) and note the wording clash.

⚡ **Quick revise** Brahmaputra — Fold ▲

Brahmaputra: Chemayungdung & Angsi glaciers, Kailash – near the **Sutlej** source. **U-turn at Namcha Barwa** → enters as **Dihang (Siang)**; **Lohit & Dibang** join; **Brahmaputra Sadiya → Dhubri**. **Name chain:** *Tsangpo – Dihang – Brahmaputra – Jamuna – Padma – Meghna.* **Majuli** island where Subansiri joins. Floods in Assam.

🔤 **Brahmaputra tributaries – "SKM-ST": S**ubansiri · **K**ameng · **M**anas · **S**ankosh · **T**eesta (**right**) | Left: **Burhi Dihing, Dhansiri, Dibang, Lohit** (+ Kolong).

Barak: Manipur hills; splits into **Surma + Kushiyara**; **Meghna** after meeting Padma; separated from Brahmaputra by **Barail Hills**; Manipur, Assam (+ Meghalaya, Nagaland, Mizoram, Tripura riparian).

Peninsular Drainage

- The peninsular drainage is approximately 200 million years old, and is in the **mature stage**
- Because of that **broad, Shallow, Largely Graded Valleys** .
- Classification based on the direction of flow:
  - **South-eastward flowing** (e.g.: Mahanadi Godavari, Krishna, Kaveri).
  - **Westward flowing** (e.g.: Narmada, Tapi).
  - **North-eastward flowing** : (e.g.: Tributaries of Ganga and Yamuna- Chambal, Ken, etc.).

⚡ **Quick revise** East-flowing — Fold ▲

East-flowing: Mahanadi, Godavari, Krishna, Kaveri · **West-flowing:** Narmada, Tapi · **NE-flowing:** Chambal, Ken etc.

South-Eastward flowing rivers:



- **Mahanadi River:**
  - **Origin: near Sihawa in Raipur district** of Chhattisgarh.
  - Runs through Odisha and **drains into the Bay of Bengal.**
  - The **river moves through CG and Odisha. Riparian States: MP** and **Chhattisgarh** (53%), and **Odisha** (47%).
  - Important Tributaries:
    - **Left Bank: Ib, Mand, Hasdo, Sheonath**
    - **Right Bank: Ong, Jonk, Tel**
  - MNEMONIC: Seonath Man Hasdo On the Joke I Told.
  - Hirakud Dam is built on it.
  - It creates a **gorge near Tikkarpara** in Odisha.
  - It makes a **delta** as it joins the Bay of Bengal.
- **Godavari River:**

  - **Longest peninsular river** , and the largest river system of Peninsular India
  - Also called **Vridha Ganga and Dakshin Ganga**

- **Origin: Trimbakeshwar,** near Nasik in Maharashtra.
- **States it moves through: Maharashtra, Telangana, Chhattisgarh, Andhra Pradesh, and Yanam (Puducherry). Riparian states include Odisha and Karnataka apart from the above.**
- Major Tributaries:



  - **Right bank: Manjara, Pravara.**
  - **Left Bank: Purna, Pranhita (Wainganga+Penganga), Indravati, Sabari.**
- **Below Rajahmundry , 2 distributaries: Gautami Godavari and Vashishtha Godavari** .
- Makes a **delta at its mouth** (KG delta)

- **Krishna River:**
  - **Originates in Mahabaleshwar** .
  - Makes a **delta near its mouth.** (KG delta)
  - Moves through: Maharashtra, Karnataka, Telangana, and Andhra Pradesh. Riparian states also same.
  - Important Tributaries:

Right Bank	Left Bank
• **Koyna** • Panchganga • Dudhganga • **Ghataprabha** • Malaprabha • **Tungabhadra**	• **Bhima** • Dindi • Halia • **Musi**

- **Kaveri River:**



  - Origin: Brahmagiri Hills, Kodagu, Karnataka.
  - **Perennial** river which gets fed by both southwest monsoon (in its upper course) and the northeast retreating monsoons (the lower course).
  - Moves through Karnataka and Tamil Nadi. Riparian States: Kerala (3%), (41%) Karnataka and 56% Tamil Nadu and Puducherry.
  - Makes a delta.
  - **Divides into 2 at Tiruchirapalli district: Kollidam & Kaveri** (both rejoin later at Srirangam, a riverine island)
  - Important Tributaries:

Right Bank	L
• Lakshmana Tirtha • Kabini • Suvarnavathi • Bhavani • **Noyall** • **Amaravati**	•

- **Other East flowing Peninsular rivers:**
  - **Brahmani:** Formed by the **confluence of S. Koel and Sankh** rivers near Rourkela. Flows into the Bay of Bengal. The **Eastern Ghats start from its South** .

- **Subarnarekha: Originates at the Ranchi plateau** ; forms the boundary between Odisha and WB; creates an estuary between Ganga and Mahanadi delta.
- **Penneru: Originates from Nandi Durg in Karnataka** ; Initially flows to the north, then attains an eastward course; forms an estuary before joining the Bay of Bengal. It makes a canyon at Gandikota.
- **Ponniyar (Karnataka & Tamil Nadu) & Vaigai (Tamil Nadu)** : Have a short course with a near coast existence only.
- Others Vamsdhara, Nagavali.

⚡ Quick revise 3.4 Peninsular rivers

Fold ▲

> 🌾 **Mahanadi left bank – "Seonath Man Hasdo On the Joke I Told" → Seonath · Mand · Hasdo** | right: **Ong · Jonk · Tel** ... and **Ib** (left). *(Handout mnemonic; "I" = Ib, "Told" = Tel.)*

River	Facts
Mahanadi	**Sihawa, Raipur (CG)**; CG + Odisha (53% / 47% as printed); **Hirakud Dam**; gorge near **Tikkarpara**; delta
Godavari	**Longest peninsular river**, **Vridha / Dakshin Ganga**; **Trimbakeshwar (Nashik)**; Maharashtra, Telangana, CG, AP, **Yanam**; **right:** Manjara, Pravara · **left:** Purna, **Pranhita (Wainganga + Penganga), Indravati, Sabari**; below **Rajahmundry** splits into **Gautami + Vashishtha**; **KG delta**
Krishna	**Mahabaleshwar**; Maharashtra, Karnataka, Telangana, AP · **right:** Koyna, Panchganga, Dudhganga, Ghataprabha, Malaprabha, **Tungabhadra** · **left:** Bhima, Dindi, Halia, Musi
Kaveri	**Brahmagiri (Kodagu)**; perennial – **SW monsoon (upper) + NE monsoon (lower)**; shares: Kerala 3%, Karnataka 41%, TN 56% (as printed); delta; splits at **Tiruchirapalli** into **Kollidam + Kaveri**, rejoin at **Srirangam island** · **right:** Lakshmana Tirtha, Kabini, Suvarnavathi, Bhavani, Noyyal, Amaravati · **left:** Harangi, Hemavati, Shimsha, Arkavati
Others (east)	**Brahmani** (S. Koel + Sankh near Rourkela; Eastern Ghats start south of it) · **Subarnarekha** (Ranchi plateau; Odisha–WB boundary; estuary) · **Penneru** (Nandi Durg; **Gandikota canyon**; estuary) · Ponniyar, Vaigai (short) · Vamsadhara, Nagavali
Narmada	**Amarkantak**; **rift valley between Vindhya & Satpura** – few tributaries (short, trellis); **estuary** at Gulf of Khambhat; Hiran, Kolar, Hatni, Choti Tawa; MP, Maharashtra, Gujarat; meander near **Bharuch**
Tapi	"Narmada's twin"; **Multai (MP)**; Purna, Girna, Panjara; Satpura north, Mahadev east, Ajanta/Satmala south
Luni	west of **Ajmer**; **Sagarmati → Luni** after Sarsuti joins; **ephemeral**; **best inland-drainage example in the world** → Rann of Kachchh; Rajasthan's biggest system
Sabarmati	Aravallis → Gulf of Khambhat; gorge at **Dharoi**
Mahi	Vindhyas (MP) → Gulf of Khambhat; **only river to cross the Tropic of Cancer twice**

River	Facts
Short west rivers	**parallel drainage** off the Western Ghats, **no deltas**: Gujarat (Auranga, Par, Daman Ganga) · Goa (**Mandovi, Zuari**, Rachol) · Maharashtra (Ulhas, Kalindi) · Karnataka (Kalinadi, **Sharavati – Jog Falls**, Tadri, Netravathi) · Kerala (Bharathapuzha, **Periyar**, Pamba)
Inland	**Ghaggar–Hakra** (ends in Thar; "Hakra" in Pakistan after Ottu barrage) · **Luni, West Banas** (→ Rann of Kachchh)

West Flowing Rivers:

- **Narmada:**
  - **Originates at the Amarkantak Plateau** , M.P. forming **a radial drainage** pattern with the river Son.
  - Makes an estuary.
  - **Mouth** at **Gulf of Khambhat** .
  - **Flows in Rift Valley** between Vindhyas and Satpuras, **so not many tributaries** . **Has short tributaries, that make a trellis pattern of drainage.**
  - **Major tributaries** include **Hiran, Kolar, Hatni; Choti Tawa** , etc.
  - Riparian states: M.P., Maharashtra & Gujarat
  - Experiences a **slight meandering near Bharuch.**
- **Tapi:**



  - Often called **Narmada's twin**

- Bounded by the Satpura range on the north, Mahadev Hills in the east, Ajanta range , and the Satmala Hills on the south.
- Riparian states MP, Maharashtra, Gujarat.
- **Origin: Multai in M.P** .
- **Main Tributary: Purna, Girna and Panjara** .

- **Luni River:**
  - Also known as **Salt (Lun) River**
  - **Source: West of Ajmer**
  - **Biggest river system of Rajasthan**
  - Called **Sagarmati** initially; and then **Luni after river Sarsuti joins it** .
  - **Ephemeral river** , i.e. it flows only when there is rain.
  - Best example of an **inland drainage system** in the world. Drains into the Rann of Kachchh which is a shoreline of emergence.
- **Sabarmati:**
  - Culturally important river of India.
  - **Origin is Aravallis** and its **mouth is the Gulf of Khambat.**
  - It forms a **Gorge in Dharoi**
- **Mahi:**
  - **Origin: Vindhyas** , M.P.
  - **Ends at the Gulf of Khambat.**
  - **Only river to cross the tropic of Cancer Twice.**
- **Other short West flowing rivers:**
  - Very short rivers, which do **not make any deltas.**
  - They are a prime example of **parallel drainage** , where these short channels originate in the Western Ghats and end in the Arabian Sea, while moving parallel to each other from West to East.
  - **Gujarat: Auranga, Par, Daman Ganga**
  - **Goa: Mandovi Zuari, Rachol**
  - **Maharashtra: Ulhas, Kalindi**
  - **Karnataka: Kalinadi, Sharavati (Jog Falls), Tadri, Netravathi**
  - **Kerala: Bharathapuzha, Periyar, Pamba**

Inland Drainage In India

- Refers to the drainage systems that **do not drain into seas/oceans.**
- **Ghaggar-Hakra river system** : It is situated in North-Western India and Eastern Pakistan. The river is known as Ghaggar in India, and as the Hakra in Pakistan, post the Ottu barrage. It ends its journey in the Thar desert.

and West Banas rivers originate in Rajasthan, enter Gujarat, and end the
e Rann of Kachchh.

Dams of India

- **Luni and West Banas** rivers originate in Rajasthan, enter Gujarat, and end their journeys in the Rann of Kachchh.

Important Dams of India

Chhattisgarh

Waterfall	Height (m)	River (including Tributary Info)
Amritdhara Falls	90	Hasdeo (Tributary of Mahanadi)
Teerathgarh	91	Munga (Tributary of Kanger which joins Mahanadi)
Chitrakote Waterfalls	29	Indravati (Tributary of Godavari)

Madhya Pradesh

Waterfall	Height (m)	River (including Tributary Info)
Dhuandhar Waterfall	30	Narmada (Independent)
Chachai	130	Bihad (Tributary of Tons, which is a tributary of Ganga)
Kapildhara Falls	30	Narmada (Independent)
Sonemuda Falls	15	Son (Tributary of Ganga)

Jharkhand

Waterfall	Height (m)	River (including Tributary Info)
Panchghagh Falls	45	Banai (Tributary of Subarnarekha)
Lodh Falls	143	Budha (Tributary of North Koel)
Hundru Falls	99	Subarnarekha (Independent)

Odisha

Waterfall	Height (m)	River (including Tributary Info)
Barehipani	399	Budhabalanga (Tributary of Subarnarekha)
Khandadhar Falls	244	Korapani Nala (Tributary of Brahmani)
Koilighughar Falls	61	Ahiraj (Tributary of Mahanadi)

Karnataka

Waterfall	Height (m)	River (including Tributary Info)
Kunchikal Falls	455	Varahi (Tributary of Krishna)
Barkana Falls	259	Sita (Tributary of Krishna)
Jog Falls	253	Sharavati
Magod Falls	198	Bedti
Shimsha Falls	46	Shimsha (Tributary of Cauvery)
Shivanasamudra Falls	98	Cauvery
Hebbe Falls	168	Thunga (Tributary of Krishna)

Other States

Waterfall	Height (m)	River (including Tributary Info)
Nohsngithiang (Seven-Sisters) Waterfall	315	Cherrapunji (Rain-fed), Meghalaya
Vantawng Falls	229	Lau (Tributary of Tlawng which joins Barak river), Mizoram
Kempty Falls	40	Yamuna, Uttarakhand
Dudhsagar	320	Mandovi, Goa
Thalaiyar Falls (Rat Tail Falls)	297	Manjalar (Tributary of Vaigai), Tamil Nadu

Difference Between Peninsular And Himalayan Drainage

Differences	Himalayan Drainage	Peni
Origin	These rivers originate from the lofty Himalayan ranges	Thes Peni
Catchment Area	Himalayan rivers have large basins and catchment areas. E.g. The total basin area of the Indus, the Ganga and the Brahmaputra is 11.78, 8.61 and 5.8 lakh square km, respectively.	Thes and God area
Valleys	The Himalayan rivers flow through deep V-shaped valleys and sometimes form gorges.	The com Thes com river

Difference Between Peninsular And Himalayan Drainage

		to p form
Drainage Type	Both antecedent and consequent drainage found	Thes cons Suba supe
Water Flow	The Himalayan rivers are perennial in nature, i.e., water flows throughout the year in these rivers. These rivers receive water from the S-W Monsoons as well as Snowmelt.	The wate flows seas river (exc
Stage	These rivers flow across the Young Fold mountains and are still in a youthful stage.	Thes one worl
Meanders	The upper reaches of the Himalayan rivers are highly tortuous. When they enter the plains, there is a sudden reduction in the speed of flow of water. Under these circumstances these rivers form meanders and often shift their beds.	The non- plate the such Plate strai
Deltas and Estuaries	The Himalayan rivers form big deltas at their mouths. The Ganga-Brahmaputra delta is the largest in the world.	Som such Tapt rest

Indian River Linking Project

History of the Project:



- The **concept originated in the 19th century** by British engineer **Sir Arthur Cotton** .
- Post-independence, the idea was revisited in the 1960s and gained momentum in the 1980s due to concerns about water scarcity and floods.
- The **National Water Development Agency (NWDA)** was **established in 1982 to study** the project's **feasibility** .
- The Supreme Court of India has intervened multiple times, directing the government to finalize plans and initiate implementation.

About the Project:

- The project aims to link numerous Indian rivers through a network of reservoirs and canals.
- **Two main components** are envisaged:
  - **Himalayan Rivers Development** : Interlinking rivers in the northern part of the country, like the Ganga and Yamuna. (14 potential links identified)
  - **Peninsular Rivers Development** : Connecting rivers in peninsular India, like the Godavari, Krishna, Kaveri, and Mahanadi. (16 potential links identified)
- Some states have also proposed intra-state river-linking projects.

Current Status:

- The project **remains under debate** .

- The **Ken-Betwa River linking project,** considered a pilot project, **received cabinet approval in 2014, but faces hurdles** related to environmental clearances and inter-state agreements.
- **Pattiseema Lift Irrigation Project** aims **to connect the Godavari and Krishna Rivers** . It was **inaugurated in 2016** . It brings **water from Godavari to Krishna** , from Godavari's **Polavaram Project.**
- The NWDA continues to study the feasibility of other potential links.

⚡ **Quick revise** River-linking project Fold ▲

River-linking project: idea of **Sir Arthur Cotton** (19th c.) · revived 1960s–80s · **NWDA 1982** · Himalayan component **14 links**, peninsular **16 links** · **Ken–Betwa** pilot (cabinet approval **2014**, hurdles) · **Pattiseema lift scheme (Godavari → Krishna), inaugurated 2016** via Polavaram.

Important Lakes in India


IMPORTANT LAKES OF INDIA
WULAR
DAL
PANGONG TSO
TSO MORORI
RENUKA
ROOPKUND
NAINITAL
BHIMTAL
GOBIND BALLABH PANT SAGAR
GURUDONGMAR
CHOLAMU
PARSHURAM KUND
LHANAK
SAMBHAR
PUSHKAR
KANWAR
DEEPOR BEEL
UDAIPUR LAKES
LOKTAK
NAL SAROVAR
BHOJTAL
LONAR
CHILIKA
SALIM ALI LAKE
HUSAIN SAGAR
KOLLERU
PULICAT
STANLEY
VEMBANAD
ASHTAMUDI

	Lake	State/UT	Characteristics
1	Wular	J&K	FW lake. Part of Jhelum
2	Dal	J&K	In Srinagar
3.	Pangong Tso	Ladakh	High altitudinal saline lake. Indo-China border
4.	Tso Morori	Ladakh	High altitudinal saline lake
5.	Renuka	HP	Largest natural lake of HP.
6.	Roopkund	Uttarakhand	Near Trishul peak. Skeletons at its bottom
7.	Nainital	Uttarakhand	Natural freshwater lake
8.	Bhimtal	Uttarakhand	Artificial FW lake
9.	Govind Ballabhpant sagar	Uttar Pradesh	Rihand dam. Largest artificial lake of India
10.	Sambhar	Rajasthan	Playa lake
11.	Pushkar	Rajasthan	Artificial lake. Religious importance
12.	Udaipur lakes	Rajasthan	Artificial lakes
13.	Kanwar	Bihar	Largest oxbow lake of Asia. By Burhi Gandak
14.	Lhonak, Cholamu.	Sikkim	High altitudinal glacial lakes
15.	Gurudongmar	Sikkim	Highest lake of India
16.	Deepor Beel	Assam	In Brahmaputra basin
17.	Loktak	Manipur	Between West & East Manipur hills. Known for Phumdis. Keibul Lamjao NP. Sangai deer.

Lake	State/UT	Characteristics
18 Nal Sarovar	Gujarat	Freshwater. Largest lake of Gujarat.
19. Bhojtal	M.P	Artificial lake
20. Lonar	Maharashtra	Meteorite crater lake
21 Salim Ali	Maharashtra	Chhatrapati Sambaji Nagar (earlier called Aurangabad)
22 Chilika	Odisha	Lagoon lake. Largest lake of India
23. Husain Sagar	Telangana	Artificial lake in Hyderabad. Gets water from Musi river
24. Kolleru	Andhra Pradesh	Was a lagoon lake. But now it is completely inland
25. Pulicat	AP-TN	At the border of Andhra Pradesh and Tamil-Nadu. Lagoon lake
26. Vembanad	Kerala	Longest lake of India. Backwater 6 rivers have a mouth here. Pamba, Periyar
27 Ashtamudi	Kerala	Octopus lake. It is also a Kayal/backwater.

PYQs:

.) The Brahmaputra, Irrawaddy and Mekong rivers originate in Tibet and flow throug arrow and parallel mountain ranges in their upper reaches. Of these rivers, Brahmaputr akes a "U" turn in its course to flow into India. This "U" turn is due to – (2011)

(a) Uplift of folded Himalayan series
(b) Syntaxial bending of geologically young Himalayas
(c) Geo-tectonic disturbance in the tertiary folded mountain chains
(d) Both (a) and (b) above

ns: B

bout 70 million years ago, there was a sea called Tethys sea where the Himalayas stan oday. The sea started to recede during the early Eocene which was marked by the comin ogether of Indian and Asian plates.

- This trend of marine regression continued until the final emergence of the Himalaya

⚡ Quick revise 3.5 Waterfalls and lakes (image pages) Fold ▲

State	Waterfall (height m) – river
Karnataka	**Kunchikal 455** (Varahi) · Barkana 259 (Sita) · **Jog 253** (Sharavati) · Magod 198 (Bedti) · Shivanasamudra 98 (Kaveri) · Hebbe 168 (Thunga) · Shimsha 46
Odisha	**Barehipani 399** (Budhabalanga) · Khandadhar 244 (Korapani Nala) · Koilighughar 61 (Ahiraj)
Kerala	**Meenmutty 300** (Kallar) · Soochipara 200 (Chaliyar) · Athirapally 25 (Chalakuddy → Periyar)
Chhattisgarh	Amritdhara 90 (Hasdeo) · Teerathgarh 91 (Munga) · Chitrakote 29 (Indravati)
MP	**Chachai 130** (Bihad → Tons) · Dhuandhar 30 & Kapildhara 30 (Narmada) · Sonemuda 15 (Son)

State	Waterfall (height m) – river
Jharkhand	**Lodh 143** (Budha) · Hundru 99 (Subarnarekha) · Panchghagh 45 (Banai)
Others	**Nohsngithiang (Seven Sisters) 315** – Meghalaya · Vantawng 229 – Mizoram · Kempty 40 – Yamuna, Uttarakhand · **Dudhsagar 320** – Mandovi, Goa · **Thalaiyar (Rat Tail) 297** – Manjalar/Vaigai, TN · Maharashtra: Thoseghar, Someshwar 10 (Godavari)

📈 **Tallest-first pegs: Kunchikal 455 (Karnataka) > Barehipani 399 (Odisha) > Dudhsagar 320 > Nohsngithiang 315 > Meenmutty 300 > Thalaiyar 297 > Jog 253.**

Lakes (handout table + map):

Lake	State/UT	Hook
Wular	J&K	freshwater, part of Jhelum
Dal	J&K	Srinagar
Pangong Tso	Ladakh	high-altitude saline; Indo-China border
Tso Moriri	Ladakh	high-altitude saline
Renuka	HP	largest natural lake of HP
Roopkund	Uttarakhand	near Trishul; skeletons at bottom
Nainital	Uttarakhand	natural freshwater
Bhimtal	Uttarakhand	artificial FW
Gobind Ballabh Pant Sagar	UP	Rihand dam; **largest artificial lake of India**
Sambhar	Rajasthan	Playa lake
Pushkar	Rajasthan	artificial; religious
Udaipur lakes	Rajasthan	artificial
Kanwar	Bihar	**largest oxbow lake of Asia**; Burhi Gandak
Lhonak, Cholamu	Sikkim	high-altitude glacial
Gurudongmar	Sikkim	**highest lake of India**
Deepor Beel	Assam	Brahmaputra basin
Loktak	Manipur	between west & east Manipur hills; **phumdis**; **Keibul Lamjao NP, sangai deer**

Also named on the map (no notes): Parshuram Kund, Nal Sarovar, Bhojtal, Lonar, Salim Ali, Husain Sagar, Kolleru, Pulicat, Stanley, Vembanad, Ashtamudi, Chilika.

PYQs:

Q.) The Brahmaputra, Irrawaddy and Mekong rivers originate in Tibet and flow through narrow and parallel mountain ranges in their upper reaches. Of these rivers, Brahmaputra makes a "U" turn in its course to flow into India. This "U" turn is due to - (2011)

(a) Uplift of folded Himalayan series

(b) Syntaxial bending of geologically young Himalayas

(c) Geo-tectonic disturbance in the tertiary folded mountain chains

(d) Both (a) and (b) above

Ans: B About 70 million years ago, there was a sea called Tethys sea where the Himalayas stand today. The sea started to recede during the early Eocene which was marked by the coming together of Indian and Asian plates.

- This trend of marine regression continued until the final emergence of the Himalayan Mountains by the end of the Tertiary Period.
- The collision of the Indian Plate with the Asian Plate retarded the pace of northerly drift of the Indian Plate.
- During the upliftment of the Himalayas, the maximum push was offered by the Aravalli mountains in the West and Shillong Plateau in the East.
- Arakanyoma and Sulaiman ranges are regarded as an offshoot of Himalaya, Arakanyoma bends near Namcha Barwa (Arunachal Pradesh East), while Sulaiman bends near Nanga Parbat.
- Both these mountains/offshoots give the appearance of a hairpin.
- Due to this structure Brahmaputra river makes U turn into the Indian subcontinent.

Q.) The Narmada river flows to the west, while most other large peninsular rivers flow to the east. Why? (2013)

1. It occupies a linear rift valley.

2. It flows between the Vindhyas and the Satpuras.

3. The land slopes to the west from Central India.

Select the correct answer using the codes given below.

(a) 1 only

(b) 1 and 3 only

(c) 2 and 3 only

(d) None

Answer: A Narmada and Tapi flow through rift valleys caused due to the upheaval of the Himalayas.

- The Narmada rises at an elevation of about 3,500 feet (1,080 metres) in the Maikala Range in eastern Madhya Pradesh state on the border with Chhattisgarh state.

- It first follows a tortuous course through the hills near Mandla and then turns northwest to pass the city of Jabalpur.
- There it turns southwest and then enters the structural trough between the Vindhya and Satpura ranges at Marble Rocks Gorge.
- Turning more westward, the river continues across Madhya Pradesh until it passes into Gujarat state.
- The Narmada enters the Gulf of Khambhat through an estuary 13 miles (21 km) wide, just below Bharuch.
- The Narmada is flowing through the area where the land is not sloping towards the west from central India, but it is flowing west because of the rift valleys only.

Q.) Consider the following rivers:

1. Barak

2. Lohit

3. Subansiri

Which of the above flows / flow through Arunachal Pradesh? (2014)

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: B The Barak is a 900 km river flowing through the states of Nagaland, Manipur and Assam in India and into the Bay of Bengal via Bangladesh.

- The Barak River flows 900 kilometres through the states of Manipur, Nagaland, Mizoram and Assam in India. Further it enters Bangladesh where it is known by the name of the Surma and the Kushiyara and later called the Meghna before receiving the combined flow of the Ganga and the Brahmaputra.
- Barak rises in the Manipur hills and enters the plains near Lakhipur. The river enters Bangladesh as Surma and Kushiyara. Later, the river is called the Meghna and receives the combined flow of the Ganga and Brahmaputra.
- It is being developed as one of the Inland waterways of India.
- The environment of Barak's basin hosts a wide variety of flora and fauna.

Q.) Consider the following rivers :

1. Vamsadhara

2. Indravati

3. Pranahita

4. Pennar

Which of the above are tributaries of Godavari? (2015)

(a) 1,2 and 3

(b) 2,3 and 4

(c) 1,2 and 4

(d) 2 and 3 only

Answer: D The Godavari is India's second longest river after the Ganga.

- Its source is in Trimbakeshwar, Maharashtra
- The major tributaries of the river can be classified as the left bank tributaries which include the Purna, Pranhita, Indravati and Sabari River covering nearly 59.7% of the total catchment area of the basin and the right bank tributaries Pravara, Manjira, Manair together contributing 16.1% of the basin.
- River Vamsadhara or River Vamsadhara is an important east flowing river between Rushikulya and Godavari, in Odisha and Andhra Pradesh states in India.
- The Penna (also known as Pennar, Penner, Penneru or North Pinākinī) is a river of southern India.
- The Penna rises in Nandi Hills in Chikballapur District of Karnataka state and runs north and east through the states of Karnataka and Andhra Pradesh to empty into the Bay of Bengal.

Q.) Which of the following is/are tributary/tributaries of Brahmaputra?

1. Dibang

2. Kameng

3. Lohit

Select the correct answer using the code given below: (2016)

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: D The Brahmaputra (meaning the son of Brahma):

- The Brahmaputra’s source is the Chemayungdung glacier in southwestern Tibet. It’s source is very close to the sources of Indus and Satluj.
- Mariam La separates the source of the Brahmaputra from the Manasarovar Lake.
- In Tibet, it passes through the depression formed by the Indus-Tsangpo Structure Zone between the Great Himalayas in the south and the Kailas Range in the north.

Brahmaputra Luit, Dilao(Basin features):

Tributaries

- **Left -** Lhasa River, Nyang River, Parlung Zangbo, Lohit River, Dhansiri River, Kolong River
- **Right -** Kameng River, Manas River, Beki River, Raidak River, Jaldhaka River, Teesta River, Subansiri River

Q.) Recently, linking of which of the following rivers was undertaken? (2016)

(a) Cauvery and Tungabhadra

(b) Godavari and Krishna

(c) Mahanadi and Son

(d) Narmada and Tapti

Answer: B

- India took a step forward in its ambitious but long-pending goal to interlink major rivers to form a national water grid. The Pattiseema project lifts flood water from the river Godavari and pumps it into the Polavaram right canal that empties into the river Krishna in Vijayawada.
- Andhra Pradesh Government is interlinking Krishna-Godavari rivers through the Pattiseema lift irrigation scheme.

Q.) With reference to river Teesta, consider the following statements:

1. The source of river Teesta is the same as that of Brahmaputra but it flows through Sikkim.

2. River Rangeet originates in Sikkim and it is a tributary of river Teesta.

3. River Teesta flows into Bay of Bengal on the border of India and Bangladesh.

Which of the statements given above is/are correct? (2017)

(a) 1 and 3 only

(b) 2 only

(c) 2 and 3 only

(d) 1, 2 and 3

Answer: B Teesta River (or Tista River) is a 315 km (196 mi) long river that rises in the eastern Himalayas, flows through the Indian states of West Bengal and Sikkim through Bangladesh and enters the Bay of Bengal

- The Teesta River originates from the Pahunri (or Teesta Kangse) glacier above 7,068 metres (23,189 ft) and flows southward through gorges and rapids in the Sikkim Himalaya.
- It is fed by rivulets arising in the Thangu, Yumthang and Donkha mountain ranges.
- The river then flows past the town of Rangpo where the Rangpo River joins, and where it forms the border between Sikkim and West Bengal up to Teesta Bazaar.
- Just before the Teesta Bridge, where the roads from Kalimpong and Darjeeling join, the river is met by its main tributary, the Rangeet River.

Q.) With reference to the Indus River system, of the following four rivers, three of them pour into one of the joins the Indus direct. Among the following, which one is such river that joins the Indus direct? (2021)

(a) Chenab

(b) Jhelum

(c) Ravi

(d) Sutlej

Answer: D

- The Indus is a transboundary river of Asia and a trans-Himalayan river of South and Central Asia. The 3,180 km river rises in Western Tibet, flows northwest through the Ladakh and Gilgit-Baltistan.
- Because Jhelum, Ravi and Chenab join theStulej, which joins the Indus directly.



Q.) Consider the following Rivers:

1. Brahmani

2. Nagavali

3. Subarnarekha

4. Vamsadhara

Which of the above rise from the Eastern Ghats? (2021)

(a) 1 and 2

(b) 2 and 4

(c) 3 and 4

(d) 1 and 3

Answer: B

- Statement 1 is incorrect. The Brahmani, known as South Koel in its upper reaches, rises near Nagri village in Ranchi district of Jharkhand at an elevation of about 600 m.
- Statement 2 is correct. The Nagavali river originate in the Eastern Ghats near Lakhbahal, located at an altitude of 1,300 metres in the Kalahandi district of the Indian state Odisha. Langulya is another name for Nagavali River.
- Statement:3 is incorrect. Subernarekha River originate in the Chhotnagpur Plateau near Nagri village in the Ranchi District of Jharkhand at an elevation of 600 m.

- Statement 4 is correct. Vamsadhara river rises in the Eastern Ghats of Orissa State and enters Srikakulam District in Bhamini Mandal and finally falls into the Bay of Bengal near Kalingapatnam.
- Vamsadhara: It is an east-flowing river that originates in the Kalahandi district of Odisha, Eastern Ghat, flows in Odisha, along its boundary with Andhra Pradesh and finally joins the Bay of Bengal at Kalingapatnam, Andhra Pradesh. Hence, 4 is correct.

Q.) Gandikota canyon of South India was created by which one of the following rivers? (2022)

(a) Cauvery

(b) Manjira

(c) Pennar

(d) Tungabhadra

Answer: C Gandikota is a small village in the Kadapa district of Andhra Pradesh. The village is majorly known for housing the spectacular gorge which is famously adjudged as the Grand Canyon of India. The stunning gorge has been created by the waters of the famous river Pennar that streams from the Erramala hills.

Q.) Consider the following statements:

1. Jhelum River passes through Wular Lake.

2. Krishna River directly feeds Kolleru Lake.

3. Meandering of the Gandak River formed Kanwar Lake.

How many of the statements given above are correct? (2023)

(a) Only one

(b) Only two

(c) All three

(d) None

Answer: A

- Wular Lake is a part of the Jhelum River Drainage.
- Kolleru lake lies between Krishna and Godavari.
- Kanwar lake is created by the meandering of Burhi Gandak (Sikrahna) River.

Q.) Identify the river of the Indian sub-continent on the basis of the following information:

1. It has an antecedent drainage system.

2. It flows through three countries.

3. It originates in the Tibetan Plateau and is an important river for irrigation.

4. It does not form distributaries.

Select the answer from the following:

(a) Brahmaputra

(b) Indus

(c) Sutlej

(d) Teesta

Answer: C Brahmaputra has distributaries. Teesta originates in Sikkim. Indus has distributaries.

⚡ **Quick revise** Drainage PYQs Fold ▲

Year	Asked	Answer / rule
2011	Why does the Brahmaputra make a U-turn?	**Syntaxial bending of young Himalaya (B)** – Arakan Yoma bends near Namcha Barwa; Sulaiman near Nanga Parbat
2013	Why does Narmada flow west?	**Linear rift valley (A)** – not "slope west" (3 wrong)
2014	Which flow through Arunachal: Barak, Lohit, Subansiri	**Lohit + Subansiri (B)**; Barak = Manipur/Nagaland/Mizoram/Assam
2015	Godavari tributaries: Vamsadhara, Indravati, Pranahita, Pennar	**Indravati + Pranahita (D)**
2016	Brahmaputra tributaries: Dibang, Kameng, Lohit	**All three (D)**
2016	Recent river linking	**Godavari–Krishna (B)** (Pattiseema)
2017	Teesta: same source as Brahmaputra? Rangeet a tributary? flows into Bay on the border?	**Only 2 (B)** – Rangeet originates in Sikkim and is the main tributary; Teesta source = Pahunri/Teesta Kangse glacier
2021	Which joins the Indus directly	**Sutlej (D)**
2021	Rise from Eastern Ghats: Brahmani, Nagavali, Subarnarekha, Vamsadhara	**Nagavali + Vamsadhara (B)**
2022	Gandikota canyon	**Pennar (C)**
2023	Jhelum through Wular ✔; Krishna feeds Kolleru ✘; Gandak meander → Kanwar ✘ (it is **Burhi Gandak**)	**Only one (A)**
–	River: antecedent, 3 countries, Tibetan origin, irrigation, **no distributaries**	**Sutlej (C)**

12 · INDIAN CLIMATE

INDIAN CLIMATE:

- Tropical conditions along the Southern coastal regions to the very cold conditions in the North, from dry desert-like conditions in the West to abundant rainfall in the East.
- Considered **Monsoonal climate** (75% rainfall in monsoon months)

⚡ **Quick revise** Indian climate — Fold ▲

> 📅 **Four seasons – "WiSAR": Wi**nter (mid-Nov–Feb) · **S**ummer/pre-monsoon · **A**dvancing monsoon (Jun–Sep) · **R**etreating monsoon (Oct–Nov).

Factors affecting Indian Climate:

- **Latitude** : India's latitudinal extent stretches from roughly **6°N to 35°N** , placing it primarily in the tropical zone. This means it receives a high amount of solar radiation throughout the year, contributing to generally warm temperatures. However, there are variations. As you move north towards higher latitudes, temperatures tend to decrease as the climate gets more temperate in nature.
- **Himalayas** : This mighty mountain range acts as a giant barrier, separating the subcontinent from the cold, dry winds blowing from Central Asia. The Himalayas play a crucial role in influencing the climate: a. **Blocking Cold Winds** : They prevent the harsh winters of Central Asia from reaching most of India. b. **Monsoons** : The Himalayas influence the formation of monsoon winds, which bring vital rainfall to the subcontinent during the summer months.
- **Pressure and Wind Systems** : The pressure and wind systems over the Indian subcontinent vary seasonally. These variations significantly impact the weather patterns: a. **Monsoon** : During summer, low pressure over the Indian landmass attracts moisture-laden winds from the southwest (Indian Ocean), bringing the monsoon season with its characteristic heavy rainfall. b. **Upper air circulation** : In winter, high pressure over the landmass leads to dry, northeasterly winds blowing from land to sea, resulting in a dry season for much of the country. But the Sub Tropical Westerly jet streams bring in the Western Disturbances, causing winter rains.
- **Ocean Currents** : India is surrounded by water bodies on three sides: the Arabian Sea to the west, the Bay of Bengal to the east, and the Indian Ocean to the south. The warm currents like the Indian Ocean Equatorial Current contribute to generally warm and humid conditions along the coastal regions.
- **Distance from the Sea (Continentality)** : As we move inland from the coast, the influence of the sea on the climate diminishes. Locations further away from the sea experience more extreme temperatures, with hotter summers and colder winters compared to coastal areas.

- **Relief Features** : India's diverse topography, including mountains, plateaus, and plains, also plays a role. Mountain ranges like the Western Ghats can act as barriers, influencing rainfall patterns and creating rain shadows in certain areas. The high altitude of the Himalayas contributes to a colder climate in the northern mountain regions.

Seasons Of India: Winter Season

- From **mid-Nov till Feb**
- **Fog formation in Northern parts** , especially in the Indo-Gangetic Region, which **can even turn into smog** . Temperature inversion can also lead to the rise of **Pollution Domes** .
- **Frost** is common **in Northern parts.**
- **Offshore Northeast trade winds** prevail.
- **Rains due to Western Disturbances** (Subtropical Westerly Jet) Winter rain " **Mewat"/ "Mahawat** " is important for Rabi crops.
- Season of **least tropical cyclones** as the land and the oceans experience high pressure due to the shift of ITCZ.
- **Cold waves are common** in the Northern parts of the country.

⚡ **Quick revise** Indian climate Fold ▲

Winter: fog/smog, inversion, pollution domes, frost, cold waves; offshore NE trades; **WD rain – "Mawath/Mahawat" – vital for Rabi**; **fewest tropical cyclones** (ITCZ shifts, high pressure).

Western Disturbances

- Western disturbances are **extratropical storms originating in the Mediterranean region** . These low-pressure systems **travel eastwards** towards India, **influenced by the Sub-Tropical Westerly jet stream** .
- As these disturbances move towards the Indian subcontinent, they encounter the Himalayas, which cause them to lose moisture, resulting in precipitation over northern India.



- **Impact:**
  - **Winter Rainfall** : Western disturbances are a major source of winter and pre-monsoon rainfall in the northwestern parts of India, including Punjab, Haryana, Uttar

Pradesh, and Delhi. Sometimes they might also bring rainfall to Peninsular and North Eastern states. They also bring snowfall to the Himalayan regions.

- **Temperature Regulation** : These disturbances help regulate winter temperatures, preventing extreme cold conditions by bringing cloud cover and moisture.
- **Water Resources** : The snowfall in the Himalayas contributes to the perennial flow of rivers such as the Ganges and Indus, which are vital for irrigation and drinking water supply. Important for Rabi crops.
- **Weather Events** : These disturbances can also lead to extreme weather events like heavy rainfall, hailstorms, and thunderstorms, which can cause damage to crops, infrastructure, and even loss of life.

- **Possible Impact of Climate Change:**
  - **Frequency and Intensity** : Climate change may alter the frequency and intensity of western disturbances. Some studies suggest that **there could be an increase in the intensity** of these disturbances due to warmer air holding more moisture.
    - Increased intensity could lead to more severe weather events, including heavier rainfall and more extensive snowfall.
  - **Shift in Patterns** : There is a possibility of a shift in the typical pathways and timing of western disturbances due to changes in global atmospheric circulation patterns.
    - Changes in the timing and pathway can affect the predictability and reliability of rainfall and snowfall during the winter months, impacting agriculture and water resources.
  - **Climatological Disasters:** Over the past few years, the WDs are being seen frequently in summers as well. They can interact with the summer cyclones in Arabian Sea and Bay of Bengal, and cause severe weather events. They can also interact with Monsoons, causing extreme precipitation.

⚡ **Quick revise**

Monsoon climate: ~75% of rain in monsoon months Fold ▲

Monsoon climate: ~75% of rain in monsoon months. Controls: latitude (~6°N–35°N) · **Himalayas** (block Central Asian cold; shape monsoon) · pressure & winds (summer low pulls SW winds; winter high → dry NE winds; **sub-tropical westerly jet → Western Disturbances**) · ocean currents · continentality · relief.

Western Disturbances: extratropical storms from the Mediterranean, steered by the **sub-tropical westerly jet**; lose moisture on the Himalaya. Give winter and pre-monsoon rain in **Punjab, Haryana, UP, Delhi**, Himalayan snow (feeds **Ganga/Indus perennial flow**), moderate winter cold; also hail/thunderstorms. *Climate change:* may be more intense (warmer air, more moisture), shifted timing/path, now **seen in summers** and **interacting with cyclones and monsoon → extreme rain**.

Seasons Of India: Summer Season

- **Northward shift of sun** , causing the **shift of ITCZ** too (around 25 degree North).

- **Loo winds and Dust storms** (Aandhi); can cause **showers** (assisted by Western Disturbances).
- **Heat waves** are a **common** feature of the season.
- **Pre-monsoon showers:** (Caused by meeting of warm continental and maritime air masses)
  - **Norwester/ Kaal baisakhi** : WB, Bihar, Jharkhand, Odisha (Rice, tea, jute)
  - **Bardoli Chheerha/Bordoisila/tea showers** : Assam
  - **Mango Showers** : Karnataka/Kerala
  - **Coffee/ Cherry Blossom Showers** : Karnataka & Kerala
- **1%** of **India's annual rainfall** in this season.
- There is a **lot of cyclonic activity** during this season.

⚡ **Quick revise** Summer: ITCZ Fold ▲

Summer: ITCZ ≈ **25°N**; **Loo**, dust storms (Aandhi); heat waves; **1% of annual rain**.

☕🥭 **Pre-monsoon showers by "what they help": Mango showers** (Karnataka/Kerala) · **Coffee / Cherry-blossom showers** (Karnataka, Kerala) · **Norwester / Kaal Baisakhi** (WB, Bihar, Jharkhand, Odisha – rice, tea, jute) · **Bardoli Chheerha / Bordoisila – tea showers** (Assam).

Cyclones In India

- India has a tropical location, with a vast coastline of over 7500 km, making it susceptible to Tropical Cyclones.
- The period between **April and December** experiences the **most significant cyclone activity** in India, with the **peak months** being **May-June and October-November** .
- **Favorable Conditions** : This period coincides with:
  - **Warm Sea Surface Temperatures** : The Bay of Bengal and the Arabian Sea reach their warmest temperatures during these months, providing the energy needed for cyclone formation.
  - **Monsoon Winds and Moisture** : The southwest monsoon brings in significant moisture, which is another crucial ingredient for cyclone formation.
- **Bay of Bengal v/s Arabian Sea:**
  - The Bay of Bengal witnesses a higher frequency of cyclones compared to the Arabian Sea, with a **ratio of approximately 4:1** . This is due to favorable conditions like **larger warm water areas** (due to continentality) and **lower wind shear** prevailing in the Bay of Bengal.
  - **On average, 5-6 tropical cyclones form annually in the North Indian Ocean (Bay of Bengal and Arabian Sea), with 2-3 being severe.**
- **Impact of Climate Change:**

- **Warmer Seas, Stronger Storms:**
  - A key factor in cyclone formation is warm sea surface temperatures. As the climate warms, both the Arabian Sea and the Bay of Bengal are experiencing rising sea surface temperatures.
  - Warmer water acts as fuel for cyclones, providing the energy they need to form, intensify, and sustain themselves. This can lead to cyclones reaching higher wind speeds and causing more extensive damage.
- **Increased Intensity, Not Necessarily Frequency:**
  - While the overall **number** of cyclones **might not** significantly increase, climate change is projected to make **existing cyclones more severe** .
  - Stronger winds, heavier rainfall, and higher storm surges are all potential consequences of climate change-induced intensification.
- **Impact on the Arabian Sea:**
  - Historically, the Arabian Sea has been less prone to cyclones compared to the Bay of Bengal. However, recent studies suggest an increase in cyclone activity in the Arabian Sea, which some experts attribute to climate change.
  - Warmer Arabian Sea waters could lead to **more frequent and intense cyclones** striking the western coast of India, posing a new threat to previously less vulnerable regions.
- **Impact on the Bay of Bengal:**
  - The Bay of Bengal is already a hotspot for cyclones. With rising sea surface temperatures, **cyclones here could become even more powerful** .
  - Increased storm surges due to rising sea levels can cause greater coastal inundation, displacing communities and damaging infrastructure.
  - Heavy rainfall associated with intensified cyclones can lead to flash floods and landslides, further exacerbating the impact on coastal populations and inland areas.
  - According to a **2022 study, numbers might decrease and intensities will increase** here.
- **Adaptation and Mitigation:**
  - **Adaptation** : Building **seawalls,** improving **early warning systems** , and promoting **disaster preparedness** measures are crucial for adapting to the changing cyclone landscape.
  - **Mitigation** : Addressing climate change at its source by **reducing greenhouse gas emissions** is essential for long-term mitigation of cyclone intensity.

- **Colour-Coded warnings:** The Indian government uses a well-defined color-coded cyclone warning system to disseminate information and prepare communities:
  - **Pre-cyclone watch (No colour):** issued **72 hours in advance** and contains an early warning about the development of a cyclonic disturbance in the north Indian Ocean.
  - **Cyclone Alert (Yellow): issued at least 48 hrs in advance** of the expected commencement of adverse weather over the coastal areas. It contains information on the location and intensity of the storm likely direction of its movement, intensification, coastal districts likely to experience adverse weather, and advice to fishermen, the general public, media, and disaster managers.

- **Cyclone Warning (Orange):** issued **at least 24 hours in advance** of the expected commencement of adverse weather over the coastal areas. The landfall point is forecast at this stage.
- **Post Landfall outlook (Red): issued at least 12 hours in advance** of the expected time of landfall. It gives the likely direction of movement of the cyclone after its landfall and adverse weather likely to be experienced in the interior areas.

- **Naming of the Cyclones in the North Indian Ocean region:** The World Meteorological Organization/Economic and Social Commission for Asia and the Pacific (WMO/ESCAP) Panel on Tropical Cyclones plays a crucial role in naming cyclones that form over the Bay of Bengal and the Arabian Sea, a **practice that began in September 2004.**
  - **13 countries** are involved in the naming process: Bangladesh, India, Iran, Pakistan, Oman, Maldives, Myanmar, Sri Lanka, Thailand, United Arab Emirates, Yemen, Qatar, and Saudi Arabia.
  - Names are chosen from a **predefined list.**
  - **Each country gives 13 names.**
  - The **RSMC** (Regional Specialized Meteorological Centre) **New Delhi Tropical Cyclone Centre** assigns names to storms reaching a designated intensity.
  - **Names are new and not used elsewhere globally** to avoid confusion.
  - **Once used, a name is retired from the list for future storms** .
  - If a cyclone originates in the South China Sea and enters the Bay of Bengal, its original name remains unchanged.

⚡ **Quick revise** Cyclones — Fold ▲

Cyclones: coast **>7,500 km** (as printed); season **April–Dec**, peaks **May–June** and **Oct–Nov**; **Bay : Arabian Sea ≈ 4 : 1** (bigger warm-water area, lower wind shear); **5–6 per year, 2–3 severe.** Climate change: *intensity up, not necessarily frequency*; Arabian Sea activity rising; a 2022 study says Bay numbers may fall but intensity rise.

IMD colour warning	Lead time
Pre-cyclone watch (no colour)	**72 h**
Cyclone Alert – Yellow	**≥48 h**
Cyclone Warning – Orange	**≥24 h** (landfall point forecast)
Post-landfall outlook – Red	**≥12 h**

(Mnemonic: 72 – 48 – 24 – 12, halving each step: "Watch → Yellow → Orange → Red".)

Naming: WMO/ESCAP Panel, since **September 2004**; **13 countries** (Bangladesh, India, Iran, Pakistan, Oman, Maldives, Myanmar, Sri Lanka, Thailand, UAE, Yemen, Qatar, Saudi Arabia) **× 13 names each**; **RSMC New Delhi** assigns; names are retired after use; a South China Sea cyclone entering the Bay keeps its name.

Urban Heat Island Effect



- The Urban Heat Island (UHI) effect refers to the phenomenon where **urban areas experience significantly higher temperatures than their rural surroundings.**
- **Causes:**
  - **Concrete and Asphalt Surfaces** : Buildings, roads, and other infrastructure absorb and retain heat, leading to higher temperatures.
  - **Reduced Vegetation** : Urban areas often have less vegetation, which would otherwise provide cooling through shading and evapotranspiration.
  - **Heat Emissions** : Vehicles, industrial activities, air conditioners, and other sources emit heat, further warming urban areas.
  - **Anthropogenic Heat** : Energy consumption for heating, cooling, and other urban needs generates additional heat.
  - **Urban Geometry** : The configuration of buildings and streets can trap heat and restrict airflow, creating "heat canyons."

⚡ **Quick revise** Urban heat island Fold ▲

Urban heat island: causes – concrete/asphalt, less vegetation, vehicle/industrial/AC heat, anthropogenic energy, urban geometry ("heat canyons").

Seasons Of India: Advancing Monsoon

- **75% rainfall.**
- Arabic word “mausim”; **seasonal reversal of winds.**
- **June-September**
- **Factors affecting Indian Monsoons:**
  - **Differential Heating of Land and Water:**
    - This is the **primary driver of the monsoon** . During summer, the Indian landmass heats up rapidly due to the high solar radiation. Conversely, the surrounding oceans (the Arabian Sea and Bay of Bengal) retain their cooler temperatures.
    - This creates a pressure difference: hot air over land rises (low pressure), while cooler air over the oceans remains denser (high pressure). This sets up a pressure

gradient that draws moisture-laden winds from the seas towards the landmass, bringing the monsoon rains.

- **Shift of the Intertropical Convergence Zone (ITCZ):**
  - The ITCZ is a belt of low pressure encircling the Earth near the equator, where warm, humid air rises and condenses to form rain.
  - **During summer, the ITCZ migrates northward** , positioning itself over the Indian subcontinent. This shift **brings the zone of heavy rainfall closer to India,** contributing significantly to the monsoon season.



- **Subtropical Westerly Jet Stream:**
  - High up in the atmosphere, the Subtropical Westerly Jet Stream is a fast-moving band of westerly winds. While its direct influence on monsoon rainfall is minimal, its position plays a role.



  - A **weaker or northward-shifted jet stream can allow more moisture-laden southerly winds to reach India** , strengthening the monsoon. Conversely, a stronger or southward-shifted jet stream can act as a barrier, hindering the monsoon winds.
- **Tibetan Plateau and Tropical Easterly Jet Stream (TEJ):**
  - The massive **Tibetan Plateau acts like a giant elevated heat source during summer.** This heating **strengthens the TEJ** , an easterly jet stream blowing over the subcontinent.
  - A **strong TEJ helps pull in moisture-laden winds** , enhancing the monsoon. Conversely, a weak TEJ can lead to a weaker monsoon.

- **Somali Jet Stream:**
  - This **low-level jet stream** blows from the southwest across the Arabian Sea during summer. It plays a crucial role in **transporting moisture from the Arabian Sea towards India** , contributing to the overall monsoon rainfall.



- **El Niño Southern Oscillation (ENSO):**
  - This **large-scale climate-ocean phenomenon** occurring in the Pacific Ocean has a significant impact on the Indian monsoon.



  - During **El Niño** events (warmer than average sea surface temperatures in the eastern Pacific compared to Western Pacific), **monsoon rainfall over India tends to be weaker.**

- Conversely, **La Niña events** (cooler than average sea surface temperatures in the Eastern Pacific) are often associated with **stronger monsoons.**



- **Indian Ocean Dipole (IOD):**



  - This climate pattern refers to the difference in sea surface temperatures between the western and eastern Indian Oceans.
  - A **positive IOD event** (warmer waters in the west, cooler in the east) can lead to **stronger monsoon** rainfall over India.
  - Conversely, a **negative IOD event** (cooler waters in the west, warmer in the east) is often associated with **weaker monsoon** activity.
- **Madden-Julian Oscillation (MJO):**



- This atmospheric phenomenon involves **eastward movement of a band of low pressure and enhanced convection** (rising air) **around the tropics** .
- The MJO can influence the intensity of the Indian monsoon rainfall on a timescale of weeks to months. A f **avorable phase of the MJO can enhance monsoon rainfall** , while an unfavorable phase can suppress it.

- **Arctic Sea Ice:** According to a **study by the National Centre for Polar and Ocean Research** , Goa, Arctic sea ice also affects the indian monsoon. What is the **exact mechanism, is yet to be identified** , but as of now, we know that reduced sea ice in the Arctic causes increased rain in central and Northeast india, and decreased rainfall in Western and southern peninsular India.

- **Monsoonal Trough:** A **low pressure region** created over India due to the above factors, that attracts the SouthWest Monsoonal winds.
- **Monsoon Burst** : **Monsoon advance with startling suddenness** accompanied by Thunder, Lightning, and heavy downpour. 1st June (Kerala).
- **2 branches** :
  - **Arabian Sea (more powerful)** & **Bay of Bengal** (Blocked by Arakan Mountains); **meet roughly around Agra & Ferozepur**
  - Arabian sea branch further divided into 3 branches (Malabar, Konkan, Kathiawar); Bay of Bengal Branch into Utkal and North Eastern Branches.
- **Very little rain in Tamil Nadu** , as it is in rainshadow of Western Ghats, and the branches are parallel to coast of Tamil Nadu.
- **Monsoon Breaks** : usually in **August** , **due to the weakening of Tibetan Low pressure** , and the **movement of monsoonal winds parallel to the Western coast of India** .
- **Flooding** can be **common** during this season, especially in the regions where rivers move.

Cloudbursts

- Cloudbursts are **intense and sudden rain events** where a significant amount of rain falls over a small area in a very short period, often leading to flash floods.
- Cloudbursts are typically defined as instances where **10 cm of rain or more falls within an hour at a particular location.**
- **Cloudburst Formation Process**

- **Rapid Upward Movement of Moist Air:** Warm, moisture-laden air rises quickly, often due to geographical factors like mountain ranges. This ascent can also be triggered by unstable atmospheric conditions.
- **Condensation at High Altitudes:** As the air rises, it cools and condenses into water droplets. In cloudburst conditions, this process happens very quickly, leading to the formation of dense, moisture-heavy clouds.
- **Sudden Rainfall Release:** Once the cloud becomes too heavy, it releases the moisture all at once, resulting in an intense and concentrated downpour. The rate of rainfall is so high that the ground cannot absorb it quickly enough, causing flash floods.
- The **release of moisture is also aided by the cold air influx into the clouds at high altitudes** (lapse rate), which **can force a downdraft** (like a piston), causing the raindrops to be pushed down.

- **Conditions Favoring Cloudbursts**
  - **Mountainous Areas:** Regions like the Himalayas and the Western Ghats are prone to cloudbursts due to orographic lifting. This is when mountains force moist air to rise rapidly, causing condensation and heavy rainfall.
  - **Monsoon Season:** Cloudbursts are most common during the Indian monsoon season (June to September), when warm, moist air from the oceans interacts with mountain ranges, creating ideal conditions for cloudburst formation.
- While cloudbursts are **more common in mountainous regions, they can also occur in plains** . However, the steep slopes of mountains make these areas particularly vulnerable to the devastating impacts of cloudbursts, such as flash floods and landslides.
- These events are **difficult to predict because they occur on such a small scale** , both in time and space. For effective prediction, we need either a high-resolution radar network or advanced weather models capable of capturing small-scale weather dynamics.
- Cloudbursts in India: Himalayas, Western Ghats.
- **Climate change is increasing the instances** of cloudbursts as warmer air can store more moisture.

⚡ **Quick revise** Advancing monsoon (Jun–Sep) Fold ▲

Drivers: land–sea differential heating (primary) · **ITCZ shifts north** · sub-tropical westerly jet (weaker/northward = stronger monsoon) · **Tibetan Plateau heating → Tropical Easterly Jet (TEJ)** · **Somali jet** (moisture across the Arabian Sea) · **ENSO** · **IOD** · **MJO** · **Arctic sea ice** (reduced ice → more rain in central & NE India, less in west & south peninsula; mechanism not yet identified).

- **Monsoon trough** = low over India. **Burst** = sudden onset with thunder/lightning, **~1 June at Kerala.**
- **Two branches: Arabian Sea** (stronger; splits into **Malabar, Konkan, Kathiawar**) and **Bay of Bengal** (blocked by **Arakan**; **Utkal and North-Eastern** branches); meet around **Agra–Ferozepur**.
- **Tamil Nadu is dry:** rain-shadow of Western Ghats, and the branches run **parallel to its coast**.

- **Break:** usually **August** – weak Tibetan Low and winds running parallel to the west coast.

Cloudbursts: ≥10 cm of rain in an hour over a small area; orographic lift + rapid condensation + cold-air downdraft; **Himalaya and Western Ghats**; monsoon; hard to predict (needs dense radar, high-res models); more with warming.

Urban Flooding In India



- Urban flooding is a phenomenon where **heavy rainfall overwhelms the drainage capacity of urban areas** , leading to water accumulation on streets, homes, and infrastructure. Unlike rural flooding, urban flooding is **primarily due to the high density of impervious surfaces** such as concrete and asphalt, which prevent water from being absorbed into the ground.
- **Causes:**
  - **Heavy Rainfall** : Intense and prolonged rainfall can exceed the capacity of urban drainage systems, leading to flooding. E.g. The Mumbai floods of July 2005 were caused by a record-breaking 944 mm of rainfall in a single day.
  - **Poor Drainage Systems** : Inadequate, poorly designed, or poorly maintained drainage infrastructure can lead to blockages and insufficient water flow. E.g. Chennai floods of 2015 highlighted the city's inadequate drainage systems and encroachments on water bodies.
  - **Unplanned Urbanization** : Rapid and unplanned urban growth reduces the amount of open land that can absorb rainwater, increasing surface runoff. E.g. Bengaluru has seen significant urban flooding due to rapid, unplanned urbanization and reduced green spaces.
  - **Improper Waste management** can choke the natural and artificial drains, exacerbating the problem.
  - **Encroachment on Natural Water Bodies** : Construction and encroachment on lakes, wetlands, and rivers disrupt natural drainage and water absorption. E.g. Hyderabad's flooding issues are partly due to encroachment on lakes and water bodies.

- **Climate Change** : Increasing frequency and intensity of extreme weather events, such as heavy rainfall, due to climate change exacerbate urban flooding. E.g. Climate change has been linked to the increased frequency of heavy rain events in cities like Mumbai and Kolkata.
- **Deforestation** : Reduction of green cover in and around cities decreases natural absorption of rainwater, leading to increased runoff. E.g. The loss of mangroves in coastal cities like Mumbai reduces natural flood buffers.

⚡ **Quick revise** Urban flooding Fold ▲

Urban flooding: impervious surfaces + drains overwhelmed. Handout examples: **Mumbai 2005 (944 mm in a day)**, **Chennai 2015** (drainage, water-body encroachment), **Bengaluru** (unplanned growth), **Hyderabad** (lake encroachment), mangrove loss in Mumbai; climate-linked heavy rain in Mumbai/Kolkata.

El Nino Southern Oscillations

- El Niño Southern Oscillation (ENSO) is a coupled **ocean-atmosphere phenomenon** that **disrupts typical weather patterns across the globe.** It's essentially a large-scale climate dance between the Pacific Ocean and the atmosphere above it, with two distinct phases: El Niño and La Niña.
- **Climatological and Oceanographic Impacts:**
  - **El Niño:** During El Niño events, **trade winds weaken in the tropical Pacific** . **Warm surface water** typically confined to the western Pacific **sloshes eastward** , leading to:
    - **Warmer sea surface temperatures in the eastern Pacific.**
    - The cold **Peruvian (Humboldt) current gets replaced by warm water** .
    - **Reduced upwelling** of cold, nutrient-rich water from the deep ocean.
    - Changes in atmospheric circulation patterns, impacting global rainfall.
  - **La Niña** : La Niña represents the opposite phase. **Trade winds intensify, pushing warm surface water westward.** This results in:
    - Cooler sea surface temperatures in the eastern Pacific.
    - **Enhanced upwelling** of cold, nutrient-rich water.
    - Altered atmospheric circulation patterns with a different global rainfall signature.
- **Global Climate Impacts of El Niño:**
  - **Warmer and Drier Conditions** : El Niño is often associated with drier-than-average conditions in many parts of the world, including:
    - **India and East Africa** : Reduced rainfall can lead to droughts and food insecurity.
    - **Australia** : Drier conditions can contribute to wildfires and heatwaves.
    - **Southeast Asia** : El Niño can suppress monsoon rainfall, impacting agriculture.
  - **Increased Rainfall and Flooding** : Conversely, El Niño can bring wetter-than-average conditions to other regions:

    - **California (USA):** Increased storminess and precipitation are common during El Niño events.
    - **Peru** : El Niño can lead to heavy rainfall and flooding along the normally dry coastal regions.
- **Global Climate Impacts of La Niña:**
    - **Cooler and Wetter Conditions** : La Niña often brings:
        - Colder-than-average temperatures in **parts of North America and Europe.**
        - **Increased precipitation** in **India, East Africa, Australia and Southeast Asia,** potentially leading to floods.
    - **Droughts** : Conversely, La Niña can cause drier conditions in some regions:
        - **Peruvian Coast** : La Niña can contribute to drier-than-average conditions and more supressed rainfall.

ELLA

What Are ELLA Years?

5. **ELLA** refers to phases when the Pacific Ocean moves between **El Niño and La Niña** conditions, neither fully active.

6. These transitional periods have distinct ocean-atmosphere characteristics, different from ENSO or El Niño Modoki. Based on the IMD-IITM Pune study, **ELLA years bring:**

- **Severe and prolonged summer heat** : with extremely high daytime and nighttime temperatures.
- **More frequent heatwaves** , particularly across northwest and central India.
- **Extended duration of extreme heat** , making summers notably harsher. **Early identification of ELLA years** could improve heatwave forecasting and preparedness in India. Recognizing this pattern adds nuance beyond standard ENSO impacts, highlighting critical transitional phases that affect summer temperatures profoundly.

⚡ **Quick revise** ENSO and the other "oscillation" family

Fold ▲

> 🥶🔥 **El Niño = El Nasty (warm east Pacific → weak SW monsoon, drought India/Australia; NE monsoon stronger). La Niña = La Nice (cold east Pacific → good monsoon, Australian floods; NE monsoon weaker).**

El Nino Modoki

- El Niño Modoki, also known as **"Central Pacific El Niño" or "dateline El Niño,"** is a variation of the traditional El Niño phenomenon.
- The term **"Modoki"** comes from a **Japanese word meaning "similar but different."**



- Unlike the conventional El Niño, where the warmest sea surface temperatures (SSTs) are found in the eastern Pacific Ocean, El Niño Modoki features **warm SST anomalies centered in the central Pacific, with cooler temperatures in both the eastern and western Pacific.**
- **Characteristics of El Nino Modoki:**
  - **Location of Warm SSTs** : In El Niño Modoki events, the central Pacific (around the dateline) experiences significant warming, while the eastern Pacific may not see as much warming and can even experience cooling.
  - **Atmospheric Patterns** : The atmospheric convection and related weather patterns shift from the eastern to the central Pacific, influencing global climate differently compared to the traditional El Niño.

Pacific Decadal Oscillation (PDO)

The **Pacific Decadal Oscillation (PDO)** is a **long-term fluctuation in sea surface temperatures (SSTs) and atmospheric circulation over the North Pacific Ocean** , with each phase typically lasting **20–30 years** . It has two phases: **positive (warm)** , when the eastern Pacific is warmer and the western Pacific cooler than average, and **negative (cool)** , with the opposite pattern. Unlike **ENSO** , which operates on a 2–7 year timescale, the PDO influences climate over **decadal scales** and can amplify or weaken El Niño and La Niña impacts. It affects **temperature, rainfall, marine ecosystems, fisheries, and storm tracks** across the Pacific basin. In India, the PDO can **modulate monsoon variability by influencing ENSO teleconnections** , thereby affecting droughts, floods, and long-term climatic trends.

Indian Ocean Dipole

- The Indian Ocean Dipole (IOD) is a climate phenomenon characterized by differences in sea surface temperatures (SSTs) between the western and eastern parts of the Indian Ocean. It has significant impacts on weather and climate patterns in the surrounding regions.
- **Characteristics of Indian Ocean Dipole**
  - **Positive IOD:** During a positive IOD event, the **Western Indian Ocean** (near the eastern coast of Africa) becomes **warmer than usual** , while the **Eastern Indian Ocean** (near Indonesia and Australia) **becomes cooler** .

- **Negative IOD** : In a negative IOD event, the **Eastern Indian Ocean warms up** while the **Western Indian Ocean cools down** , creating the opposite dipole pattern.

- **Climatological Impacts of Indian Ocean Dipole**
  - The IOD has wide-ranging impacts on weather and climate across the Indian Ocean rim and beyond:
  - **Indian Subcontinent:**
    - **Monsoon Variability** : A **positive IOD often enhances the Indian monsoon** , leading to above-average rainfall in the Indian subcontinent.
    - Conversely, a **negative IOD can suppress the monsoon** , causing drought conditions in the region.
  - **Australia** : **Positive IOD** events are typically associated with **drier conditions** in Australia, particularly in the southeastern parts, leading to droughts and increased bushfire risk. **Negative IOD** events bring **wetter conditions** , increasing the likelihood of floods.
  - **East Africa** : **Positive IOD** events can lead to **increased rainfall** in East Africa, resulting in wetter conditions and sometimes causing floods. **Negative IOD** can lead to **drier conditions** and droughts in the region.
  - **Indonesia and Southeast Asia** : During a **positive IOD** , Indonesia and parts of Southeast Asia can experience **drier conditions** , affecting agriculture and water resources. **Negative IOD** usually brings **wetter conditions** to these areas.
- **Global Climate Teleconnections** : El Niño Interaction: The IOD **often interacts with El Niño-Southern Oscillation** (ENSO) phenomena.

Atlantic Ocean Dipole

- The Atlantic Ocean Dipole (AOD), **also known as the Atlantic Zonal Mode** , refers to the climate variability pattern characterized by sea surface temperature (SST) anomalies between the northern tropical Atlantic (NTA) and the southern tropical Atlantic (STA).
- **Characteristics of Atlantic Ocean Dipole**
  - **Positive Phase** : During the positive phase, the **northern tropical Atlantic experiences warmer SSTs, while the southern tropical Atlantic sees cooler SSTs** .
  - **Negative Phase** : In the negative phase, the **northern tropical Atlantic has cooler SSTs** , and the **southern tropical Atlantic is warmer** .

Atlantic Nino & Nina

- **Atlantic Nina:** The Atlantic Ocean cousin of La Nina. When the **eastern equatorial Atlantic cools down more than normal** , western equatorial Atlantic warms up.
- **Atlantic Nino: warming of eastern equatorial Atlantic** and **cooling of Western equatorial Atlantic.**
- Do **not have as wide climatic impacts as Pacific** El Nino and la nina.

Arctic Ocean Dipole

- The Arctic Ocean Dipole (AOD) or **Arctic Dipole Anomaly** refers to the sea level pressure pattern in the Arctic.

Madden Julian Oscillation

- The Madden-Julian Oscillation (MJO) is a major atmospheric phenomenon that significantly influences global weather patterns and climate. It is characterized by a **large-scale coupling between atmospheric circulation and tropical deep convection, primarily affecting the Indian and Pacific Oceans.**



- **Characteristics of the MJO**
  - Propagation: The MJO is an **eastward-moving disturbance of clouds, rainfall, winds, and pressure that typically recurs every 30 to 60 days.**
  - It **travels from the Indian Ocean across the Pacific Ocean and sometimes continues into the Atlantic.**
  - Phases: The MJO consists of two primary phases:
    - **Active (Convective) Phase:** Characterized by **enhanced cloudiness and precipitation.** This phase is associated with increased convection and storm activity.
    - **Inactive (Suppressed) Phase:** Marked by **reduced cloudiness and precipitation** , leading to drier and more stable conditions.
- **Climatological Impacts of the MJO**
  - **Tropical Regions:**

- **Monsoon Variability:** The MJO affects monsoon systems in Asia, Africa, and Australia. During the **active phase, it can enhance monsoon rainfall** , while the **inactive phase can suppress it** . For instance, the Indian monsoon can see increased rainfall during the MJO's active phase. The shorter the cycle, the better the impact on the monsoon.
- **Cyclone Formation:** The MJO can modulate the formation of tropical cyclones. **Enhanced convection during the active phase increases the likelihood of cyclone development** in the Indian Ocean, Pacific Ocean, and Atlantic Ocean.

⚡ **Quick revise** ENSO and the other "oscillation" family

Fold ▲

Phenomenon	What it is	India link
El Niño	trade winds weaken; warm water sloshes east; **Humboldt (Peruvian) current replaced**, less upwelling	**weak SW monsoon**; Australia drought/fire; Peru floods; California storms
La Niña	trades strengthen; cold east Pacific; more upwelling	**better monsoon**; floods in Australia, E. Africa, SE Asia
ELLA years	**transition** between El Niño and La Niña (IMD–IITM Pune study)	**severe, prolonged summer heat; more heatwaves in NW & central India**
El Niño Modoki	"similar but different": **warmth in the central Pacific (dateline)**, cooler east and west	different global pattern
PDO	North Pacific, each phase **20–30 yrs** (ENSO = 2–7 yrs); positive = warm east	modulates ENSO links to monsoon
IOD	west–east SST gap in the Indian Ocean	**Positive IOD (warm west near Africa, cool east) → stronger monsoon**, even in El Niño years; negative → weaker
Atlantic Niño/Niña, Atlantic dipole	Atlantic equivalents	not as wide-ranging
Arctic Ocean Dipole	Arctic sea-level pressure pattern	–
MJO	**eastward** cloud-rain pulse, **30–60 days**; active (wet) vs suppressed (dry)	active phase enhances monsoon; modulates **cyclone formation**; *shorter cycle = better monsoon impact*

Seasons Of India: Retreating Monsoon Season

- **Shift of ITCZ** and increase in pressure over Northern India.



Heavy thunder clouds hovered above parts of sea over the Bay of Bengal and east and central India as part of a broad but weak circulation that set up a fresh low-pressure area, ahead of forecasts, on Tuesday afternoon.

- **October Heat** : **High Temperature and Humidity.** Its causes include:
  - As the southwest monsoon retreats, there is a **transition period where the atmospheric moisture remains high, but the cooling effect of the rains is absent** , leading to increased humidity and heat.
  - With the **reduction in cloud cover** post-monsoon, there is **increased solar radiation** , which raises daytime temperatures.
- **Northeast Monsoon** experienced in this season in **coastal Andhra Pradesh and a major part of Tamil Nadu** , due to movement of the Northeastern trade winds from over the Bay of Bengal.
- During the El Nino years, the SW Monsoons get weaker while the NE Monsoons get stronger.
- During the La Nina years, the NE Monsoons get weaker while the SW Monsoons get stronger.
- Season of **most severe cyclones** , especially in the Bay of Bengal (high sea surface temperature; southward shift of monsoon trough; change in wind direction)

⚡ **Quick revise** Retreating monsoon (Oct–Nov) Fold ▲

Retreating monsoon (Oct–Nov): ITCZ shifts, pressure rises over north India; **"October heat"** (humid, clear skies); **NE monsoon rain in coastal Andhra and most of Tamil Nadu**; El Niño years: NE stronger; **most severe cyclones** (Bay: hot sea, trough moves south, winds change).

Spatial Distribution Of Rainfall In India

- **High Rainfall Regions (>200 cm annually)**
  - **Windward side of Western Ghats:** This includes the western coastal strip and the Western Ghats, extending from southern Gujarat through Maharashtra, Goa, Karnataka, and Kerala.

- **Northeastern India:** Primarily Assam, Meghalaya, and parts of Arunachal Pradesh. Notable places include Cherrapunji and Mawsynram, which receive some of the highest rainfall in the world. **Funneling effect:** Clouds get channeled into constricted space between mountains. Happens in NE India.



- **Andaman and Nicobar Islands:** Island location. Experience both South-West and North-East Monsoons.

- **Medium Rainfall Regions (100-200 cm annually)**
  - **Foothills of the Himalayas:** This includes regions in Uttarakhand, Himachal Pradesh, and northern West Bengal (like Darjeeling).
  - **Eastern India:** Odisha, Jharkhand, Bihar, Eastern MP, parts of North East India. East Tamil Nadu (NE Monsoon).
- **Low Rainfall Regions (50-100 cm annually)**
  - **Parts of Peninsular India:** Parts of Deccan plateau in Maharashtra and Karnataka.
  - **North-Western India:** Punjab, Haryana, western Uttar Pradesh, Gujarat, and Eastern Rajasthan. Parts of Jammu Kashmir.
- **Very Low Rainfall Regions (<50 cm annually)**
  - **Rajsthan:** This includes the semi-arid regions of Rajasthan, particularly the Western part.
  - **Rain Shadow Regions:** The eastern slopes of the Western Ghats, including parts of Maharashtra, Karnataka, and Tamil Nadu. Also Ladakh (rainshadow of the Himalayas).

⚡ **Quick revise** Rainfall map — Fold ▲

Rainfall map: >200 cm – windward W. Ghats (S. Gujarat → Kerala), NE (Assam, Meghalaya, Arunachal; **funnelling**: Cherrapunji, Mawsynram), Andaman & Nicobar · **100–200** – Himalayan foothills, east India, east TN

· **50–100** – parts of the Deccan, NW India · **<50** – west Rajasthan, W. Ghats rain-shadow, **Ladakh**.

PYQs

Q.) La Nina is suspected to have caused recent floods in Australia. How is La Nina different from El Nino?

1. La Nina is characterised by unusually cold ocean temperature in the equatorial Indian Ocean whereas El Nino is characterised by unusually warm ocean temperatures in the equatorial Pacific Ocean.

2. El Nino has an adverse effect on the south-west monsoon of India, but La Nina has no effect on monsoon climate.

Which of the statements given above is/are correct? (2011)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: D La Niña is a climate pattern that describes the cooling of surface ocean waters along the tropical west coast of South America (It means when Peru/Chile cold current reestablished itself).

- La Nina is considered to be the counterpart to El Nino, which is characterised by unusually warm ocean temperatures in the equatorial region of the Pacific Ocean.
- Over the Indian subcontinent, El Nino during winter results in the development of warm conditions. During summer, it leads to dry conditions and deficient monsoon.
- It also leads to drought in Australia.
- On the other hand, La Nina results in better than normal monsoon in India.
- At the same time, in Australia, it has caused floods.
- La Nina is characterised by unusually cold ocean temperature in the equatorial Indian Ocean whereas, El Nino is characterised by unusually warm ocean temperature in the equatorial Pacific Ocean.El Nino has an adverse effect on the south-west monsoon of India, but La Nina has no effect on the monsoon climate.

Q.) With reference to ‘Indian Ocean Dipole (IOD)’ sometimes mentioned in the news while forecasting Indian monsoon, which of the following statements is/are correct?

1. IOD phenomenon is characterized by a difference in sea surface temperature between tropical Western Indian Ocean and tropical Eastern Pacific Ocean.

2. An IOD phenomenon can influence El Nino’s impact on the monsoon.

Select the correct answer using the code given below: (2017)

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: B The Indian Ocean Dipole (IOD) is defined by the difference in the sea surface temperature between the two equatorial areas of the Indian Ocean – a western pole near the Arabian Sea (in western Indian Ocean) and an eastern pole closer to the Bay of Bengal (in the eastern Indian Ocean).

- The IOD affects the climate of Southeast Asia, Australia and other countries that surround the Indian Ocean Basin. The Indian Monsoon is invariably influenced by the IOD.
- A positive IOD occurs when the sea surface temperatures are greater than normal in the Arabian Sea and less than normal in the tropical eastern Indian Ocean.
- When the reverse is the case, a negative IOD is said to have developed.
- A positive IOD leads to greater monsoon rainfall and more active (above normal rainfall) monsoon days while negative IOD leads to less rainfall and more monsoon break days (no rainfall).
- El-Nino conditions normally result in diminished monsoonal rainfall.
- However, if there is a positive IOD, it can bring good rains to India despite an El Nino year.

Q.) With reference to the climate of Andaman and Nicobar Islands, which of the following statements is/are correct?

1. The climate can be defined as a humid, tropical coastal climate.

2. It receives rainfall from both South-west monsoon and North-east monsoon.

3. Maximum precipitation is between December and May.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Answer: B The Andaman and Nicobar Islands experience a humid tropical maritime/coastal climate. In Ocean, so both SW and NE winds are onshore. Maximum rainfall generally occurs during the South-West Monsoon months, especially May to September. (Heat=LP=rain, and NH has heat in March-Sept)

⚡ **Quick revise** Climate PYQs Fold ▲

Year	Asked	Answer
2011	La Niña vs El Niño: (1) La Niña = cold equatorial **Indian** Ocean? (2) La Niña has	**D – neither** (La Niña is cold *Pacific* water; it

Year	Asked	Answer
	no effect on monsoon?	helps the monsoon)
2017	IOD: (1) difference between W Indian Ocean and **Eastern Pacific**? (2) IOD can influence El Niño's effect	**B – only 2** (IOD = W vs **E Indian Ocean**)
–	Andaman & Nicobar climate: humid tropical coastal ✔; SW + NE monsoon ✔; max rain Dec–May ✘	**B (1 and 2)**; max rain May–Sept (SW monsoon)

13 · FORESTS OF INDIA

India boasts of a **variety of vegetation** , owing to the **diverse climatic conditions** .
Important forest types in India include:

- Tropical Evergreen and Semi Evergreen forests.
- Tropical Deciduous forests.
- Tropical Thorn forests.
- Montane forests.
- Littoral and Swamp forests.



Tropical Evergreen Forests:

- The **trees do not shed their leaves** due to optimal conditions all year round.
- Due to high canopy density, they **lack an undergrowth.**
- Depending upon the precipitation levels, they can be further divided into the following types:
- **Wet Evergreen:**
  - **Rainfall >200cm; Temperature: 25-27 degrees C**
  - Found in Western Ghats (south of Mumbai), Northeast India, Andaman & Nicobar
  - Jackfruit, Rosewood, Mahogany, Aini, ebony, Agarwood (Oudh) etc
- **Semi-Evergreen:**
  - **Bordering wet evergreen**
  - **Rainfall: 175-225cm; Temperature: 24-27 degrees C**
  - Found in the Western Coast, Assam, the lower slope of Eastern Ghats, Odisha, and parts of Andamans.
  - **Transition between wet evergreen and moist deciduous** .
  - Have a **mix of trees from both** the above forests.
  - **Evergreen due to undergrowth** .

© NCERT not to be republished

- Kadam, Cashew, cedar, chestnut, thorny bamboo, etc.

- **Dry Evergreen**
  - About **100 cm of rain** spread throughout the year.
  - Found on the **Tamil Nadu coast** .
  - They have **short trees** as the amount of rainfall here is low.
  - Tamarind, Neem, toddy palm, etc.

⚡ **Quick revise**

Andaman & Nicobar forests (for the 2015 PYQ) Fold ▲

Andaman & Nicobar forests (for the 2015 PYQ): ~2,200 plant varieties, 200 endemic, 1,300 not on the mainland · South Andaman epiphytes (ferns, orchids) · **Middle** moist deciduous · **North** wet evergreen + woody climbers · **North Nicobar** has **no evergreen forest**; central/southern Nicobars mostly evergreen · **grasslands only in Nicobars** · forest cover **86.2%** · mangroves **1,136 km² (Andaman) + 35 km² (Nicobar)**.

Tropical Deciduous Forests

- The trees in these forests **shed their leaves once a year.**
- Based on the amount of precipitation received, can be classified into:
- **Moist Deciduous:**
  - **Rainfall: 100-200cm; Temperature: about 27 degrees C**
  - Found in parts surrounding western ghats, NE India, Shiwalik range (along Terai & Bhabher belt); hills of Eastern MP, CG, Chotanagpur plateau; most of Odisha, parts of WB & Andaman and Nicobar.
  - They have a **shrubby undergrowth.**
  - Valuable trees like teak, sal, mahua, tendu, bamboo, etc.
- **Dry Deciduous:**
  - **Rainfall: 75-100 cm.**
  - **Transition between moist deciduous and thorny forests** .
  - Found in the rainier parts of the Peninsula, plains of UP, Bihar, and Madhya Pradesh.
  - Some parts of Rajasthan, also show these type of forests, but with scantier growth.
  - Teak, tendu, Red Sanders, sal, Khair, etc.

Tropical Thorny Forests:

- **Found in Arid locations.**
- **Rain <50cm; 25-30 degree C**
- **Low, widely scattered trees**
- In India, they are present in Arid, Semi-arid Northwestern India: Rajasthan, SW Punjab, Western Haryana, Kachchh, Saurashtra, some parts of UP; lee-side of Western Ghats (MH, Karnataka, Telangana, Andhra Pradesh, Tamil Nadu)

- Acacia, Babul, Neem, Khair, Cacti. In wetter margins, Indian wild date and some grasses.

Littoral & Swamp

- **In and around deltas, estuaries, and creeks.**
- Highly developed in the Andaman and Nicobar Islands and Sunderbans (Sundri tree) of West Bengal. Also seen in Mahanadi, the Godavari and the Krishna deltas.
- These are the **mangrove forests** , which are important for **protecting the coastline from erosion, storm surges, and tsunamis,** but their capacity to adapt to rising sea levels is being compromised. Mangroves play a crucial role in coastal protection, biodiversity, and carbon sequestration, but their capacity to adapt to rising sea levels is being compromised.
- **India has about 7% of the world's Mangrove forests** .
- **Survive and grow in both fresh & Brackish water.**
- Have **special adaptations** like **Viviparity and Pneumatophores** .
- Sundri, Keora, Agar, Rhizophora, etc.

⚡ **Quick revise** Mangroves protect Fold ▲

Mangroves protect coast from erosion, storm surge, tsunami; trap sediment (protect coral and seagrass); fisheries nurseries; timber, fodder, medicine; tourism. **Handout alert: capacity to adapt to rising seas is being compromised.**

Alpine:

- **In the mountain regions, depending on the altitude** , different types of forests are found.
- **In the Northern Mountains:** The **base region** has **moist deciduous forests** , **as the height increases, the forests become temperate.** At **further heights, beyond the tree line, up to the snow line, there is alpine tundra vegetation** .
  - At 1000-2000m we get wet temperate forests with Pine trees. British replaced Oaks with Pine in these forests, and used it for laying Railways.
  - Above this upto 3000 m (approx) are Blue Pine, Spruce and some Grasslands.
  - 3000-4000m: Alpine forests (Silver firs, Junipers, Rhodohenderon), and Grasslands (Bugyals)
  - At further heights, beyond the tree line, up to the snow line, there is alpine tundra vegetation.
  - The **Northern slopes of the Himalayas do not have much vegetation** , as this region continuously remains devoid of sunlight.
- **Southern Mountains (Western Ghats):**
  - **On their western slopes** , at the **base** they have **evergreen forests** which eventually **convert into temperate evergreen forests known as sholas** , as the height increases. Found in Nilgiris, Annmalai and Palani hills mainly.

- **On the eastern slopes** there are **dry evergreen or deciduous forests** , which again **change into temperate sholas** as the height increases.
- Due to **height limitation, the southern forests do not have alpine tundra type of vegetation.**

⚡ **Quick revise** Forests of India Fold ▲

🌳 **Rainfall ladder (cm): >200 wet evergreen → 175–225 semi-evergreen → 100–200 moist deciduous → 75–100 dry deciduous → ~100 spread through year: dry evergreen (TN coast) → <50 thorn**. *Wetter = evergreener, taller; drier = shorter, thornier.*

Forest	Climate	Where	Trees / traits
Wet evergreen	>200 cm; 25–27 °C	W. Ghats (south of Mumbai), NE, A&N	**no undergrowth** (dense canopy); jackfruit, rosewood, mahogany, aini, ebony, agarwood
Semi-evergreen	175–225; 24–27 °C	W. coast, Assam, lower E. Ghats, Odisha, Andamans	wet evergreen + moist deciduous mix; evergreen due to undergrowth; kadam, cashew, cedar, chestnut, thorny bamboo
Dry evergreen	~100 cm all year	**TN coast**	short; tamarind, neem, toddy palm
Moist deciduous	100–200; ~27 °C	around W. Ghats, NE, Shiwalik (Terai–Bhabar), E. MP, CG, Chotanagpur, Odisha	shrubby undergrowth; **teak, sal, mahua, tendu, bamboo** (also shisham, sandalwood, khair, kusum, arjun, mulberry)
Dry deciduous	75–100	rainier peninsula, plains of UP, Bihar, MP, Rajasthan	transition to thorn; teak, tendu, **red sanders**, sal, khair
Thorn	<50; 25–30 °C	Rajasthan, SW Punjab, W. Haryana, Kachchh, Saurashtra, lee of W. Ghats	low scattered; acacia, babul, neem, khair, cacti, wild date
Littoral & swamp (mangroves)	brackish water	**Sundarbans (Sundri)**, A&N, Mahanadi–Godavari–Krishna deltas	**viviparity, pneumatophores**; sundri, keora, agar, rhizophora; **~7% of world mangroves**

Mountain forests. *Himalaya (north):* base moist deciduous → **1,000–2,000 m wet temperate with pine (British replaced oak with pine for railway sleepers)** → up to ~3,000 m **blue pine, spruce, grassland** → **3,000–4,000 m silver fir, juniper, rhododendron + bugyals** → tree line → alpine tundra → snow line. Northern Himalayan slopes carry little vegetation. *Western Ghats (south):* evergreen at base → **temperate**

evergreen "sholas" (Nilgiris, Anamalai, Palani) on both slopes; **no alpine tundra** (height limit).

Status of Indian Forests:

- **Key Points:**
  - Total forest and tree cover is 8,27,357 sq km (25.17% of India's geographical area).
  - Forest cover is 7,15,343 sq km (21.76%), and tree cover is 1,12,014 sq km (3.41%).
  - As compared to 2021, forest and tree cover increased by 1445 sq km (156 sq km increase in the forest cover and 1289 sq km increase in tree cover)
  - However, this is still below the recommended 33% for ecological stability.
  - The total mangrove cover in the country is 4,992 sq km.
  - The bamboo-bearing area has increased compared to the 2021 assessment by 5227 sq. km.
  - Forest cover in hill districts increased by 234.14 sq.km.
  - 19 states/UTs have over 33% forest cover, with 8 states/UTs—Mizoram, Lakshadweep, Andaman & Nicobar Islands, Arunachal Pradesh, Nagaland, Meghalaya, Tripura, and Manipur—having over 75% forest cover.
  - Among the six megacities—Bengaluru, Chennai, Delhi, Kolkata, Hyderabad, and Mumbai—Delhi has the highest forest cover

Category	Details
Largest Total Forest and Tree Cover	Madhya Pradesh, Arunacha
Largest Forest Cover	Madhya Pradesh, Arunacha
Highest Percentage of Forest Cover	Lakshadweep (91.33%), Mi Nicobar Islands (81.62%)
Maximum Increase in Total Cover	Chhattisgarh, Uttar Pradesh
Maximum Increase in Forest Cover	Mizoram, Gujarat, Odisha
Maximum Increase in Tree Cover	Gujarat, Bihar, Kerala
Maximum Bamboo-Bearing Area	Madhya Pradesh, Arunacha

- **Concerns:**

- Total cover grew only by 1445 sq.km, compared to 1540 sq.km in 2021. Notable decrease in Andhra Pradesh and Maharashtra.
- Decline in total mangrove cover by 7.43 sq.km.
- In Western Ghats Ecologically Sensitive Area, an overall loss of 58.22 km² in forest cover in the past 10 years.
- In North-East India, decline in forest cover by 327.30 sq. km

- **Difference between 2021 & 2023 reports:**
  - Separate chapter on Agroforestry.
  - Re-introduction (after 10yrs) of chapter on "Forest characteristics" which talks about soil health, grazing instances etc.
  - Bamboo cover estimates done and added under tree cover.
  - For the first time, an estimate of tree cover contributed by smaller trees with a diameter of 5 to 10 cm at breast height has been provided. Reason: smaller trees like Sababul are being planted for commercial purposes.
  - Gives estimates for 751 districts compared to 636 in 2021.
- **FAO Report:**
  - According to the latest report (**State of World's Forests Report**) **by FAO** (released July 22, 2024), **India gained 2,66,000 hectares of forest area annually from 2010 to 2020** , securing the **third spot** (after China and Australia) **among the top 10 countries with the most significant forest area gains during this period.**
  - FAO **appreciated Indian efforts in Agroforestry and reclamation of degraded lands** for forestry.
  - **FAO's Global Forest Resources Assessment (GFRA) 2025** report indicates that **India has risen to 9th globally** in total forest area and **maintains its 3rd position for annual net forest area gain** . The report also highlights that global deforestation rates have slowed but remain high, and **Asia is the only continent to have seen an increase in forest area since 1990** , thanks to contributions from countries like India and China.

⚡ **Quick revise** Forest status (handout numbers) Fold ▲

Forest status (handout numbers).

Item	Figure
Forest + tree cover	**8,27,357 km² = 25.17%** of area (target **33%** for stability)
Forest cover	**7,15,343 km² = 21.76%** · Tree cover **1,12,014 km² = 3.41%**
Change vs 2021	**+1,445 km²** (forest +156, tree +1,289) – lower than **+1,540** in 2021
Mangrove	**4,992 km²** (**−7.43 km²**)
Bamboo area	**+5,227 km²**
>33% forest	**19** states/UTs; **>75%: 8** – Mizoram, Lakshadweep, A&N, Arunachal, Nagaland, Meghalaya, Tripura, Manipur
Highest % cover	**Lakshadweep 91.33 · Mizoram 85.34 · A&N 81.62**

Item	Figure
Largest forest + tree cover	**MP, Arunachal, Maharashtra** · largest *forest* cover: **MP, Arunachal, Chhattisgarh**
Max increase total	**Chhattisgarh, UP, Odisha** · forest: **Mizoram, Gujarat, Odisha** · tree: **Gujarat, Bihar, Kerala**
Biggest bamboo area	**MP, Arunachal, Maharashtra**
Megacities	**Delhi has the highest forest cover** among the six
Worries	decline in **Andhra Pradesh, Maharashtra**; **Western Ghats ESA −58.22 km² in 10 years**; **NE −327.30 km²**
New in 2023 report	agroforestry chapter; **"forest characteristics" chapter back after 10 years**; bamboo in tree cover; first estimate of **5–10 cm DBH small trees**; **751 districts** (636 in 2021)
FAO	India **+2,66,000 ha/year, 2010–2020, 3rd** (after China, Australia); **GFRA 2025: 9th in total forest area, 3rd in annual gain**; Asia the only continent with forest gain since 1990

Social Forestry

- Social forestry is a **land management approach that aims to increase tree cover and improve the livelihoods of rural communities** through the involvement of local people.
- The emphasis is on planting trees that meet the social and economic needs of local communities, while also providing environmental benefits.
- **The term "social forestry" was first used in India in 1976 by the National Commission on Agriculture.**
- Its types include:
  - **Community Forestry:**
    - This involves local communities actively participating in the management of **forests on degraded or wasteland.**
  - **Urban Forestry:**
    - This type of social forestry involves **planting and managing trees in urban areas** , including parks, streetscapes, and institutional lands.
    - **Miyawaki method:** Japanese method. Growing native trees in high density in a region.
  - **Extension Forestry:**
    - **Extension of forests in previously non-forested regions.** This approach involves **educating and empowering rural communities about the benefits of trees** and **providing technical assistance** for tree planting and management.
  - **Agroforestry:**
    - This **combines trees and shrubs with crop production** on the same land. (**Not necessarily by a single farmer**).
- **Farm Forestry:**
  - **Focus:** This approach **encourages farmers to integrate trees into their existing agricultural lands** for commercial as well as protection purposes.

- **Benefits:** Trees planted along field borders or in small woodlots provide **wind protection for crops, reduce soil erosion, and offer additional income** from timber, fruits, or other non-timber forest products (NTFPs).

- **Advantages:**
  - **Social Benefits:**
    - Creates **rural employment** opportunities through tree planting and maintenance.
    - Provides **fuelwood, fodder, and other forest products for local communities** , improving livelihoods.
    - **Empowers local communities through participation in decision-making and management of forest resources.**
  - **Economic Benefits:**
    - Provides a **source of income** for rural communities through the sale of timber, fruits, and other non-timber forest products (NTFPs) like medicinal plants and honey.
  - **Environmental Benefits:**
    - Increased tree cover helps **mitigate climate change** by absorbing carbon dioxide.
    - **Reduces soil erosion** and **promotes land reclamation** .
    - Provides **habitat** for diverse species, promoting **biodiversity** .
    - **Reduces dependence on traditional forests** , potentially leading to less deforestation.
    - Supports **sustainable agriculture** by improving soil quality and providing wind protection.

⚡ **Quick revise** Social forestry Fold ▲

Social forestry – term first used in India in **1976 by the National Commission on Agriculture.** Types: **community, urban (Miyawaki = dense native planting, Japanese), extension, agroforestry** (trees with crops, not necessarily one farmer), **farm forestry.** Benefits: jobs, fuel/fodder, NTFPs, carbon, erosion control, habitat, less pressure on natural forests.

14 · SOILS OF INDIA

Multiple factors affect the formation of soils:

- **Relief (Topography):**
  - **Slope:** Steeper slopes experience faster erosion, leading to thinner and less developed soils. Gentle slopes allow for better weathering and accumulation of organic matter, promoting richer soil development.
  - **Drainage:** Well-drained areas allow for water to percolate through the soil, influencing nutrient leaching and mineral composition. Poorly drained areas can lead to waterlogging, affecting oxygen availability and plant growth.

- **Aspect:** South-facing slopes in the Northern Hemisphere receive more sunlight, leading to faster decomposition of organic matter and potentially drier soils. North-facing slopes may have slower decomposition and cooler, moister soils.

- **Parent Material (Bedrock):**
  - **Texture:** The type of rock the soil forms from determines its initial mineral content. For example, granite weathers into sandier soils, while basalt weathers into clay-rich soils.
  - **Chemical composition:** Parent material can influence soil fertility. Rocks rich in certain minerals like calcium or potassium can contribute to a more fertile soil.
- **Climate:**
  - **Temperature:** Warmer temperatures accelerate the rate of chemical weathering of rocks, releasing minerals for soil formation. Cold temperatures slow down the process.
  - **Precipitation:** Rainfall patterns directly impact soil development. High rainfall can leach nutrients from the upper layers, while low rainfall can lead to salinization in arid regions.
- **Vegetation and Other Forms of Life (Biota):**
  - **Plants:** Plant roots help break down rocks, contributing to physical weathering. Decomposing plant matter adds organic material to the soil, enhancing fertility and soil structure.
  - **Microorganisms:** Bacteria and fungi in the soil break down organic matter and rock minerals, releasing nutrients for plants and influencing soil structure.
  - **Animals:** Earthworms and burrowing animals improve aeration and drainage by creating tunnels in the soil.
- **Time:**
  - Soil formation is a slow and continuous process. Over time, the interaction of the above factors leads to the development of distinct soil horizons with varying characteristics. The **older the soil, the more developed and mature its profile will be** .
  - Time helps in the further disintegration of the rocks into finer material and the addition of minerals to the soil via various biotic and abiotic activities.
- **Anthropogenic Factors:**



The Indian Council of Agriculture Research Defines 8 types of soils in India:

- Alluvial soil
- Red and Yellow soil
- Black soil
- Arid Desert soil
- Laterite soil
- Saline soil
- Peaty / marshy soil
- Forest & Mountain soil

Descriptions of the soils:

- **Alluvial:**
  - The **most common soil of India** , **covering >40% Indian area.**
  - Found in Northern Plains, Coastal Plains.
  - **Types include Bhangar, i.e. the less fertile older alluvium, and Khadar, i.e. the more fertile newer alluvium.**
  - **Rich** in **Potash, and Lime, Phosphoric acid** .
  - **Poor** in **Nitrogen, Humus** .
  - Ideal for the growth of **sugarcane, paddy, wheat, and other cereal and pulse crops** .
- **Black (Regur):**
  - **3rd common soil of India.**
  - Plateaus of Maharashtra, Saurashtra, Madhya Pradesh, and Chhattisgarh; also extend in the Southeast direction along the Godavari and the Krishna valleys.

- **Rich: Magnesium, potash, iron, lime and alumina.**
- **Poor: Phosphoric contents, Nitrogen, and organic matter.**
- When dry, it develops cracks which gives it **self-ploughing ability.**
- **High water retention** , making it an ideal soil for rain-fed agriculture.
- **Important crops: Cotton and sugarcane, Pulses**
- Very ideal for the growth of cotton, and often **called Cotton soil too.**

- **Red (& Yellow) soil**
  - Develops on **crystalline igneous rocks, rich in Iron, giving it a red hue.**
  - Becomes **yellow** when iron occurs in **hydrated form.**
  - Red soil is found in areas of low rainfall in the eastern and southern parts of the Deccan Plateau and also along the Piedmont zone of the Western Ghats.
  - Yellow in parts of Odisha & Chhattisgarh.
  - **Rich in Iron, Potash; Poor in Nitrogen, Phosphorous, and Humus**
  - **Fertile if fine-grained.**
  - **Important Crops: Wheat, cotton, pulses, tobacco, oilseeds, potato.**
- **Laterite Soil**
  - **In the wetter margins, the red soil converts into Laterite soil.**
  - It is an **acidic** soil, making it unsuitable for cultivation.
  - **Poor: organic matter, nitrogen, phosphate, and calcium.**
  - **Rich: iron oxide and potash are in excess.**

⚡ **Quick revise** Formation factors — Fold ▲

Formation factors: relief · parent rock · climate · biota · time · humans.

Types of Soils in India

⚡ **Quick revise** Soils of India — Fold ▲

> 🎨 **ICAR's 8 soils – "A Red Black Arid Laterite Saline Peaty Forest"** → **A**lluvial · **R**ed & yellow · **B**lack · **A**rid desert · **L**aterite · **S**aline · **P**eaty/marshy · **F**orest & mountain. **Memory pairs: Black = cotton + cracks (self-ploughing); Red = iron; Laterite = tea-coffee-cashew (acidic, leached); Alluvial = the breadbasket.**

Descriptions of the soils:

- Found in Southern states, the Western Ghats region of Maharashtra, Odisha, some parts of West Bengal, and the North-east regions.
- **Tea, coffee, Cashew**

- **Arid Soils**

- **Sandy & Saline** soil
- **Lack of moisture, Humus, and Nitrogen; Phosphorous is adequate.**
- **Include Calcium Kankar nodules** which reach the topsoil layers by the capillary action.
- **Can be cultivated if irrigation is available** . E.g. The Indira Gandhi Canal project has helped cultivate these soils.
- Found in Rajasthan, in and around the Thar desert.
- **Millets, Fodder crops, some beans (Guar Beans), wheat if irrigated** .

- **Saline Soils**
  - Also called **Usara soil.**
  - **Infertile, with a large proportion of salts** of Sodium, Potassium, Magnesium, etc.
  - Found in arid and semi-arid parts of Rajasthan, Punjab, Uttar Pradesh, Haryana, Gujarat (Kachchh), and Maharashtra.
  - **In some parts, coconuts, and salt resistant fodder crops** being grown (**White Revolution Impact**).
- **Peaty/Marshy**
  - In the **areas of heavy rainfall and high humidity.**
  - **Rich in humus** , and **organic content may go even up to 40-50 percent** .
  - **If properly drained and fertilized** , can be used for cultivating **crops like rice** .
  - In Odisha, the Sundarbans in West Bengal, parts of Bihar, Uttar Pradesh, and parts of Kerala.
- **Forest & Mountain Soil**
  - In **Forest areas, with adequate rainfall** .
  - In Forest regions of Himalayas, Western and Eastern Ghats.
  - They **vary in their structure and texture depending upon the mountain environment where they are formed.**
  - **Valley sides- loamy and silty** soil (can be used to grow rice and wheat)
  - **Upper slopes- coarse-grained** soil (can be used for Horticulture)

⚡ **Quick revise** Soils of India — Fold ▲

Soil	Where	Rich / poor	Crops & hooks
Alluvial	N. plains, coastal plains; **>40% of India**, most common	rich: **potash, lime, phosphoric acid**; poor: **nitrogen, humus**	sugarcane, paddy, wheat, pulses; **Bhangar / Khadar**
Black (Regur)	Maharashtra, Saurashtra, MP, CG; along Godavari–Krishna; **3rd most common**	rich: **Mg, potash, iron, lime, alumina**; poor: phosphoric, N, organic	**cotton**, sugarcane, pulses; **cracks when dry**; **high water retention**; **volcanic (Deccan/Rajmahal trap lava)**
Red & yellow	low-rain **E and S Deccan**; Western	rich: iron, potash; poor: N,	crystalline igneous rocks; yellow = **hydrated iron**;

Soil	Where	Rich / poor	Crops & hooks
	Ghat piedmont; yellow in parts of Odisha/CG	phosphorus, humus	wheat, cotton, pulses, tobacco, oilseeds, potato
Laterite	wetter margins; south, W. Ghats of Maharashtra, Odisha, WB, NE	iron oxide + potash high; poor in organic, N, phosphate, calcium	**acidic**; **tea, coffee, cashew** (tapioca – PYQ)
Arid	Rajasthan/Thar	little humus, N; **phosphorus adequate**; **kankar**	millets, fodder, guar; **Indira Gandhi Canal**
Saline (Usara)	arid/semi-arid Rajasthan, Punjab, UP, Haryana, **Kachchh**, Maharashtra	salts of Na, K, Mg	some coconut, salt-resistant fodder (White Revolution)
Peaty/marshy	heavy rain: Odisha, **Sundarbans**, Bihar, UP, Kerala	**humus/organic 40–50%**	rice if drained
Forest & mountain	Himalaya, W & E Ghats	varies	valley sides loamy-silty (rice, wheat); upper slopes coarse (horticulture)

Soil Degradation

- Soil degradation is the **decline in quality and health of soil** . It's a gradual process caused by various factors that diminish the soil's ability to support life. Excessive degradation can cause desertification.
- According to a UNCCD Report, deforestation, urbanization, and unsustainable farming practices are causing land degradation at an unprecedented scale, affecting 15 million square kilometers—an area larger than Antarctica. The report emphasizes that the global area impacted by land degradation is expanding by about 1 million square kilometers each year, posing significant threats to biodiversity, climate stability, and human livelihoods.
- **Causes of Soil Degradation:**
  - **Soil Erosion:** Wind and water can wear away topsoil, the most fertile layer. This happens due to:
    - **Deforestation:** Lack of trees exposes soil to wind and heavy rain.
    - **Unsustainable farming practices:** Tilling practices and leaving fields bare leave soil vulnerable.
    - **Overgrazing:** Excessive grazing by livestock removes vegetation and compacts soil.
  - **Chemical Degradation:**
    - **Excessive use of fertilizers and pesticides** can disrupt the natural balance of nutrients and soil organisms.

- **Improper use of chemicals** can lead to salinization, where salts accumulate in the soil, making it unsuitable for plant growth.
- **Improper irrigation practices** can lead to an increase in the acidity of soils.

- **Physical Degradation** : Activities like mining, construction, and heavy machinery traffic can damage soil structure and reduce fertility.

⚡ **Quick revise** Soil degradation — Fold ▲

Soil degradation: erosion (deforestation, bad tillage, overgrazing) · chemical (fertiliser/pesticide, **salinisation**, irrigation acidity) · physical (mining, construction, machinery). **UNCCD: land degradation affects 15 million km² (larger than Antarctica), growing ~1 million km²/year.**

INDIA: A SUB-CONTINENT

- Has **natural borders** , separating it from the rest of Asia (The Himalayas).
- Sits on a **separate plate** which has a geological history separate from the rest of Asia (rest of Asis in Eurasian plate, which was part of Laurasia; India was a part of Gondwanaland).
- Within itself has **almost all the major global climate and forest types** , ranging from deserts to Tundra.
- Because of its Physical separation from the rest of Asia, has a **distinct history, cultures, traditions** , etc.

⚡ **Quick revise** "India: a sub-continent" — Fold ▲

"India: a sub-continent": Himalaya as natural border; **own plate** (Gondwana, vs Eurasian/Laurasia); almost all major climate and forest types (desert to tundra); distinct history/culture.

PYQ

Q.) Consider the following statements: Statement-I : The soil in tropical rain forests is rich in nutrients. **Statement-II** : The high temperature and moisture of tropical rain forests cause dead organic matter in the soil to decompose quickly.

Which one of the following is correct in respect of the above statements? (2023)

- Both Statement-I and Statement-II are correct and Statement-II is the correct explanation for Statement-I
- Both Statement-I and Statement-II are correct and Statement-II is not the correct explanation for Statement-I
- Statement-I is correct but Statement-II is incorrect
- **Statement-I is incorrect but Statement-II is correct**

Answer: D

- The soil of Tropical Rainforests is highly leached and poor in nutrients. Though there is a high availability of leaf litter and high decomposing activity, the trees in the region consume these nutrients as fast as they are produced. This leaves the soil devoid of nutrients.

Q.) The 2004 Tsunami made people realise that mangroves can serve as a reliable safety hedge against coastal calamities. How do mangroves function as a safety hedge? (2011)

- The mangrove swamps separate the human settlements from the sea by a wide zone in which people neither live nor venture out\
- The mangroves provide both food and medicines which people are in need of after any natural disaster
- The mangrove trees are tall with dense canopies and serve as art excellent shelter during a cyclone or tsunami
- The mangrove trees do not get uprooted by storms and tides because of their extensive roots.

Answer: D Benefits of mangroves:

- Coastal protection: The dense root systems of mangrove forests trap sediments flowing down rivers and off the land.
- This helps stabilize the coastline and prevents erosion from waves and storms.
- In areas where mangroves have been cleared, coastal damage from hurricanes and typhoons is much more severe.
- By filtering out sediments, the forests also protect coral reefs and seagrass meadows from being smothered in sediment.
- Fisheries: Mangrove forests are home to a large variety of fish, crab, shrimp, and mollusk species.
- These fisheries form an essential source of food for thousands of coastal communities around the world.
- The forests also serve as nurseries for many fish species, including coral reef fish.
- This makes mangrove forests vitally important to coral reef and commercial fisheries as well.
- Timber and plant products: Mangrove wood is resistant to rot and insects, making it extremely valuable.
- Many coastal and indigenous communities rely on this wood for construction material as well as for fuel.
- These communities also collect medicinal plants from mangrove ecosystems and use mangrove leaves as animal fodder. Recently, the forests have also been commercially harvested for pulp, wood chip, and charcoal production.
- Tourism: Given the diversity of life inhabiting mangrove systems, and their proximity in many cases to other tourist attractions such as coral reefs and sandy beaches, it is perhaps surprising that only a few countries have started to tap into the tourism potential of their mangrove forests.

Q.) Which of the following statements regarding laterite soils of India are correct?

1. They are generally red in colour.

2. They are rich in nitrogen and potash.

3. They are well-developed in Rajasthan and UP.

4. Tapioca and cashew nuts grow well on these soils.

Select the correct answer using the codes given below. (2013)

(a) 1 only

(b) 1 and 2 only

(c) 1 and 4 only

(d) 1, 2, 3 and 4

Ans: C Laterite is soil and rock type **rich in iron and aluminium** but **poor in nitrogen, potash, potassium, lime and organic matter** and is commonly considered to have **formed in hot and wet tropical areas.**

- Nearly all laterites are of **rusty-red colouration** , because of high iron oxide content.
- They develop by intensive and prolonged weathering of the underlying parent rock.
- Tropical weathering (alteration) is a prolonged process of chemical weathering which produces a wide variety in the thickness, grade, chemistry and ore mineralogy of the resulting soils.
- The majority of the land area containing laterites is **between the tropics of Cancer and Capricorn.**
- Laterite soil is commonly found in areas of high altitude and heavy rainfall in Karnataka, TN, MP, Bihar, Orissa, Assam and Meghalaya.
- Plants like Cashew grow well along with Tapioca in this type of soil.

Q.) Which one of the following regions of India has a combination of mangrove forest, evergreen forest and deciduous forest? (2015)

(a) North Coastal Andhra Pradesh

(b) South-West Bengal

(c) Southern Saurashtra

(d) Andaman and Nicobar Islands

Answer: D

- The Andaman and Nicobar Islands have a tropical rainforest canopy, made of a mixed flora with elements from Indian, Myanmar, Malaysian and endemic floral strains. So far, about 2,200 varieties of plants have been recorded, out of which 200 are endemic and 1,300 do not occur in mainland India.
- The South Andaman forests have a profuse growth of epiphytic vegetation, mostly ferns and orchids.
- The Middle Andamans harbours mostly moist deciduous forests.
- North Andamans is characterised by the wet evergreen type, with plenty of woody climbers.

- The North Nicobar Islands (including Car Nicobar and Batti Malv) are marked by the complete absence of evergreen forests, while such forests form the dominant vegetation in the central and southern islands of the Nicobar group.
- Grasslands occur only in the Nicobars, and while deciduous forests are common in the Andamans, they are almost absent in the Nicobars. The present forest coverage is claimed to be 86.2% of the total land area.
- The Andamans and Nicobars have nearly 1136 Sq. Km. and 35 Sq. Km. (450 and 13.8 square miles) respectively of mangrove forests.
- They develop in areas covered by brackish water at high tide, such as the mud flats along tidal streams.

Q.) Consider the following States:

1. Arunachal Pradesh

2. Himachal Pradesh

3. Mizoram In which of the following states do ‘Tropical Wet Evergreen Forests’ occur? (2015)

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: C In India, evergreen forests are found on the eastern and western slopes of the Western Ghats in such states as Tamil Nadu, Karnataka, Kerala and Maharashtra.

- Tropical Wet Evergreen Forests are found in Assam Arunachal Pradesh Meghalaya Nagaland Tripura West Bengal and Andaman and Nicobar Island and on the eastern and western slopes of the Western Ghats in such states as Tamil Nadu Karnataka Kerala and Maharashtra.
- Tropical wet evergreen forests: these forests occur in the states of Maharashtra, Karnataka, Tamilnadu, Kerala, Andaman and Nicobar islands in the south and in the entire north-eastern region including the sub-mountain division of West Bengal.



Figure 5.3 : Natural Vegetation

Q.) In India, in which one of the following types of forests is teak a dominant tree species? (2015)

(a) Tropical moist deciduous forest

(b) Tropical RainForest

(c) Tropical thorn scrub forest

(d) Temperate forest with grasslands

Answer: A Tropical moist deciduous forests are found in areas of moderate rainfall of 100 to 200 cm per year, mean annual temperature of about 27°C and the average annual relative humidity of 60% to 75%.

- The trees of these forests drop their leaves for about 6 to 8 weeks during the spring and early summer when sufficient moisture for the leaves is not available.
- Teak trees are the most dominant species of trees found in these forests.
- Bamboos, sal, shisham, sandalwood, khair, Kusum, Arjun, mulberry are some of the other commercially important species found here.

Q.) With reference to 'Red Sanders', sometimes seen in the news, consider the following statements:

1. It is a tree species found in a part of South India.

2. It is one of the most important trees in the tropical rainforest areas of South India.

Which of the statements given above is/are correct? (2016)

(a) 1 only

(b) 2 Only

(c) Both 1 and 2

(d) Neither 1 nor 2

Answer: A Pterocarpus santalinus, with the common names red sanders, red sandalwood, and sanders wood, is a species of Pterocarpus endemic to the southern Eastern Ghats mountain range of South India.

- This tree is valued for the rich red colour of its wood.
- The wood is not aromatic.
- The tree is not to be confused with the aromatic Santalum sandalwood trees that grow natively in South India.
- It's found in Tropical Dry Deciduous Forests.

Q.) The black cotton soil of India has been formed due to the weathering of (2021)

(a) Brown forest soil

(b) Fissure volcanic rock

(c) Granite and schist

(d) Shale and limestone

Answer: B

- Most pedologists believe that these black soils have been formed due to the solidification of lava spread over large areas during volcanic activity in the deccan plateau, thousands of years ago.
- Black soil is of volcanic origin. These soils have been formed by the solidification of lava that has been spread over a large area during volcanic activities in the Deccan plateau (Deccan and the Rajmahal trap) occurred over thousands of years ago.

Q.) Consider the following trees:

1. Jackfruit (Artocarpus heterophyllus)

2. Mahua (Madhuca indica)

3. Teak (Tectona grandis)

How many of the above are deciduous trees? (2023)

(a) Only one

(b) Only two

(c) All three

(d) None

Answer: B

- Jackfruit is a tree of Tropical Evergreen Forests. It is an evergreen tree. Mahua and Teak are deciduous trees.

Q.) Which of the following best explain(s) the rationale for protecting mangrove ecosystems in the context of climate resilience?

1. Mangroves reduce tidal energy and store freshwater, making them ideal sites for paddy cultivation in saline estuarine belts.

2. Their salt-sensitive roots filter seawater, making mangroves key to converting coastal land into freshwater aquaculture zones.

3. By withstanding tidal surges and offering biomass resources, mangroves function both as natural bio-shields and livelihood bases for rural communities.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Answer: D Do not store salt water. Roots are salt tolerant not salt sensitive.

⚡ **Quick revise** Forest and soil PYQs Fold ▲

Year	Asked	Answer / rule
2023	**Tropical rainforest soil is nutrient-rich** because of fast decomposition	**I false, II true (D)** – soil is **leached and poor**; trees take nutrients as fast as they are produced
2011	How do mangroves act as a safety hedge (2004 tsunami)?	**D – extensive roots hold against storms and tides**
2013	Laterite: red ✔; rich in N & potash ✘; Rajasthan/UP ✘; tapioca & cashew ✔	**C (1 and 4)**
2015	Region with mangrove + evergreen + deciduous	**Andaman & Nicobar (D)**
2015	Tropical wet evergreen in: Arunachal, HP, Mizoram	**Arunachal + Mizoram (C)**
2015	**Teak** is dominant in	**Tropical moist deciduous (A)**
2016	**Red Sanders**: found in a part of S. India ✔; important in tropical rainforest ✘	**A (1 only)** – endemic to southern Eastern Ghats; **not aromatic**, not Santalum sandalwood; dry deciduous
2021	Black cotton soil forms from weathering of	**Fissure volcanic rock (B)**
2023	Jackfruit, Mahua, Teak – how many deciduous?	**Two (B)** – jackfruit evergreen
–	Mangrove rationale for climate resilience	**D** – "bio-shield + livelihood"; roots are **salt-tolerant, not salt-sensitive**; they do not store freshwater

Recall sheet

1. **82°30′E:** *U C My Old Aunt* (UP, CG, MP, Odisha, AP).
2. **Ten – Nine – Six channels:** Andaman|Nicobar, Lakshadweep|Minicoy, Great Nicobar|Sumatra.
3. **Highest:** Kanchenjunga (India's Himalayan peak) · Nanda Devi (entirely in India) · Kamet (Uttarakhand) · Kalsubai (Maharashtra) · Anaimudi (W. Ghats/Kerala) · Doddabetta (TN) · Gurudongmar (lake) · Kunchikal (waterfall).
4. **Panch Prayag:** Vishnu–Nand–Karna–Rudra–Dev. **Left Ganga tributaries:** Ram-Gomti-Ghaghara-Gandak-Kosi-Mahananda.
5. **Mahi** crosses Tropic of Cancer twice · **Luni** = inland-drainage best example · **Kosi** = Sorrow of Bihar · **Kanwar Tal** = Asia's biggest oxbow (Burhi Gandak).
6. **Godavari** longest peninsular river; **Narmada** in a rift – estuary; **Kaveri** gets both monsoons.
7. **Cyclone alerts:** 72 – 48 – 24 – 12. **Season:** Oct–Nov and May–June.
8. **IOD +ve = good monsoon; El Niño = weak SW monsoon, stronger NE. ELLA = heatwave years.**
9. **Soils:** Black=cotton, Alluvial >40%, Laterite=tea-coffee-cashew.
10. **Forest cover: 21.76% + tree 3.41% = 25.17%** (33% target).

Traps

- **Peak "highest of ..." lists are state-specific** – Karoh (Haryana), Amsot (UP), Someshwar (Bihar) are **Shiwalik** peaks; **Kangto** is Arunachal; **Shillong Peak** is Meghalaya.
- **Narmada flows west because of the rift, not the slope** (2013). **Kolleru** lies between Krishna and Godavari (not fed directly by Krishna).
- **Nokrek = Meghalaya, not Sikkim; Namcha Barwa ≠ Garhwal** (2022). **Amarkantak = Vindhya + Satpura** (2023).
- **IOD is Indian Ocean W–E, not the Pacific** (2017). **La Niña is Pacific; it helps the monsoon** (2011).
- **Rainforest soil is poor** despite lush trees (2023). **Laterite is poor in N and potash… but rich in iron** (the handout's two lists differ slightly – laterite "rich in iron oxide and potash" in the notes vs "poor in nitrogen, potash" in the 2013 explanation) → *rule: PYQ asks N-rich? say no.*
- **Jackfruit evergreen; mahua/teak deciduous. Red Sanders is not aromatic sandalwood.**
- **Mahananda / last-tributary wording; UP/Rajasthan/Sikkim PYQ answer conflict** – see alerts above.

Connects to: *Geomorphology* (Tethys, plate collision, drainage patterns, antecedent/superimposed) · *Climatology* (jet streams, ITCZ, cyclones) · *Oceanography* (ENSO, IOD) · *Economic Geography* (soils → crops, minerals of the plateau).

GEOGRAPHY · GS FOUNDATION · SUBJECT 5

Human Geography

Handout + mind maps: no class slides | Handout pages 13

MEDIUM · 4 Prelims

The whole Human Geography handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 6 mind-map pages are in Chapter 11.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: MEDIUM (short chapter, concept-heavy). 4 Prelims PYQs** printed: **2013** (tribe–state pairs) · **2014** (Changpa) · *two undated* (definition of **Total Fertility Rate**; countries known for **low birth rate / ageing**). The handout also reprints five "missed" Indian-Geography PYQs at the end – they are covered in the Indian chapter. **Mains quoted:** none quoted in the material.
- **Pattern:** *definitions* (TFR, sex ratio, dependency) and *where-is-which-tribe*. Plus theory: **Malthus, Demographic Transition**.
- **What to prepare:** Prelims – learn the five DTT stages as a **birth-rate / death-rate table**; learn state-wise tribes (table figure below). Mains hooks – demographic dividend, ageing, urbanisation, gentrification, climate-driven settlement shifts.
- *No teacher ink in this PDF; priority = PYQ count in the handout.*

1 · DEMOGRAPHY

Demography is the study of populations, including their size, structure, distribution, and dynamics. It involves analyzing various demographic factors such as birth rates, death rates, migration patterns, age distribution, and fertility rates to understand population trends and processes.

Factors Affecting Population Growth

- **Birth Rate:** The number of **live births per thousand people in a given population over a specific period** . High birth rates contribute to population growth.
- **Death Rate:** The **number of deaths per thousand people in a given population over a specific period** . Low death rates can contribute to population growth by reducing population loss.

- **Migration:** The **movement of people from one place to another** . **Immigration (inward migration)** can increase population size, while **emigration (outward migration)** can decrease it.
- **Fertility Rate:** The **number of live births per 1000 women of childbearing age in a population** . The **total fertility rate** is the **average number of children born to women of childbearing age in a population.** High fertility rates lead to population growth, while low fertility rates can lead to population decline.
- **Age Structure** : The **distribution of individuals in different age groups within a population** . A youthful age structure (with a high proportion of young people) can lead to rapid population growth, while an aging population structure may lead to slower growth or decline.
- **Economic and Social Factors** : Socioeconomic conditions, access to healthcare, education, employment opportunities, and cultural norms can influence fertility rates, mortality rates, and migration patterns, thereby affecting population growth.

⚡ **Quick revise** Demography Fold ▲

Demography = size, structure, distribution, dynamics of population.
Drivers: birth rate, death rate, migration, fertility, age structure, social-economic factors.

Temporal Distribution of Population



According to the UN World Population Prospects report, the global population will peak at 10.3 billion people around the mid-2080s.

⚡ **Quick revise** World: UN – global population peaks at Fold ▲

- **World:** UN – global population peaks at **10.3 billion in the mid-2080s. 2025 shares: Asia 62.18% · Africa 19.36% · Europe 9.3% · N. America 7.71% · S. America 5.47% · Oceania 0.58%.**
- **India:** most populous country; **1.46 billion (UN)**; UN says it **peaks in the early 2060s at 1.7 billion**; latest **SRS: TFR 1.9**.

> ⚠️ **Handout alert:** the handout quotes **"1.31 billion according to the 2011 census"** next to UN's 1.46 billion. Check that census figure against your notes before quoting it.

Spatial Distribution of Population

- **Factors affecting population distribution:**
  - **Geography and Climate:** Areas with favorable climates, such as temperate regions, tend to have higher population densities. Harsh climates, such as deserts or polar regions, generally have lower population densities. Example: The Indo-Gangetic Plain in India is densely populated due to its fertile land and favorable climate.
  - **Resources:** Availability of natural resources like water, minerals, and fertile land attracts populations. Example: River valleys like the Nile in Egypt and the Ganges in India are densely populated due to their water resources and fertile land.
  - **Economic Opportunities:** Regions with more economic opportunities, such as jobs and industries, attract people. Example: Urban areas like Mumbai, New York, and Tokyo have high population densities due to abundant job opportunities and economic activities.
  - **Political and Social Factors:** Stability, government policies, and social services also influence population distribution. Example: Countries with political stability and good social services, like Canada and Australia, attract migrants and have specific regions with high population densities.
  - **Historical Factors:** Historical events, colonization, trade routes, and historical settlements impact population distribution. Example: Coastal regions and ancient trade routes often have higher population densities due to historical trade and settlement patterns.
- **Patterns of Population Distribution**
  - **Urban vs. Rural:** Urban areas tend to have higher population densities due to concentrated economic activities, better infrastructure, and more services. Rural areas generally have lower population densities. Example: Metropolitan cities like Delhi, Mumbai, and Kolkata in India are densely populated compared to the rural hinterlands.
  - **Continental Distribution:** Population density varies significantly across different continents. Example: Asia is the most populous continent, home to major population centers like China and India, while continents like Australia and Antarctica have very low population densities.

- **Coastal vs. Inland:** Coastal areas are often more densely populated than inland areas due to access to trade, transportation, and resources. Example: Coastal cities like Mumbai, Chennai, and New York are densely populated compared to many inland areas.

- **In 2025, Asia was home to about 62.18% of the world's population, followed by Africa (19.36%), Europe (9.3%; reduction of 0.09% compared to 2024), North America (7.71%), South America (5.47%). Oceania had the smallest population, with just 0.58%** of the world's total population.

Indian Population

(a) India is the **most populous country** in the world.

(b) According to UN Population data, India's population is **1.46 billion** (1.31 billion according to 2011 census.)

(c) UN World Population Prospect report has stated that Indian population **will peak in early 2060s, at 1.7 billion, and then will start reducing.**

(d) However, the latest SRS report, released last month by India's Office of the Registrar General and Census Commissioner, said that India's Total Fertility Rate (TFR) had dropped to 1.9 children per woman.

⚡ **Quick revise** Distribution drivers Fold ▲

Distribution drivers: geography/climate (Indo-Gangetic plain, temperate zones), resources (river valleys – Nile, Ganga), economic opportunity (Mumbai, New York, Tokyo), political/social stability (Canada, Australia), history (coasts, trade routes). **Patterns:** urban > rural; Asia most populous; coast > inland.

Important Terminologies

- **Total Fertility Rate: average number of children a woman is expected to have during her reproductive years** , based on the current Birth Rate. 2.1 is the replacement rate.
  - NFHS 5 data: TFR of India is 2.0
  - Only 5 states TFR > 2.1; Bihar, Meghalaya, Uttar Pradesh, Jharkhand and Manipur
  - India's fertility rate is projected to decline to 1.29 by 2050, significantly below the replacement level of 2.1,
- **IMR (Infant Mortality Rate): deaths of infants under one year of age per 1,000 live births** in a **given year.** In India: 25.5 in 2021
- **Sex Ratio: Males per 1000 females** . In India calculated by females per 1000 males. 2011: 943; NFHS-5: 1020;
- **Child Sex Ratio:** aka **Juvenile Sex Ratio; 0-6 yrs age group** . 2011: 914; NFHS-5: 920
- **Life Expectancy: Average number of years a person is expected to live if current mortality rates continue.** 70.19 yrs in 2022

- **Child Wasting** (low weight for height); **Child Stunting** (low height for age); **Child Underweight** (low weight for age).
- **Dependency Ratio** : Ratio of dependent (0-14 yrs & >65 yrs) to total working age population
- **Working age population** : 15-64 yrs
- **Demographic Dividend:** When **working age population is much higher than the dependent population.**
  - India: 56% in 2022-23
  - Will peak at 68.9% in 2030.
- **Second Demographic Dividend** : **Lower fertility and increased life expectancy** generally lead to greater wealth. Older people with fewer dependents and large savings can improve the money available for investment in an economy, thereby boosting growth.

⚡ **Quick revise** Demography in numbers Fold ▲

👶 **Replacement = 2.1 ("two parents → two kids + a little extra").** India's TFR: **NFHS-5 = 2.0 → SRS = 1.9 → projected 1.29 by 2050.** Only **5 states have TFR > 2.1: "Big Mummy's Up-to-Juggle Mania" = Bihar, Meghalaya, UP, Jharkhand, Manipur.**

Term	Meaning / India number
TFR	average children a woman has in her reproductive years at present birth rates (**PYQ: "average live births a woman would have by end of child-bearing age"**)
IMR	infant deaths (<1 yr) per 1,000 live births · India **25.5 (2021)**
Sex ratio	India counts **females per 1,000 males** · **2011: 943** · **NFHS-5: 1020**
Child sex ratio	age **0–6** · **2011: 914** · **NFHS-5: 920**
Life expectancy	**70.19 yrs (2022)**
Wasting / Stunting / Underweight	low **weight-for-height** / low **height-for-age** / low **weight-for-age**
Dependency ratio	dependants (**0–14 and >65**) ÷ working-age (**15–64**)
Demographic dividend	working-age ≫ dependants · **56% in 2022-23 → peak 68.9% in 2030**
Second dividend	lower fertility + longer life → older people with few dependants and big savings fund investment

🍼🍎🍃 **Stunting = short for age; Wasting = thin for height; Underweight = light for age** – *"Stunt = Short, Waste = Weight-for-height, Under = Weight-for-age".*

Population Pyramids



(a) A population pyramid is a **graphical representation of the age and sex distribution of a population.**

(b) It is a **useful tool for understanding the demographic structure of a population** , which can provide insights into the population's growth patterns, economic development, and potential future challenges.

(c) Population pyramids are typically divided into **different shapes** based on the characteristics of the population they represent.

(d) Expansive Pyramid (i) **Broad base** , indicating a high number of young people. (ii) **Narrow top** , indicating a lower number of older people. (iii) **Rapid population growth** . (iv) Example: **Nigeria** 's population pyramid has a wide base, showing a high proportion of children and young people. This reflects high birth rates and a high dependency ratio. The pyramid narrows sharply at older ages, indicating lower life expectancy. (v) **Indian Pyramid is expansive but with a stabilizing base** . **(e) Constricted Pyramid** (i) **Narrow base** , indicating lower birth rates. (ii) **Bulging middle** , indicating a higher proportion of working-age individuals. (iii) **Narrow top** , reflecting fewer elderly people compared to the working-age population.

(iv) **Population growth slowing down** . (v) Example: **Japan** 's population pyramid is more constricted at the base, showing lower birth rates. The middle section is wider, indicating a large working-age population. The top of the pyramid is wider compared to the base, reflecting an aging population with longer life expectancy. **(f) Stationary Pyramid** (i) **Relatively equal distribution across all age groups** , except for the very old. **(ii) Population growth is stable or very slow.** (iii) Example: The **U.S** . population pyramid shows a more balanced distribution across different age groups. The birth and death rates are low, leading to a relatively stable population size.

⚡ **Quick revise** Population pyramids Fold ▲

> 🔺🏺▮ **Shape = story. Expansive 🔺 (broad base, narrow top)** = Nigeria, rapid growth, high dependency; **India = expansive with a stabilising base. Constricted 🏺 (narrow base, bulging middle)** = Japan, slowing growth, ageing. **Stationary ▮ (even, narrow only at the very old)** = USA, stable.

Population Theory of Malthus

- **Thomas Robert Malthus** , an English cleric and scholar, proposed one of the most influential theories on population growth in his **work "An Essay on the Principle of Population** ," first published in **1798** .



- Malthus's theory **focuses on the relationship between population growth and resources** , particularly food supply.
- **Key Principles of Malthus's Theory**
  - **Population Growth** : Malthus asserted that the **population grows geometrically** (exponentially), meaning it doubles at regular intervals (e.g., 2, 4, 8, 16, etc.).
  - **Food Supply** : He argued that **food supply grows arithmetically** (linearly), meaning it increases by a fixed amount each period (e.g., 1, 2, 3, 4, etc.).
  - **Imbalance** : According to Malthus, **over time** , the **potential for population growth outstrips the potential for food production** , leading to a situation where the population exceeds the food supply.
  - **Checks on Population** : Malthus proposed two types of checks to keep the population in balance with the food supply:
    - **Preventive Checks** : These are **measures that reduce the birth rate** , such as moral restraint (delaying marriage and celibacy), and contraceptive methods. **These methods are under our control.**
    - **Positive Checks** : **Occur after the Malthusian catastrophe** takes place. They are **not under our control.** These **increase the death rate** through famine, disease, war, and other calamities that reduce the population.
- **Criticisms**
  - **Technological Advancements** : Critics argue that **Malthus underestimated human ingenuity and the capacity for technological advancements to increase food production.** The Industrial Revolution and later the Green Revolution significantly increased food supply, countering Malthus's predictions.

- **Economic and Social Factors** : Malthus **did not fully account for the complex social and economic factors** that influence population growth and resource distribution. He equated sociological and biological needs as the same.

- **Modern interpretations** of Malthusian theory consider the **impact of population growth on environmental sustainability and resource depletion beyond just food supply.**

⚡ **Quick revise**

Malthus (1798, Essay on the Principle of Population) Fold ▲

Malthus (1798, *Essay on the Principle of Population*): population grows **geometrically (2-4-8-16)**, food **arithmetically (1-2-3-4)** → population outruns food. **Checks:** *Preventive* (moral restraint, delayed marriage, celibacy, contraception – **under our control**) vs *Positive* (famine, disease, war – **after the catastrophe, not in our control**). **Critics:** technology (Industrial and **Green Revolution**), ignores social/economic complexity; "equated sociological and biological needs". Modern reading: environmental sustainability and resource depletion, not just food.

> 🥖 **Malthus = Multiply vs Add.** *Babies multiply (×2), bread adds (+1).* **Preventive = "Pre" = before birth; Positive checks = "Pos-t-mortem", death does the checking.**

Demographic Transition Theory

- The Demographic Transition Theory (DTT) is a model that **describes the transformation of a country's population structure over time.**
- This theory outlines the **transition from high birth and death rates to low birth and death rates as a country develops from a pre-industrial to an industrialized economic system.**
- Assumption: Fertility and mortality rates reduce over time. Former before latter
- The theory is divided into four to five stages, each characterized by distinct demographic patterns.



- **Stage 1: High Stationary Stage (Pre-Industrial Stage)**
  - Birth Rates: High
  - Death Rates: High
  - **Population Growth: Slow and fluctuating**
  - Characteristics: This stage is marked by high fertility and high mortality due to limited access to medical care, poor nutrition, and lack of sanitation. Populations remain relatively stable but can fluctuate due to famines, diseases, and wars.
  - Example: Most societies before the 18th century. Primitive tribal societies still display this stage.
- **Stage 2: Early Expanding Stage (Transitional Stage)**
  - Birth Rates: High
  - Death Rates: Declining
  - **Population Growth: Rapid**
  - Characteristics: Improved healthcare, sanitation, and food supply lead to a significant decline in mortality rates, particularly among infants and children. Birth rates remain high, resulting in rapid population growth.
  - Example: Many developing countries in Africa.
- **Stage 3: Late Expanding Stage (Industrial Stage)**
  - Birth Rates: Declining
  - Death Rates: Low
  - **Population Growth: Slowing**
  - Characteristics: Birth rates begin to decline due to changes in social norms, increased access to contraception, improved women's education and employment opportunities, and urbanization. Population growth rates begin to slow down.
  - Example: Countries like India, and Brazil.
- **Stage 4: Low Stationary Stage (Post-Industrial Stage)**
  - Birth Rates: Low
  - Death Rates: Low

- **Population Growth: Stable or declining**
- Characteristics: Both birth and death rates are low, leading to a stable or slowly growing population. Societies in this stage often face challenges related to aging populations and the economic impacts of a shrinking workforce.
- Example: Most developed countries, such as the United States, and much of Europe.

- **Stage 5: (Proposed) Decline Stage**
  - Birth Rates: Very low
  - Death Rates: Low
  - **Population Growth: Declining**
  - Characteristics: In this speculative stage, birth rates fall below death rates, leading to a decline in the overall population. This stage is associated with highly developed economies, where the cost of raising children is high and social attitudes towards family size have changed significantly.
  - Example: Some demographers suggest that countries like Japan, South Korea, Germany, and Italy are entering this stage.
- **Significance of the Demographic Transition Theory**
  - **Economic Development** : The DTT suggests that economic development is associated with lower birth and death rates. As countries develop, they move through the stages, which has implications for planning and policy-making.
  - **Population Aging** : In the later stages, countries face challenges related to aging populations, such as increased healthcare costs and the need for pension systems.
  - **Urbanization** : The transition often accompanies increased urbanization, changing patterns of settlement and resource use.
  - **Education and Women's Rights** : The decline in birth rates is closely linked to improvements in women's rights, education, and employment opportunities.
  - **Environmental Impact** : Population dynamics affect environmental sustainability. Rapid population growth can lead to increased pressure on natural resources while declining populations might lead to labor shortages and economic challenges.

Demographic Transition in India

1. Till 1921: Stagnant population

2. 1921-1951: Increasing population

3. 1951-1981: Population explosion

4. (1921-81: Early Expanding stage)

5. 1981- till date: Late Expanding Stage (Growth with stability)

MAJOR TRIBES OF INDIA

Table No. 1A: State-wise major tribes of India		
Sl_No.	**States/UTs**	**Major Tribes**
1	Andaman and Nicobar Islands	Jarawa , Oraons, Onges, Sentinelese, Shompens
2	Andhra Pradesh	Andh, Sadhu Andh, Bhagata, Bhil, Chenchus (Chenchawar), Gadabas, Gond, Goundu, Jatapus, Kammara, Kattunayakan, Kolawar, Kolam, Konda, Manna Dhora, Pardhan, Rona, Savaras, Dabba Yerukula, Nakkala, Dhulia, Thoti, Sugalis.
3	Arunachal Pradesh	Apatanis, Abor, Dafla, Galong, Momba, Sherdukpen, Singpho.
4	Assam	Chakma, Chutiya, Dimasa, Hajong, Garos, Khasis, Gangte.
5	Bihar	Asur, Baiga, Birhor, Birjia, Chero, Gond, Parhaiya, Santhals, Savar.
6	Chhattisgarh	Agariya, Bhaina, Bhattra, Biar, Khond, Mawasi, Nagasia.
7	Goa	Dhodia, Dubia, Naikda, Siddi,Varli.
8	Gujarat	Barda, Bamcha, Bhil, Charan, Dhodia, Gamta, Paradhi, Patelia.
9	Himachal Pradesh	Gaddis, Gujjars, Khas, Lamba, Lahaulas, Pangwala, Swangla.
10	Jammu and Kashmir	Bakarwal, Balti, Beda, Gaddi, Garra, Mon, Purigpa, Sippi.
11	Jharkhand	Birhors, Bhumij, Gonds, Kharia, Mundas, Santhals, Savar.
12	Karnataka	Adiyan, Barda, Gond, Bhil, Iruliga, Koraga, Patelia, Yerava.
13	Kerala	Adiyan, Arandan, Eravallan, Kurumbas, Malai arayan, Moplahs, Uralis.
14	Madhya Pradesh	Baigas, Bhils, Bharia, Birhors, Gonds,Katkari, kharia, Khond, Kol, Murias.
15	Maharashtra	Bhaina, Bhunjia, Dhodia, Katkari, Khond, Rathawa, Warlis.
16	Manipur	Aimol, Angami, Chiru, Kuki, Maram, Monsang, Paite, Purum, Thadou.
17	Meghalaya	Chakma, Garos, Hajong, Jaintias Khasis, Lakher, Pawai, Raba.
18	Mizoram	Chakma, Dimasa, Khasi, Kuki, Lakher, Pawai, Raba, Synteng.
19	Nagaland	Angami, Garo, Kachari, Kuki, Mikir, Nagas, Sema.
20	Odisha	Gadaba, Ghara, Kharia, Khond, Matya, Oraons, Rajuar, Santhals.
21	Rajasthan	Bhils, Damaria, Dhanka, Meenas(Minas), Patelia, Sahariya.
22	Sikkim	Bhutia, Khas, Lepchas.
23	Tamil Nadu	Adiyan, Aranadan, Eravallan, Irular, Kadar, Kanikar, Kotas, Todas
24	Telangana	Chenchus.
25	Tripura	Bhil, Bhutia, Chaimal, Chakma, Halam, Khasia, Lushai, Mizel, Namte.
26	Uttarakhand	Bhotias, Buksa, Jannsari, Khas, Raji, Tharu.
27	Uttar Pradesh	Bhotia, Buksa, Jaunsari, Kol, Raji, Tharu.
28	West Bengal	Asur, Khond, Hajong, Ho, Parhaiya, Rabha, Santhals, Savar.
Source: Indian Census		

⚡ Quick revise Demographic Transition Theory (DTT) Fold ▲

Demographic Transition Theory (DTT): birth and death rates fall from high to low as a country industrialises.

Stage	Name	Birth	Death	Growth	Example (handout)
1	High stationary (pre-industrial)	High	High	slow, fluctuating	tribal societies, pre-18th c.
2	Early expanding	High	**Falling**	**rapid**	many African countries
3	Late expanding (industrial)	**Falling**	Low	slowing	**India, Brazil**
4	Low stationary (post-industrial)	Low	Low	stable/declining	USA, much of Europe
5	Decline (proposed)	Very low	Low	**declining**	Japan, South Korea, Germany, Italy

🌈 **"Death Drops first, Birth Boards later."** Stage 1: both high. **Stage 2: death falls → boom. Stage 3: births fall → boom slows.** Stage 4: both low. Stage 5: births **below** deaths.

⚠️ **Handout alert:** the assumption line reads *"Fertility and mortality rates reduce over time. Former before latter"* – but the stage table (and every textbook-style description in the same handout) shows **mortality falling first (Stage 2), fertility next (Stage 3).** *Exam rule: follow the stage table.*



Original sketch of the handout's stage table (not to scale): **dashed = death rate**, **solid = birth rate**. The gap between them is the population boom.

India's transition: till **1921 stagnant** → **1921–51 increasing** → **1951–81 explosion** (1921–81 = **early expanding**) → **1981 onwards late expanding** ("growth with stability").

Why DTT matters: development ↔ lower birth/death; ageing and pensions; urbanisation; **women's education and rights lower fertility**; environmental pressure vs labour shortages.

PYQ

Q.) Consider the following pairs :

1. Limboo (Limbu): Sikkim

2. Karbi: Himachal Pradesh

3. Dongaria: Odisha

4. Bonda: Tamil Nadu

Which of the above pairs are correctly matched? (2013) 1 and 3 only 1 and 2 only 2 and 3 only 1, 2, 3 and 4

Answer: A The **Limbu are Kirati people** indigenous and native to the **Himalayan Limbuwan region** of the Indian subcontinent, what is now modern-day Eastern Nepal,

Northern Sikkim, India and Western Bhutan.

- The **Niyamgiri hill range in Odisha** state, eastern India, is **home to the Dongria Kondh tribe** .
- Niyamgiri is an area of densely forested hills, deep gorges and cascading streams. To be a Dongria Kondh is to farm the hills' fertile slopes, harvest their produce, and worship the mountain god Niyam Raja and the hills he presides over, including the 4,000 metre Mountain of the Law, Niyam Dongar.

Q.) With reference to 'Changpa' community of India, consider the following statement:

- They live mainly in the State of Uttarakhand.
- They rear the Pashmina goats that yield fine wool.
- They are kept in the category of Scheduled Tribes.

Which of the statements given above is/are correct? (2014)

(a) 1 only

(b) 2 and 3 only

(c) 3 only

(d) 1, 2 and 3

Answer: B The Changpa or Champa are a semi-nomadic Tibetan people found mainly in the Changtang in Ladakh and in Jammu and Kashmir.

- The Changpa or Champa are a semi-nomadic Tibetan people found mainly in the Changtang in Ladakh, India. A smaller number resides in the western regions of the Tibet Autonomous Region and were partially relocated for the establishment of the Changtang Nature Reserve.
- Most of the regions that the Changpas reside in is quite inhospitable and unsuitable for farming. Thus, the primary occupation for the tribe is rearing of yaks, horses and sheep. Cashmere, also known as pashmina, is majorly sourced from the pashmina sheep that the tribe rears.
- A smaller number resides in the western regions of the Tibet Autonomous Region and were partially relocated for the establishment of the Changtang Nature Reserve.
- As of 1989, there were half a million nomads living in the Changtang area.

Q.) The total fertility rate in an economy is defined as:

- The number of children born per 1000 people in the population in a year.
- The number of children born to a couple in their lifetime in a given population.
- The birth rate minus death rate.
- The average number of live births a woman would have by the end of her child-bearing age.

Answer:

(d) The average number of live births a woman would have by the end of her child-bearing age.

Q.) Consider the following countries:

- Italy
- Japan
- Nigeria
- South Korea
- South Africa **Which of the above countries are frequently mentioned in the media for their low birth rates, or ageing population or declining population?**
- (a) 1, 2 and 4
- (b) 1, 3 and 5
- (c) 2 and 4 only
- (d) 3 and 5 only

Answer: A MISSED QNS OF INDIAN GEOGRAPHY

Q.) With reference to the climate of Andaman and Nicobar Islands, which of the following statements is/are correct?

1. The climate can be defined as a humid, tropical coastal climate.

2. It receives rainfall from both South-west monsoon and North-east monsoon.

3. Maximum precipitation is between December and May.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Ans: (b)

Explanation: The Andaman and Nicobar Islands experience a humid tropical maritime/coastal climate. It is in the ocean, so both SW and NE winds are onshore. Maximum rainfall generally occurs during the South-West Monsoon months, especially May to September. (Heat=LP=rain, and NH has heat in March-Sept)

Q. Which of the following geographical features or phenomena is/are associated with the Peninsular Block of India?

1. Submergence of parts of the western coast due to tectonic activity

2. Presence of residual mountain ranges such as the Veliconda Hills and Mahendragiri Hills

3. Deep, V-shaped river valleys formed by fast-flowing rivers

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Ans: (b)

Explanation: The Peninsular Block is an ancient stable cratonic landmass, but parts of the western coast experienced submergence due to tectonic movements. Eastern Ghats are denuded mountains. Most rivers of Peninsular India are mature rivers flowing over old hard rocks.

Q. Which of the following best explain(s) the rationale for protecting mangrove ecosystems in the context of climate resilience?

1. Mangroves reduce tidal energy and store freshwater, making them ideal sites for paddy cultivation in saline estuarine belts.

2. Their salt-sensitive roots filter seawater, making mangroves key to converting coastal land into freshwater aquaculture zones.

3. By withstanding tidal surges and offering biomass resources, mangroves function both as natural bio-shields and livelihood bases for rural communities.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Ans: (d)

Explanation: Do not store salt water. Roots are salt tolerant not salt sensitive. (Played with words)

Q. Identify the river of the Indian sub-continent on the basis of the following information:

1. It has an antecedent drainage system.

2. It flows through three countries.

3. It originates in the Tibetan Plateau and is an important river for irrigation.

4. It does not form distributaries.

Select the answer from the following:

(a) Brahmaputra

(b) Indus

(c) Sutlej

(d) Teesta

Ans: (c)

Explanation: Brahmaputra has distributaries. Teesta originates in Sikkim. Indus has distributaries.

Q. Which of the following with reference to Indian States is/are not correct?

1. Uttar Pradesh shares its boundary with the highest number of other Indian States.

2. Rajasthan shares the longest international border among all Indian States.

3. Sikkim is the only State that shares its boundary with just one other Indian State.

Select the answer using the code given below:

(a) 1 only

(b) 1 and 2

(c) 2 and 3

(d) 3 only

Ans: (a)

Explanation: West Bengal has longest international border. Meghalaya also shares border with only 1 state

⚡ **Quick revise** Tribes (state-wise table from the handout)

Fold ▲

Source figure: Census table printed in the handout (p. 7). Study the states in your PYQs first.

Table No. 1A: State-wise major tribes of India		
Sl_No.	States/UTs	Major Tribes
1	Andaman and Nicobar Islands	Jarawa , Oraons, Onges, Sentinelese, Shompens
2	Andhra Pradesh	Andh, Sadhu Andh, Bhagata, Bhil, Chenchus (Chenchawar), Gadabas, Gond, Goundu, Jatapus, Kammara, Kattunayakan, Kolawar, Kolam, Konda, Manna Dhora, Pardhan, Rona, Savaras, Dabba Yerukula, Nakkala, Dhulia, Thoti, Sugalis.
3	Arunachal Pradesh	Apatanis, Abor, Dafla, Galong, Momba, Sherdukpen, Singpho.
4	Assam	Chakma, Chutiya, Dimasa, Hajong, Garos, Khasis, Gangte.
5	Bihar	Asur, Baiga, Birhor, Birjia, Chero, Gond, Parhaiya, Santhals, Savar.
6	Chhattisgarh	Agariya, Bhaina, Bhattra, Biar, Khond, Mawasi, Nagasia.
7	Goa	Dhodia, Dubia, Naikda, Siddi,Varli.
8	Gujarat	Barda, Bamcha, Bhil, Charan, Dhodia, Gamta, Paradhi, Patelia.
9	Himachal Pradesh	Gaddis, Gujjars, Khas, Lamba, Lahaulas, Pangwala, Swangla.
10	Jammu and Kashmir	Bakarwal, Balti, Beda, Gaddi, Garra, Mon, Purigpa, Sippi.
11	Jharkhand	Birhors, Bhumij, Gonds, Kharia, Mundas, Santhals, Savar.
12	Karnataka	Adiyan, Barda, Gond, Bhil, Iruliga, Koraga, Patelia, Yerava.
13	Kerala	Adiyan, Arandan, Eravallan, Kurumbas, Malai arayan, Moplahs, Uralis.
14	Madhya Pradesh	Baigas, Bhils, Bharia, Birhors, Gonds,Katkari, kharia, Khond, Kol, Murias.
15	Maharashtra	Bhaina, Bhunjia, Dhodia, Katkari, Khond, Rathawa, Warlis.
16	Manipur	Aimol, Angami, Chiru, Kuki, Maram, Monsang, Paite, Purum, Thadou.
17	Meghalaya	Chakma, Garos, Hajong, Jaintias Khasis, Lakher, Pawai, Raba.
18	Mizoram	Chakma, Dimasa, Khasi, Kuki, Lakher, Pawai, Raba, Synteng.
19	Nagaland	Angami, Garo, Kachari, Kuki, Mikir, Nagas, Sema.
20	Odisha	Gadaba, Ghara, Kharia, Khond, Matya, Oraons, Rajuar, Santhals.
21	Rajasthan	Bhils, Damaria, Dhanka, Meenas(Minas), Patelia, Sahariya.
22	Sikkim	Bhutia, Khas, Lepchas.
23	Tamil Nadu	Adiyan, Aranadan, Eravallan, Irular, Kadar, Kanikar, Kotas, Todas
24	Telangana	Chenchus.
25	Tripura	Bhil, Bhutia, Chaimal, Chakma, Halam, Khasia, Lushai, Mizel, Namte.
26	Uttarakhand	Bhotias, Buksa, Jannsari, Khas, Raji, Tharu.
27	Uttar Pradesh	Bhotia, Buksa, Jaunsari, Kol, Raji, Tharu.
28	West Bengal	Asur, Khond, Hajong, Ho, Parhaiya, Rabha, Santhals, Savar.
Source: Indian Census		

PYQ	Asked	Answer
2013	Limboo–Sikkim ✔ · Karbi–Himachal ✘ · Dongaria–Odisha ✔ · Bonda–Tamil Nadu ✘	**A (1 and 3)** – **Dongria Kondh** of **Niyamgiri**, Odisha (god Niyam Raja)
2014	**Changpa:** live mainly in Uttarakhand ✘; rear pashmina goats ✔; Scheduled Tribe ✔	**B (2 and 3)** – semi-nomadic in **Changtang, Ladakh**; yaks, horses, sheep; pashmina/cashmere

Changpa = "Chang-Pa-shmina" – Ladakh's high plateau herders (Changtang). **Dongria = "Dongar" (hill)** in Niyamgiri hills.

2 · SETTLEMENTS

3 · RURAL SETTLEMENTS

Factors Affecting Settlements

Rural settlements are influenced by a variety of factors that determine their location, size, structure, and pattern.

- **Physical Factors**
  - **Topography:**
    - Flat and fertile plains are more conducive to agriculture and human habitation.
    - Hilly and mountainous regions often lead to dispersed settlements due to difficult terrain and limited arable land.
    - Example: The Ganges Plains in India have dense rural settlements due to flat terrain and fertile soil.
  - **Climate:**
    - Areas with moderate climates are more likely to have higher population densities.
    - Extreme climates (too hot, too cold, or too wet) can hinder large-scale settlements.
    - Example: The Thar Desert in Rajasthan, India, has sparse rural settlements due to harsh climatic conditions.
  - **Water Supply:**
    - Proximity to rivers, lakes, or groundwater sources is crucial for drinking water, irrigation, and other daily needs.
    - Settlements are often concentrated near reliable water sources.
    - Example: The rural settlements along the Nile River in Egypt depend heavily on the river for water.
  - **Soil Fertility:**
    - Fertile soils support agriculture, which is the primary livelihood in rural areas.
    - Areas with rich, alluvial soils are densely populated.
    - Example: The Indo-Gangetic Plain in India has high rural population density due to its fertile soil.
  - **Natural Resources:**
    - Availability of resources like minerals, forests, and fisheries can attract settlements.
    - Regions rich in natural resources often see clustered settlements.
    - Example: The mining regions in Jharkhand, India, have developed settlements around mining activities.
- **Economic Factors**
  - **Agricultural Productivity:**
    - High agricultural productivity supports larger populations and dense settlements.
    - Regions with better irrigation, technology, and crop yields tend to have more stable settlements.
    - Example: The Punjab region in India has high rural population density due to its advanced agricultural practices.
  - **Employment Opportunities:**
    - The availability of non-agricultural jobs (e.g., cottage industries, mining) can attract and retain population.
    - Economic diversification can influence the settlement pattern.

        - Example: The rural areas near industrial towns often have higher population densities.
    - **Accessibility and Transportation:**
        - Proximity to roads, railways, and other transport networks facilitates trade, and access to markets, and services.
        - Better connectivity generally leads to more populated and prosperous settlements.
        - Example: Villages near major highways or railway lines tend to be larger and more developed.
- **Social and Cultural Factors**
    - **Community and Kinship:**
        - Social bonds and kinship ties can influence the location and growth of rural settlements.
        - Communities with strong social networks often cluster together.
        - Example: Traditional villages in India often have extended families living in close proximity.
    - **Cultural Practices:**
        - Cultural and religious practices can dictate settlement patterns, such as the layout and structure of homes.
        - Specific cultural requirements (e.g., space for religious rituals) influence the design of rural settlements.
        - Example: Some rural areas in Rajasthan, India, have unique architectural styles due to cultural influences.
    - **Land Ownership Patterns:**
        - The distribution of land ownership can affect the settlement pattern.
        - Areas with fragmented landholdings may have dispersed settlements.
        - Example: The zamindari system in India historically influenced the pattern of rural settlements.
- **Political Factors**
    - **Government Policies:**
        - Policies related to land use, agriculture, infrastructure development, and rural development impact settlement patterns.
        - Programs aimed at rural development can lead to the establishment of new settlements or the growth of existing ones.
        - Example: The Green Revolution in India led to significant changes in rural settlement patterns in regions like Punjab and Haryana.
    - **Security and Stability:**
        - Political stability and security influence where people choose to settle.
        - Areas prone to conflict or instability may see depopulation or fragmented settlements.
        - Example: Rural areas in conflict zones may have sparse and dispersed settlements.
- **Environmental Factors**
    - **Natural Disasters:**

- Areas prone to natural disasters (floods, earthquakes, hurricanes) may have fewer stable settlements.
- Repeated exposure to natural disasters can lead to migration and changes in settlement patterns.
- Example: Coastal villages in Odisha, India, are often affected by cyclones, influencing their settlement structure and resilience measures. **(e) Climate Change** (i) In future, climate change might force the coastal settlements to be removed to more interior locations. (ii) Some villages might get displaced due to extreme drought of flood conditions. (iii) E.g. some regions of Brazil and Burkina Faso have started to feel these problems.

⚡ **Quick revise** What decides where villages stand? Fold ▲

What decides where villages stand? *Physical* (relief, climate, **water**, soil, resources) · *Economic* (farm productivity – Punjab; jobs; roads/rail) · *Social-cultural* (kinship, ritual, land ownership – zamindari) · *Political* (policies, security; Green Revolution reshaped Punjab/Haryana) · *Environmental* (floods, cyclones – coastal Odisha) · **climate change** (coastal villages shift inland; drought/flood displacement – parts of Brazil and Burkina Faso).

Types of Rural Settlements

- **Compact/Nucleated Settlements:**
  - **Favourable relief + favourable Climate** .
  - Houses and buildings are **closely packed** .
  - **Common in fertile plains, valleys, and areas with abundant natural resources.**
  - Sometimes in **locations where resources are concentrated or for security reasons.**
  - North Indian Plains, Wet points in Rajasthan, regions of Bundelkhand and Nagaland.
- **Semi-Compact Settlements:**
  - **Favourable relief + Less favourable climate**
  - **Transitional** form between compact and dispersed settlements.
  - Houses are somewhat clustered but with more space between them compared to compact settlements.
  - Found in areas with **moderate agricultural productivity** .
  - Indus Basin; Gujarat Plains, parts of Rajasthan Plains.
- **Hamleted Settlements:**
  - **Fragmentation of settlements in socially/ethically motivated separate units.**
  - Often known by common names like **panna, para, palli, nagla, dhani,** and others.
  - Northern India: Ganga plains, lower Himalayan valleys; Chhattisgarh Plains (tribal clustering); parts of South India
- **Dispersed Settlements:**
  - Houses and buildings are **spread out over a large area** .
  - **Bad climate & Relief**

- Himalayan Region: Uttarakhand, Himachal Pradesh, Ladakh; Meghalaya, etc.

⚡ **Quick revise** Rural settlements — Fold ▲

> 🏘️ **Good relief + good climate → huddled together; bad → scattered. Compact** 🧱 (north Indian plains, wet points of Rajasthan, Bundelkhand, Nagaland) · **Semi-compact** 🏠🏠 (Indus basin, Gujarat plains, parts of Rajasthan) · **Hamleted** 🧩 (socially separate units – **panna, para, palli, nagla, dhani**; Ganga plains, lower Himalayan valleys, **Chhattisgarh plains (tribal)**, parts of south) · **Dispersed** 🏔️ (Uttarakhand, HP, Ladakh, Meghalaya).

Patterns of Rural Settlements

- **Linear** : Common-most pattern; Along highways, forest pathways, and rivers; at cross-sections, can make T-shaped or star-shaped patterns or Checkerboard Pattern.
- **Oval/Circular** : around a nodal point of attraction. E.g. a religious place, oasis, pond, etc.
- **Triangular** : found in the deltas and doabs
- **Rectangular (Grid pattern)** : plains
- **Step:** hilly regions

⚡ **Quick revise** Patterns (shape) — Fold ▲

Patterns (shape): Linear (most common – roads, rivers; T/star/checkerboard at crossings) · **Oval/circular** (around temple, oasis, pond) · **Triangular** (**deltas, doabs**) · **Rectangular/grid** (plains) · **Step** (hills).

4 · URBAN SETTLEMENTS

Terminologies associated with Urban Settlements:

- **City:** A large and **significant urban area with a high population density and infrastructure** of built environment.
  - High population density.
  - Diverse economic activities.
  - Advanced infrastructure and services.
  - Example: Mumbai, Lucknow, Guwahati.
- **Town:** An **urban area that is larger than a village but smaller than a city** , often with local administrative and commercial functions.

- **Megacity:** An **urban area with a population of more than 10 million people** . Due to its population, it often has a significant economic, cultural, and political influence.
  - Example: Tokyo, Japan; Greater Mumbai, Greater Chennai, India.
- **Metropolis** : **A large and densely populated urban area that serves as a significant economic, political, and cultural center of a country or region.**
  - Major urban center.
  - Advanced infrastructure and services.
  - High levels of cultural and economic activities.
  - Example: New York City, USA; Delhi, India.
- **Conurbation:** An extended urban area formed by the merging of several towns or cities and their suburbs.
  - Large, continuous urban area.
  - Multiple urban centers within a single metropolitan area.
  - Integrated economic and social networks.
  - Example: Delhi NCR Region.
- **Megalopolis** : **A region in which several large cities and surrounding areas grow together, forming a massive, interconnected urban complex.**
  - Extensive urbanized region.
  - Multiple large cities (often metropolises) in close proximity.
  - Integrated transportation and economic systems.
  - **Labour market separation, due to different specialization and large distances** (due to the large size).
  - Example: The BoshWash corridor (Boston to Washington D.C.) in the USA.
- **Suburb: A residential area on the outskirts of a city** , typically less densely populated than the urban core.
  - Primarily residential.
  - Lower population density compared to the city center.
  - Dependence on the city for employment and services.
  - Example: earlier Gurugram was a suburb of Delhi (now a part of the conurbation). Bandra, Mumbai.
- **Urban Sprawl** : The **uncontrolled expansion of urban areas into the surrounding rural land.**
  - Low-density, automobile-dependent development.
  - Expansion into rural areas.
  - Often leads to environmental and infrastructural challenges.
- **Slum** : A **densely populated urban area characterized by substandard housing and squalor.**
  - Poor housing conditions.
  - Inadequate access to basic services (water, sanitation, electricity).
  - High population density and poverty levels.
  - Example: Dharavi in Mumbai, India.
- **Central Business District (CBD):** The **commercial and business center of a city** , often referred to as the "downtown" area.

- High concentration of retail and office buildings.
- High land values and rent prices.
- Major transportation hubs.
- Example: Chandni Chowk and Connaught Place in Delhi.

- **Edge City:** A **concentration of business, shopping, and entertainment outside a traditional downtown (CBD) in what had previously been a suburban residential or rural area** .
  - Significant commercial and business activity.
  - Located on the outskirts of larger cities.
  - Often characterized by new office buildings, shopping malls, and entertainment facilities.
  - Example: Gurgaon for Delhi
- **Gentrification:** The **process by which urban areas undergo redevelopment and revitalization, often resulting in increased property values and the displacement of lower-income residents.**
  - Upgrading of housing and infrastructure.
  - Influx of higher-income residents.
  - Potential displacement of long-term, lower-income residents
  - E.g. Dharavi Re-development **Satellite City:** A planned city established to reduce the population pressure on a major city.

⚡ **Quick revise** Urban settlements – the size ladder Fold ▲

🪜 **Climb the ladder: Town → City → Metropolis → Megacity (>10 million) → Conurbation → Megalopolis.**

Term	Meaning	Handout example
Town	bigger than village, smaller than city	–
City	dense, diverse economy, advanced infrastructure	Mumbai, Lucknow, Guwahati
Metropolis	major economic/political/cultural centre	New York, **Delhi**
Megacity	**> 10 million**	Tokyo, **Greater Mumbai, Greater Chennai**
Conurbation	several towns/cities + suburbs merged	**Delhi NCR**
Megalopolis	several big cities grown together; labour-market separation	**BosWash** (Boston–Washington)
Suburb	residential, depends on city	Bandra; Gurugram earlier
Urban sprawl	uncontrolled low-density, car-based spread into rural land	–

Term	Meaning	Handout example
Slum	substandard housing, no basic services	**Dharavi**
CBD	commercial centre; high land values	**Chandni Chowk, Connaught Place**
Edge city	business/shopping outside the CBD in former suburb/rural land	**Gurgaon** (for Delhi)
Gentrification	redevelopment → higher values → **lower-income residents displaced**	**Dharavi redevelopment**
Satellite city	planned city to relieve a major city	–

> 🛍️ **CBD = the city's wallet (downtown). Edge city = the mall that moved to the edge. Gentrification = the café that raised the rent.**

Recall sheet

1. **Replacement TFR 2.1** · India NFHS-5 **2.0**, SRS **1.9** · only **5 states** above 2.1 (Bihar, Meghalaya, UP, Jharkhand, Manipur).
2. **Sex ratio 2011: 943 · Child sex ratio 914** (NFHS-5: 1020 / 920).
3. **Working age 15–64**; dependants **0–14 & >65**; dividend **56% → 68.9% (2030)**.
4. **Pyramids:** Expansive (Nigeria/India), Constricted (Japan), Stationary (USA).
5. **DTT:** Stage 2 death falls first; **India in stage 3**.
6. **Malthus:** geometric vs arithmetic; preventive vs positive.
7. **Settlements:** compact → semi-compact → hamleted → dispersed.
8. **Urban ladder:** town, city, metropolis, megacity (>10 m), conurbation, megalopolis.

Traps

- **TFR ≠ birth rate** (PYQ): TFR is *average live births per woman over her child-bearing years.*
- **Changpa ≠ Uttarakhand** – Ladakh (Changtang). **Karbi ≠ Himachal; Bonda ≠ Tamil Nadu.**
- **Low-birth-rate PYQ (undated):** answer **A = Italy, Japan, South Korea**; Nigeria and South Africa are not in it.
- **Sex ratio** in India = *females per 1,000 males* (the world often counts the other way).
- **DTT "former before latter" wording** is misleading – death rate falls first.

Connects to: *Indian Geography* (Changpa and Ladakh, Dongria and Niyamgiri; Thar/Himalaya settlements) · *Economic Geography* (labour, agriculture, urban industry).

GEOGRAPHY · GS FOUNDATION · SUBJECT 6

Economic Geography

Handout + mind maps: no class slides | Handout pages 54

VERY HIGH · 31 Prelims

The whole Economic Geography handout in its original order with every picture, table and highlight, an exam lens on top, with my memory cards, PYQs and alerts placed right inside the topic they belong to (⚡ Quick revise boxes; tap the title to fold them), then a recall sheet and traps. The 21 mind-map pages are in Chapter 12.

Exam lens and priority

🎯 **Exam lens (from the handout only)**

- **Priority: HIGH. 31 Prelims PYQs** printed (counted from the handout text): **agriculture 19** (2011×4, 2012×2, 2013, 2014×2, 2015, 2017, 2018, 2019, 2020, 2021×2, 2022, 2024, 2025) · **minerals & energy 9** (2012, 2013×2, 2016, 2018, 2022, 2023×2, one undated on rare earths) · **transport 3** (2014, 2015, 2023) · **industries 0 – none quoted in the material**. **Mains quoted:** none quoted in the material.
- **Pattern:** *which crop where, which mineral where, which state does what.* Plus "statement" questions on fertiliser, salinisation, permaculture, conservation agriculture, and current affairs (cocoa, turmeric, REE, NCMM).
- **Heavy handout, thin on teacher stress.** *No teacher ink in this PDF; priority = PYQ count in the handout.* Two image pages (crop table, waterways) were checked visually.

1 · AGRICULTURE

- India boasts the **largest area of arable land globally** , at around 1.597 million square kilometers (394.6 million acres).

Cropping Seasons of India:

- **Kharif Season**
  - **Time Period** : **June to October** .
  - **Sowing:** Beginning with the **onset of the southwest monsoon** in June.
  - **Harvesting: September to October** .

- **Major Crops: Paddy** (rice), **maize, sorghum** (jowar), millet (bajra), pulses (such as moong, urad, and arhar), cotton, groundnut, soybean, and sugarcane.
- **Climatic Conditions:** Requires **hot and humid weather with substantial rainfall** . Millets, pulses, and oilseeds can grow even in less humid areas.

- **Rabi Season**
  - **Time Period: October/November to March/April** .
  - **Sowing:** After the monsoon rains, from **October to December.**
  - **Harvesting: March to April.**
  - W **inter rainfall from Western disturbances** supports growth.
  - **Major Crops: Wheat, barley, oats, chickpea** (gram), mustard, peas, and linseed.
  - **Climatic Conditions:** Requires **cool and dry weather for growth and warmer weather during the harvesting period** .
- **Zaid Season**
  - **Time Period: March to June** .
  - **Sowing:** Between the **end of the Rabi season and the beginning of the Kharif season** .
  - **Harvesting: June to July** .
  - **Major Crops: Watermelon, muskmelon, cucumber** , vegetables, fodder crops, and some pulses (such as moong and urad).
  - **Climatic Conditions:** Requires **warm and dry weather** with the **availability of irrigation** .

⚡ **Quick revise** 1.1 Seasons — Fold ▲

> 🌦️ **Kharif = Kar-Rain (monsoon crops). Rabi = Rest-of-winter (WDs help). Zaid = Zara-garmi (summer gap).**

	Kharif	Rabi	Zaid
Time	**Jun–Oct** (sow with SW monsoon; harvest Sep–Oct)	**Oct/Nov–Mar/Apr** (sow Oct–Dec; harvest Mar–Apr)	**Mar–Jun** (harvest Jun–Jul)
Weather	hot, humid, heavy rain	cool and dry growth, warm at harvest; **Western Disturbance rain** helps	warm, dry; **irrigation needed**
Crops	**paddy, maize, jowar, bajra, pulses (moong, urad, arhar), cotton, groundnut, soybean, sugarcane**	**wheat, barley, oats, gram, mustard, peas, linseed**	**watermelon, muskmelon, cucumber, vegetables, fodder, some moong/urad**

⚡ **Quick revise** 1.2 Crop table (read from the image) — Fold ▲

🗺️ **Pairing pegs: Cotton + black soil (Maharashtra, Gujarat, Telangana) · Jute + West Bengal (wet, 120–200 cm) · Tea = acidic slopes (Assam, WB, TN, Kerala) · Coffee = Karnataka (shade, Arabica & Robusta) · Rubber = Kerala · Coconut = Kalpavriksha (Kerala) · Bajra = toughest (tolerates 42 °C), Rajasthan.**

Crop	Temp / rain	Soil	Top Indian states	Top world	Remarks
Rice	20–32 °C · 150–300 cm	deep clayey, alluvial	UP, Telangana, WB	India, China, Bangladesh, Indonesia, Vietnam	staple; **Aus, Aman, Boro** in WB; high humidity
Wheat	10–25 °C · 50–100 cm	loamy, well-drained	UP, MP, Punjab	China, EU, India, Russia, USA	winter crop; bright sun at ripening
Jowar	20–27 · 50–75	light sandy–loamy	Maharashtra, Karnataka, MP	India, Niger, China, Mali, Nigeria	Kharif & Rabi; fodder + biofuel
Bajra	20–27 · 50–75	sandy loam	Rajasthan, UP, Gujarat	India, Niger, China, Nigeria	**drought-resistant, up to 42 °C**
Millets (general)	20–27 · 50–75	wide range	Rajasthan, Karnataka, MP	India, Niger, China, Mali, Nigeria	antioxidants, diabetes/heart, protein; **low cost; marginal farmers**; jowar, bajra, ragi
Maize	15–27 · 50–80	well-drained loam	MP, Karnataka, Bihar	USA, China, Brazil, EU, Argentina	food, **alcohol, feed, biofuel**
Cotton	21–30 · 50–100	**black (regur)**	Maharashtra, Gujarat, Telangana	China, India, Brazil, USA, Australia	cash crop; **≥ 210 frost-free days**
Jute	24–35 · 120–200	alluvial	WB, Bihar, Assam	India, Bangladesh, China	**retting**; plastic alternative
Sugarcane	20–27 · 100–150	loamy	UP, Maharashtra, Karnataka	Brazil, India, China,	**very water-intensive**; ethanol + sugar

Crop	Temp / rain	Soil	Top Indian states	Top world	Remarks
				Thailand, Pakistan	
Pulses	15–27 · low rain	well-drained	MP, Maharashtra, UP	India, Canada, Myanmar, Brazil, Ethiopia	**nitrogen-fixing**; intercropping; **India 766 kg/ha vs world 1,015**
Oilseeds	15–30 · 30–75	loamy–sandy	Rajasthan, MP, Maharashtra	USA, China, Brazil, India	soybean, mustard, groundnut
Palm oil	25–28 · >200	tropical well-drained	AP, Telangana, Kerala	Indonesia, Malaysia, Thailand, Colombia, Nigeria	deforestation concern; push in NE
Tea	16–32 · 150–300	acidic, slopes	Assam, WB, TN, Kerala (+ Tripura, HP)	China, India, Kenya, Sri Lanka, Vietnam	plantation; labour-intensive
Coffee	15–28 · 150–200	acidic, fertile	Karnataka, Kerala, TN	Brazil, Vietnam, Colombia, Ethiopia, Indonesia	shade-grown; Arabica & Robusta
Cocoa	18–32 · 100–250	laterite	AP, Kerala, Karnataka, TN	Ivory Coast, Ghana, Indonesia, Nigeria, Ecuador	shade (under coconut/areca); **~70% of world beans from W. Africa (Ivory Coast, Ghana, Nigeria, Cameroon)**
Rubber	25–35 · 200–450	laterite, acidic	Kerala, TN	Thailand, Indonesia, Vietnam, India, China	latex
Cashew	20–30 · moderate	sandy loam	Maharashtra, AP, Odisha, Karnataka, Kerala	Vietnam, India, Ivory Coast, Brazil	export cash crop
Turmeric	20–30 · >150	sandy loam	Maharashtra, Telangana, TN, Karnataka	India, China, Myanmar, Bangladesh	**>30 varieties, >20 states**; **National Turmeric Board HQ Nizamabad (Telangana), inaugurated June 2025**

Crop	Temp / rain	Soil	Top Indian states	Top world	Remarks
Mango	24–30 · 75–250	deep loamy	UP, AP, Telangana, Karnataka, Bihar	India, China, Thailand, Indonesia, Mexico	**India is largest producer**; Alphonso, Dasheri, Langra
Coconut	20–32 · 100–200	sandy, coastal alluvial, laterite	Kerala, TN, Karnataka, AP	Indonesia, Philippines, India, Sri Lanka, Brazil	**Kalpavriksha**; humid coast
Red gram (tur)	20–30 · 60–100	loam–black	Maharashtra, Karnataka, UP, MP	India, Myanmar, Malawi, Tanzania, Kenya	Kharif; deep-rooted; N-fixing
Mustard	10–25 · 25–75	alluvial–loamy	Rajasthan, UP, Haryana, MP	Canada, China, India, Germany, France	Rabi; frost-sensitive; **sarson, rai**

Arable land: India has the largest arable area in the world, **≈ 1.597 million km².**

Sugarcane moved south: tropical climate, **higher sucrose**, **longer crushing season (Oct–May vs Nov–Feb)**, **cooperatives more successful in the south.**

Types of Farming in India

- **Irrigated:**
  - **Protective:** Irrigation is done to **protect the crops from drying** up.
  - **Productive:** Irrigation is done to **increase the produce harvested from the crops** .
- **Dry farming:** It is done in **places where rainfall is <75 cm and the growing season is <75 days** . Focus needs to be on moisture conservation and drought-resistant crops.
- **Dryland farming:** It is done where the **rainfall** is **75-115 cm** , and the **growing season is between 75-120 days** . Focus on s oil moisture retention, drought-tolerant crops, and adaptive strategies.
- **Wetland farming:** Practiced in locations where the **rainfall is >115cm** , and the **growing season is >120 days** . Focus on rainwater harvesting, moisture conservation, and suitable crop selection.

- **In India, almost 80 percent of maize, 90 percent of Bajra, approximately 95 percent of pulses, and 75 percent of oilseeds are obtained from dry or dryland agriculture.**

Micro Irrigation

- Use of methods to **improve the water use efficiency of agriculture** .
- Includes use of Drip Irrigation, Sprinklers, Foggers, etc.
- **Advantages:**
  - Water Conservation by increasing water use efficiency.
  - **Prevention of soil erosion** as the fields are not flooded with water.
  - **No need to level the fields** .

  - Can be **used in all types of soil** .
  - **Uniform irrigation** and **reduced nutrient loss** , thereby increasing crop productivity.
  - **Fertilizers/Pesticides etc can be added** to the water and provided to the plants.
  - Drip irrigation: Needs lesser pressure to operate, reducing energy costs.
- **Disadvantages:**
  - **High Investment**
  - **Clogging of pipes** can render certain regions devoid of water.
  - **Drip irrigation needs defined spacing between crops** , so not feasible in regions where
  - **Technical installation** : needs expertise, otherwise waste of resources (time, money, pipes, etc.)

Important Crops across the World

Crop	Temperature	Precipitation	Soil	Major Producing States (India)	M Pro Cou (W
Rice	20–32 °C	150–300 cm	Deep clayey & alluvial	Uttar Pradesh, Telangana, West Bengal	India, Bangl Indon Vietn
Wheat	10–25° C	50–100 cm	Loamy, well-drained	Uttar Pradesh, Madhya Pradesh, Punjab	China India, USA
Jowar (Sorghum)	20–27° C	50–75 cm	Light sandy to loamy	Maharashtra, Karnataka, Madhya Pradesh	India, China Niger
Bajra (Pearl Millet)	20–27° C	50–75 cm	Sandy loam	Rajasthan, Uttar Pradesh, Gujarat	India, China Niger
Millets (General)	20–27° C	50–75 cm	Wide range (clay to sandy)	Rajasthan, Karnataka, Madhya Pradesh	India, China Niger

MY NOTES

Maize	15–27° C	50–80 cm	Well-drained loamy	Madhya Pradesh, Karnataka, Bihar	USA, Brazil Arger
Cotton	21–30° C	50–100 cm	Black soil (regur)	Maharashtra, Gujarat, Telangana	China Brazil Austr
Jute	24–35 °C	120–200 cm	Alluvial	West Bengal, Bihar, Assam	India, Bang China
Sugarcane	20–27° C	100–150 cm	Loamy	Uttar Pradesh, Maharashtra, Karnataka	Brazil China Thaila Pakis
Pulses	15–27° C	Low rainfall	Well-drained	Madhya Pradesh, Maharashtra, Uttar Pradesh	India, Cana Myan Brazil Ethio
Oilseeds	15–30° C	30–75 cm	Loamy to sandy	Rajasthan, Madhya Pradesh, Maharashtra	USA, Brazil
Palm Oil	25–28 °C	>200 cm	Well-drained tropical	Andhra Pradesh, Telangana, Kerala	Indon Malay Thaila Color Niger
Tea	16–32° C	150–300 cm	Acidic, well-drained (slopes)	Assam, West Bengal, Tamil Nadu, Kerala. Minor states: Tripura, Himachal	China Kenya Lanka Vietn

				Pradesh	
Coffee	15–28° C	150–200 cm	Acidic, fertile	Karnataka, Kerala, Tamil Nadu	Brazil Vietn Color Ethio Indon

MY NOTES

Cocoa	18–32	100–250	Well-drained laterite soil	Andhra Pradesh, Kerala, Karnataka, Tamil Nadu	Ivory Ghan Indon Niger Ecua
Rubber	25–35 °C	200–450 cm	Laterite, acidic	Kerala, Tamil Nadu	Thaila Indon Vietn India,
Cashew	20–30 °C (tropical)	Moderate rainfall	Sandy loam	Maharashtra, Andhra Pradesh, Odisha, Karnataka, Kerala	Vietn India, Coas
Turmeric	20–30 °C	>150 cm	Sandy loam	Maharashtra, Telangana, Tamil Nadu, Karnataka	India, Myan Bang
Mango	24–30	75–250	Deep,	Uttar	India,

	°C (frost-free)	cm	well-drained loamy (can tolerate laterite &	Pradesh, Andhra Pradesh, Telangana, Karnataka, Bihar	Thaila Indon Mexic

⚡ **Quick revise** Farming by water — Fold ▲

Farming by water.

> 💧 **"75 – 115 – 120" ladder. Dry farming:** rain **< 75 cm**, season **< 75 days** (moisture conservation, drought-resistant crops). **Dryland: 75–115 cm**, **75–120 days**. **Wetland: > 115 cm**, **> 120 days** (rainwater harvesting). **Dry/dryland farming gives ≈ 80% of maize, 90% of bajra, ~95% of pulses, 75% of oilseeds.**

Irrigated: **protective** (saves crops from drying) vs **productive** (raises yield).

Micro-irrigation (drip, sprinkler, foggers): saves water, stops soil erosion, no levelling, any soil, uniform watering, **fertigation**, drip needs low pressure. **Costs:** high investment, **pipe clogging**, spacing needed, technical installation. *PYQ 2011 (ans C):* reduces nutrient loss ✔; checks falling groundwater table in some areas ✔; **"the only means of irrigation in dryland farming" ✘.**

System of Rice Intensification

			alluvial)		
Coconut	20–32 °C	100–200 cm (well distributed)	Sandy loam, coastal alluvial, laterite (well-drained)	Kerala, Tamil Nadu, Karnataka, Andhra Pradesh	Indon Philip India, Lanka
Red Gram (Tur / Pigeon Pea)	20–30 °C	60–100 cm (can tolerate low rainfall)	Well-drained loamy to black soils	Maharashtra, Karnataka, Uttar Pradesh, Madhya Pradesh	India, Myan Malaw Tanza Kenya
Mustard (Rapeseed & Mustard)	10–25° C	25–75 cm (low rainfall)	Alluvial to loamy soils (well-drained)	Rajasthan, Uttar Pradesh, Haryana, Madhya Pradesh	Canad China Germ Franc

System of Rice Intensification

- Originated in Madagascar in 1980s.
- Increase yields while reducing resources used.
- Instead of continuous flooding, used AWD (Alternate Wetting and Drying) technique for rice nurseries.
- Encourages use of organic manures.
- Known to improve yields by 30-50%.
- Reduces Methane emissions.

⚡ **Quick revise** SRI (System of Rice Intensification) Fold ▲

SRI (System of Rice Intensification): Madagascar, 1980s; **AWD (alternate wetting and drying)**; organic manure; yields **+30–50%**; **less methane**.

The shift of sugarcane industry to the South

- Tropical Climate better for sugarcane yield

- Higher sucrose content
- Longer crushing season (does not get extremely hot in the summers): only Nov-Feb in North; October to May in South
- Cooperatives more successful in South

Agriculture Revolutions in India

- **Green Revolution**
  - **Period: Late 1960s to early 1980s** .
  - **Key Figure: Dr. M.S. Swaminathan.**
  - **Main Focus** : Introduction of **high-yielding varieties (HYVs) of seeds, increased use of fertilizers, pesticides, and irrigation.**
  - **Successes:**
    - Significant **increase in food grain production** , particularly wheat and rice.
    - Achieved self-sufficiency in food grains.
    - **Reduced the import dependency** for grains.
  - **Drawbacks:**
    - **Regional disparities** , with major benefits concentrated in Punjab, Haryana, and Western Uttar Pradesh.
    - **Social disparities** , as only the farmers with significant land holdings benefitted.
    - **Environmental degradation** due to overuse of chemical fertilizers and pesticides.
    - **Water table depletion** due to intensive irrigation.
- **White Revolution (Operation Flood)**
  - **Period: 1970s onwards** .
  - **Key Figure: Dr. Verghese Kurien.**
  - **Main Focus:** Development of the **dairy industry** .
  - **Successes:**
    - India became the **largest producer of milk** in the world.
    - **Improved rural incomes and employment** .
    - Created a **robust cooperative network** (e.g., Amul).
  - **Drawbacks:**
    - Challenges in **maintaining quality standards** .
    - **Inadequate cold chain infrastructure** in many parts of the country.
    - **Inadequate processing** capabilities.
- **Blue Revolution**
  - **Period: 1980s onwards** .
  - **Main Focus:** Development of the **fisheries sector** .
  - **Successes:**
    - **Increase in fish production** , making India one of the top fish-producing countries.
    - **Enhanced export** opportunities. Around 17% of India’s agricultural exports in 2021-22.

    - **Improved nutritional standards and employment** in coastal regions.
  - **Drawbacks:**
    - **Overfishing and depletion of marine resources** .
    - **Environmental issues** such as water pollution and habitat destruction.
- **Yellow Revolution**
  - **Period: 1980s onwards** .
  - **Main Focus:** Increase in **oilseed** production.
  - **Successes:**
    - Significant **increase in the production of oilseeds** like mustard, sunflower, and soybean.
    - **Reduced dependency** on edible oil **imports** .
  - **Drawbacks:**
    - **Inconsistent growth and production challenges** due to climatic variability.
    - **Market fluctuations** affecting farmer incomes.
- **Evergreen Revolution**
  - **Proposed by: Dr. M.S. Swaminathan** .
  - **Main Focus: Sustainable agriculture** through a blend of traditional and modern practices.
  - **Successes:**
    - Emphasis on **ecological sustainability** .
    - Focus on **genetic diversity and conservation of natural resources** .
  - **Drawbacks:**
    - **Implementation challenges** on a large scale.
    - Need for **significant policy support and farmer education** .
- **Other Revolutions:**
  - Brown Revolution: Coffee/Leather/Cocoa
  - Golden Revolution: Fruits/Honey/Horticulture (Nirpakh Tutaj)
  - Golden Fibre: Jute
  - Silver Revolution: Eggs
  - Silver Fibre Revolution: Cotton
  - Pink Revolution: Onion/Pharma/Prawn
  - Red Revolution: Tomato/Meat
  - Round Revolution: Potato
  - Evergreen Revolution: Ecology with technology (M.S. Swaminathan)
  - Grey Revolution: Fertilizers
  - Protein Revolution: encourages farmers to adopt scientific agricultural techniques to boost food grain production and reduce the import of protein-rich lentils.

⚡ **Quick revise** 1.3 Agricultural revolutions Fold ▲

🌈 **Colour = crop. Green** = grains · **White** = milk · **Blue** = fish · **Yellow** = oilseeds · **Golden** = fruit/honey/horticulture · **Golden fibre** = jute · **Silver** = eggs · **Silver fibre** = cotton · **Pink** = onion/pharma/prawn · **Red** = tomato/meat · **Round** = potato · **Brown** = coffee/leather/cocoa · **Grey** = fertiliser · **Protein** = lentils · **Evergreen** = ecology + technology.

Revolution	When · who	Win	Cost
Green	late 1960s–early 1980s · **M.S. Swaminathan**	HYV seeds + fertiliser + irrigation; wheat/rice jump; **food self-sufficiency**	regional (Punjab, Haryana, W. UP), social (big farmers), pollution, **water-table fall**
White (Operation Flood)	1970s · **Verghese Kurien**	**world's largest milk producer**; cooperatives (**Amul**)	quality, cold chain, processing
Blue	1980s	fish output; **~17% of agri exports (2021-22)**	overfishing, pollution
Yellow	1980s	oilseeds; fewer edible-oil imports	climate swings, price swings
Evergreen	Swaminathan's proposal	sustainable mix of traditional + modern	scale, policy support

Agricultural Regions of the World

Developing World:

- **Nomadic Herding:**
  - Mainly 4th world communities in unfavourable envt. Regions.
  - Camels (Bedouins), Cattle (Masais), Horses (Kazaks), Reindeers (Chukchis)
- **Livestock Ranching:**
  - Subsistence level: African Savannah
  - Commercial level: Australia (sheep, cattle); Brazil (beef)
- **Shifting Agriculture:**
  - Mostly by 4th world community
- **Sedentary Agriculture:**
  - Rotation of crops instead of fields
  - Transition between shifting & subsistence.
- **Intensive Subsistence**
  - Small land holding; intensive use of resource; mostly for self-consumption.
  - Multiple cropping seasons
  - With Paddy: Wetter margins
  - Without Paddy: drier climates

- Primitive type: true subsistence; absolute dependence on manual labour; no tech; no surplus
- Non-Primitive: Use of tech; Significant surplus

- **Mixed Subsistence:**
  - Evolving
  - In Savannah type climate

Temperate Developed:

- **Extensive Commercial Grain Farming:**
  - Breadbasket of Anglo-America & Wheat Triangle of Eurasia.
  - Temperate Grasslands
  - Intensive commercial development in India & China
- **Extensive Commercial Grain Farming:**
  - Dual benefit
  - Largest geographical expanse of temperate regions
- **Commercial Dairy Farming:**
  - Intensive Commercial: needs manual + machine labour.
  - Wetter temperate margins
  - India: tropical, but leading milk producer

Specialized Agriculture

- **Plantation:**
  - Extensive in the tropics
  - Same stand gives yield for years, so special care
  - Specialized harvesting
  - Coffee (Brazil); Cocoa (Ghana), Rubber (Indonesia, Malaysia), Spices (India), Tea (China)
  - Non-traditional plantation: 2-3 yrs; Cotton & Cane (India, Brazil, Egypt, etc)
- **Mediterranean:**
  - Intensive Commercial, sometimes mixed culture
  - Gardens & Vineyards of the world
- **Horticulture:**
  - Truck farming
  - Intensive: manual labour
  - Traditionally in temperate regions [Bulgaria (roses), Netherlands (tulips)], now in the tropics too [India (marigolds, roses, lilies, brinjals, etc), China (Potatoes, tulips)]
  - Special fertilizers and harvesting
- **Cooperative Farming:**
  - Farmers come together to form cooperatives through resource pooling (except land).
  - Cooperatives help farmers get access to farm inputs at affordable rates.
  - Increases the negotiating powers of the farmers.

- Successful in many West European nations like Belgium, Netherlands, and especially Denmark.
- Eg in India: IFFCO (Indian Farmers Fertilisers Cooperative Limited); Amul.

- **Collective Farming:**
  - In USSR: Kolkhoz
  - State ownership of land and other resources.
  - Farmers work together on collective lands.
  - Small land parcels provided to farmers only for their personal needs.
  - Problem: Free-riding

⚡ **Quick revise** 1.4 World agricultural types Fold ▲

Type	Key points (as printed)
Nomadic herding	4th-world communities; **camels – Bedouins · cattle – Masai · horses – Kazakhs · reindeer – Chukchis**
Livestock ranching	subsistence – African savannah; commercial – **Australia (sheep, cattle), Brazil (beef)**
Shifting agriculture	mostly 4th-world communities; **sedentary** = rotation of crops, not fields
Intensive subsistence	small holdings, multiple seasons; **with paddy – wetter margins; without paddy – drier**; *primitive* (manual, no surplus) vs *non-primitive* (tech, surplus)
Extensive commercial grain	**breadbasket of Anglo-America, Wheat Triangle of Eurasia**; temperate grasslands
Commercial dairy	wetter temperate margins; India = tropical yet leading milk producer
Plantation	tropics; years of yield from the same stand – **coffee (Brazil), cocoa (Ghana), rubber (Indonesia, Malaysia), spices (India), tea (China)**; non-traditional (2–3 yrs): cotton, cane
Mediterranean	gardens and vineyards
Horticulture / truck farming	**Bulgaria (roses), Netherlands (tulips)**; now India (marigold, rose, lily, brinjal)
Cooperative farming	pool resources **except land**; Belgium, Netherlands, **Denmark**; **IFFCO, Amul**
Collective farming	USSR **Kolkhoz**; state owns land; **free-riding** problem

Agricultural Terminologies

- **Organic Farming** : Growing crops without synthetic pesticides or fertilizers.
- **Permaculture** : Designing agricultural systems that mimic natural ecosystems.
- **Agroforestry** : Integrating trees and shrubs into agricultural landscapes.
- **Regenerative Agriculture** : Focus on renewing soil health and ecosystems.

- **Conservation Agriculture** : Practices that preserve and improve soil health.
- **Hydroponics** : Growing plants in nutrient-rich water without soil.
- **Aquaponics** : Combining aquaculture (raising fish) with hydroponics.
- **Aeroponics** : Growing plants in an air or mist environment without soil.
- **Vertical Farming** : Growing crops in stacked layers, often indoors.
- **Precision Farming** : Using technology (like GPS and IoT) to optimize field-level management.
- **Smart Farming** : Incorporating modern technologies like AI, drones, and sensors to enhance productivity and efficiency.
- **Crop Rotation** : Alternating crops in the same field to maintain soil fertility.
- **Intercropping** : Growing different crops in proximity for pest control and better resource use.
- **Cover Cropping** : Growing specific crops to protect and enrich soil between periods of regular crop production. E.g. Mustard, Clover, Radish, etc.
- **Silviculture** : Managing forest plantations for timber and other forest products.
- **Viticulture** : Cultivation of grapevines for winemaking.
- **Apiculture** : Beekeeping for honey and other bee products.
- **Sericulture** : Raising silkworms for silk production.
- **Horticulture** : Cultivating fruits, vegetables, flowers, and ornamental plants.
- **Floriculture** : Growing flowers and ornamental plants for the floral industry.
- **Climate-Smart Agriculture:** Practices that increase productivity, resilience, and reduce emissions.
- **Biopharming** : Using genetically modified plants and animals to produce pharmaceuticals.
- **Agrivoltaic Farming:** Combinind solar energy production with farming, to maximize land use and rural energy access.

⚡ **Quick revise** Terms in one line each Fold ▲

Terms in one line each: Organic (no synthetic inputs) · **Permaculture** (mimic natural ecosystems; mulching; multicropping) · **Agroforestry** · **Regenerative** (renew soil) · **Conservation agriculture** (minimum tillage, residue cover, rotations) · **Hydroponics** (nutrient water) · **Aquaponics** (fish + hydroponics) · **Aeroponics** (mist) · **Vertical farming** · **Precision** (GPS, IoT) · **Smart** (AI, drones) · **Intercropping** · **Cover cropping** (mustard, clover, radish) · **Silviculture** (forest plantations) · **Viticulture** (grapes) · **Apiculture** (bees) · **Sericulture** (silk) · **Floriculture** · **Climate-smart** · **Biopharming** · **Agrivoltaic** (solar + farming).

Natural Farming

- **Natural farming** (also known as " **Fukuoka Method"**) is an agricultural philosophy and practice that **emphasizes minimal human intervention and working in synergy with nature** .

- It **avoids synthetic chemicals, fertilizers, pesticides, and excessive tilling** .
- The concept **promotes biodiversity, soil health, and sustainable farming practices** using locally available natural resources.
- It was **established by Japanese farmer and philosopher Masanobu Fukuoka** in his 1975 book The One-Straw.
- **Zero Budget Natural Farming (ZBNF) is a low-cost approach to natural farming** that relies on locally made preparations and avoids external inputs like pesticides and fertilizers. It does not treat weeds, insects and animals as enemies, but tries to emulate the natural conditions.
- Key concepts:
  - **Minimal Human Intervention** : The core idea is to let nature follow its course as closely as possible.
  - **Use of Natural Inputs** : Reliance on natural fertilizers and pest control methods, such as compost, manure, and biological pest control.
  - **Soil Health** : Maintaining healthy soil ecosystems through practices like crop rotation, cover cropping, and reduced tillage.

⚡ **Quick revise** Natural farming (Fukuoka) Fold ▲

Natural farming (Fukuoka): minimal intervention; **Masanobu Fukuoka, *The One-Straw Revolution*, 1975**; **ZBNF (Zero Budget Natural Farming)** = low-cost, local preparations, no weeds-as-enemies.

Land Use Categories of India

- According to the **Land Revenue records** , land use in India is mainly of 9 types:
  - **Forests:** Area identified by the government as forest.
  - **Barren & Wasteland:** Area that cannot be brought under cultivation on the basis of the available technologies.
  - **Land put to Non-Agri Use:** Settlements, Commercial centres, Industries, Infrastructure etc.
  - **Area under Permanent Pastures and Grazing Lands:** Most of such land is owned by the Panchayat (Common Property Resource) or government. (Rarely Private)
  - **Area under Miscellaneous Tree Crops and Groves:** Orchards and fruit trees. (Mostly Private)
  - **Culturable Wasteland:** Land left fallow for >5yrs. Can be reclaimed.
  - **Current Fallow:** Land left without cultivation for less than or equal to 1 agricultural year. (Land resting for re-couping)
  - **Fallow other than Current Fallow:** Left fallow for less than 5 yrs but more than and year.
  - **Net Sown Area:** Land used for cultivation of crops
- **Land Use Change:**
  - **Changes due to changes in the economic activities.**

- Over time, land under cultivation reduces (improvement of agricultural production, diversion of land for non-agricultural activities)
- Land under forests can also reduce due to diversion.
- As land is a finite resource, wastelands are also either reclaimed for agriculture or other works. Similarly Barren lands start getting used for non-agricultural activities.
- **In India, these land use records are maintained by Land Revenue Department** (exercise by Survey of India)
- **Digital India Land Records Modernization Program** is working to digitize land records in India.

- Land Degradation: Covered in Soils of India

⚡ **Quick revise**

Land use (9 categories, land-revenue records) Fold ▲

Land use (9 categories, land-revenue records): Forests · Barren & wasteland · Non-agricultural use · Permanent pastures (mostly Panchayat/government) · Misc. tree crops and groves (mostly private) · Culturable wasteland (fallow > 5 yrs) · Current fallow (≤ 1 year) · Fallow other than current (1–5 yrs) · Net sown area. Records kept by Land Revenue Department (Survey of India); **Digital India Land Records Modernization Programme.**

PYQ

Q.) With reference to micro-irrigation, which of the following statements is/are correct?

1. Fertiliser/nutrient loss can be reduced.

2. It is the only means of irrigation in dryland farming.

3. In some areas of farming, the receding of the groundwater table can be checked.

Select the correct answer using the codes given below: (2011)

(a) 1 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: C Micro-irrigation is a modern method of irrigation; by this method, water is irrigated through drippers, sprinklers, foggers and by other emitters on the surface or subsurface of the land.

- Major components of a micro irrigation system are as follows.
  - Water source, pumping devices (motor and pump), ball valves, fertigation equipment, filters, control valves, PVC joining accessories (Main and sub-main) and emitters.

- In this system, water is applied drop by drop nearer the root zone area of the crop.
- The drippers are fixed based on the spacing of the crop.
- Many different types of emitters are available in the market. They are classified as Inline drippers, online drippers, Microtubes, and Pressed compensated drippers.
- Drip irrigation is most suitable for wider spacing crops.
- The micro sprinkler irrigation system is mostly followed in sandy or loamy soils.
- This system is most suitable for horticultural crops and small grasses.
- In this method, water is sprinkled at a lower height in various directions.
- It is used for preparing nursery and lawns in soils with low water holding capacity.

Advantages of drip irrigation system-

- Water saving and higher yield
- High quality and increased fruit size
- Suitable for all types of soil
- The easy method of fertigation and chemigation
- Saving in labour and field preparation cost

Q.) The lower Gangetic plain is characterised by a humid climate with high temperatures throughout the year. Which one among the following pairs of crops is most suitable for this region? (2011)

(a) Paddy and cotton

(b) Wheat and Jute

(c) Paddy and Jute

(d) Wheat and cotton

Answer: C The lower deltaic plains of the Ganges and Brahmaputra rivers that are Characterised by swamps and Sundarbans.

1. The suitable climate for growing jute is a warm and wet climate, which is offered by the monsoon climate during the fall season, immediately followed by summer. Temperatures ranging to more than 25 °C and relative humidity of 70%–90% are favourable for successful cultivation.

2. Rice crops need a hot and humid climate. It is best suited to regions which have high humidity, prolonged sunshine and an assured supply of water. The average temperature required throughout the life period of the crop ranges from 21 to 37º C. Maximum temp which the crop can tolerate is 400C to 42 0C.

Q.) A state in India has the following characteristics:

1. Its northern part is arid and semi-arid.

2. Its central part produces cotton.

3. Cultivation of cash crops is predominant over food crops.

Which one of the following states has all of the above characteristics? (2011)

(a) Andhra Pradesh

(b) Gujarat

(c) Karnataka

(d) Tamil Nadu

Answer: B

1. Gujarat possesses varied landforms with the dry deciduous forests, majestic grasslands, marine ecosystems, wetlands and rich moist deciduous forests.

2. These habitats are home to rare wildlife inhabitants.

3. The three main sources of growth in Gujarat's agriculture are from cotton production, the rapid growth of high-value foods such as livestock, fruits and vegetables, and from wheat production, which saw an annual average growth rate of 28% between 2000 and 2008 (According to the International Food Policy Research Institute).

Q.) Salinization occurs when the irrigation water accumulated in the soil evaporates, leaving behind salts and minerals. What are the effects of salinization on the irrigated land? (2011)

(a) It greatly increases the crop production

(b) It makes some soils impermeable

(c) It raises the water table

(d) It fills the air spaces in the soil with water

Answer: B Soil salinization occurs when water-soluble salts accumulate in the soil to a level that impacts on agricultural production, environmental health, and economics.

- Salt at higher levels causes soil aggregates to break down, collapsing soil pores and reducing the movement of water and air.
- In the early stages, salinity affects the metabolism of soil organisms and reduces soil productivity, but in advanced stages it destroys all vegetation and other organisms living in the soil, consequently transforming fertile and productive land into barren and desertified lands

Q.) Which of the following is the chief characteristic of ‘mixed farming’? (2012)

(a) Cultivation of both cash crops and food crops

(b) Cultivation of two or more crops in the same field

(c) Rearing of animals and cultivation of crops together

(d) None of the above

Answer: C Mixed farming is a type of agriculture which involves both crops and livestock.

(a) This is not to be confused with mixed cultivation, implying merely a series of different crops.

(b) It is found throughout Europe— from Ireland in the west through Central Europe to Russia.

(c) It is also found in North America, east of the 98° meridian, in the ‘pampas’ of Argentina, southeast Australia, South Africa and New Zealand.

(d) Mixed farming is essentially associated with the densely populated, urbanised and industrialised societies, dependent upon high incomes for the sale of its products and upon manufacturing in dustry for the provision of its inputs.

Q.) Consider the following crops of India :

1. Groundnut

2. Sesamum

3. Pearl millet

Which of the above is/are predominantly rainfed crop/crops? (2012)

(a) 1 and 2 only

(b) 2 and 3 only

(c) 3 only

(d) 1, 2 and 3

Answer: D Sesamum is mainly a rainfed crop and is grown on marginal, dry and sloppy lands.

- Nearly 80% of the area sown to groundnuts in India is rain-fed and relies entirely on summer monsoon rainfall.
- Among cereals, the main rainfed cereals grown in India are sorghum, pearl millet (Bajra), Finger Millet (Ragi).

Q.) Consider the following crops

1. Cotton

2. Groundnut

3. Rice

4. Wheat

Which of these are Kharif crops? (2013)

(a) 1 only

(b) 1, 2 and 3 only

(c) 2 and 3 only

(d) 1, 2, 3 and 4

Answer: B The word “Kharif” is Arabic for autumn since the season coincides with the beginning of autumn or winter.

- Kharif crops also are known as monsoon crops. These are the crops that are cultivated in the monsoon season.
- The Kharif season differs in every state of the country but is generally from June to September.
- These crops are usually sown at the beginning of the monsoon season around June and harvested by September or October.
- Rice, maize, bajra, ragi, soybean, groundnut, cotton are all Kharif types of crops.

Q.) Region Well-known for the production of

1. Kinnaur (Himachal) Areca nut

2. Mewat (Haryana) Mango

3. Coromandel (TN-Andhra) Soya bean

Which of the above pairs is/are correctly matched? (2014)

(a) 1 and 2 only

(b) 3 only

(c) 1, 2 and 3

(d) None

Answer: D Portions of Kinnaur are situated high in the Himalaya, where vegetation is sparse and consists primarily of hardy grasses. Alpine species such as juniper, pine, fir, cypress, and rhododendron can be found at elevations between 3,500 and 5,000 metres, primarily in Middle Kinnaur. At lower altitudes, temperate-climate trees are found, including oak, chestnut, maple, birch, alder, magnolia, apple, and apricot.

- Yaks and dzos are reared by local farmers in the higher areas. Scattered populations of the Himalayan black bear and small ponies may also be found.
- The main occupation in the Mewat district is agriculture along with allied and agro-based activities.
- The Meos are the predominant population group and are all agriculturists.
- Agriculture is mostly rain-fed except in small pockets where canal irrigation is available.
- Agriculture production measured in terms of crop yield per hectare is low in comparison to the other districts of the State.
- Animal husbandry, particularly dairy, is a secondary source of income for the people and those who live closer to the hilly ranges of Aravali also keep sheep and goats.
- MP is famous for the production Soybean

Q.) Consider the following statements :

1. Maize can be used for the production of starch.

2. Oil extracted from maize can be a feedstock for biodiesel.

3. Alcoholic beverages can be produced by using maize.

Which of the statements given above is/are correct? (2014)

(a) 1 only

(b) 1 and 2 only

(c) 2 and 3 only

(d) 1, 2 and 3

Answer: D In Central America, the alcoholic beverage brewed from malted maize is known as chica fuerte.

- Corn starch has a wide variety of industrial applications. Biodiesel can be produced from corn oil.
- Maize can be used for the production of starch.Oil extracted from maize can be a feedstock for biodiesel. Alcoholic beverages can be produced by using maize.
- The type of alcohol in the alcoholic drinks we drink is a chemical called ethanol. To make alcohol, you need to put grains, fruits, or vegetables through a process called fermentation. The common source of ethanol is maize and other grains.

Q.) What can be the impact of excessive/inappropriate use of nitrogenous fertilisers in agriculture?

1. The proliferation of nitrogen-fixing microorganisms in the soil can occur.

2. Increase in the acidity of soil can take place

3. Leaching of nitrate to the ground-water can occur.

Select the correct answer using the code given below. (2015)

(a) 1 and 3 only

(b) 2 only

(c) 2 and 3 only

(d) 1,2 and 3

Answer: C Farmers occasionally need to add nitrogen fertiliser to their farms and gardens to make available just the precise nutrients for their plants' growth.

- The applications of inorganic nitrogen fertilisers to various crops have been continuously increasing since the last many decades globally.
- Excessive use of N (urea) encourages climate change (when lost through denitrification) and groundwater pollution (when lost through leaching).
- Increase in nitrate content of groundwater in some intensively-cropped areas has been reported, which is obviously due to leaching of nitrates beyond the crop root zone. Increase in nitrate content of groundwater is potentially harmful, as it is used for drinking purposes in most of the rural areas.
- Although nitrogen fertiliser contributes substantially to yield enhancement, excessive use of this manure has posed serious threats to the environment and human health.
- The rate of nitrogen fertiliser application has a close relationship with nitrate accumulation in the surrounding environment, groundwater, as well as leafy and root vegetables.

- Excessive use of urea leads to several adverse implications on soil, crop quality and overall ecosystem.
- Nitrogen applied in excess of crop demand is lost through volatilization, acidification, denitrification and leaching.

Q.) With reference to the circumstances in Indian agriculture, the concept of "Conservation Agriculture" assumes significance. Which of the following fall under the Conservation Agriculture?

1. Avoiding monoculture practises

2. Adopting minimum tillage

3. Avoiding the cultivation of plantation crops

4. Using crop residues to cover the soil surface

5. Adopting spatial and temporal crop sequencing/crop rotations

Select the correct answer using the code given below: (2017)

(a) 1, 3 and 4

(b) 2, 3, 4 and 5

(c) 1, 2, 3 and 5

(d) 2, 4 and 5

Answer: D

- Conservation agriculture (CA) technologies involve minimum soil disturbance, permanent soil cover through crop residues or cover crops, and crop rotations for achieving higher productivity.
- Main features of Conservation agriculture (CA) are; Least interference with natural processes, No-till or drastically reduced tillage (biological tillage, Low wind and soil erosion, Surface retention of residues (permanently covered), Infiltration rate of water is high, Use of in-situ organics/composts, Brown manuring/cover crops (surface retention),Diversified and more efficient rotations etc.
- The principal indicators of non-sustainability of agricultural systems include soil erosion, soil organic matter decline, salinization. These are caused mainly by
  - intensive tillage-induced soil organic matter decline, soil structural degradation, water and wind erosion, reduced water infiltration rates, surface sealing and crusting, soil compaction,
  - insufficient return of organic material, and
  - Monocropping.

Q.) With reference to agricultural soils, consider the following statements :

1. The high content of organic matter in soil drastically reduces its water holding capacity.

2. Soil does not play any role in the sulphur cycle.

3. Irrigation over a period of time can contribute to the salinization of some agricultural lands.

Which of the statements given above is/are correct? (2018)

(a) 1 and 2 only

(b) 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: B

- Plants obtain nutrients from two natural sources: organic matter and minerals. Organic matter includes any plant or animal material that returns to the soil and goes through the decomposition process. In addition to providing nutrients and habitat to organisms living in the soil, organic matter also binds soil particles into aggregates and improves the water holding capacity of the soil.
- The sulphur cycle is the collection of processes by which sulphur moves to and from rock, waterways and living systems. The isotopic composition of sedimentary sulphides provides primary information on the evolution of the sulphur cycle.
- Too much salt can reduce crop production and water infiltration on soils that have been irrigated for 20 years or more. Too little salt can also result in a chemically compacted soil. The level of sodium (Na) in the soil can result in soil particles or even layers of differing soil profiles forming a compacted layer that the roots do not penetrate.

Q.) Consider the following statements:

1. Agricultural soils release nitrogen oxides into the environment.

2. Cattle release ammonia into the environment.

3. Poultry industry releases reactive nitrogen compounds into the environment.

Which of the statements given above is/are correct? (2019)

(a) 1 and 3 only

(b) 2 and 3 only

(c) 2 only

(d) 1, 2 and 3

Answer: D Statement 1 is correct:

- Nitrous oxide is emitted into the atmosphere as a result of biomass burning, and biological processes in soils. Biomass burning is not only an instantaneous source of nitrous oxide, but it results in a longer term enhancement of the biogenic production of this gas.
- The current belief is that 90% of the emissions come from soils. Nitrous oxide is formed in soils during the microbiological processes of nitrification and denitrification. Because nitrous oxide is a gas it can escape from soil during these transformations.

Statement 2 is correct:

- Cattle release both ammonia and methane into the atmosphere. Ruminant animals do not efficiently utilise dietary nitrogen.

Statement 3 is correct:

1. Nitrogen emissions in chicken production occur in several forms but mainly ammonia can contribute directly or indirectly to several environmental and public health hazards.

2. Chicken production also contributes to some extent to climate change through the emissions of nitrous oxide, fine particulate matter, and methane.

Q.) How is permaculture farming different from conventional chemical farming ?

1. Permaculture farming discourages monocultural practices but in conventional chemical farming, monoculture practices are predominant.

2. Conventional chemical farming can cause increase in soil salinity but the occurrence of such phenomenon is not observed in permaculture farming.

3. Conventional chemical farming is easily possible in semi-arid regions but permaculture farming is not so easily possible in such regions.

4. Practice of mulching is very important in permaculture farming but not necessarily so in conventional chemical farming.

Select the correct answer using the code given below. (2020)

(a) 1 and 3

(b) 1, 2 and 4

(c) 4 only

(d) 2 and 3

Answer: B

- Permaculture farming discourages monocultural practices but in conventional chemical farming, monoculture practices are predominant.
- 2. Conventional chemical farming can cause an increase in soil salinity but the occurrence of such phenomenon is not observed in permaculture farming.
- Statement 1 is correct. Permaculture is a totally integrated design system that’s modelled on nature. Permaculture farming promotes multicropping and integrated farming systems. Agroforestry is its example.
- Chemical farming is more suitable for monoculture cropping due to use of crop specific inputs like irrigation, chemical fertiliser and harvesting methods.
- Statement 2 is correct. Chemical fertilizers will make the soil less fertile over time, which can lead to serious problems like soil salinisation. But such problems are not seen in permaculture farming as it relies on organic fertilizers.
- Statement 3 is incorrect. Chemical farming is based on intense use of inputs like Chemical fertilizers and irrigation that make the soil less fertile over time, thus restricting its practise in semi-arid region. Thus, conventional farming is not suitable for semi-arid regions.

- The concept of permaculture involves well-designed systems don't produce waste and permaculture tries to mimic well-designed systems. Thus, permaculture tries to take into account the local conditions like arid climate. This helps in developing a suitable system to ensure sustainable production.
- Statement 4 is correct. Mulching is an important component of Permaculture systems for promoting maximum efficiency.

Q.) With reference to 'palm oil', consider the following statements:

1. The palm oil tree is native to Southeast Asia.

2. The palm oil is a raw material for some industries producing lipstick and perfumes.

3. palm oil can be used to produce biodiesel.

Which of the statements given above are correct? (2021)

(a) 1 and 2 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

Answer: B

- The palm oil tree is native to hills of guinea.
- It is used in cosmetics products like lipstick and perfumes and cleaning products. Palm oil is one of the best options and most cost-effective cosmetics ingredients. It is versatile. Due to its high fatty acid content, it can be used to make surfactants, emulsifiers and emollient
- Palm oil can become a sustainable option for biodiesel production because it is a bye product of palm oil (Pannilawithana and Pathirana, 2017). Both unrefined palm oil (UPO) and palm kernel oil are primary oil sources from oil palm trees.

Q.) Among the following, which one is the least water-efficient crop? (2021)

(a) Sugarcane

(b) Sunflower

(c) Pearl millet

(d) Red gram

Answer: A

- Option a is correct. Sugarcane is the least water efficient crop. Among the given options, its water requirement is 1800-2200 mm/season which is highest.
- Option b is incorrect. Sunflower's water requirement was estimated at 672.4 mm/season.
- Option c is incorrect. Pearl millet is a drought tolerant crop.
- Water requirement: 350 mm/season.

Q.) Consider the following States:

1. Andhra Pradesh

2. Kerala

3. Himachal Pradesh

4. Tripura

How many of the above are generally known as tea-producing States? (2022)

(a) Only one State

(b) Only two States

(c) Only three States

(d) All four States

Answer: D

- All of these states produce tea. Kerala is a major, while the rest are minor tea producing states.

Q.) Which of the following countries are well known as the two largest cocoa producers in the world? (2024)

(a) Algeria and Morocco

(b) Botswana and Namibia

(c) Côte d'Ivoire and Ghana

(d) Madagascar and Mozambique

Answer: D

Explanation: These two countries produce about 70% of world's cocoa.

Q.) Consider the following statements about turmeric during the year 2022-23: I. India is the largest producer and exporter of turmeric in the world. II. More than 30 varieties of turmeric are grown in India. III. Maharashtra, Telangana, Karnataka and Tamil Nadu are major turmeric producing States in India.

Which of the statements given above are correct? (2025)

(a) I and II only

(b) II and III only

(c) I and III only

(d) I, II and III

Answer: D

Explanation: India is the largest turmeric producer in the world, we also produce the maximum varieties.

⚡ Quick revise Agriculture PYQs

Fold ▲

Year	Asked	Answer / rule
2011	Micro-irrigation statements	**C (1 and 3)**
2011	Lower Gangetic plain, humid and hot all year: best crop pair	**Paddy and jute (C)**
2011	State: arid north, cotton in centre, cash crops > food crops	**Gujarat (B)**
2011	Effect of salinisation	**Makes some soils impermeable (B)** – salt breaks aggregates, pores collapse
2012	Chief feature of **mixed farming**	**Crops + livestock together (C)**
2012	Rainfed: groundnut, sesamum, pearl millet	**All three (D)**
2013	Kharif: cotton, groundnut, rice, wheat	**1, 2, 3 – not wheat (B)**
2014	Kinnaur–areca? Mewat–mango? Coromandel–soya?	**None (D)**
2014	Maize: starch, biodiesel oil, alcohol	**All three (D)**
2015	Excess nitrogen fertiliser	**Acidity ↑ and nitrate leaching (C)** – not "more N-fixing microbes"
2017	Conservation agriculture	**Minimum tillage, residue cover, rotations (D: 2, 4, 5)**
2018	Soil organic matter ↓ water holding? Soil no role in sulphur cycle? Irrigation salinises?	**Only 3 (B)**
2019	Agri soils emit nitrogen oxides; cattle ammonia; poultry reactive N	**All three (D)**
2020	Permaculture vs chemical farming	**1, 2, 4 (B)** – chemical farming is **not** easy in semi-arid areas
2021	Palm oil native to SE Asia ✘; lipstick/perfume raw material ✔; biodiesel ✔	**B (2 and 3)** – native to the hills of Guinea
2021	Least water-efficient crop	**Sugarcane (A)**
2022	Tea-producing: AP, Kerala, HP, Tripura	**All four (D)** – Kerala major; others minor
2024	Two largest cocoa producers	**Côte d'Ivoire and Ghana (C)** (~70% of world)
2025	Turmeric 2022-23: India largest producer/exporter; >30 varieties; Maharashtra, Telangana, Karnataka, TN	**All three (D)**

2 · WORLD RESOURCES

Resources are essentially **anything that we can use from our environment to fulfill our needs and wants** . **An object/entity should fulfill these 2 criteria to be called a resource** :

- **Availability:** The resource needs to be accessible somehow. This can involve technology to extract it, economic feasibility to develop it, and cultural acceptance of its use.
- **Utility:** It must be something that can be used for some kind of benefit.

Types of Resources:

Resources can be classified based on various parameters:

- **Based on Origin**
  - **Natural Resources:** Resources that exist without human intervention. E.g. Water, minerals, forests, and air.
  - **Man-made (Human-made) Resources:** Resources created by humans using natural resources. Buildings, machinery, infrastructure.
- **Based on Exhaustibility**
  - **Renewable Resources:** Resources that can be replenished naturally over time. E.g. Solar energy, wind energy, biomass, hydropower.
  - **Non-renewable Resources:** Resources that are finite and cannot be replaced within a human timescale once depleted. E.g. Fossil fuels (coal, oil, natural gas), minerals (gold, iron ore).
- **Based on Origin (Biotic and Abiotic)**
  - **Biotic Resources:** Resources derived from living organisms. E.g.: Forests, animals, plants, and fossil fuels (originated from ancient biological matter).
  - **Abiotic Resources:** Resources derived from non-living elements. E.g. Water, air, minerals, metals.
- **Based on Ownership**
  - **Individual Resources:** Resources owned by private individuals or companies. E.g. Private land, personal vehicles, privately owned mines.
  - **Community Resources:** Resources accessible to all members of a community. E.g. Public parks, community forests, common grazing lands.
  - **National Resources:** Resources owned by the nation and managed by the government. E.g. National highways, public sector industries, forests.
  - **International Resources:** Resources that are regulated by international bodies and are beyond the jurisdiction of any one country. E.g. International waters (High Seas), outer space.



Mining:

- Extracting Resources from the ground.
- Condition for a mine: **The cost of extraction and purification of the ore should be less than the value of the final product.**
- **Types** of Mines:
  - **Open Cast:** for resources near the surface
  - **Shaft:** for resources at a deeper level

Mineral	Major Ores	Uses	Top producers and reserves in world	Top producers In
Iron ore	Hematite, Magnetite, Limonite, Siderite	It is mainly used **for the production of Steel**.	**Producers:** Australia, Brazil, India, China, Russia; **Reserves:** Australia, Brazil, Russia, China, Mauritania.	**Reserves:** 95% i Jharkhand, Chh Karnataka, Goa, Andhra Pradesh **Producers / maj** (Mayurbhanj, Jh

				Jharkhand (East Singhbhum), Ch (Durg, Dantewa Karnataka (Balla Kudremukh etc. Telangana, Andh Tamil Nadu, Goa
Manganese	Pyrolusite, Psilomelane, Rhodochrosite	Manganese, a critical industrial metal, is an essential component in the **manufacturing of steel, batteries, and various alloys.**	**Producers:** South Africa, Gabon, Ghana, Australia, Brazil; **Reserves:** South Africa, Australia, Brazil, China, Gabon.	**Reserver**: Odish Madhya Pradesh Maharashtra. **Producers:** • Odisha: near th Kendujhar, Sundergarh, Kor • Madhya Prade Balaghat-Chhin Nimar-Mandla a districts • Karnataka: Dha Belagavi, North Canara, Chikkm Shivamogga, Chitradurg and • Maharashtra: N Bhandara and Ratnagiri Minor Producers Goa, Jharkhand
Copper	Chalcopyrite, Bornite, Malachite, Cuprite	**Extensive applications** in industries ranging from **construction to electronics**, is a crucial resource in the global economy	**Producers:** Chile, DR Congo, Peru, China, Russia; **Reserves:** Chile, Australia, Peru, DR Congo, Russia.	**Reserves:** • Jhunjhunu, Kh districts in Rajasthan (52.2 deposits) • Balaghat distri Pradesh (23.28 per cent • Singhbhum dis Jharkhand (15.14 per cent) • Minor in Andh (Guntur), Karnataka (Chit & Tamil Nadu (South Arc in very small quantities Haryana, Maharashtra, Me Nagaland, Odisha, Sikkim, Uttarakhand, and West Benga **Production:** Madhya Pradesh Jharkhand
Bauxite (Aluminium)	Gibbsite, Boehmite, Diaspore	Crucial mineral supporting modern	**Producers:** Guinea, Australia, China, Brazil, India; **Reserves:** Guinea,	**Reserves:** Odisha, Andhra Gujarat,

		infrastructure, transportation, and various industrial applications	Australia, Indonesia, Jamaica, Brazil.	Jharkhand, Mah Chhattisgarh. Recently, the Ge of India (GSI) has identi deposits Bauxite deposit **Production:** • Odisha: Kalaha Bolangir, Koraput • Jharkhand: Lol Patlands • Gujarat: Bhavr • Chhattisgarh: plateau • Madhya Prade Katni–Jabalpur Balaghat • Maharashtra: K Ratnagiri, Satara Pune Minor Producer Karnataka, Goa
Gold	Native gold, Auriferous quartz veins, Placer deposits	Jewellery, Financial Systems, Electronics.	**Producers:** China, Russia, Australia, Canada, USA; **Reserves:** Australia, Russia, Indonesia, China, Canada.	**Reserves:** Bihar possibly very lar Chhattisgarh, Ra Pradesh; recent Odisha (Sundar Nabarangpur, K Deogarh). **Produ** Karnataka (Kola mines in Jharkh (U.P.), Jonnagiri
Silver	Argentite, Galena (by-product), Native silver	Silver's **industrial applications**, including solar panels, electronics, and medical devices, drive a significant portion of its demand.	**Producers:** Mexico, Peru, China, Bolivia, Chile; **Reserves:** Peru, Russia, Australia, China, Poland.	**Reserves:** limite deposits in Raja reserves in Karn Jharkhand, Andl **Producers:** Raja contributes abo production, esp mines (Udaipur) Agucha; also pr Andhra Pradesh Karnataka as a b zinc/lead/coppe
Uranium	Pitchblende (Uraninite), Carnotite	Uranium is a crucial element used primarily as a fuel in **nuclear reactors for electricity generation.**	**Producers:** Kazakhstan, Canada, Namibia, Australia, Uzbekistan; **Reserves:** Australia, Kazakhstan, Canada, Namibia, Russia.	**Potential reserv** (Udaipur, Alwar, Chhattisgarh (D Maharashtra (Bh Himachal Prade Hamirpur, Una); reserves in Andl Meghalaya, Karr **Producers / min** at Jaduguda, Jh

Thorium	Monazite (important in India), Thorite	Much cleaner than Uranium. More energy-efficient. Very crucial for Clean Energy Production. Also, U233 produced via Thorium is more fissile than U235.	**Production data:** not publicly available; **Reserves:** India, Brazil, USA, Australia, China.	**Reserves:** India of global deposi monazite depos and Kollam (Ker Visakhapatnam Mahanadi delta **Producer:** Kerala producer.
Tin	Cassiterite	Tin is a silvery-white metal revered for its **high malleability and corrosion resistance**, embarks on a captivating journey across the globe. Its is **used to make alloys** like Bronze.	**Producers:** China, Indonesia, Peru, Brazil, DR Congo; **Reserves:** Indonesia, China, Brazil, Myanmar.	**Reserves:** Chha (Dandkaranya); deposits in Odis **Producer:** Chha (Dandkaranya).
Lead	Galena	Used in batteries, radiation shielding, ammunition, cable sheathing, and some construction materials.	**Producers:** China, Australia, Peru, USA, Russia; **Reserves:** Australia, China, Russia, Mexico, Peru.	**Reserves:** prima also Andhra Pra Madhya Pradesh in Bihar, Mahara Meghalaya, Odis Tamil Nadu, Utta Bengal. **Produce** the largest lead- state.
Zinc	Sphalerite (Zinc blende), Smithsonite	Primarily used in **galvanization**, where it protects iron and steel from corrosion. Key **component in alloys like brass** and is essential for making	**Producers:** China, Peru, Australia, India, Mexico; **Reserves:** Australia, China, Russia, Peru, Mexico.	**Reserves:** Gujar Odisha, Sikkim, Uttarakhand, W **Producers:** mair HZL mines—Ran Sindesar Khurd, Zawar, Kayad; o include Andhra Madhya Pradesh Maharashtra.

		batteries, fertilizers, paints, and pharmaceuticals		
Nickel	Pentlandite, Garnierite	Key ingredient in stainless steel and rechargeable batteries.	**Producers:** Indonesia, Philippines, Russia, Canada, New Caledonia; **Reserves:** Indonesia, Australia, Brazil, Russia, New Caledonia.	**Reserves:** about reserves; mainly (Sukinda Valley, Mayurbhanj, Ked Jharkhand (Sing Nagaland (Kiphi associated with deposits at Jadu **Producer:** no ac
Cobalt	Cobaltite, Smaltite	Powering our electronics and enabling the clean energy transition.	**Producers:** DR Congo, Indonesia, Russia, Madagascar, Philippines, Australia; **Reserves:** DR Congo, Australia, Russia, Indonesia, Philippines.	**Potential reserv** (Jharkhand), Ke Jajpur (Odisha), (Rajasthan), Tue (Nagaland), Jha Hoshangabad (M no domestic pro currently.
Lithium	Spodumene, Lepidolite, Petalite, Brines	Powering our electronics and enabling the clean energy transition.	**Producers:** Australia, China, Chile, Argentina, Brazil; **Reserves:** Chile, Australia, China, USA. The text snippet repeats Australia once; I have removed the duplicate in the final rendering.	**Reserves:** Jamm Rajasthan, Karna Yadgiri); recent Meghalaya and **Producer:** no op currently; first m Korba, Chhattisg
Titanium	Ilmenite, Rutile	Used in aircraft, spacecraft, medical implants, paints, and corrosion-resistant equipment due to its light weight and strength.	**Producers:** China, Japan, Russia, Kazakhstan, Saudi Arabia; **Reserves:** Australia, China, South Africa, Canada, Mozambique.	**Indian occurren** Kerala, Tamil Na Pradesh, Odisha rutile, brookite, **Production:** Ker Odisha
Rare Earth Elements	Monazite, Bastnäsite, Xenotime	After the table	**Producers:** China, USA, Australia, Myanmar, Thailand; **Reserves:** China, Brazil, Australia, Russia, Vietnam, Greenland (Denmark).	**Reserves:** about tonnes of in-situ resource; 5th la deposits; coasta in Kerala, Tamil I Andhra Pradesh Gujarat, and inla Uttar Pradesh, J Bengal, Tamil Na Odisha, Kerala,

				Andhra Pradesh Gujarat.
Mica	Muscovite, Biotite, Phlogopite	Mica is a group of silicate minerals prized for their unique properties like flexibility, and heat resistance, which play a vital role in various industries. From **electronics and construction to cosmetics and pharmaceuticals**.	**Producers:** India, China, Russia, Brazil, USA; **Reserves:** India, China, Republic of Korea, Turkey.	**Reserves / occu** Jharkhand (Haz Koderma), Andh (Nellore), Rajast (Jaipur–Bhilwar Karnataka (Mysu Tamil Nadu (Co Tiruchirapalli, M Kanyakumari), K Maharashtra, W **Top Producer:** J
Tantalum	Columbite-Tantalite (Coltan)	Used for **electronics & semiconductors**, from computers to ACs (for making capacitors mainly)	**Top Producers** Democratic Republic of Congo, Nigeria, Rwanda, Brazil & China **Top Reserves** Australia, China, Brazil, Russia. For the rest of the countries, data is not available	Potential found sands in Punjab. No Production i
Silicon	Quartz, Silica sand	Widely used in **electronics** (as a semiconductor in chips and solar cells) and in **construction** (as silica in glass, cement, and ceramics).It is also crucial in **metallurgy (ferrosilicon), silicones (lubricants, sealants)**, and in **cosmetics and medical implants**	**Top producers:** Silicon metal-China, Brazil, Norway; Ferrosilicate-China, Russia, Norway, Brazil; **Reserves:** abundantly available in most countries.	**Reserves / sour** from quartz; sig quartzite/quartz Haryana, Rajasth Pradesh, Tamil N Maharashtra, Jh Pradesh. **Produc** Rajasthan, Andh

Tungsten	Wolframite, Scheelite	**Dense, hard metal** known for its high **melting point**, which is the **highest of all metals.** **Production of hard materials**, including tungsten carbide, used for cutting tools and wear-resistant applications. **Used in the aerospace and electronics industries**, including for filaments in light bulbs and electrodes	**Producers:** China, Vietnam, Kazakhstan, Russia, North Korea; **Reserves:** China, Australia, Russia, Vietnam, Spain.	**Reserves:** Karna Andhra Pradesh smaller resource Tamil Nadu, Utta Bengal; notable Degana, Rajasth Rajasthan and K
Antimony	Stibnite	Used to create many compounds and alloys, which are then used to make flame-retardant materials, paints, enamels, glass, and pottery	**Top Producers:** China, Russia, Tajikistan, Australia, Bolivia **Top Reserves:** China, Russia, Bolivia, Kyrgyzstan, Myanmar	Minor deposits i Spiti district of HP and Damo Complete impo
Gallium	By-product of Bauxite (Aluminum ores)	**Soft, silvery-white strategic metal** predominantly used in electronics	**Top Producers:** China (>90 per cent), Russia, Japan, South Korea, Ukraine **Top Reserves**: Data on Global reserves not available in public domain	NALCO planning gallium extracti plant at its Dam refinery. Used to be proc Hindalco Industr and National Alu (NALCO), but not anymore.
Germanium	By-product of Zinc ores (Sphalerite)	**Electronics and solar applications**, **fiber-optic systems, infrared**	**Top Producers:** The producers do not publicly give data regarding their production. But the biggest producer is	No known reser cent import dep China, Germany & Sou

		optics, and polymerization catalysts. Other uses included chemotherapy, metallurgy, and phosphors.	China.Other producers include the USA, Belgium, Canada, Russia, South Africa, and Germany. **Top Reserves:** Data on Global reserves not available in public domain	
Tellurium	By-product of Copper refining	Used primarily in **thermoelectric devices**, solar panels (CdTe solar cells), and as an additive in **steel and copper alloys.** Also in **optical discs, semiconductors**, and **vulcanization of rubber**.	**Producers:** China, Russia, Japan, Sweden, Canada; **Reserves:** China, Russia, Japan, Sweden, Uzbekistan.	Both Tellurium a are rarely found in their na are typically extracted as byp other mining operations. **Occurrence / pr** as by-product c gold ores; Ghats smelter, Jharkha tellurium from D no recent produ cited.
Selenium	By-product of Copper refining	Used in **photocells, solar panels, and glass manufacturing** (to decolorize or tint glass). It is also important in **electronic rectifiers**, and in **agriculture and animal nutrition** as a micronutrient.	**Top Producers:** China, Japan, Russia, Belgium, Canada **Top Reserves:** China, Japan, Russia, Belgium, Mexico	
Indium	By-product of Zinc ores	Mainly used in **Indium Tin Oxide (ITO)** for making **touchscreens, LCDs, and solar panels** due to its transparency and conductivity. It is also used in	**Top Producers:** China, South Korea, Japan, Canada, France **Top Reserves:** Detailed Reserves data not available publicly	Complete impo NIL production.

		semiconductors, solders, and alloys that require low melting points and good thermal properties.		
Helium	Natural gas fields (radioactive decay origin)	Helium is used in **balloons/airships, cryogenic cooling of MRI machines, deep-sea diving gas mixtures, welding, leak detection, and as a protective gas in rockets and scientific instruments**.	**Top Producers:** USA, Qatar, Russia, Algeria, Australia, Canada **Top Reserves:** USA, Qatar, Algeria, Russia	No major reserv dependency
Diamonds	Kimberlite pipes, Alluvial deposits	Used in jewellery, cutting/drilling tools, abrasives, and precision instruments due to its hardness.	**Top Producers:** Russia, Botswana, Canada, Angola, Democratic Republic of Congo, South Africa **Top Reserves:** Russia, Botswana, Democratic Republic of Congo, Angola	Approximately 9 concentrated in Pradesh, specifi Panna belt. Othe diamond-bearir include Andhra (Krishna River b Chhattisgarh. Production only
Crude Oil	Sedimentary basins (organic origin)	Used as fuel, lubricants, asphalt, and raw material for petrochemicals like plastics, fibres and fertilizers.	**Top Producers:** USA, Saudi Arabia, Russia, Canada, China **Top Reserves:** Venezuela, Saudi Arabia, Canada, Iran, Iraq	Major reserves a centres: Mumba Rajasthan (Barm Gujarat (Camba (Assam-Arakan The Krishna-Go reserves, but no Recent discover Andaman Sea re evaluated.
Natural Gas	Associated & non-associated gas fields	Used for cooking, power generation, fertilizers, industries, and as CNG/PNG fuel.	**Top Producer:** USA, Russia, Iran, China, Canada **Top Reserves:** Russia, Iran, Qatar, USA, Saudi Arabia	Assam (Digboi a Rudrasagar), Mu Gujarat, Rajasth Basin. Reserves Andaman Sea a

- **Applications of REE:**

- **Electronics:** Rare earth metals are essential components in the production of electronics, including smartphones, computers, and TVs.
- **Magnets:** Neodymium magnets, which contain rare earth elements, are crucial for various applications, such as electric motors in electric vehicles and wind turbines.
- **Catalysis** : Some rare earth metals serve as catalysts in chemical reactions, impacting processes like refining petroleum and automobile emissions control.
- **Lighting** : Phosphors containing rare earth metals are used in the production of energy-efficient lighting, including fluorescent lamps and LED lights.
- **Medical Imaging** : Gadolinium is used in contrast agents for magnetic resonance imaging (MRI) in medical diagnostics.
- **Defense Technologies** : Rare earth elements are used in the manufacturing of components for defense technologies, including guided missiles and radar systems.

⚡ **Quick revise** Resource = availability + utility Fold ▲

Resource = availability + utility. Classes: **natural / human-made · renewable / non-renewable · biotic / abiotic · individual / community / national / international (high seas, outer space).**

Mining: a mine pays only if **extraction + purification cost < value of product. Open-cast** (near surface), **shaft** (deep).

Limestone

- Limestone, a sedimentary rock **composed primarily of calcium carbonate (CaCO3)** , is a ubiquitous and versatile resource with immense economic and geologic significance.
- From building materials and cement production to agriculture, soil amendment, and environmental applications, limestone underpins numerous industries and plays a crucial role in shaping landscapes and ecosystems worldwide.
- **Top Producers: China, India, USA, Russia, Brazil**
  - China: Hubei, Guangxi, and Shandong.
  - India: Rajasthan, Madhya Pradesh, Andhra Pradesh, and Gujarat
  - USA: Texas, Florida, and Kentucky
  - Russia: Caucasus Mountains and the Ural Mountains
  - Brazil: Minas Gerais and Bahia
- **India:**
  - **Rajasthan** : Jaisalmer, Jodhpur, Nagaur, and Bikaner
  - **MP** : Satna, Katni, and Damoh
  - **Andhra Pradesh** : Kurnool and Anantapur
  - **Gujarat** : Bhavnagar, Amreli, and Kachchh

Some Important Ores and their Products

S.No	Ore	Product	Prese
1	Apatite and Rock Phosphate	Phosphorous, Fluorine, Uranium (secondary product)	Apati Megh Rock
2	Borax and Sassolite	Boron	No e Small
3	Chromite	Chromium	>96% Megh P.
4	Cryolite	Fluorine, Sodium (very less)	Nil
5	Diatomite (Dead Diatom Deposits)	Silica	Gujar
6	Fluorite	Fluorine	GJ, R

⚡ **Quick revise** World resources and minerals Fold ▲

⛏️ **State-hotspot pegs: Iron = the Odisha–Jharkhand–Chhattisgarh–Karnataka–Goa belt · Copper = Rajasthan (Khetri) · Bauxite = Odisha · Gold = Kolar & Hutti (Karnataka) · Diamond = Panna (MP) only · Mica = Jharkhand · Monazite/Thorium = Kerala · Uranium = Jaduguda (Jharkhand) · Lead–zinc–silver = Rajasthan (Zawar, Rampura Agucha) · Chromite = Odisha > 96%.**

Mineral	Where in India (handout)	Hooks
Iron ore	Odisha (Mayurbhanj, Keonjhar, Sundargarh); Jharkhand (Singhbhum); Chhattisgarh (Durg, Dantewada, **Bailadila**); Karnataka (Ballari, Hospet, Kudremukh); Goa, Maharashtra, Telangana, AP, TN	hematite, magnetite, limonite, siderite; **steel**
Manganese	Odisha, Karnataka, MP, Maharashtra	steel, batteries
Copper	**Rajasthan 52.25%** (Jhunjhunu, Khetri, Alwar) · **MP 23.28%** (Balaghat) · **Jharkhand 15.14%** (Singhbhum)	chalcopyrite; world leaders Chile, DR Congo, Peru
Bauxite	Odisha (Kalahandi, Koraput), Gujarat, Jharkhand (Lohardaga), Maharashtra, Chhattisgarh (Amarkantak), MP; new finds in Meghalaya	aluminium smelters sit near **power plants** (huge electricity)

Mineral	Where in India (handout)	Hooks
Gold	Karnataka (**Kolar, Hutti**), Jharkhand, Chhattisgarh, Rajasthan, AP (Jonnagiri); Bihar preliminary	Kolar to restart 24 yrs after closure
Silver	by-product; **Rajasthan ≈ 60%** (Zawar, Rampura Agucha)	lead–zinc by-product
Uranium	Jharkhand (UCIL **Jaduguda**); potential Rajasthan, Chhattisgarh, Maharashtra, HP, Andhra…	natural U contains **0.72% U-235**; **U-235 fissile, U-238 not**; enrichment raises concentration
Thorium (monazite)	**India ≈ 25% of global deposits**; Palakkad & Kollam (Kerala), Visakhapatnam, Mahanadi delta	cleaner; **U-233**; thorium reactors can use up to 99% of fuel
Tin	Chhattisgarh (Dandkaranya)	bronze
Lead / Zinc	**Rajasthan** largest lead and zinc state (HZL mines: Rampura Agucha, Sindesar Khurd, Rajpura Dariba, Zawar, Kayad)	zinc – galvanising; Zn–S batteries as lithium alternative
Nickel	Odisha (**Sukinda**), Jharkhand, Nagaland; ~0.2% of global reserves; **no active mine**	stainless steel, batteries
Cobalt	potential – Jharkhand, Odisha, Rajasthan, Nagaland, MP; **no domestic production**	
Lithium	J&K, Rajasthan, Karnataka (Mandya, Yadgiri); first mine expected in **Korba, Chhattisgarh**; **no operating mine**	
Titanium	Kerala, TN, AP, Odisha (**ilmenite, rutile**)	**2023 PYQ**
Rare earths	**monazite, bastnäsite, xenotime**; coastal placers in Kerala, TN, Odisha, AP; **13.07 million t monazite (5th largest)**	**17 elements = 15 lanthanides + scandium + yttrium**; China's near-monopoly → **Minerals Security Partnership**
Mica	**Jharkhand** (Hazaribagh, Koderma), AP (Nellore), Rajasthan	**Feb 2025: classified Major Mineral** with feldspar, barytes, quartz
Tantalum, silicon	no production; silicon from quartz in Haryana, Rajasthan, AP	
Tungsten	Karnataka, Rajasthan (Degana)	**Madurai mining scrapped after protest**
Antimony, germanium, indium, helium	**~100% import dependent** (antimony "complete"; germanium 100%; indium complete; helium 100%)	China limited exports (antimony – July 2024; gallium – July 2023)
Gallium, tellurium, selenium	by-products of bauxite/copper	ALCO plans 10 tpy gallium plant at Damanjodi

Mineral	Where in India (handout)	Hooks
Diamond	**~90% Panna (MP)**; AP (Krishna basin), Chhattisgarh	**production only in Panna**
Limestone	Rajasthan, MP (Satna, Katni), AP, Gujarat	world: China, India, USA, Russia, Brazil
Others	**Chromite** (> 96% Odisha) · **Pyrite** (no production since 2003) · **Borax** (small deposits, Puga valley) · **Cryolite** (nil)	

Coal

7	Ilmenite, Rutile, Brookite, Leucoxene	Titanium	Kerala Odish
8	Kyanite, Sillimanite and Andalusite	Aluminium	Kyani OD, reser
9	Magnesite	Magnesium	UK, T
10	Monazite	Thorium, Lanthanum, Cerium	130 c place Odish Maha inlanc Jhark Nadu
11	Pyrite	Iron, Sulphur	No pr
12	Vermiculite	Similar to Mica	TN, A
13	Zircon	Zirconium	State Occu Rutile

Coal

- Coal, a cornerstone of the global energy sector for centuries, plays a crucial role in powering industries and generating electricity.



- Coal is primarily **formed from the remains of plants** that lived and died millions of years ago by the **process called Coalification.**
- **Indian coal is of non-coking grade** (non-metallurgical coal, has higher impurities, so cannot be converted into coke).
- **Top Producers: China, India, USA, Australia, Indonesia.**
  - China: Shanxi, Inner Mongolia (mainly Anthracite)
  - India
  - USA: Appalachians (mainly Anthracite)
  - Australia: Queensland, New South Wales
  - Indonesia: Kalimantan, Sumatra
- **India:**
  - Production less than requirement. 75% of electricity produced via coal based thermal power plants.
  - **Gondwana** (**Bituminous)** : **River Valleys** (Damodar, Mahanadi, Godavari, Son)
    - **Jharkhand-Bengal Coal Belt** : Raniganj, Jharia, Bokaro, Giridih, Karanpura.
    - **Madhya Pradesh** : Singrauli (part in UP, but CIL plant in MP)
    - Korba in **Chhattisgarh**
    - Talcher and Rampur in **Odisha**
    - Chanda–Wardha, Kamptee and Bander in **Maharashtra**
    - Singareni in Telangana and Pandur in **Andhra Pradesh** .
  - **Tertiary Coal** (Can be Anthracite, bituminous, sub-bituminous or lignite):
    - **Anthracite** : **Kalakot,** JnK
    - **Sub-Bituminous:**
- **Assam** (Makum, Jaipur and Nazira)
- **Arunachal Pradesh** (Namchik – Namphuk)
- **Meghalaya** (Darangiri, Cherrapunji, Mewlong and Langrin)

- **Nagaland** (minor rat-hole mining allowed despite a ban by the supreme court, under article 371A)
    - **Lignite: Tamil Nadu** (Neyveli), **Gujarat, Puducherry** .

Shale Oil And Gas

- Shale oil (tight oil) and shale gas, **unconventional oil and gas resources trapped within shale rock formations** (less permeable and difficult to drill through), have revolutionized the energy landscape in recent decades.
- These resources, previously inaccessible, are now **extracted through** innovative technologies like **hydraulic fracturing** , unlocking vast new energy reserves and impacting global energy markets significantly.
- **Top Producers** : **USA, China, Argentina, Canada, Russia**



- **India** : **Promising potential** in:
    - Cambay Basin
    - Ganga Valley Basin
    - Assam and Arakan Basin
    - KG Basin
    - Kaveri Basin
- **Problems related to shale oil & gas mining** : Shale oil and gas extraction, particularly through **hydraulic fracturing (fracking),** faces several environmental and public health concerns:
    - **Water Contamination** : Fracking uses large volumes of water mixed with chemicals, raising concerns about potential contamination of groundwater resources through spills or improper disposal of wastewater. The underground aquifers might also get contaminated.
    - **Induced Seismicity** : The process of injecting water underground during fracking can trigger earthquakes in some areas.

- **Land Disturbance** : Shale oil and gas development requires significant land use for well pads, pipelines, and infrastructure, disrupting ecosystems and impacting local communities.
- **Air and Climate Impacts** : Methane leaks during extraction and processing contribute to greenhouse gas emissions. Additionally, air pollution from drilling and associated activities can harm local air quality.

⚡ **Quick revise** Coal: from plant remains by Fold ▲

- **Coal:** from plant remains by **coalification**. **Indian coal is non-coking, high ash (up to 40–45%), low sulphur (~0.5–1%, ~0.7%), high ash-fusion temperature (~1,500 °C). ~75% of electricity comes from coal-based thermal plants.** World leaders: China (Shanxi, Inner Mongolia), India, USA, Australia, Indonesia.
- **Gondwana (bituminous):** river valleys – **Damodar, Mahanadi, Godavari, Son.** *Jharkhand–Bengal belt:* **Raniganj, Jharia, Bokaro, Giridih, Karanpura** · Singrauli (MP/UP) · **Korba (CG)** · Talcher, Rampur (Odisha) · Chanda–Wardha, Kamptee, Bander (Maharashtra) · Singareni (Telangana), Pandur (AP).
- **Tertiary coal:** anthracite – **Kalakot (J&K)**; sub-bituminous – Assam (Makum, Jaipur, Nazira), Arunachal (Namchik–Namphuk), Meghalaya (Darangiri, Cherrapunji, Mewlong, Langrin), Nagaland (rat-hole mining, Art. 371A); **lignite – Neyveli (TN), Gujarat, Puducherry.**
- **Why does India import coal?** *(2012 PYQ, ans B)* – power plants cannot get enough domestic coal **and** steel needs **coking coal**; **not** a policy to "save reserves".
- **Crude oil (India): Mumbai High**, Barmer (Rajasthan), Cambay (Gujarat), **Assam–Arakan**; **KG belt has reserves but no production**; new find in the Andaman Sea region (being evaluated). **Natural gas:** Assam (**Digboi, Rudrasagar**), Mumbai High, Gujarat, Rajasthan, Tripura, KG; also Andaman Sea.
- **Shale oil and gas:** unconventional, fracking; **USA, China, Argentina, Canada, Russia**. India's basins: **Cambay, Ganga valley, Assam–Arakan, KG, Kaveri** (handout PYQ lists **Cambay, Gondwana, KG, Cauvery onshore, Indo-Gangetic**). **Problems:** water contamination, **induced seismicity**, land disturbance, methane leaks.

Non-Conventional Energy Resources

- **India's total Renewable Energy Generation: 283.46 GW** upto march 2026

Solar Energy

- **Thermal & Electric Energy** both can be **harnessed** from it.
- **Top Producers** : **China, USA, India, Japan, Germany**
- India being a tropical nation has a huge solar energy potential.

- **India: target of 280 GW capacity by 2030**
- **By May 2026, total installed capacity 157 GW**
- **Top Producers: Rajasthan, Gujarat, Maharashtra, Tamil Nadu, Karnataka**
- **Problems of covering sub-tropical deserts with Solar Panels:**
  - Covering just 1.2% of Sahara desert can produce enough energy to the world. But why is this not being done?
  - **Solar panels** , which are darker in colour compared to the sands, **may absorb more heat** -> increase in temperature -> changes in rainfall and wind patterns in surrounding areas.
  - **High Logistic costs**
  - **High maintenance costs** (sands will cover the solar panels)

Wind Energy

- **Onshore and Offshore**
- **In offshore locations,** due to the **lack of frictional forces, the speed of the winds is higher** . However, the cost of these plants with the current technology nullifies the increased electricity production due to higher wind speeds.
- **Top Producers: China, USA, India, Germany, Brazil**
- **India installed:** 56 GW.
- **Top Producers in India: Gujarat, Tamil Nadu, Karnataka, Rajasthan, Maharashtra**
- **No operational offshore** wind farm in India.

Bio-Energy

- **Waste to Energy:** Waste to Energy (WtE) is the process of generating energy in the form of electricity or heat from the primary treatment of waste. It typically involves incineration, where municipal solid waste is burned to produce steam that drives turbines to generate electricity.
- **Biogas:** Biogas is produced through the anaerobic digestion of organic materials such as agricultural waste, manure, and food scraps. The process results in a mixture of methane and carbon dioxide, which can be used as a renewable energy source for heating, electricity, and as a vehicle fuel.
- **Top Producers: USA, Brazil, Indonesia, China, India**
- **In India: Maharashtra, Karnataka, Uttar Pradesh, Punjab, and Bihar**
- India total production: 10.75 GW

Geothermal Energy

- **USA, Indonesia, Philippines, Turkiye, New Zealand.**



- Indian potential: 10600 MW
- **India plans to harness 10 GW by 2030 especially at Manikaran, HP.**
- **Currently not tapped.**
- **Singareni Collieries Company Limited** (SCCL) has commissioned a **20 kW pilot geothermal power plant in Manuguru area of Bhadradri Kothagudem district in Telangana.**

Tidal Energy

- **Top Producers: S. Korea, France, Canada, China, Russia**
- **In India:** We had **started building** plants in Gujarat (50MW in Kachchh) and West Bengal (3.75 MW at Durgaduani Creek). **But stopped due to exorbitant prices.**

Otec (Ocean Thermal Energy Conversion)

- **Only 2 global operating (pilot-sized) plants:** Japan (Okinawa) & USA (Makai, Hawaii)
- **In India, a 65-kW plant at Kavaratti, Lakshadweep** , to power a desalination plant.

Nuclear Energy



- Includes **Fusion and Fission.**

Currently, only Fission is used for energy generation by humans.

- **Top Producers (2021): USA, China, France, Russia, South Korea.**
- According to Government of India, the present installed nuclear power capacity is set to increase from 8180 MW to 22480 MW by 2031-32.
- There has been a **70% Surge in India's Nuclear Power Capacity in the last 10 years** .
- **3 stages of India’s Nuclear Power Program** :
  - **Natural uranium** fuelled Pressurized Heavy Water Reactors (PHWRs) (U235)
  - **Fast Breeder Reactors** (FBRs) **utilizing plutonium** -based fuel. (Pu239): Kalpakkam, TN (achieved criticality)
  - **Advanced nuclear power systems for utilization of thorium** . (U233 from Thorium): KAMINI-Kalpakkam Mini Reactor, Kalpakkam, TN (experimental)
- Map shows the locations of stage 1 reactors in India:

⚡ **Quick revise** Non-conventional energy Fold ▲

Source	India facts	Notes
All renewables	283.46 GW up to March 2026	
Solar	157 GW by May 2026; target 280 GW by 2030; top: Rajasthan, Gujarat, Maharashtra, TN, Karnataka	desert panels warm the area (darker than sand), logistics + maintenance costs, sand cover

Source	India facts	Notes
Wind	**56 GW**; top: **Gujarat, TN, Karnataka, Rajasthan, Maharashtra**; **no offshore farm**	offshore winds faster (no friction) but cost cancels the gain
Bio-energy	**10.75 GW**; Maharashtra, Karnataka, UP, Punjab, Bihar	waste-to-energy, biogas
Geothermal	potential **10,600 MW**; **Manikaran (HP)**; target 10 GW by 2030; **SCCL 20 kW pilot, Manuguru (Telangana)**	not yet tapped
Tidal	Kachchh (Gujarat) 50 MW and **Durgaduani Creek (WB) 3.75 MW** started, **stopped due to cost**	
OTEC	**only 2 pilot plants in the world (Okinawa, Makai–Hawaii)**; **65 kW at Kavaratti (Lakshadweep)** for desalination	
Nuclear	**8,180 MW → 22,480 MW by 2031-32**; **+70% in 10 years**	**3 stages:** (1) **PHWR**, natural uranium (U-235) → (2) **Fast Breeder, plutonium (Pu-239) – Kalpakkam** → (3) **thorium, U-233 – KAMINI**, Kalpakkam

⚠️ **Source conflict (handout vs mind map):** the Economic-Geography **mind map** prints older figures – **renewables 220.10 GW** and **solar 105.65 GW (31 Mar 2025, MNRE)** – while the handout text says **283.46 GW (Mar 2026)** and **157 GW (May 2026)**. *Rule: use the handout's newer numbers; the 280 GW-by-2030 target is the same in both.*

🔋 **Nuclear 3 stages = "U → Pu → Th":** *Use uranium, Produce plutonium, Then thorium.* All three are Kalpakkam-linked after stage 1.

3 · PYQ

Q.) Despite having large reserves of coal, why does India import millions of tonnes of coal ?

1. It is the policy of India to save its own coal reserves for the future, and import it from other countries for the present use.

2. Most of the power plants in India are coal-based and they are not able to get sufficient supplies of coal from within the country.

3. Steel companies need a large quantity of coking coal which has to be imported.

Which of the statements given above is/are correct? (2012)

(a) 1 Only

(b) 2 & 3 Only

(c) 1 & 3 Only

(d) 1,2, & 3

Ans. B Coal is and will continue to be the backbone of India's economic and development growth. India has the world's fifth largest coal reserves and is also the world's third largest producer and importer of coal.

- India's total thermal coal imports in 2021 may have edged higher from 156.09mn t in 2020. India imported 116.57mn t of thermal coal in January-September 2021, according to customs data, and going by the average fourth-quarter imports in the last five years, the receipts could have been slightly over 160mn t in 2021.
- The lack of coking coal reserves that are used as a raw material in steelmaking and allied industries is the main reason India imports millions of tonnes of coal. Coal in India is used either from domestic sources, mostly mined by coal India, or is imported.
- The issue for India is that the gap between domestic supply and demand is growing at a phenomenal rate.This gap is being filled by imported coal.

Q.) Consider the following statements:

1. Natural gas occurs in the Gondwana beds.

2. Mica occurs in abundance in Kodarma.

3. Dharwars are famous for petroleum.

Which of the statements given above is/are correct? (2013)

(a) 1 only

(b) 2 only

(c) 2 and 3 only

(d) None

Ans. B

- KG basin, Assam, Gulf of Khambhat, Cuddalore district of Tamil Nadu, Barmer in Rajasthan etc. are the areas where natural gas can be found.Nearly 80% of the world's total proven natural gas reserves are located in ten countries. Russia tops the list, holding about a quarter of the world's total gas reserves, followed by Iran and Qatar in the Middle East.
- 4. Kodarma mines in Jharkhand are one of the important mica producing areas. Kodarma district of Jharkhand state has been famous worldwide for mica

mining,especially for ruby mica. The district had the monopoly in producing mica for years and themica was exported to many countries worldwide. The main reserve of mica is found under the forest of mild life sanctuary of Kodarma.

- 5. The Dharwar rocks are rich in iron ore, manganese, lead, zinc, gold, silver etc. The Champions series containing gold mines lie within this system. This Champion system is named after the Champion reef in the Kolar Gold Fields. The Kolar Gold Fields contain one of the deepest gold mines of world
- 6. The Dharwar rocks are rich in iron ore, manganese, lead, zinc, gold, silver etc. The Champions series containing gold mines lie within this system.

Q.) Which of the following is/are the characteristics/ characteristics of Indian coal?

1. High ash content

2. Low sulphur content

3. The low ash fusion temperature

Select the correct answer using the codes given below. (2013)

(a) 1 only

(b) 1 and 2 only

(c) 2 and 3 only

(d) 1, 2 and 3

Ans. B Indian coals like any other Gondwana coals are of high ash content.

- Indian coals by their very nature are high in ash content but low in sulphur content. Indian coal has a high ash fusion temperature of about 1,500°C.
- India's domestic coal reserves have high ash content—up to 40 to 45 percent. As a result more coal is required to generate one kilowatt hour of electricity. This in turn means higher emissions.
- Sulphur content of Indian coal is, in general, around 0.5 – 1%. Studies show that the sulphur content in Indian coal is hardly 0.7%, which is not uniformly distributed in the coal but concentrated in the pyritic bands which are visible to the naked eye. The best solution to get rid of sulphur from Indian coal is to wash it
- Ash fusion temperature of Indian coal is generally high.

Q.) In which of the following regions of India are shale gas resources found ?

1. Cambay Basin

2. Cauvery basin

3. Krishna-Godavari Basin

Select the correct answer using the code given below: (2016)

(a) 1 and 2 only

(b) 3 only

(c) 2 and 3 only

(d) 1, 2 and 3

Ans. D To strengthen its hydrocarbon resource base, India has identified six basins as areas for shale gas exploration: Cambay (Gujarat), Assam-Arakan (North East), Gondwana (Central India), Krishna Godavari onshore (East Coast), Cauvery onshore, and Indo-Gangetic basins.

- Shale Gas can emerge as an important new source of energy in the country. India has several Shale formations which seem to hold Shale Gas and oil. The Shale Gas formations are spread over several sedimentary basins such as Cambay, Gondwana, Krishna-Godavari and Kaveri on land.
- Shale gas is an unconventional source of energy found in non-porous rocks. It has become an important source of natural gas in the US.

Q.) Consider the following statements:

1. In India, State Governments do not have the power to auction non-coal mines.

2. Andhra Pradesh and Jharkhand do not have gold mines.

3. Rajasthan has iron ore mines.

Which of the statements given above is/are correct? (2018)

(a) 1 and 2

(b) 2 only

(c) 3 only

(d) 1 and 3

Ans. C

- Statement 1 is incorrect: State governments can auction Non-coal mines.
- Statement 2 is incorrect: Both Jharkhand and Andhra Pradesh have gold mines.
- Statement 3 is correct: Bhilwara in Rajasthan has an iron ore mine.
- In India, State Governments do not have the power to auction non-coal mines. As per the new mining law — Mines and Minerals (Development and Regulation) Amendment Act, 2015 — which came into effect from January 2015, the non-coal mines have to be auctioned by the respective state governments. The first ever non-coal auction in the country began in December, 2015.
- Andhra Pradesh and Jharkhand do not have gold mines. Rajasthan has iron ore mines.
- Rajasthan is the richest state in terms of availability and variety of minerals in the country and produces about 57 different minerals. Rajasthan is the sole producer of lead & zinc ores, selenite and wollastonite.
- In the State copper deposits are located in the districts of Jhunjhunu, Sikar, Sirohi, Udaipur, Bhilwara, Ajmer, Alwar, Bharatpur, Chittorgarh, Rajsamand and Dungarpur. With resources of 809.09 million tonnes, the State possesses a substantial share of 54% of the total resources of the country. The State is the second leading producer of copper concentrate contributing 41% of the nation’s output.

Q.) With reference to India, consider the following statements?

1. Monazite is a source of rare earths.

2. Monazite contains thorium.

3. Monazite occurs naturally in the entire Indian coastal sands in India.

4. In India, Government bodies only can process or export monazite.

Which of the statements given above are correct? (2022)

(a) 1, 2 and 3 only

(b) 1, 2 and 4 only

(c) 3 and 4 only

(d) 1, 2, 3 and 4

Ans. (b)

- Monazite is a mineral mainly containing rare earths and thorium - a prescribed substance to be handled by the Department of Atomic Energy (DAE).
- Accordingly, Indian Rare Earths Ltd. (IREL) which is wholly owned by the Govt. of India, under the administrative control of the Dept. of Atomic Energy (DAE), utilises monazite mainly for production of rare earth compounds, and thorium, as needed in the Department of Atomic Energy.
- In-situ monazite established by AMD so far are present in Odisha, Andhra Pradesh, Tamil Nadu, Kerala, West Bengal, and Jharkhand.

Q.) Ilmenite and rutile, abundantly available in certain coastal tracts of India, are rich sources of which one of the following? (2023)

(a) Aluminium

(b) Copper

(c) Iron

(d) Titanium

Ans: (d) Ilmenite is represented chemically as FeO.TiO2 and rutile as TiO2. They are both ores of Titanium.

Q.) Consider the following statements: Statement-I: India, despite having uranium deposits, depends on coal for most its electricity production. **Statement-II:** Uranium, enriched to the extent at least 60%, is required for the production of electricity.

Which one of the following is correct in respect of the above statements? (2023)

(a) Both Statement-I and Statement-II are correct and Statement-II is the correct explanation for Statement-I

(b) Both Statement-I and Statement-II are correct and Statement-II is not the correct explanation for Statement-1

(c) Statement-I is correct but Statement-II is incorrect

(d) Statement-I is incorrect but Statement-II is correct

Ans: C

- India gets around 75% of its power from thermal power plants.
- Uranium enriched to 5% is enough for nuclear power production.

Q.) Which of the following statements about Rare Earth Elements (REEs) and Critical Minerals is/are correct?

1. Modern technological innovations including Artificial Intelligence, robotics and space exploration extensively utilise Rare Earth Elements (REEs).

2. China has the highest share in mining of REEs followed by India.

3. The Government of India launched the National Critical Mineral Mission (NCMM) in 2025 to establish a robust framework for self-reliance in the critical mineral sector.

4. Rare Earth Elements are a set of 13 metallic elements.

Select the answer using the code given below:

(a) 1 and 3 only

(b) 3 only

(c) 1, 3 and 4

(d) 1, 2 and 4

Ans: (a)

Explanation: India isnt 2nd highest. REE=17 elements.

⚡ **Quick revise** Minerals and energy PYQs Fold ▲

Year	Asked	Answer / rule
2012	Why does India import coal?	**Power plants short + steel needs coking coal (B)**
2013	Natural gas in Gondwana beds ✘ · Mica abundant in Kodarma ✔ · Dharwars famous for petroleum ✘	**B (2 only)** – Dharwars: iron, manganese, lead, zinc, gold (**Kolar**)
2013	Indian coal: high ash ✔, low sulphur ✔, **low** ash-fusion ✘	**B (1, 2)**
2016	Shale gas in Cambay, Cauvery, KG	**All three (D)**
2018	States cannot auction non-coal mines ✘ · AP & Jharkhand have no gold ✘ · Rajasthan has iron ore ✔	**C (3 only)** – Bhilwara has iron ore
2022	Monazite: rare earths ✔, thorium ✔, entire coast ✘, only government bodies process/export ✔	**B (1, 2, 4) – IREL, under DAE**
2023	Ilmenite and rutile are ores of	**Titanium (D)**
2023	India depends on coal despite uranium (✔); uranium "enriched ≥ 60%" for power (✘)	**C – ~5% enrichment is enough**

Year	Asked	Answer / rule
–	REEs: AI/robotics use ✔; China first, **India second ✘**; **NCMM 2025 ✔**; "REE are **13** elements" ✘	**A (1 and 3)** – REE = **17 elements**

⚠️ **Handout alert (2018 mines PYQ):** the explanation prints both *"State governments can auction non-coal mines"* and *"State governments do not have the power to auction"* and says AP/Jharkhand "do not have gold mines" while the table lists Jharkhand and AP among gold states. **Take the printed answer (C) and the gold table.**

4 · INDUSTRIES

Types of Industries:

- **Primary Industries:** These industries involve the extraction and harvesting of natural resources.
  - **Agriculture:** Farming, horticulture, and animal husbandry.
  - **Fishing:** Commercial fishing and aquaculture.
  - **Forestry:** Logging, timber production, and forest management.
  - **Mining and Quarrying:** Extraction of minerals, ores, and fossil fuels.
- **Secondary Industries:** These industries involve the processing and manufacturing of raw materials into finished goods.
  - **Manufacturing:** Production of goods using raw materials. It includes:
  - **Light Industry:** Consumer goods like clothing, electronics, and household items.
  - **Heavy Industry:** Capital goods like machinery, steel production, and shipbuilding.
  - **Construction:** Building infrastructure such as roads, bridges, buildings, and other structures.
- **Tertiary Industries (Service Sector):** These industries provide services rather than goods.
  - **Retail and Wholesale Trade:** Selling goods directly to consumers and businesses.
  - **Hospitality and Tourism:** Hotels, restaurants, travel agencies, and tour operators.
  - **Transport and Logistics:** Transportation services (road, rail, air, sea), warehousing, and logistics management.
  - **Financial Services:** Banking, insurance, investment, and real estate.
  - **Healthcare:** Hospitals, clinics, pharmaceuticals, and medical services.
  - **Education:** Schools, colleges, universities, and vocational training.
  - **Information Technology:** Software development, IT services, and telecommunications.
  - **Professional Services:** Legal, accounting, consulting, and advertising.

- **Quaternary Industries**
  - These industries involve the **intellectual activities** often associated with technological innovation and knowledge-based services.
  - **Research and Development (R&D):** Scientific research and technological innovation.
  - **Information Services:** Data processing, information management, and media.
  - **Education and Training:** Advanced education and specialized training programs.
- **Quinary Industries**
  - These industries include the **highest levels of decision-making and societal advancement** .
  - **Government:** Public administration and defense.
  - **Nonprofit Organizations:** Charities, NGOs, and other community services.
  - **Culture and Arts:** Museums, theaters, and cultural heritage management.

⚡ **Quick revise** Industries — Fold ▲

Sector	What	Examples
Primary	extraction/harvest	agriculture, fishing, forestry, mining
Secondary	processing/manufacture	light (consumer goods), heavy (capital goods – steel, ships), construction
Tertiary	services	trade, tourism, transport, finance, health, education, IT, professional
Quaternary	knowledge	R&D, information services, advanced training
Quinary	top decisions	government, NGOs, culture and arts

🏭 **P-S-T-Q-Q: Dig → Make → Serve → Think → Decide.**

Other Classifications of Industries

- **Based on Size:**
- **Micro:** Very small businesses, often family-run. **Investment <2.5 crore and turnover <10 crore** .
- **Small:** Small-scale enterprises with limited employees and capital. **Investment between 2.5-25 crore; turnover between 10-100 crore.**
- **Medium:** Mid-sized businesses with moderate resources. **Investment between 25-125 crore; turnover between 100-500 crore.**
- **Large:** Large corporations with **substantial resources and workforce** .
- **Based on Ownership:**
- **Public Sector:** Owned and operated by the government.
- **Private Sector:** Owned and operated by private individuals or companies.

- **Joint Sector:** Collaboration between government and private entities.
- **Cooperative Sector:** Owned and operated by a group of individuals for mutual benefit.
- **Based on Market Orientation:**
- **Consumer Goods:** Products sold directly to consumers (e.g., food, clothing, electronics).
- **Capital Goods:** Products used in the production of other goods (e.g., machinery, tools, equipment).

⚡ **Quick revise** By size (investment / turnover) Fold ▲

By size (investment / turnover): Micro < ₹2.5 cr / < ₹10 cr · Small ₹2.5–25 cr / ₹10–100 cr · Medium ₹25–125 cr / ₹100–500 cr · Large above. **By ownership:** public, private, joint, cooperative. **By use:** consumer goods vs capital goods.

Factors Affecting the Location of Industries:

The location of industries depends on various factors that influence their operational efficiency, cost-effectiveness, and overall success.

- **Raw Materials**
  - **Proximity to Raw Materials:** Industries that rely heavily on raw materials, such as mining and agriculture-based industries, often locate near their sources to minimize transportation costs. On this basis, industries can be called:
    - **Weight losing:** Raw materials bulky, and finished product lighter
    - **Weight gaining:** Raw materials lighter, and finished products bulky.
    - **Footloose:** No dependency on the weight of the raw materials or the finished product.
  - **Availability and Quality of Raw material:** The type, quality, and availability of raw materials can significantly impact the location decision.
- **Labor**
  - **Availability of Labor:** Access to a sufficient and skilled labor force is crucial for industries requiring intensive manual or specialized labor.
  - **Labor Costs:** Regions with lower labor costs may attract labor-intensive industries.
  - **Labor Laws and Conditions:** Favorable labor laws and conditions can also influence industrial location.
- **Market Access**
  - **Proximity to Markets:** Industries producing perishable goods or those with high transportation costs often locate near their primary markets to reduce distribution expenses.
  - **Market Size and Demand:** Larger markets with higher demand can attract industries looking to maximize their sales and distribution.
- **Transport and Infrastructure**

- **Transportation Facilities:** Good connectivity through roads, railways, ports, and airports facilitates the efficient movement of raw materials and finished goods.
- **Infrastructure Availability:** Adequate infrastructure, including power supply, water supply, and communication networks, is essential for industrial operations.

- **Energy Supply**
  - **Access to Power:** Reliable and affordable energy sources are crucial for energy-intensive industries like manufacturing and mining.
  - **Types of Energy:** The availability of specific types of energy, such as electricity, natural gas, or renewable sources, can influence industrial location.
- **Capital and Financial Factors**
  - **Access to Capital:** Proximity to financial institutions and access to credit can impact the decision on where to locate an industry.
  - **Investment Incentives:** Government incentives, such as tax breaks and subsidies, can attract industries to specific regions.
- **Government Policies and Regulations**
  - **Industrial Policies:** Government policies, including zoning laws, environmental regulations, and industrial promotion policies, play a significant role in industrial location decisions.
  - **Taxation:** Favorable tax policies and lower tax rates can attract industries to particular areas.
- **Climate and Environmental policies**
  - **Climate Suitability:** Certain industries, such as agriculture and tourism, depend on specific climatic conditions.
  - **Environmental Regulations:** Stringent environmental regulations can deter industries from locating in certain regions.
  - **Circular Economy Push** : The push for sustainability forces new industries to co-locate near major recycling centers to utilize secondary raw materials (urban mining).
- **Social and Cultural Factors**
  - **Community Attitudes:** Local community support or opposition can influence industrial location.
  - **Quality of Life:** Areas offering a higher quality of life, including good housing, education, and healthcare, may attract industries that value employee well-being.
- **Availability of Technology and Innovation**
  - **Access to Technology:** Proximity to technology hubs and research institutions can benefit high-tech industries.
  - **Innovation Ecosystem:** A robust ecosystem that supports innovation and research and development can attract industries looking to stay competitive.
  - **Reshoring:** Advanced robotics negates the need for cheap human labor. This causes industries to "reshore" back to high-income countries closer to the final consumer.
- **Global and International Factors**
  - **Trade Agreements:** International trade agreements and export incentives can influence the location of industries targeting global markets.

- **Political Stability:** Regions with stable political environments are more attractive to industries, reducing the risk of disruptions.
- **Nearshoring and Friendshoring:** As the cost of production in a particular country increases, the industry tries to relocate to a nearby country, so that the markets are still nearby. This is called nearshoring. If two countries sign FTAs, etc., then companies of one country are free to establish their plants in the other country, if the costs of production increase in the former (friend-shoring)

- **Ecological and Sustainability Considerations**
  - **Sustainability Practices:** Industries increasingly consider ecological footprints and sustainability practices, favoring locations that support green initiatives and offer renewable resources.
- **Weight-Gaining and Weight-Losing Industries:**
  - Weber classified industries into two main types: weight-gaining and weight-losing. **Weight-gaining** industries are those where the **final product is heavier than the raw materials** , such as manufacturing industries.
  - **Weight-losing** industries are those where the **final product is lighter than the raw materials** , such as mining or industries producing bulky raw materials.
  - The location of an industry depends on whether it is weight-gaining or weight-losing, as **transportation costs are influenced by the weight and bulk of goods** .
- **Isotim:**
  - An isotim is a **line that connects all locations that have the same transportation cost from a specific source of raw materials or market.**
  - It represents the boundaries of equal transport costs for a specific commodity.
  - Isotims form concentric circles around the source, with each circle representing a higher transportation cost.
- **Isodapane:**
  - **An isophane is a line that connects all locations that have the same total cost (transport cost for both raw materials and finished products + labour cost) for both raw materials and finished goods.**
  - It depicts the combined impact of transportation costs for both inputs and outputs on a firm's total cost.
  - The optimal location for an industry typically lies on the lowest-cost isodapane.

⚡ **Quick revise** Location factors — Fold ▲

Location factors: raw material · labour · market · transport/infrastructure · energy · capital · government policy · climate/environment (**circular-economy push**: co-locate near recycling centres, "urban mining") · social/cultural · technology (**reshoring** – robots replace cheap labour, plants return to high-income countries) · global factors (**nearshoring, friend-shoring**) · sustainability.

⚖️ **Weber's weight rule: weight-losing** (bulky raw material → light product, e.g., mining/ore) → **sit at the raw material. Weight-gaining** (product heavier/bulkier than inputs) → **sit near the**

market. Footloose = weight does not matter. **Isotim** = equal **transport cost** line (rings round a source). **Isodapane** = equal **total cost** line (transport of inputs + outputs + labour); the best site lies on the **lowest isodapane**.

Important Industrial Regions of the World

North America

- **North-Eastern United States:**
  - Regions: New York, New Jersey, Pennsylvania, Ohio, Illinois, and Michigan.
  - Industries: Steel, automobiles, chemicals, machinery, textiles, and electronics.
  - Key Cities: New York City, Chicago, Detroit, Pittsburgh, and Cleveland.
- **West Coast of the United States:**
  - Regions: California, Oregon, and Washington.
  - Industries: Technology, aerospace, entertainment, electronics, and biotechnology.
  - Key Cities: San Francisco, Los Angeles, Seattle, and San Diego.
- **Southern United States:**
  - Regions: Texas, Florida, Georgia, and North Carolina.
  - Industries: Oil and gas, aerospace, textiles, automobiles, and information technology.
  - Key Cities: Houston, Dallas, Atlanta, Miami, and Charlotte.
- **Ontario and Quebec in Canada:**
  - Regions: Toronto, Montreal, and Ottawa.
  - Industries: Automobile manufacturing, aerospace, information technology, and machinery.

Europe

- **Western Europe:**
  - Regions: Ruhr Valley (Germany), Alsace-Lorraine (France), and Benelux countries (Belgium, Netherlands, Luxembourg).
  - Industries: Steel, automobiles, chemicals, machinery, and textiles.
  - Key Cities: Berlin, Hamburg, Paris, Brussels, and Amsterdam.
- **Northern Europe:**
  - Regions: United Kingdom, particularly the Midlands and Northern England.
  - Industries: Pharmaceuticals, aerospace, chemicals, and electronics.
  - Key Cities: London, Manchester, Birmingham, and Glasgow.
- **Eastern Europe:**
  - Regions: Poland, Czech Republic, Slovakia, and Hungary.
  - Industries: Heavy machinery, steel, chemicals, and automotive.
  - Key Cities: Warsaw, Prague, Bratislava, and Budapest.

Asia

- **East Asia:**
  - Regions: Eastern China, Japan, South Korea, and Taiwan.
  - Industries: Electronics, automobiles, shipbuilding, textiles, and heavy machinery.
  - Key Cities: Shanghai, Beijing, Tokyo, Seoul, and Taipei.
- **South Asia:**
  - Regions: India, particularly Maharashtra, Gujarat, Tamil Nadu, and Karnataka.
  - Industries: Information technology, textiles, automobiles, chemicals, and pharmaceuticals.
  - Key Cities: Mumbai, Bangalore, Chennai, Delhi, and Ahmedabad.
- **Southeast Asia:**
  - Regions: Singapore, Malaysia, Thailand, and Indonesia.
  - Industries: Electronics, petrochemicals, textiles, and food processing.
  - Key Cities: Singapore, Kuala Lumpur, Bangkok, and Jakarta.

Africa

- **North Africa:**
  - Regions: Egypt, Morocco, and Tunisia.
  - Industries: Textiles, food processing, chemicals, and petroleum refining.
  - Key Cities: Cairo, Casablanca, and Tunis.
- **Southern Africa:**
  - Regions: South Africa, particularly Gauteng and Western Cape.
  - Industries: Mining, automobile manufacturing, textiles, and food processing.
  - Key Cities: Johannesburg, Cape Town, and Durban.
- **West Africa:**
  - Regions: Nigeria and Ghana.
  - Industries: Oil and gas, food processing, textiles, and chemicals.
  - Key Cities: Lagos, Accra, and Abuja.

Australia

- **Eastern Australia:**
  - Regions: New South Wales, Victoria, and Queensland.
  - Industries: Mining, machinery, food processing, and chemicals.
  - Key Cities: Sydney, Melbourne, Brisbane, and Adelaide.

South America

- **Brazil:**
  - Regions: São Paulo, Rio de Janeiro, and Minas Gerais.
  - Industries: Automobiles, steel, petrochemicals, and textiles.
  - Key Cities: São Paulo, Rio de Janeiro, and Belo Horizonte.
- **Argentina:**
  - Regions: Buenos Aires and Córdoba.

- Industries: Food processing, automobiles, textiles, and chemicals.
- Key Cities: Buenos Aires and Córdoba.

- **Chile:**
  - Regions: Santiago and Valparaíso.
  - Industries: Mining, food processing, and chemicals.
  - Key Cities: Santiago and Valparaíso.

⚡ **Quick revise**

World industrial regions (handout, one line each) Fold ▲

World industrial regions (handout, one line each). *N. America:* NE USA (steel, autos), West Coast (tech, aerospace), South USA (oil, aerospace), Ontario–Quebec. *Europe:* **Ruhr, Alsace-Lorraine, Benelux**; UK Midlands; Poland–Czechia–Slovakia–Hungary. *Asia:* East Asia (China, Japan, S. Korea, Taiwan), **South Asia (Maharashtra, Gujarat, TN, Karnataka)**, SE Asia (Singapore, Malaysia, Thailand, Indonesia). *Africa:* N. Africa, **Gauteng & Western Cape**, W. Africa (Nigeria, Ghana). *Australia:* east. *S. America:* Brazil (São Paulo–Rio–Minas Gerais), Argentina, Chile.

Important Industrial Regions of India



Fig. 27.18. India : Industrial Regions

- **Mumbai-Pune Industrial Region**
  - Industries: Textiles, automobiles, chemicals, engineering goods, pharmaceuticals, and information technology.
  - Key Cities: Mumbai, Pune, Thane, Nashik.
- **Kolkata-Hooghly Industrial Region**

- Industries: Jute, textiles, chemicals, engineering, iron and steel, and paper.
- Key Cities: Kolkata, Howrah, Durgapur, Haldia.

- **Delhi-NCR Industrial Region**
  - Industries: Information technology, electronics, automobiles, consumer goods, and textiles.
  - Key Cities: Delhi, Gurugram, Noida, Faridabad.
- **Ahmedabad-Vadodara Industrial Region**
  - Industries: Tourism, Textiles, chemicals, pharmaceuticals, petrochemicals, and engineering.
  - Key Cities: Ahmedabad, Vadodara, Surat, Bharuch.
- **Chennai-Bengaluru Industrial Region**
  - Industries: Automobiles, electronics, information technology, textiles, and heavy engineering.
  - Key Cities: Chennai, Bengaluru, Hosur, Coimbatore.
- **Kollam-Thiruvananthapuram Industrial Region**
  - Industries: Cashew processing, coir manufacturing, software development, biotechnology, and electronics.
  - Key Cities: Kollam, Thiruvananthapuram.
- **Chota Nagpur Industrial Region**
  - Industries: Iron and steel, coal mining, heavy engineering, and chemicals.
  - Key Cities: Jamshedpur, Ranchi, Bokaro, Dhanbad.
- **Visakhapatnam-Guntur Industrial Region**
  - Industries: Shipbuilding, petrochemicals, fertilizers, steel, and electronics.
  - Key Cities: Visakhapatnam, Guntur.

⚡ **Quick revise** Indian industrial regions (8) Fold ▲

Indian industrial regions (8).

Region	Industries	Key cities
Mumbai–Pune	textiles, autos, chemicals, engineering, pharma, IT	Mumbai, Pune, Thane, Nashik
Kolkata–Hooghly	jute, textiles, chemicals, engineering, iron & steel, paper	Kolkata, Howrah, Durgapur, Haldia
Delhi–NCR	IT, electronics, autos, consumer goods	Delhi, Gurugram, Noida, Faridabad
Ahmedabad–Vadodara	textiles, chemicals, pharma, petrochemicals	Ahmedabad, Vadodara, Surat, Bharuch
Chennai–Bengaluru	autos, electronics, IT, heavy engineering	Chennai, Bengaluru, Hosur, Coimbatore
Kollam–Thiruvananthapuram	**cashew, coir**, software, biotech	Kollam, Thiruvananthapuram

Region	Industries	Key cities
Chotanagpur	**iron & steel, coal**, heavy engineering	Jamshedpur, Ranchi, Bokaro, Dhanbad
Visakhapatnam–Guntur	shipbuilding, petrochemicals, fertiliser, steel	Visakhapatnam, Guntur

5 · TRANSPORT AND COMMUNICATION

Means Of Transportation



6 · Land Transport

Roadways

- **India's roadway** network is **one of the largest in the world** , spanning over 63 lakh kilometers (Jan 2025).
- It plays a crucial role in the country's transportation system, connecting remote areas with cities, facilitating trade, and contributing to economic growth.
- The roadways in India can be categorized into various types based on their importance and the authority responsible for their construction and maintenance.
- **Types of Roadways in India**



- **National Highways (NH)**
  - These are the **primary roadways connecting major cities, ports, state capitals, and large industrial areas.** They facilitate long-distance travel and play a vital role in the country's economic integration.
  - **Only 2% of the total road network** .
  - **Authority:** Constructed and maintained by the **National Highways Authority of India (NHAI)** , under the Ministry of Road Transport and Highways (MoRTH).
  - Examples: NH44 (previously NH7, the longest highway running from Srinagar to Kanyakumari), NH48 (previously NH8, connecting Delhi and Mumbai).
- **State Highways (SH)**
  - These roads **connect important cities within a state and provide links to the national highways and other state highways.**
  - **4% of the total Road network.**
  - **Authority:** Constructed and maintained by the respective **State Public Works Departments (PWDs).**
  - Examples: Maharashtra State Highway 4 (connecting Pune and Ratnagiri), Karnataka State Highway 1 (connecting Bangalore and Mysore).
- **District Roads**
  - These roads **connect district headquarters with other places in the district.** They are further **classified into Major District Roads (MDR) and Minor District Roads** .
  - **14% of the total road network** .
  - **Authority:** Constructed and maintained by the **Zilla Parishads (District Councils) and the District PWDs.**
  - Examples **:** MDRs in Tamil Nadu and Uttar Pradesh that connect rural areas to district centers.

- **Rural Roads**
  - These are **roads connecting rural areas and villages to nearby towns or district roads.** They are crucial for local connectivity and agricultural activities.
  - **Authority:** Constructed and maintained by the **local bodies (Gram Panchayats) and under the Pradhan Mantri Gram Sadak Yojana (PMGSY).**
  - Examples: Roads constructed under the PMGSY program in various states, improving rural connectivity.
- **Urban Roads**
  - These roads are **found within the limits of towns and cities.** They include arterial roads, sub-arterial roads, and local streets.
  - **Authority:** Constructed and maintained by **Urban Local Bodies (Municipal Corporations or Municipalities).**
  - Examples: Major city roads like Marine Drive in Mumbai, and Rajpath in Delhi.

- **Naming of National Highways in India:**
  - **Up until 2010** , all of our National Highways were numbered **on the basis of the National Highways Act of 1956.**
  - However, these numbers of National Highways **did not give any indication of its location and direction.** Hence, a new system was brought into place by the Ministry of Road Transport and Highways.
  - **North-South Highways:** These highways have **even numbers** . They run primarily in a north-south direction.
    - The **numbering increases from east to west** . Highways close to the eastern part of India have smaller even numbers, and the numbers increase as one moves westward. For example, NH2 is in the eastern part, while NH68 is further west.
  - **East-West Highways:** These highways have **odd numbers** . They run primarily in an east-west direction.
    - The **numbering increases from north to south** . Highways closer to the northern part of India have smaller odd numbers, and the numbers increase as one moves southward. For example, NH1 is in the northern part, while NH87 is further south.
  - **Sub-Highways:**
    - For **highways that branch off from primary highways** , a **prefix is added** to the primary highway number. For example, NH101 would be a branch of NH01.
- April 1, 2018: a **new way to calculate roads** . Lane system: each lane's length is calculated separately.

⚡ **Quick revise** 4.1 Roads Fold ▲

Network > 63 lakh km (Jan 2025). NH = 2% (NHAI, MoRTH) · **State highways = 4%** (state PWDs) · **District roads = 14%** · rural roads (Gram Panchayats, **PMGSY**) · urban roads (municipalities). **NH44** (earlier NH7) **Srinagar–Kanyakumari – longest**; **NH48** (earlier NH8) **Delhi–Mumbai**. Since **1 April 2018** road length counts **each lane separately**.

> 🔢 **NH numbers – start in the north-east corner and walk away. Even = North–South** (numbers rise **east → west**: NH2 east, NH68

west). **Odd = East–West** (numbers rise **north → south**: NH1 north, NH87 south). **Sub-highway = prefix**: NH101 branches from NH01. *(New scheme after 2010; the old 1956-Act numbers told you nothing about location.)*

Railways

- India has **one of the largest and most complex railway networks in the world** , spanning over 68,000 km, playing a critical role in the country's transportation system.
- The Indian Railways network is extensive and serves both passenger and freight needs across diverse geographical regions.
- **Types of Railways in India**
  - **Broad Gauge (BG)**
    - The **most widely used** gauge in India, with a **track width of 1,676 mm (5 ft 6 in)** .
    - Predominantly used for long-distance and heavy freight traffic, as well as passenger services.
  - **Meter Gauge (MG)**
    - A narrower gauge with a **track width of 1,000 mm (3 ft 3 3⁄8 in).**
    - Mostly found in some rural and less developed areas, but is gradually being converted to broad gauge under the Unigauge project.
  - **Narrow Gauge (NG)**
    - The narrowest gauge in India, with a **track width of 762 mm (2 ft 6 in) or 610 mm (2 ft).**
    - Used in hilly and mountainous regions where constructing broad gauge is difficult. Also being phased out or converted to broad gauge.
  - **Standard Guage**
    - A railway track with a gauge of **1,435 mm (4 feet 8 1/2 inches)** .
    - Primarily used for urban rail transit systems like **metro, monorail, and tram** .



  - **Project Unigauge** , launched on **April 1, 1992,** is an **ongoing initiative** by Indian Railways aimed at **converting and standardizing all rail gauges in India** to the

1,676 mm (5 ft 6 in) **broad gauge.**

- **Suburban Railways**
  - **Serve the metropolitan and larger urban areas** , facilitating daily commutes.
  - Mumbai Suburban Railway, Delhi Suburban Railway.
- **Metro Railways**
  - Urban transit systems operating in major cities to reduce congestion and provide efficient public transportation.
  - Delhi Metro, Kolkata Metro, Bengaluru Metro.
- **High-Speed Railways**
  - **Proposed and under-construction** projects aimed at providing high-speed travel between major cities.
  - Mumbai-Ahmedabad High-Speed Rail Corridor.

- **Construction and Maintenance Authorities**
  - **Indian Railways (IR)**
    - A state-owned national railway company, responsible for the construction, maintenance, and operation of the majority of the rail network in India.
    - Divided into 19 zones, each headed by a General Manager. These zones are further divided into divisions.
    - Under the Ministry of Railways, Government of India.
  - **Dedicated Freight Corridor Corporation of India Limited (DFCCIL)**
    - A public sector unit under the Ministry of Railways, responsible for planning, construction, and operation of dedicated freight corridors to improve the efficiency of freight movement.
  - **Metro Rail Corporations**
    - Autonomous bodies or special purpose vehicles (SPVs) formed to plan, construct, and operate metro rail systems in cities.
    - Delhi Metro Rail Corporation (DMRC), Bangalore Metro Rail Corporation Limited (BMRCL).
- **Dedicated Freight Corridors:** Dedicated freight corridors (DFCs) in India are **specialized railway tracks designed exclusively for freight transportation.** These corridors aim to improve the efficiency, speed, and capacity of freight movement across the country, reducing congestion on the existing railway network and facilitating economic growth.



- **Eastern Dedicated Freight Corridor (EDFC)**
  - **Route: Ludhiana** in Punjab **to Dankuni** near Kolkata in West Bengal.
  - **Length:** Approximately **1,839 kilometres.**
  - **Purpose:** To facilitate the movement of coal, steel, and other bulk goods from eastern India to northern India, improving the efficiency of goods transportation in the region.
  - **Status: Under construction** , with **several sections already operational** .
- **Western Dedicated Freight Corridor (WDFC)**
  - **Route: Dadri** in Uttar Pradesh **to Jawaharlal Nehru Port Trust (JNPT)** near Mumbai in Maharashtra.
  - **Length:** Approximately **1,506 kilometres** .
  - **Purpose:** To enhance the movement of goods from the industrial heartland of Delhi-NCR to the major port in Mumbai, thus facilitating export-import trade and reducing logistics costs.
  - **Status: Under construction** , with **significant sections already operational.**
- **Future Projects:**
  - **East-West Corridor** : Connecting Kolkata to Mumbai.
  - **North-South Corridor** : Connecting Delhi to Chennai.
  - **East Coast Corridor** : Connecting Kharagpur to Vijayawada.
  - **Southern Corridor** : Connecting Chennai to Goa.

⚡ **Quick revise** 4.2 Railways Fold ▲

> 68,000 km.

Gauge	Width	Where
Broad	**1,676 mm**	most widely used; long-distance and freight
Metre	**1,000 mm**	rural; being converted (**Unigauge, launched 1 April 1992**)
Narrow	**762 mm or 610 mm**	hills; being phased out

Gauge	Width	Where
Standard	**1,435 mm**	metro, monorail, tram

> 📏 **Metre = 1,000 mm (name = number). Broad = the biggest, 1,676. Standard 1,435 = city metros.**

Indian Railways: state-owned, **19 zones** (General Manager each), Ministry of Railways. **DFCCIL** runs **dedicated freight corridors: Eastern DFC – Ludhiana (Punjab) → Dankuni (near Kolkata), ~1,839 km** (coal, steel) · **Western DFC – Dadri (UP) → JNPT (Mumbai), ~1,506 km**. Future: East–West (Kolkata–Mumbai), North–South (Delhi–Chennai), East Coast (Kharagpur–Vijayawada), Southern (Chennai–Goa). Also suburban (Mumbai, Delhi), metro (DMRC, BMRCL), proposed **Mumbai–Ahmedabad high-speed rail.**

Pipelines



- Pipelines in India form a crucial part of the country's infrastructure for transporting a variety of commodities, including crude oil, natural gas, petroleum products, and water. They **provide a safe, efficient, and cost-effective means of transportation over long distances.**
- Their **limiting factor** is that they **can only transport liquids or gases** , and they **have less flexibility compared to railways and roadways.**
- **What They Transport**
  - **Crude Oil:** Transported from oil fields to refineries.
  - **Natural Gas:** Transported from gas fields to processing plants and consumers.
  - **Petroleum Products:** Transported from refineries to distribution points, including petrol, diesel, kerosene, and aviation fuel.

- **Ores:** Mixed with water to allow them to flow.
- **Water:** Transported from water sources to urban, rural, and industrial areas.
- **Milk:** in very local locations in the world as milk is perishable

- **Important Examples**
  - **Naharkatia-Noonmati-Barauni Pipeline**
    - Initially till Noonmati, later extended to Barauni.
    - Crude oil
  - **Mumbai High-Mumbai-Ankaleshwar-Koyali Pipeline**
    - Crude oil & Natural Gas
  - **Salaya-Koyali-Mathura**
    - Crude oil to Koyali & Mathura.
    - Extended from Mathura to Panipat & later Jalandhar.
  - **Hazira-Bijapur-Jagdishpur Gas Pipeline**
    - By GAIL
    - Largest underground pipeline of world
    - Extended to Delhi via Dadri

⚡ **Quick revise** Pipelines Fold ▲

Pipelines (liquids and gases only; less flexible): **Naharkatia–Noonmati–Barauni** (crude) · **Mumbai High–Mumbai–Ankleshwar–Koyali** (crude + gas) · **Salaya–Koyali–Mathura** (crude; extended to Panipat, Jalandhar) · **Hazira–Bijapur–Jagdishpur (GAIL)** – **largest underground pipeline** in the world; extended to Delhi via Dadri. Also carry ores (slurry), water, local milk.

Ropeways: first major passenger ropeway **Auli, 1969**; **Parvatmala Scheme** (Budget 2022-23, PPP, MoRTH; 8 projects ~60 km initially; **Jan 2024: 400 projects, ₹1.25 lakh crore in 5 yrs**); **Varanasi urban ropeway; Gaurikund–Kedarnath (9.7 km, 3,584 m)**.

Ropeways

- Ropeways **use suspended cables to transport goods and passengers, typically over difficult terrain.** They are especially **popular in** hilly, mountainous, and remote **areas where conventional transportation** modes like roads and railways **are challenging to construct** .
- Ropeways in India **date back to the early 20th century** , with industrial applications in mining and logging. The **first major passenger ropeway** in India was established in 1969 in **Auli** , Uttarakhand, which opened the door for further developments across the country. Since then, several ropeways **have been established in popular tourist destinations** , such as in Darjeeling, Gulmarg, and Vaishno Devi.
- **PARVATMALA SCHEME**
  - The **Parvatmala Scheme** , announced in the 2022-23 Union Budget by India's Union Finance Minister, is a **National Ropeways Development Programme**

focused on improving connectivity in India's hilly regions.

- The program, launched under a **Public-Private Partnership (PPP)** model, offers an eco-friendly alternative to conventional roadways in difficult terrains. It aims to enhance commuter convenience, promote tourism, and potentially relieve congestion in crowded urban areas.
- The scheme **initially targets states like Uttarakhand, Himachal Pradesh, Manipur, Jammu & Kashmir, and other Northeastern regions, with eight ropeway projects** covering 60 kilometers planned for 2022-23.
- The **Ministry of Road Transport and Highways (MORTH)** oversees the program, supported by a regulatory framework to ensure standards and efficiency.
- The scheme is **significant for its economic benefits** (lower land acquisition costs due to direct routes), **speed** (aerial transport bypasses road obstacles), **environmental friendliness** (reduced dust emissions), and potential for **last-mile connectivity** .
- In **January 2024** , it was announced that the **number of projects** under the scheme would be **increased to 400** , with an expense of Rs. 1.25 lakh crore in 5 years.
- Ropeways under construction:
  - Varanasi Urban Ropeway
  - Gaurikund-Kedarnath Ropeway (9.7 km, 3,584m altitude)

7 · WATER TRANSPORT

- **Why** is water transport **cheap** ?
  - Water does not provide as much friction as land, so **fuel efficiency is higher** .
  - There is **no need to create separate infrastructure** for sea travel **except the terminals** . E.g. roads, rails, and pipes are needed for land-based transportation. This adds to the savings.

Inland Waterways

- Inland waterways in India have a long history dating back to ancient times, when rivers were utilized as crucial arteries for trade and transportation.
  - **Ancient Importance:** Rivers like the Ganges, Brahmaputra, and Godavari have historically served as major trade routes, facilitating commerce and cultural exchange.
  - **Colonial Legacy:** The British colonial administration developed some inland waterways for transportation of goods during the colonial period, recognizing the strategic importance of river navigation.
- **Construction and Development**
  - **Inland Waterways Authority of India (IWAI):** Established in 1986, IWAI is the primary authority responsible for the regulation and development of inland waterways in India.
  - **National Waterway Act:** Enacted in 1986, this legislation identified specific stretches of rivers, canals, and backwaters as national waterways for development

and regulation.

- **Modernization Efforts:** Recent years have seen increased investment in the modernization and expansion of inland waterways infrastructure, including dredging, construction of terminals, and installation of navigation aids.

- There are 111 officially notified Inland Waterways in the country. Some Important Examples **:**



THE LINKS AND THE LENGTHS

NW-1	Ganga–Bhagirathi–Hooghly (Haldia-Allahabad)	1,620 km
NW-2	Brahmaputra river	891 km
NW-3	West Coast Canal–Champakara Canal–Udyogmandal Canal	205 km
NW-4	Krishna (Muktiyala–Vijayawada)	82 km
NW-10	Amba river	45 km
NW-83	Rajpuri Creek	31 km
NW-85	Revadanda Creek–Kundalika river	31 km
NW-91	Shastri river-Jaigad Creek System	52 km
NW-68	Mandovi river (Usgaon Bridge–Arabian Sea)	41 km
NW-111	Zuari river (Sanvordem Bridge–Marmugao Port)	50 km
NW-73	Narmada river	226 km
NW-100	Tapi river	436 km
NW-97	Sunderbans Waterways	172 km

⚡ **Quick revise** Why cheap? Fold ▲

Why cheap? low friction → fuel efficient; **no separate infrastructure except terminals. IWAI established 1986; National Waterway Act 1986; 111 notified inland waterways.**

Waterway	Route	Length
NW-1	Ganga–Bhagirathi–Hooghly (**Haldia–Allahabad/Prayagraj**)	**1,620 km**
NW-2	Brahmaputra (**Dhubri–Sadiya**)	**891 km**
NW-3	West Coast Canal – Champakara – Udyogmandal	**205 km**
NW-4	Krishna (Muktiyala–Vijayawada)	**82 km**
NW-73 / NW-100	Narmada / Tapi	226 / 436 km
NW-97	Sunderbans	172 km
NW-10, 68, 83, 85, 91, 111	Amba 45 · Mandovi 41 · Rajpuri creek 31 · Revdanda–Kundalika 31 · Shastri–Jaigad 52 · Zuari 50	km

THE LINKS AND THE LENGTHS

NW-1	Ganga–Bhagirathi–Hooghly (Haldia-Allahabad)	1,620 km
NW-2	Brahmaputra river	891 km
NW-3	West Coast Canal–Champakara Canal–Udyogmandal Canal	205 km
NW-4	Krishna (Muktiyala–Vijayawada)	82 km
NW-10	Amba river	45 km
NW-83	Rajpuri Creek	31 km
NW-85	Revadanda Creek–Kundalika river	31 km
NW-91	Shastri river-Jaigad Creek System	52 km
NW-68	Mandovi river (Usgaon Bridge–Arabian Sea)	41 km
NW-111	Zuari river (Sanvordem Bridge–Marmugao Port)	50 km
NW-73	Narmada river	226 km
NW-100	Tapi river	436 km
NW-97	Sunderbans Waterways	172 km

Handout figure: the 13 national waterways in operation and their lengths.

Open Ocean Transport

- India's ocean-based water transport has a **rich history** and plays a vital role in the country's economy and connectivity.
- Approximately **95 percent of India's foreign trade by volume and 70 percent by value** moves through ocean routes.
- **History:**

- **Ancient Times:** India's maritime history **dates back to the Indus Valley Civilization** , with evidence of trade links with Mesopotamia and Egypt. Ancient ports like Lothal were significant centers of commerce.

- **Medieval Period:** During the medieval period, Indian ports facilitated **trade with Africa, the Middle East, and Southeast Asia** . Notable ports included Calicut, Surat, and Cochin.
- **Colonial Era:** The colonial period saw the establishment of **modern ports by the British** , such as Mumbai (Bombay), Kolkata (Calcutta), and Chennai (Madras), to facilitate trade and control.
- **Post-Independence:** After independence, India **expanded and modernized its port infrastructure** to support economic growth and international trade.

- India has **13 major ports** (newest being Galathea Bay at Great Nicobar) **and over 200 minor and intermediate ports** . The 14th Port has been commissioned at Vadhavan, near Mumbai, but it is yet to be constructed.

Major Port	Location	Important facts /
Deendayal Port	Kandla, Gulf of Kachchh, Gujarat	Formerly **Kandla** west-coast port chemicals, salt, Strategically serv
Mumbai Port	Mumbai Harbour, Maharashtra	One of India's ol deep-water harb western India's t cargo.
Jawaharlal Nehru Port Authority (JNPA)	Nhava Sheva, Navi Mumbai, Maharashtra	India's leading c decongest Mum containerised tra India.
Vadhavan Port	Palghar district, Maharashtra	**Notified/approve** planned as a d port. The Gover 2024 at an estim
Mormugao Port	Mormugao, Goa	Important wes significant for irc handles coal, pet cargo.

New Mangalore Port	Panambur, Mangaluru, Karnataka	Only major port petroleum produ fertilisers, iron ore
Cochin Port	Kochi, Kerala	Major port on the **International Terminal**; impor petroleum, ferti maritime trade of
V.O. Chidambaranar Port	Thoothukudi/Tuticorin, Tamil Nadu	Important port in coal, salt, fertili cargo. It became **implement a Di** management.
Chennai Port	Chennai, Tamil Nadu	One of India's o coast; important petroleum proc historically a ma Nadu.
Kamarajar Port	Ennore, near Chennai, Tamil Nadu	India's first corpo for coal, automob and bulk cargo. Chennai Port.
Visakhapatnam Port	Visakhapatnam, Andhra Pradesh	Natural harbour important for fertilisers and ste linked with easter
Paradip Port	Paradip, Odisha	Major bulk-carg important for c fertilisers. It has bulk-cargo perfo
Syama Prasad Mookerjee Port	Kolkata-Haldia, West Bengal	Formerly **Kolkat** riverine port sys of **Kolkata Doc Complex**; crucial India.
Galathea Bay Port	Great Nicobar Island, Andaman & Nicobar Islands	**Notified major pc Container Transs** Nicobar project; east-west interna

- **Port Management**
  - **Ministry of Ports, Shipping, and Waterways:** The central government body responsible for formulating policies and overseeing the development and regulation of ports in India.

- **Port Trusts:** Major ports are managed by autonomous Port Trusts established under the Major Port Trusts Act, of 1963. Each Port Trust operates under the supervision of a Board of Trustees appointed by the government.
- **State Maritime Boards:** Minor and intermediate ports are managed by State Maritime Boards or State Port Departments. These entities oversee the development, regulation, and management of ports within their respective states.
- **Private Sector Participation:** In recent years, there has been significant private sector participation in port development and operations through Public-Private Partnerships (PPPs). Private companies invest in port infrastructure, container terminals, and logistics services.

⚡ **Quick revise** 4.4 Water transport Fold ▲

Ocean: ~95% of foreign trade by volume and 70% by value. 13 major ports + Vadhavan (14th, approved June 2024, ₹76,220 crore, Palghar). Newest major port: **Galathea Bay (Great Nicobar)** – transshipment. Managed by **Port Trusts (Major Port Trusts Act 1963)**; minor ports by **State Maritime Boards.**

⚓ **West coast, north → south: "Kind Men Join Very Many New Clubs"** = **K**andla (Deendayal) · **M**umbai · **J**NPA (Nhava Sheva – leading container port) · **V**adhavan · **M**ormugao (Goa, iron ore) · **N**ew Mangalore (only major port of Karnataka) · **C**ochin (Vallarpadam transshipment). **East coast, south → north: "Very Clever Kids Visit Pretty Kolkata"** = **V**.O. Chidambaranar (Tuticorin) · **C**hennai · **K**amarajar (Ennore) · **V**isakhapatnam (natural harbour) · **P**aradip (bulk) · **K**olkata–Haldia (**Syama Prasad Mookerjee Port – only riverine major port**, Hooghly).

8 · AIR TRANSPORT

Airways in India

- **History:**
  - **Early Beginnings:** Air transport in India began in the early 20th century. The **first commercial flight took place on February 18, 1911, from Allahabad to Naini,** covering a distance of about 6 miles.
  - **Post-Independence Era:** After independence, India **established its own national carrier** , which played a key role in developing the country's aviation sector. In **1948, the government acquired a 49% stake in Tata-owned Air India**
  - **Nationalization:** Done in **1953** , when all Private airlines were merged to form a national state carrier, i.e. the **Indian Airlines for all domestic flights, and Air India for international services.**

- **Liberalization and Growth:** The aviation sector was **liberalized in the 1990s** , leading to the entry of private airlines and significant growth in the industry. This period saw the emergence of several domestic carriers, including Jet Airways, IndiGo, SpiceJet, and others.
- **Modern Era:** Today, India boasts a rapidly growing aviation market, with a large number of domestic and international flights connecting major cities and regions. The sector continues to expand with increasing passenger traffic and investment in airport infrastructure.
- **State exit:** A proposal for the sale of **Air India** was floated in 2017. After multiple failures and delays, it was **finally sold to the Tata Group in 2021** (officially handed over in Jan 2022).
- >90% of Indian air traffic handled by 2 companies. Concerns of Duopoly.

- India has **157** operational airports of which 133 are overseen by Airports Authority of India.
- **Airport Management in India:**
  - **Airports Authority of India (AAI):** AAI is the statutory body responsible for creating, upgrading, maintaining, and managing civil aviation infrastructure in India. It operates over 100 airports, including international, domestic, and civil enclaves.
  - **Private Sector Participation:** In recent years, several major airports have been developed and managed through Public-Private Partnerships (PPPs). Private entities such as GMR Group, Adani Group, and GVK Group operate major airports like Indira Gandhi International Airport in Delhi, Chhatrapati Shivaji Maharaj International Airport in Mumbai, and Kempegowda International Airport in Bengaluru, in conjunction with the AAI (except Mumbai).
  - **State Governments:** Some state governments also play a role in managing smaller airports within their territories, often in collaboration with AAI or private players.

⚡ **Quick revise** Air: first commercial flight Fold ▲

Air: first commercial flight **18 Feb 1911, Allahabad–Naini**; govt took **49%** of Tata-owned Air India **1948**; **nationalised 1953** (Indian Airlines domestic, Air India international); liberalised **1990s**; **Air India sold to Tata Group 2021 (handed over Jan 2022)**; **>90% traffic with two companies (duopoly concern)**; **157 operational airports, 133 with AAI.** Private operators: **GMR, Adani, GVK** (Delhi, Mumbai, Bengaluru).

Post/telecom timeline: Indian Post Office Act 1837 → Railway Mail → Act 1854 (uniform rates) → Speed Post **1986** · **Telegraph: Calcutta–Diamond Harbour 1851; ended 2013** · **First telephone exchange Kolkata 1881** · **Radio:** clubs 1920s, **Indian Broadcasting Company 1927**, **All India Radio 1936** (Quit India used it) · **Doordarshan 1959; colour 1982** · **Internet 1995** · **Digital India 2015** · fast 5G rollout · **ISRO** communication satellites.

9 · COMMUNICATION

- **Prior to the introduction of the Postal System in India, communication was mostly in conjunction with transportation.** Either the person who wanted to communicate travelled to the receiver, or they sent some messengers.
- The British colonial administration recognized the importance of an efficient communication system for governance and trade. The **first organized postal system was established in the late 18th century.**

Postal System

- **Indian Post Office Act of 1837:** This act laid the foundation for the modern postal system in India. It standardized postal services and rates across the country.
- **Railway Mail Service:** With the expansion of the railway network in the **mid-19th century** , the Railway Mail Service was introduced, significantly speeding up mail delivery.
- **Indian Post Office Act of 1854:** This act reorganized the postal system, making it more efficient and widespread. It **introduced uniform postage rates** and **established post offices in urban and rural areas.**
- **Expansion and Modernization: After independence in 1947** , the Indian postal system expanded rapidly to cover rural and remote areas. The **postal network became one of the largest in the world.**
- **Introduction of Speed Post:** In **1986** , India Post introduced Speed Post, a high-speed postal service, **to compete with private courier services** .

Telegraph

- **Introduction of Telegraph:** The telegraph was introduced to India in the **mid-19th century.** The **first telegraph line was laid between Kolkata (then Calcutta) and Diamond Harbour in 1851.**
- **Expansion: By 1854** , telegraph lines **connected major cities** like Kolkata, Mumbai (then Bombay), Chennai (then Madras), and Peshawar. The telegraph network **expanded rapidly** , covering vast distances and enhancing communication across the British Empire in India.
- **End of Telegraph Services:** With the rise of digital communication, the use of telegraph declined. **In 2013, India officially ended its telegraph services.**

Telephones

- The **first telephone exchange** in India was established in **Kolkata in 1881** , marking the beginning of telephone services in the country.
- **Expansion:** Telephone services gradually **expanded to other major cities** , but their reach remained limited primarily to urban areas and government offices.
- **Liberalization and Privatization:** The liberalization of the Indian economy **in the 1990s** led to the **entry of private telecom companies** , which transformed the sector. This period saw the launch of mobile phone services and significant growth in the number of subscribers.

- **Mobile Revolution:** The **early 2000s** marked the mobile revolution, with affordable handsets and competitive pricing making mobile phones accessible to a large population.
- **Currently,** India has become **one of the fastest countries to roll out 5G** .

Radio

- **Introduction of Radio Broadcasting:** Radio broadcasting began in India with the **establishment of private radio clubs in the early 1920s.** The **first commercial radio station, Indian Broadcasting Company, started operations in Mumbai and Kolkata in 1927.**
- **All India Radio (AIR):** The **British government took over radio broadcasting in 1936** , establishing **All India Radio (AIR) as the national broadcaster** . AIR played a key role in disseminating news and information during World War II.
- The **Quit India Movement was also energised by the use of this mass media.**
- All India Radio **expanded its reach post-independence** , with more radio stations set up across the country, broadcasting in multiple languages.

Television:

- **Introduction of Television: Doordarshan** , India's public service broadcaster, **began television services in 1959** . Television rapidly gained popularity, especially with the **introduction of color broadcasts in 1982.**
- **Private Channels:** The **1990s liberalization policy** led to the entry of private television channels, breaking the monopoly of Doordarshan and diversifying content.

Digital Communication

- **Introduction of the Internet:** The Internet was introduced in India in 1995. The subsequent years saw a rapid increase in internet users, driven by affordable data plans and smartphones.
- **Growth of Digital Services:** The government launched various initiatives to promote digital communication, including e-governance services, online education, and telemedicine.
- **Digital India Initiative:** Launched in 2015, the Digital India initiative aimed to transform India into a digitally empowered society and knowledge economy. It focused on improving digital infrastructure, enhancing internet connectivity, and providing digital services to citizens.

Satellite Communication:

- Satellite communication in India has significantly enhanced the country's capabilities in broadcasting, telecommunications, and internet services.
- **ISRO, has been instrumental in developing and deploying a series of communication satellites** that cater to various needs, including television broadcasting, telephony, data transmission, and internet services. More details in Science & Tech.

10 · PYQ

Q.1) Consider the following pairs :

National Highway	Cities conr
NH4	Chennai an
NH6	Mumbai an
NH15	Ahmedaba

Which of the above pairs is/are correctly matched ? (2014)

(a) 1 and 2 only

(b) 3 only

(c) 1, 2 and 3

(d) None

Answer: D National Highway 4 or NH 4, It is of 230.7 km of length.

- This highway is known as Andaman Trunk Road (The Great Andaman Trunk Road).
- A stretch of national highway from Mumbai to Pune to Hubli to Bangalore to Chennai was earlier called NH 4 before renumbering of national highways in the year 2010.
- The former NH 4 is now renumbered as NH 48.
- National Highway 6, is a national highway in India running from Jorabat in Meghalaya to Selling in Mizoram.
- The highway passes through the Indian states of Meghalaya, Assam, and Mizoram.
- National Highway 15 (NH 15) starts from Baihata in Assam and terminates at Wakro in Arunachal Pradesh.



Q.2) In which of the following activities are Indian Remote Sensing (IRS) satellites used?

1. Assessment of crop productivity

2. Locating groundwater resources

3. Mineral exploration

4. Telecommunications

5. Traffic studies

Select the correct answer using the code given below (2015)

(a) 1, 2 and 3 only

(b) 4 and 5 only

(c) 1 and 2 only

(d) 1, 2, 3, 4 and 5

Answer: A India's remote sensing programme under the Indian Space Research Organisation (ISRO) started off in 1988 with the IRS-1A, the first of the series of indigenous state-of-art operating remote sensing satellites, which was successfully launched into a polar sun-synchronous orbit on March 17, 1988, from the Soviet Cosmodrome at Baikonur.

- Data from Indian Remote Sensing satellites are used for various applications of resources survey and management under the National Natural Resources Management System (NNRMS). Following is the list of those applications:
  - Space Based Inputs for Decentralised Planning (SIS-DP)
  - National Urban Information System (NUIS)

- ISRO Disaster Management Support Programme (ISRO-DMSP)
- Biodiversity Characterization at landscape level- http://bis.iirs.gov.in
- Preharvest crop area and production estimation of major crops.
- Drought monitoring and assessment based on vegetation condition.
- Flood risk zone mapping and flood damage assessment.
- Hydro-geomorphological maps for locating underground water resources for drilling wells.
- Irrigation command area status monitoring
- Snow-melt run-off estimates for planning water use in downstream projects
- Land use and land cover mapping
- Urban planning
- Forest survey
- Wetland mapping
- Environmental impact analysis
- Mineral Prospecting
- Coastal studies

Q.3) Consider the following pairs:

Port	Well known as
Kamarajar Port	First major port in India registere
Mundra Port	Largest privately owned port in In
Visakhapatnam	Largest container port in Port Ind

How many of the above pairs are correctly matched? (2023)

(a) Only one pair

(b) Only two pairs

(c) All three pairs

(d) None of the pairs

Answer: B

- Kamarajar port at Tuticorin is the first corporate port of India, registered under the Companies Act. It started in 2001.
- The Mundra Port owned by Adani Ports is the largest private port of India. It has also overtaken JNPT port to become the largest container port of India.

⚡ **Quick revise** Transport PYQs Fold ▲

Year	Asked	Answer / rule
2014	NH4 Chennai–Hyderabad? NH6 Mumbai–Kolkata? NH15 Ahmedabad–Jodhpur?	**None (D)** – old NH4 became **NH48**; NH6 (Jorabat–Selling) is in the NE; NH15 starts at Baihata, Assam
2015	**IRS satellites** used for: crop productivity, groundwater, mineral exploration (✔); telecom, traffic (✘)	**A (1, 2, 3)**
2023	Kamarajar = first major port registered as a company ✔ · Mundra = largest privately owned ✔ · Visakhapatnam = largest container port ✘	**B (two pairs)** – Mundra also overtook JNPT as largest container port

⚠️ **Handout alert:** the 2023 PYQ explanation says **"Kamarajar Port at Tuticorin"**, but the port table puts **Kamarajar at Ennore, near Chennai** (and **V.O. Chidambaranar at Tuticorin**). Use the table.

Recall sheet

1. **Kharif Jun–Oct** (paddy, cotton, maize…) · **Rabi Oct–Mar** (wheat, gram, mustard) · **Zaid Mar–Jun** (melons).
2. **75–115–120**: dry <75 cm/75 days; dryland 75–115/75–120; wetland >115/>120.
3. **Black soil + cotton; sugarcane = most water; bajra survives 42 °C; pulses fix nitrogen (766 vs 1,015 kg/ha).**
4. **Revolutions:** Green (Swaminathan), White (Kurien), Blue, Yellow; Round = potato; Silver fibre = cotton.
5. **Weber:** weight-losing → raw material; weight-gaining → market. **Isotim = transport cost; isodapane = total cost.**
6. **Copper:** Rajasthan 52.25 / MP 23.28 / Jharkhand 15.14. **Diamond:** Panna only. **Mica:** Jharkhand (Koderma). **Thorium:** Kerala, 25% of world.
7. **Indian coal: non-coking, high ash, low sulphur.** Gondwana belts: Damodar–Mahanadi–Godavari–Son.
8. **Renewables 283.46 GW; solar 157→280 GW (2030); wind 56 GW; nuclear 8,180→22,480 MW.**
9. **Gauges:** 1,676 / 1,000 / 762–610 / 1,435. **NH:** even N–S, odd E–W.
10. **Ports:** 13 major + Vadhavan; **Galathea Bay** newest; **only riverine major port = Kolkata–Haldia (SPM)**.

Traps

- **Wheat is Rabi (not Kharif)** – 2013 PYQ. **Groundnut, sesamum, bajra are rainfed.**
- **Palm oil is native to Guinea hills, not SE Asia** (2021). **Excess N → acidity + nitrate leaching, not more N-fixing microbes** (2015).
- **Chemical farming is hard in semi-arid regions** (2020); **permaculture mimics local conditions**.

- **Uranium for power needs ~5% enrichment, not 60%** (2023). **REE = 17 elements; China first, not India second.**
- **Mundra/JNPT, not Visakhapatnam, are container leaders**; **Kamarajar = Ennore** (table) vs "Tuticorin" (PYQ text).
- **Old NH numbers:** NH4 → NH48; NH44 is the long north–south road; **even = N–S, odd = E–W**.
- **Coal import reason = coking coal + short supply**, not "saving reserves".

Connects to: *Indian Geography* (soils/crops, Gondwana coal, rivers/waterways, ports) · *Climatology* (monsoon and crop seasons, Western Disturbances) · *Human Geography* (settlements and industrial regions).

GEOGRAPHY · MIND MAPS · SUBJECT 1

Mind maps · Geomorphology

36 mind-map pages | Tap a map to zoom

Every mind-map page of the Geomorphology section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.72



Mind map · PDF p.73



Mind map · PDF p.74



Mind map · PDF p.75



Mind map · PDF p.76



Mind map · PDF p.77



Mind map · PDF p.78



Mind map · PDF p.79



Mind map · PDF p.80



Mind map · PDF p.81



Mind map · PDF p.82



Mind map · PDF p.83



Mind map · PDF p.84



Mind map · PDF p.85



Mind map · PDF p.86



Mind map · PDF p.87



Mind map · PDF p.88



Mind map · PDF p.89



Mind map · PDF p.90



Mind map · PDF p.91



Mind map · PDF p.92

Mind map · PDF p.93



Mind map · PDF p.94



Mind map · PDF p.95



Mind map · PDF p.96



Mind map · PDF p.97



Mind map · PDF p.98



Mind map · PDF p.99



Mind map · PDF p.100



Mind map · PDF p.101



Mind map · PDF p.102



Mind map · PDF p.103



Mind map · PDF p.104



Mind map · PDF p.105



Mind map · PDF p.106



Mind map · PDF p.107


LAKES
Body of water of considerable size, localized in a basin, surrounded by land apart from a river or other outlet that feeds/drains it.
Most; have at least one natural outflow
Endorheic Lakes do not have a natural outflow; lose water solely by evaporation or underground seepage or both
Formation of lakes
earth movements Tectonic lakes; Rift valley lakes
By glaciation ; Tarns, Rock hollow lakes
By Volcanic Activities Crater and Caldera lakes
By Erosion Karst, Playa, Wind deflated
By Deposition Ox-Bow lakes, Lagoons; Glacial lakes
Man-made lakes

GEOGRAPHY · MIND MAPS · SUBJECT 2

Mind maps · Climatology

21 mind-map pages | Tap a map to zoom

Every mind-map page of the Climatology section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.161



Mind map · PDF p.162



Mind map · PDF p.163



Mind map · PDF p.164



Mind map · PDF p.165



Mind map · PDF p.166



Mind map · PDF p.167



Mind map · PDF p.168



Mind map · PDF p.169



Mind map · PDF p.170



Mind map · PDF p.171



Mind map · PDF p.172



Mind map · PDF p.173



Mind map · PDF p.174



Mind map · PDF p.175



Mind map · PDF p.176



Mind map · PDF p.177



Mind map · PDF p.178



Mind map · PDF p.179



Mind map · PDF p.180



Mind map · PDF p.181



GEOGRAPHY · MIND MAPS · SUBJECT 3

Mind maps · Oceanography

18 mind-map pages | Tap a map to zoom

Every mind-map page of the Oceanography section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.216



Mind map · PDF p.217



Mind map · PDF p.218



Mind map · PDF p.219



Mind map · PDF p.220



Mind map · PDF p.221



Mind map · PDF p.222



Mind map · PDF p.223



Mind map · PDF p.224



Mind map · PDF p.225



Mind map · PDF p.226



Mind map · PDF p.227



Mind map · PDF p.228



Mind map · PDF p.229



Mind map · PDF p.230



Mind map · PDF p.231



Mind map · PDF p.232



Mind map · PDF p.233


OCEAN RESOURCES
Biotic Resources
Edible Resources
Fish
Shellfish
Seaweed
Non-Edible Resources
Marine Pharmaceuticals
Coral Reefs
Pearls
Seal and Whale Products
Non-Biotic Resources
Energy Resources
Oil and Gas
Renewable Energy
Mineral Resources
Dissolves salts
Manganese Nodules (Poly-Metallic Nodules)
Phosphorite Deposits
Intangible Resources
Regulating Services (Environmental Balance)
Climate regulation; ~90% of the excess heat
Carbon sink
Heat & circulation regulation
Weather & water cycle regulation
Coastal protection
Cultural & Recreational Services
Tourism & recreation
Cultural & spiritual values
Education & research
Aesthetic & inspiration
Supporting Services (Underlying Systems)
Oxygen production; 50–60% of Earth's oxygen
Biodiversity hotspots
Nutrient cycling
Habitat provision
LAW OF SEA
Body of international law that governs the use of the world's oceans and seas.
United Nations Convention on the Law of the Sea (UNCLOS): 1982
Components
Territorial Sea: 12 nautical miles (22.2 kilometres
Contiguous Zone; 24 nautical miles (44.4 kilometres)
Exclusive Economic Zone (EEZ): 200 nautical miles (370.4 kilometres)
High Seas

GEOGRAPHY · MIND MAPS · SUBJECT 4

Mind maps · Indian Geography

29 mind-map pages | Tap a map to zoom

Every mind-map page of the Indian Geography section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.318



Mind map · PDF p.319



Mind map · PDF p.320



Mind map · PDF p.321



Mind map · PDF p.322



Mind map · PDF p.323



Mind map · PDF p.324



Mind map · PDF p.325



Mind map · PDF p.326



Mind map · PDF p.327



Mind map · PDF p.328



Mind map · PDF p.329



Mind map · PDF p.330



Mind map · PDF p.331



Mind map · PDF p.332



Mind map · PDF p.333



Mind map · PDF p.334



Mind map · PDF p.335



Mind map · PDF p.336



Mind map · PDF p.337



Mind map · PDF p.338



Mind map · PDF p.339



Mind map · PDF p.340



Mind map · PDF p.341



Mind map · PDF p.342



Mind map · PDF p.343



Mind map · PDF p.344



Mind map · PDF p.345



Mind map · PDF p.346


SOILS OF INDIA
Soil Degradation
Decline in quality and health of soil
Causes of Soil Degradation
Soil Erosion
Deforestation
Unsustainable farming practices
Overgrazing
Chemical Degradation
Excessive use of fertilizers and pesticides
Improper use of chemicals
Improper irrigation practices
Physical Degradation
INDIA: A SUB-CONTINENT
Natural borders
Separate plate
Almost all the major global climate and forest types
Distinct history, cultures, traditions

GEOGRAPHY · MIND MAPS · SUBJECT 5

Mind maps · Human Geography

6 mind-map pages | Tap a map to zoom

Every mind-map page of the Human Geography section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.365



Mind map · PDF p.366



Mind map · PDF p.367



Mind map · PDF p.368



Table No. 1A: State-wise major tribes of India

Sl_No.	States/UTs	Major Tribes
1	Andaman and Nicobar Islands	Jarawa , Oraons, Onges, Sentinelese, Shompens
2	Andhra Pradesh	Andh, Sadhu Andh, Bhagata, Bhil, Chenchus (Chenchawar), Gadabas, Gond, Goundu, Jatapus, Kammara, Kattunayakan, Kolawar, Kolam, Konda, Manna Dhora, Pardhan, Rona, Savaras, Dabba Yerukula, Nakkala, Dhulia, Thoti, Sugalis.
3	Arunachal Pradesh	Apatanis, Abor, Dafla, Galong, Monba, Sherdukpen, Singpho.
4	Assam	Chakma, Chutiya, Dimasa, Hajong, Garos, Khasis, Gangte.
5	Bihar	Asur, Baiga, Birhor, Birjia, Chero, Gond, Parhaiya, Santhals, Savar.
6	Chhattisgarh	Agariya, Bhaina, Bhattra, Biar, Khond, Mawasi, Nagasia.
7	Goa	Dhodia, Dubia, Naikda, Sidhi, Varli.
8	Gujarat	Barda, Bamcha, Bhil, Charan, Dhodia, Gamta, Paradhi, Patelia.
9	Himachal Pradesh	Gaddis, Gujjars, Khas, Lambu, Lahaulas, Pangwala, Swangla.
10	Jammu and Kashmir	Bakarwal, Balti, Beda, Gaddi, Garra, Mon, Purigpa, Sippi.
11	Jharkhand	Birhors, Bhumij, Gonds, Kharia, Mundas, Santhals, Savar.
12	Karnataka	Adiyan, Barda, Gond, Bhil, Iruliga, Koraga, Patelia, Yenava.
13	Kerala	Adiyan, Arnadan, Eravallan, Kurumbas, Mala arayas, Moplahs, Uralis.
14	Madhya Pradesh	Baigas, Bhils, Bharia, Birhors, Gonds,Katkari, kharia, Khond, Kol, Murias.
15	Maharashtra	Bhuina, Bhunjia, Dhodia, Katkari, Khond, Rathawa, Warlis.
16	Manipur	Aimol, Anganni, Chiru, Kuki, Marram, Meisang, Paite, Purum, Thadou.
17	Meghalaya	Chakma, Garos, Hajong, Jaintias Khasis, Lakher, Pawai, Raba.
18	Mizoram	Chakma, Dimasa, Khasi, Kuki, Lakher, Pawai, Raba, Synteng.
19	Nagaland	Angami, Garo, Kachari, Kuki, Mikir, Nagas, Sema.
20	Odisha	Gadaba, Ghara, Kharia, Khond, Matya, Oraons, Rajuar, Santhals.
21	Rajasthan	Bhils, Damaria, Dhanka, MeenasMinasi, Patelia, Sahariya.
22	Sikkim	Bhutia, Khas, Lepchas.
23	Tamil Nadu	Adiyan, Arnadan, Eravallan, Irular, Kadar, Kanikar, Kotas, Todas
24	Telangana	Chenchus.
25	Tripura	Bhil, Bhutia, Chaimal, Chakma, Halam, Khasia, Lushai, Mizel, Namte.
26	Uttarakhand	Bhotias, Buksa, Jaunsari, Khas, Raji, Tharu.
27	Uttar Pradesh	Bhotia, Buksa, Jaunsari, Kol, Raji, Tharu.
28	West Bengal	Asur, Khond, Hajong, Ho, Parhaiya, Rabha, Santhals, Savar.

Source: Indian Census

Mind map · PDF p.369



Mind map · PDF p.370



GEOGRAPHY · MIND MAPS · SUBJECT 6

Mind maps · Economic Geography

21 mind-map pages | Tap a map to zoom

Every mind-map page of the Economic Geography section of your PDF, in the PDF's order. Mark each one read as you revise it.

Mind map · PDF p.431



Mind map · PDF p.432



Mind map · PDF p.433



Mind map · PDF p.434



Mind map · PDF p.435



Mind map · PDF p.436



Mind map · PDF p.437



Mind map · PDF p.438



Mind map · PDF p.439



Mind map · PDF p.440



Mind map · PDF p.441



Mind map · PDF p.442



Mind map · PDF p.443



Mind map · PDF p.444



Mind map · PDF p.445



Mind map · PDF p.446



Mind map · PDF p.447



Mind map · PDF p.448



Mind map · PDF p.449



Mind map · PDF p.450



Mind map · PDF p.451



Priority and PYQ index

H = 4 chapters · M = 8 · L = 0. Prelims years as quoted in the handout and class slides; a number in brackets is the count in that year.

Ch	Chapter	Priority	Prelims PYQs
1	Geomorphology	H	HIGH · 19 Prelims, 2 Mains
2	Climatology	H	VERY HIGH · 32 Prelims, 1 Mains
3	Oceanography	M	MEDIUM · 6 Prelims, 1 Mains
4	Indian Geography	H	VERY HIGH · 40 Prelims
5	Human Geography	M	MEDIUM · 4 Prelims
6	Economic Geography	H	VERY HIGH · 31 Prelims
7	Mind maps · Geomorphology	M	none quoted in the material
8	Mind maps · Climatology	M	none quoted in the material
9	Mind maps · Oceanography	M	none quoted in the material
10	Mind maps · Indian Geography	M	none quoted in the material
11	Mind maps · Human Geography	M	none quoted in the material
12	Mind maps · Economic Geography	M	none quoted in the material