Shrinkage test: a real donor faded towards the dataset mean, blend = mean + lambda (donor - mean), lambda = r of the rung (repeat 0.85, mirror 0.24). R: 0 = repeat, 1 = unrelated person; lower = judged closer.

metric	repeat scan R donor	R blend	blend - donor [95% CI]	verdict	other hemisphere R donor	R blend	blend - donor [95% CI]	verdict
MSE	0.00	-0.01	-0.01 [-0.03, -0.00]	prefers the faded donor	0.72	0.37	-0.35 [-0.40, -0.28]	prefers the faded donor
RMSE	0.00	-0.01	-0.01 [-0.03, -0.00]	prefers the faded donor	0.77	0.49	-0.27 [-0.32, -0.22]	prefers the faded donor
MAE	0.00	-0.01	-0.01 [-0.03, +0.00]	indifferent	0.70	0.50	-0.21 [-0.26, -0.15]	prefers the faded donor
PSNR	0.00	-0.01	-0.01 [-0.04, +0.00]	indifferent	0.81	0.62	-0.20 [-0.24, -0.15]	prefers the faded donor
PSNR, fixed range	0.00	-0.01	-0.01 [-0.03, +0.00]	indifferent	0.82	0.62	-0.20 [-0.23, -0.15]	prefers the faded donor
SSIM	0.00	0.00	+0.00 [-0.02, +0.02]	indifferent	0.82	0.70	-0.13 [-0.16, -0.09]	prefers the faded donor
Pearson r	0.00	-0.01	-0.01 [-0.01, -0.00]	prefers the faded donor	0.82	0.56	-0.26 [-0.29, -0.23]	prefers the faded donor
ACC vs dataset mean	0.00	-0.00	-0.00 [-0.00, +0.00]	indifferent	0.75	0.75	-0.00 [-0.00, +0.00]	indifferent
ACC vs sex-age mean	0.00	-0.00	-0.00 [-0.00, +0.00]	indifferent	0.75	0.74	-0.02 [-0.02, -0.01]	prefers the faded donor
lDDT-like	0.00	-0.07	-0.07 [-0.09, -0.05]	prefers the faded donor	0.89	0.39	-0.50 [-0.56, -0.45]	prefers the faded donor

Intervals: bootstrap over families and positions (random cubes, both hemispheres). pre-QC: affine registration only, MRI registration gate not met (results/registration/status.md)