

# Sulfate-reducing bacteria unearthed: ecological functions of the diverse prokaryotic group in terrestrial environments

Konstantin A. Demin,<sup>1</sup> Evgeniya V. Prazdnova,<sup>1</sup> Tatiana M. Minkina,<sup>1</sup> Andrey V. Gorovtsov<sup>1</sup>

**AUTHOR AFFILIATION** See affiliation list on p. 15.

**ABSTRACT** Sulfate-reducing prokaryotes (SRPs) are essential microorganisms that play crucial roles in various ecological processes. Even though SRPs have been studied for over a century, there are still gaps in our understanding of their biology. In the past two decades, a significant amount of data on SRP ecology has been accumulated. This review aims to consolidate that information, focusing on SRPs in soils, their relation to the rare biosphere, uncultured sulfate reducers, and their interactions with other organisms in terrestrial ecosystems. SRPs in soils form part of the rare biosphere and contribute to various processes as a low-density population. The data reveal a diverse range of sulfate-reducing taxa intricately involved in terrestrial carbon and sulfur cycles. While some taxa like *Desulfitobacterium* and *Desulfosporosinus* are well studied, others are more enigmatic. For example, members of the Acidobacteriota phylum appear to hold significant importance for the terrestrial sulfur cycle. Many aspects of SRP ecology remain mysterious, including sulfate reduction in different bacterial phyla, interactions with bacteria and fungi in soils, and the existence of soil sulfate-reducing archaea. Utilizing metagenomic, metatranscriptomic, and culture-dependent approaches will help uncover the diversity, functional potential, and adaptations of SRPs in the global environment.

**KEYWORDS** sulfate reduction, uncultured microorganisms, soil microbiology, sulfur cycle, rare biosphere, terrestrial environment

Existing trends in the study of certain taxa of microorganisms in modern science can lead to biases that distort our understanding of their place in nature. These biases can be associated with the location or ecological niche where representatives of the group were first discovered. An example of such bias can be seen in research on archaea, which for a long time were solely considered through the prism of extremophilic lifestyles. However, recent studies indicate the need to study mesophilic archaeal species and acknowledge the group's ubiquitous nature (1) and even its potential involvement in human pathological processes (2).

We believe that the study of sulfate-reducing prokaryotes (SRPs) has encountered a similar issue. Abundant new data, including metagenomic analyses, indicate that the involvement of SRPs in ecological cycles in soil habitats may be underestimated.

SRPs have been extensively studied since the 19th century, and these works concentrated predominantly on specific habitats where this group plays crucial roles, such as structuring of microbial community in paddy soils and wetlands, driving the carbon flux in the global marine environment, and the role in pathogenic processes in the intestinal tract. Moreover, the biotechnological importance of SRPs was realized, either as the primary corrosion agent (3, 4) or essential part of bioreactors and activated sludges (5, 6). However, the substantial amount of new information collected about this group for the last two decades allows one to suggest even more ecological functions mediated by SRPs in nature. It includes SRP contribution to microbial communities at different levels, existence of a large proportion of uncultured and unclassified taxa

**Editor** Gemma Reguera, Michigan State University, East Lansing, Michigan, USA

Address correspondence to Konstantin A. Demin, i@kincognita.ru.

The authors declare no conflict of interest.

See the funding table on p. 15.

**Published** 29 March 2024

Copyright © 2024 American Society for Microbiology. All Rights Reserved.

related to SRPs, and, most importantly, the ecology of SRP in soils. The latter is the most important gap in studying the microbial turnover of sulfur: SRPs were reported in soils multiple times, but the actual role of this group is poorly understood.

The main reasons for this bias are as follows:

1. SRPs are more commonly studied in the context of marine ecosystems.
2. While planning research, a wrong assumption can emerge that anaerobic SRPs are exclusive for anaerobic environments, not aerated soils.
3. Sulfate-reducing bacteria in terrestrial environments do not represent a cohesive taxonomic group, which impedes comprehensive study of whole terrestrial SRP diversity.
4. This group encompasses many non-culturable forms, whose physiological understanding has only recently become accessible with the advancement of metagenomic, metatranscriptomic, and predictive metabolic pathway analysis methods.

Considering all the aforementioned points, we strongly believe that the current perspective on SRPs needs to be revised based on the latest data, which will enable the emergence of new trends in the study of the environmental role of SRPs.

Therefore, the primary aim of this review is to examine the present status of understanding regarding SRP ecology within terrestrial ecosystems and to bring together new and overlooked data. Additionally, we propose a model highlighting the role of SRPs in terrestrial environments, particularly in soils.

## WHAT DO WE KNOW ABOUT SULFATE-REDUCING PROKARYOTES?

The ability to reduce sulfur compounds, especially using sulfate anions as the terminal electron acceptors coupled to sulfide generation [dissimilatory sulfate reduction (DSR)], is not connected to a specific taxonomic group. This is due to the lateral gene transfer events and the unique evolutionary trajectory of SR metabolism. Species of the same taxonomic unit (genus or family) may be able to reduce (i) sulfate solely, (ii) sulfate and other S compounds such as elemental sulfur or thiosulfate, (iii) only some compounds except sulfate, or (iv) even disproportionate S compounds. There is no strict linkage between phylogeny and physiology. Moreover, the preference of electron acceptor depends on the environmental conditions, which results in using ferrous iron, nitrates, or even oxygen and organics. These considerations make it impossible to limit the group of bacteria under discussion only to complete reducers to sulfide or only to those which reduce sulfate solely, not to mention specific taxa. Here, we will use “sulfate reducers” or “SRPs” as generalizing terms for microbes with dissimilatory metabolism of sulfur compounds.

Sulfate reduction is one of the most critical processes in known ecosystems since it is the main way sulfur compounds are reduced in modern ecosystems. DSR is often coupled with the oxidation of many organic substrates, from volatile fatty acids to recalcitrant aromatic hydrocarbons (7). Some estimates indicate that sulfate is the second most abundant anion in marine environments after chlorides (8). With low oxygen levels and high concentrations of dissolved organic matter (9), water environments provide preferable conditions for sulfate-reducing prokaryotes to flourish, which is why microorganisms capable of DSR are well studied in marine and freshwater environments. Sulfate reducers also populate flooded soils and deep soil horizons (10, 11). In some cases, microbial reduction of sulfur compounds can be one of the major drivers of biogeochemical turnover, which is especially true for wetlands (12), peatlands (13), paddy soils (14), and soils undergoing flooding-evaporation cycles. Polluted and harsh environmental sites with active soil formation processes, such as sludges and acid mines, also comprise niches for SRPs (15–17). At the same time, the role of SRPs in natural ecosystems is not restricted solely to participation in the sulfur cycle. Microbial metabolism intensely connects the sulfur and carbon cycle (18). For example, sulfate

reducers are responsible for a significant flux of CO<sub>2</sub> from organics, which is especially important in marine ecosystems (19), where they can oxidize up to 30% of dissolved organics. As for the nitrogen cycle, the most critical SRP process is the ability to reduce nitrites and nitrates if the sulfate anions are absent (20). Moreover, data that sulfate reduction can be coupled with nitrogen fixation by some SRPs existed for more than 70 years (21) and are confirmed by recent works (22, 23).

Furthermore, according to some estimations, SRPs can act as primary producers in crust microbiomes without relying on photosynthetic processes (24). Another remarkable SRP characteristic is their interaction with compounds that are often considered xenobiotics or presented in xenobiotic concentrations. Many pollutants of significant environmental concern can be metabolized or immobilized by sulfate reducers. It is well known that these microorganisms can degrade mono- and polyaromatics (7, 25–28) and chlorinated organics (29) and precipitate heavy metals (such as Zn, Ni, and Cu) in the form of poorly soluble sulfides or enhance precipitation indirectly, so they gain a selective advantage in ecosystems with polymetallic or combined pollution (30, 31). Sulfate reducers are specifically known for sphalerite (ZnS) and related minerals precipitation (32, 33). Moreover, SRP-immobilized zinc was suggested to be more stable than its abiogenic alternative (34). One unique group of SRPs, known as dissimilatory metal-reducing bacteria, can reduce most heavy metal (HM) ions, including uranium (35, 36). Currently, many sulfate reducers are known to be able to reduce extracellular metals directly or via shuttle molecules. Overall, sulfate-reducing bacteria are essential in metal biogeochemistry (37–39). The intricate cell biology and physiology of SRP, however, are beyond the scope of the review and can be explored in other recent publications (40).

Due to the ubiquitous distribution of SRPs in natural ecosystems, there are numerous gaps in our understanding of their ecology. This includes the phenomenon of dormancy in SRPs, their role in the rare biosphere, and the degree to which SRP metabolic capacities go beyond the reduction of sulfur species. The most prominent gaps are visible when considering terrestrial microbial communities with SRPs.

## EVIDENCE FOR SRPs AS A VITAL PART OF TERRESTRIAL ECOSYSTEMS

Marine and aquatic conditions are favorable for SRPs, which is why our knowledge of this large and diverse group is limited to water habitats (41). To date, SRP population dynamics, diversity, and physiology are poorly understood in pristine and anthropogenically impacted terrestrial ecosystems, especially in soils (28, 42). Collected data are mostly restricted to areas where anaerobic conditions are sustained most of the time, such as paddy soils and subsurface water horizons. Despite the recognition of the urgent need for studies addressing the biogeochemical role of sulfate reducers in aerobic or hypoxic terrestrial zones (41, 42), little progress has been made in the last years. One can presume that the strictly anaerobic group of microorganisms will not be an active part of environments with the regular influx of oxygen (such as topsoils) or that its influence will be negligible and limited to anaerobic microniches. However, there are multiple aerotolerant and oxygen-respiring SRP genera, such as *Desulfovibrio*, *Desulfobulbus*, *Desulfomicrobium*, and *Desulfofaba* (43, 44), and even proliferation in aerated topsoil was reported (28). The understanding slowly emerges that DSR can substantially affect ecosystem productivity in terrestrial environments (12, 13, 41, 45). Moreover, low sulfate concentrations in soils, peatlands, and rice paddy fields are not constraints for either high sulfate reduction rates or sulfate reducer diversity (46).

Except for the extensively studied *Desulfovibrio* and *Desulfotomaculum*, plenty of other important SRPs inhabit soils, including those with occurring aerobic conditions (Fig. 1). The most demonstrative in-land sulfate reducers belong to the Bacillota phylum. The first example is *Desulfitobacterium* genus. More than 20 years ago, scientists showed that pollutant-degrading *Desulfitobacterium* species, initially isolated from the environment contaminated with chlorinated aromatics, are widespread throughout different soils with various levels of anthropogenic impact (47). The authors hypothesized that the studied *Desulfitobacterium* are a part of diverse natural microbial communities with




FIG 1 Summary of different ecological functions mediated by SRPs in soils.

the ability to degrade chlorinated organics as an accessory. Further studies reported the active role of SRP species in HM attenuation in contaminated soils (48–50), with *Desulfitobacterium* among such species (50, 51), because of their ability to form poorly soluble metal sulfides (52). The ubiquity of the genus was confirmed by the study of Mingo et al. (53), who showed that different phenyl methyl ether-demethylating *Desulfitobacterium* strains were isolated in great quantity from top horizons of different soils, and 16S rRNA gene copies were observed in each soil sample studied. In another study, deoxynivalenol was significantly de-epoxidized by a microbial consortium isolated from wheat topsoil, with *Desulfitobacterium* sp. as a significant actor in deoxynivalenol detoxification (54). Interestingly, phenyl methyl ethers and deoxynivalenol are products of fungal metabolism, suggesting another role of *Desulfitobacterium* spp. in soils, aside from the sulfur and iron reduction.

Recently, petroleum-degrading, thiosulfate-, and iron-reducing microbe was isolated from contaminated soil, which genetically clustered with *Desulfitobacterium aromaticivorans* but showed distinct physiological and phylogenetical traits (55). It was suggested to classify this strain to the new *Paradesulfitobacterium* genus and reclassify *Desulfitobacterium aromaticivorans* to it. An interesting study was performed by Huang et al. (56), where soil incubation experiments with glucose, lignin, and hemicellulose amendments and hypoxic conditions resulted in the enrichment of *Desulfitobacterium* and *Desulfosporosinus* genera. Furthermore, these two genera and *Desulfotomaculum* were the significant taxa emerging from approximately 60-cm depth of cryosols during long-term thawing experiments, which confirms that they are adapted to altering conditions in soils (57). All these findings widen our perspective on the SRP diversity and ecology in soils.

*Desulfosporosinus* genus appears to be another SRPs frequently met in hypoxic/oxic environments. It is usually captured from water-saturated soils such as paddy fields (56, 58), wetlands, peatlands, and other soils with anoxic conditions. However, the presence of the genus has been shown in more air-saturated soils too. These include enrichment cultures from As-Pb-contaminated soils near smelters in China (49), microcosms with alkane-contaminated soils (59), and natural communities of benzene-toluene-xylene-contaminated soils (60). A study of Gagliano et al. (61) describes a topsoil (0–3 cm) microbial community near the terrestrial hydrothermal vents with harsh conditions, including low pH and temperature >60°C. *Desulfosporosinus*, followed by *Desulfotomaculum*, were among the major dominating taxa. The authors did not specifically discuss that observation but reported a high level of water saturation in this soil and attributed it to the active condensation of the hydrothermal water vapor from below. It presumably should create hypoxic conditions, but intense water evaporation in hydrothermal vents and the vicinity to the surface air make this case especially interesting in the scope of this review.

Genera *Desulfosporosinus* and *Desulfotomaculum*, along with *Desulfovermiculus*, again have been reported as dominating groups in the deep horizon of colluvial soils (62). The authors mentioned that anoxic conditions for studied soils, even at the deep level, are rare and transient. Furthermore, a study on more temperate soils revealed the presence and the dominance of *Desulfosporosinus* operational taxonomic unit (OTU) (followed by *Desulfobulbus*) among SRPs in areas of intensive land use, such as arable soils and grasslands (42). One more SRP genus from Bacillota for which soil as an isolation source and habitat was reported is *Desulfonispota*. Iron-rich soils (63), soils from ice-free zones in the Arctic (64), and ornithogenic arctic soils (65) have been stated as the source where *Desulfonispota* was found to be isolated or detected in metagenomic studies. Although *Desulfonispota* instead metabolizes organosulfonates than actively reduces sulfate (66), its publicly available genome annotation contains *DsrAB* genes, suggesting the ability to reduce sulfite (BioProject accession number [PRJEB20235](#)).

Other phyla, such as Thermodesulfobacteriota, Acidobacteriota, Campylobacterota, and Thermotogota, also provide SRPs found in land ecosystems. A compelling and counterintuitive case is the metagenomic study undertaken in China wetlands, where the *Desulfuromonadaceae* family was significantly enriched in sediments during the dry season (67). Based on the network analysis, authors suggested this family to be one of the putative keystones shaping the dry season microbial community. Sequences affiliated with *Desulfuromonas*, a typical member of *Desulfuromonadaceae* family, were found to be abundant in an amplicon metagenomic survey of soils near the Roopkund Glacier, India, and the already mentioned *Desulfosporosinus* and *Desulfitobacterium* genera were present as well (68). Like *Desulfuromonadaceae* (67), *Desulfurellaceae* family was among the significant and active members of the soil microbiome, as reported by two independent studies. The first denotes the increased abundance of the family because of farm activities (69), and the second addresses the effects of tree species on adjusting soil's microbial communities (70). The microbial community of the latter study was completely in-land without flooding, and the authors managed to uncover significant differences in *Desulfurellaceae* abundance between soils under different tree coverage. The study by Miletto et al. (71) found that in experimentally flooded grassland soil, there was an increase in the abundance of *Desulfobulbaceae* family members compared to salt marsh soil, where *Desulfobacteraceae* members were the predominant sulfate-reducing bacteria. Finally, *Mesotoga* genus, which is the only mesophilic member of Thermotogota phylum, can reduce thiosulfate and molecular sulfur and can be found in soils (72, 73). Moreover, it was hypothesized that the genus is spreading across various terrestrial biomes as a result of human technogenic activity, since its presence is often accompanied by high content of polyaromatic hydrocarbons (PAHs) (73).

In contrast to soils, deep terrestrial subsurface is well recognized as a habitat for SRPs. These bacteria were repeatedly reported as the active members of deep crystalline bedrock microbial community (74–79) and can be found at more than 2-kilometer depth



(80). In deep subsurface, SRPs contribute to symbiosis with archaea or to metal corrosion. The latter is specifically studied in the context of radioactive waste disposal, since SRPs immobilize and absorb radioactive elements in deep fractures (81–85) as well as mobilize radioactive metals such as plutonium both directly (86) or indirectly by copper containers corrosion (87).

## BIOTIC RELATIONS OF SRPs IN SOILS

### Rhizosphere

Sulfate reducers are present in diverse ecological niches and habitats, including plant rhizosphere. However, most studies focus on the microbial community of seagrass and other plants inhabiting soil-water interfaces. Many papers are studying the SRP communities in paddy soils. A rather diverse SRP community characterizes rice fields. Among the genera identified in such soils are *Desulfobacca*, *Desulfovibrio*, *Syntrophobacter*, *Desulforhopalus*, *Desulfarculus*, and *Desulfobulbus* (88). The SRPs play a vital role in the rhizosphere of plants grown in soils contaminated by potentially toxic elements (PTEs). In paddy soils located in the zone adjacent to thallium sulfide utilization plant, thallium's geochemical fractionation is largely reliant on the activity of Fe-reducing *Geothrix* and sulphate-reducing *Anaerolinea* (89). Sulfur-cycling microorganisms also contributed to the mitigation of chromium toxicity by reduction of Cr(VI) to less toxic Cr(III). A recent study showed that in the rice rhizosphere, *Desulfovibrio* species reduced sulfate to sulfide utilizing citric acid from root exudates as electron donors. *Thiobacillus* used the resulting sulfide to reduce Cr(VI) (90). It has been shown that the addition of sulphate reduced Cd's translocation from soil to rice grain by 53.2%–63.2%, depending on moisture conditions. The mechanism involved CdS precipitation by *Desulfovibrio*, promoted by flooding (91). As for the influence of SRPs on the bioavailability of arsenic, the evidence is contradictory (92). It has been shown that SRPs can take part in direct and indirect reduction of As, leading to increased bioavailability (93). At the same time, other studies suggest that the microbially mediated formation of secondary mackinawite (FeS) leads to trapping As in its structure, decreasing the mobility of this toxic element (94).

SRPs were also found in other highly flooded yet in-land niches, such as coastal silt soils and salt marshes. In coastal silts, it was shown that functional bacteria enriched in *S. glauca* rhizosphere can enhance the driving effect of the sulfur cycle and promote the transformation and circulation of Fe, N, and C in the soil by enhancing the degradation of organic matter and sulfate reduction (95). Sulfate-reducing bacteria were thus thought to be particular hub bacteria for the growth of salt-tolerant plants. For salt marshes, a recent study revealed that the core rhizosphere microbiome of a typical salt marsh plant, *Spartina alterniflora*, is highly enriched with sulfate-reducing genera, such as *Desulfatiglans*, *Desulfocarbo*, *Desulfatitalea*, *Desulfobulbus*, *Desulfopila*, and *Desulfosarcina* (96). Thus, significantly higher rates of enzymatic organic matter decomposition as well as the relative abundances of putative sulfur (S)-oxidizing, sulfate-reducing, and nitrifying prokaryotes correlated with *S. alterniflora* productivity (96). Moreover, prokaryotic species from the *Desulfobulbus*, *Desulfatitalea*, *Desulfovibrio*, and *Sulfurospirillum* genera were shown to couple sulfate or sulfur respiration with nitrogen fixation (96–98), which expands their role in rhizosphere communities.

In the rhizosphere of plants in paddy soils, there are several driving factors determining the numbers and activity of the SRPs. Main factors include the availability of sulphates, dissolved organic matter, and their ratio. More than 30 years ago, it was shown that the optimal ratio of sulphates to available carbon sources is 2:1, and the maximum numbers of SRPs are observed in the rhizosphere during the tillering phase (99). This can be attributed to root exudation activity which differs significantly during plant development. Another important factor is the oxygen level, which should limit SRP growth. Studies of rice rhizosphere have shown that the conditions in rhizosphere soil are oxic and would theoretically be unsuitable for SRP activity. However, the presence of anoxic microsites ensures their survival and participation in geochemical processes (100). Furthermore, among SRPs, several species are tolerant to the presence of oxygen

to some degree (101, 102). Thus, we should expect SRPs to be present in the rhizosphere of plants growing in well-drained soils that are not subjected to regular flooding.

However, there are very few publications dedicated to SRP communities of such soils. In 2006, the black layer formation in the rootzone of *Agrostis stolonifera* was attributed to active biologically induced sulfate reduction (103). The studies of oak rhizosphere revealed that the abundance of particular genera was different in rhizosphere and bulk soil. The abundance of *Desulfacinum* was higher in bulk soil compared to oak rhizosphere. The abundance of *Desulfuromonas* was low in both rhizosphere and bulk soils (104). The influence of tobacco plants on the abundance of SRPs was different and dependent on the Cd contamination level. High Cd treatment ( $5.8 \text{ mg kg}^{-1}$ ) tended to enrich sulfate-reducing bacteria (such as *Desulfarculaceae*). In comparison, low Cd treatment ( $0.2 \text{ mg kg}^{-1}$ ) led to the enrichment of plant growth-promoting bacteria (*Bacillus*, *Dyadobacter*, *Virgibacillus*, and *Lysobacter*) (105). This observation demonstrates the potential role of SRPs in decreasing the bioavailability of PTEs in soils with normal water regimes. The SRP communities are also active in the rhizosphere of crop plants. For example, sulfate reducers are present in the rhizosphere of maize, and their abundance is more stable when the plants are colonized by mycorrhizal fungi (106).

To conclude, among the SRPs, the role in terrestrial plants' rhizosphere is shown for *Desulfovibrio*, *Syntrophobacter*, *Desulfobacca*, *Desulforhopalus*, *Desulfarculus*, *Desulfacinum*, *Desulfuromonas*, *Anaerolinea*, and *Desulfobulbus*. The representatives of the mentioned genera can take part in regulating PTE bioavailability and depend on plant root exudates as electron donors. Despite the potentially oxic conditions, their ecological niches are not limited to flooded soils and should be studied in more detail in all terrestrial ecosystems.

### SRPs as part of the soil rare biosphere and the role of phages in their life strategy

The most demonstrative and cited studies about the SRP impact on an ecosystem concerning their low relative abundance have been conducted by Pester et al. (13). Results showed that SRPs, being the "rare" part of the particular microbial community, can still considerably contribute to the biogeochemical flow of carbon, with the effects going way beyond the community itself. Furthermore, the intriguing observation was made that sulfate reducers can increase ribosomal count and sustain some transcriptional activity even at zero-growth, comprising less than 0.01% of the microcosm community (107). Finally, the group showed that a small population of *Desulfosporosinus* in microcosms does not multiply its biomass by growth or proliferation but remains transcriptionally and metabolically active for more than a month (108). Such zero growth coupled with cellular activity was hypothesized to be the meaningful strategy in highly fluctuating niches, i.e., peat soils. Unfortunately, not so many studies of SRPs combining community profiling and cultural data have been conducted so far.

A survey of George et al. (42) has shown noticeable diversity of sulfate reducers in different soils at low levels of absolute abundance, which peaked in grassland soils and was not correlated strictly with the acidity or moisture level. Again, the most abundant SRP taxa were *Desulfosporosinus* and *Desulfobulbus*. The authors hypothesized that the low observable relative abundance of sulfate reducers can be explained by both SRP dormancy/rarity and the inappropriateness of general prokaryotic primers to amplify SRP ribosomal genes. This, in turn, can shed light on the other study of (109), where the amplicon metagenomic approach failed to reveal the presence of sulfate reducers in the flooded coal mine. However, plenty of sulfur oxidizers were found, hinting at some source of reduced sulfur compounds. Indeed, subsequent enrichment culturing revealed the presence of *Desulfotomaculum* and *Desulfomicrobium* species, and similarly, it was then hypothesized that they are rare biosphere members. It is the question of whether such results are caused by actual rarity or insufficient primers. It is perhaps both, especially since it is reported that microbial groups can be enriched from a sample inoculate while not showing in its metagenomic profile (109).

An interesting case was reported by Karnachuk et al. (110). The study focuses on the thermophilic *Desulfofundulus* species, which was isolated from soils and rocks near burning coal seams. This species produces spores capable of germinating into metabolically active sulfate-reducing cells, even in a mesophilic environment. However, these cells are unable to divide under such conditions. Based on this discovery, authors challenge the existing explanation that abundant thermophiles in mesophilic conditions are merely metabolically inactive biomass. It can be further hypothesized that at least part of reductive sulfur metabolism taking place in moderate soils is attributed to thermophilic bacteria, like *Desulfofundulus*, *Thermotoga*, *Caldisericum*, and many others.

Based on the above, we should consider it essential to conduct culture-dependent studies to characterize an SRP community. These observations are also interesting in the context of SRP virome. Viruses of sulfate reducers, presented especially as prophages, are numerous (111). They also can coinfect their metabolic and syntrophic partners (112) and contribute to the ecological role of SRPs (113, 114). For example, based on an analysis of soil-recovered SRP genomes, it was suggested by Li et al. (41) that SRP phages can participate in cell life by providing protection from oxidative stress and facilitating glycoside hydrolysis. Such participation can be achieved if SRP viruses are present as prophages, which, in turn, can be facilitated by a life strategy with low population density, rare cell division events, and high transcriptional activity.

## Fungi

Fungal-SRP interactions are an even rarer topic in literature. Anaerobic fungi in deep granite fractures were hypothesized to be in syntrophic relationships with SRPs by supplying the latter with hydrogen and acetate (77). Both fungi and SRPs are known for colonization and degradation of metal or concrete materials, such as rocks, ceramics, metal statues, and several built environments (115), where they can coexist in a biofilm. This view is supported by the study of Nuppenen-Puputti et al. (116), where cross-kingdom interactions between fungi and SRPs were detected on schist surfaces, and by the survey of Edgcomb et al. (117), found collective hydrocarbon degradation by fungi and sulfate-reducing bacteria (SRB) in hydrothermal vents. Again, possible metabolic interaction can exist between SRB and fungi in soils, given the ability of *Desulfotobacterium* spp. to transform fungal metabolites (53, 54). Thus, arid soils, rocks, granite intrusions, and ore deposits provide potential spots for interactions between sulfate reducers and fungal species in terrestrial ecosystems. More studies are required for understanding fungal-SRP interactions.

## Syntrophic interactions with other bacteria

Despite the long research history, little is known about syntrophic interactions between SRPs and other bacteria, especially in soils. It has been known for a long time (118) that sulfate reducers naturally develop relationships with sulfur-oxidizing (both lithoautotrophic and photosynthesizing) bacteria by providing them with reduced sulfur species such as sulfide or thiosulfate. Their co-occurrence can be found in multiple areas (96, 101, 119). However, the relationships between sulfate-reducing and sulfur-oxidizing bacteria (SOB) in soils are poorly understood. Since the high diversity of both SRPs and SOB groups as well as different sulfur compounds readily available in natural conditions, we should expect complex cooperative-competitive interactions between different members of sulfur-transforming community. This picture became even more sophisticated if oxidative-type Dsr pathway in SOB and sulfide oxidation metabolism in SRPs [like in *Desulfobulbus*, *Desulfovibrio*, see references (120, 121)] are considered. Many studies undertaken so far describe reducer-oxidizer relationship only in marine sediments (122, 123) or artificial systems, such as municipal wastewater treatment plants (124–127) and laboratory co-cultures (128). Direct syntrophic association was described by Yamamoto-Ikemoto et al. (125), where sulfur bacteria filamentous bulking and increased sulfide oxidation rates were observed after the elevation of microbial sulfate-reducing activity in an activated sludge system. Fukui et al. (122) reported close natural associations between



gliding filamentous sulfate-reducing *Desulfonema* spp. and giant filamentous sulfide-oxidizing *Tioplota* and *Beggiatoa* species in coastal and freshwater pond's sediments. The authors hypothesized in the first place that the capacity for gliding motility of *Desulfonema* is hindered in sediments and that *Tioplota* sheaths provide another substrate for gliding in such conditions. Recently, a close association between sulfide-oxidizing cable Desulfobulbaceae bacteria and *Desulfatiglans* was confirmed to take place in sediments suboxic zone (129). These interactions should take place in soils too, since such electroactive Desulfobulbaceae genera as *Electronema*, *Electrothrix*, and *Desulfobulbus* can be found in temperate and anaerobic soils (39, 42, 71). On the other hand, soils, as aquatic sediments, are solid-state heterogenic media, where filamentous species can provide not only nutritional but also spatial advantage for motile sulfate metabolizing bacteria. SRP potential interactions with fungal hyphae discussed above also hint on such type of symbiosis. Xia et al. (130), reported co-occurrence of SRPs and SOB in landfill cover soil layers. Uranium-bearing sandstone was also documented as a source with an active sulfate-reducing and sulfur-oxidizing microbial community (131). In the rhizosphere zone, the existence of SRPs depends on the activity with which aerobic species remove oxygen, although it is not a specialized type of interaction. More close interactions were reported multiple times in between SRPs and methanogens or methane oxidizers, as reviewed in references (132, 133). It was the subject of recent investigations (133–135). Almost all comprehensively studied syntrophy includes archaea in subseafloor, except for Candidatus *Acidiflora* and methanogens in thawing permafrost (136), and the case of *Mesotoga prima*, a sulfur and thiosulfate-reducing microorganism that can establish interspecies hydrogen transfer with hydrogenotrophic sulfate reducers in oil reservoirs and, possibly, soils (137).

Finally, microbial interactions between SRPs and other representatives of natural microbial communities, particularly sulfur compound oxidizers, have been studied for many years. However, the studies have been heavily focused on aquatic or artificial ecosystems, while terrestrial habitats have largely been overlooked. The diversity of SRP morphology and life strategies (motile and attached, filamentous and single-celled, actively dividing and capable of maintaining activity at zero growth, microaerophiles and obligate anaerobes), metabolism types, available sulfur compounds, and potential symbiotic partners suggests that we have only scratched the surface in studying their relationships with other bacteria.

## UNCULTURED AND UNCLASSIFIED REALM OF SULFATE-REDUCING MICROORGANISMS

Most of the knowledge about sulfate reducers was obtained by studying culturable representatives of the group. However, a vast gap exists between the cultured and observed diversity of dissimilatory sulfite reductase (*Dsr*) genes. As far back as in 2008 (11), unique phylotypes deeply divergent from common cultured sulfate-reducing bacteria were found. Since then, novel *Dsr* variants continued to emerge from multiple studies. In wetlands, for example, most sulfate reducers belong to unclassified and rare taxa (12) as well as in peatlands (138). The distribution of *Dsr* genes across various genomes suggests the presence of full gene clusters for sulfate, sulfite, or sulfur reduction in very different phyla (19). Interestingly, almost full clusters were found in typically aerobic bacteria such as Actinomycetota. More recently, the diversity of *Dsr* genes was estimated in arsenic-contaminated soils, and again, most variants were new at the genus and family levels (139). In the same year as the study of Anantharaman group, a high-coverage metabarcoding survey with more than 200 samples representing 14 habitats revealed even more hidden diversity of *Dsr* genes, with an amount of new species-level OTU reaching more than 150,000 (140). The authors stated that the *Dsr* copies of already cultivated sulfate reducers comprise less than 1% of the discovered quantity, even when we consider the existence of inactive copies of *Dsr* genes and those which are working in the opposite direction in the sulfur-oxidizing taxa. It was concluded that a "rare sulfate-reducing microorganisms' biosphere" exists that can serve as a seed

bank triggered by environmental changes. From the perspective of the current review, it should be noted that a limited amount of temperate terrestrial ecosystems has been taken into the study and no highly aerated representatives at all. Can it then be that even such high previously undetected diversity is underrated? The answer is likely yes, given that, in the study of Vigneron et al. (140), nearly 30% of uncovered gene copies were attributed to unknown and unclassified taxa and affiliated with different uncultured environmental superclusters. Finally, no *Dsr* clusters or sulfate-reducing taxa were found to be ubiquitous, which again shows how diverse and specialized SRPs can be, in terms of metabolic capacities and role in a community. The point is also supported by the variety of SRP stress coping mechanisms, defense systems, and syntrophic interactions (141) and by the reports of niche partitioning between different SRPs at the microscale in microbial mats (142) and macroscale in sediments (143). Considering all of this, one should ask what a rarefaction curve would look like for all known *Dsr* variants thus far. Are we even close to a plateau?

If such an enormous diversity of sulfate reducers exists in all environments and is hard to culture, we should rely on metatranscriptomic data. A limited number of studies have considered sulfate reduction taxa and activity based on metatranscriptomic data so far: in microbial fuel cells (144), in waters (145–147), and in artificial systems (148, 149); only a few has been soil oriented. Surprisingly, Acidobacteriota was reported as the active sulfate reducers in terrestrial ecosystems. To our knowledge, the first report of Acidobacteria contribution to the dissimilatory sulfur cycle was made by Pester et al. (13), where a minor but significant share of peatland sulfate reduction was affiliated with Acidobacteria subgroup 3. In recent years, acidobacterial sulfate reducers appeared periodically in studies of the marine environment (150, 151), while on the surface, only peatlands have shown a full set of SR genes in Acidobacteria genomes. A remarkable study of Li et al. (41) follows metatranscriptomic data obtained from revegetated acid soil near the acid mines. The authors recovered multiple high-quality metagenome-assembled genomes (MAGs) of *Deltaproteobacteria* and *Acidobacteria* lineages, which carry complete and actively expressing DSR pathways; revealed the role of viruses in their life strategy; and concluded that acidobacterial SRPs dominate over canonical SR *deltaproteobacteria* in studied sulfate-rich soils. Hausmann et al. (152) recovered MAGs of sulfite- and sulfate-respiring Acidobacteria from peatlands and confirmed the activity of DSR pathways both in soils and microcosm experiments.

Finally, MAGs of similar bacteria were obtained in seasonally thawed Sweden peatlands and was named Candidatus *Acidiflorens* (136). Although the study did not discuss a complete set of necessary genes for DSR, the MAG did have it. It later clustered with other SR acidobacteria in visualization by Flieger et al. (151). All four studies indicate usage of plant-derived oligo- or polysaccharides, namely, xylanes, by these Acidobacteria. Acidobacteria in peatlands perform a unique process of coupling complex organic matter's degradation to sulfate reduction. Besides that, a few studies were conducted. A clear increase in *Desulfobacterota* transcripts was observed for acid sulfate soils in unoxidized deep soil horizons (153). Contradictory metatranscriptomic results were reported regarding the effect of the increase in atmospheric CO<sub>2</sub> content on soil microbial communities and S cycle specifically. In the study of Padhy et al. (154), dissimilatory and assimilatory S pathways were upregulated as well as the abundance of sulfate-reducing genera, while in reference (155), sulfate reduction was significantly downregulated.

To sum up, metagenomics constantly reveals an unexpected diversity of sulfate reducers in all Earth habitats. At the same time, limited environmental transcriptomic studies show that this diversity can extend beyond the canonical SR taxa. Table 1 summarizes the data on SRP diversity in terrestrial habitats obtained with different omics approaches. More metatranscriptomic and culture-based studies are required to characterize such SRP diversity fully.

TABLE 1 SRPs that have been identified in different terrestrial ecosystems in the last 10 years and their contribution to sulfur cycling

Material	Habitat	Identified SRPs	Sulfur cycle contribution	Approach	Study
Soil	Revegetated acidic mine wasteland	Acidobacteria sb. 1, Deltaproteobacteria ( <i>Desulfovibrio</i> , Syntrophobacteraceae)	Sulfate, sulfite reduction	Shotgun metagenomics, metatranscriptomics	(41)
Peat, soil	Thawing palsas and bog	Acidobacteria, Acidobacteriaceae family, Candidatus <i>Acidifluores</i> spp.	Sulfate, sulfite reduction	Shotgun metagenomics, metatranscriptomics, metaproteomics	(136)
Soil	Acid sulfate soils	Desulfobacteria	Sulfate, sulfite, sulfur reduction	Shotgun metagenomics, 16S rRNA amplicon sequencing, metatranscriptomics	(153)
Peat, soil	Schlöppnerbrunnen II acidic peatland	<i>Desulfosporosinus</i> , <i>Syntrophobacter</i> , <i>Desulfobacca acetoxidans</i> , Acidobacteria sb. 1 and 3, Verrucomicrobia	Sulfate, sulfite reduction	Shotgun metagenomics, metatranscriptomics, stable-isotope probing	(152)
Rock and soil	Chagan-Uzun lignite open mountain pit	<i>Desulfofundulus</i> sp., <i>Thermanaeromonas</i> sp.	Sulfate, sulfite, thiosulfate reduction	Shotgun metagenomics, 16S rRNA amplicon sequencing, pure culture isolation	(110)
Soil	Soil contaminated with chloroethene near the Alameda Naval Air Station	<i>Mesotoga</i> spp.	Sulfur, thiosulfate reduction	Shotgun metagenomics	(73)
Sediments	Salt marsh sediments	<i>Desulfatitalea</i> , <i>Desulfopila</i> , <i>Desulfosarcina</i> , <i>Sulfurospirillum</i> ( <i>S. alterniflora</i> rhizosphere)	Sulfate, sulfite, sulfur reduction	16S rRNA amplicon sequencing	(96)
Soil	Wales grasslands with different degree of land use	<i>Desulfosporosinus</i> , <i>Desulfobulbus</i> , <i>Desulfobacca</i> , <i>Desulfomonile</i>	Sulfate, sulfite reduction	16S rRNA amplicon sequencing	(42)
Soil	Hainich National Park Forest soil	Desulfurellaceae	Sulfate, sulfite reduction	16S rRNA amplicon sequencing	(70)
Dried out coastal sediments	Subtropical intertidal wetland	Desulfuromonadaceae	Sulfate, sulfite, sulfur reduction	16S rRNA amplicon sequencing	(67)
Soil	South Moravia colluvial chernozems	<i>Desulfosporosinus</i> , <i>Desulfotomaculum</i> , <i>Desulfovermiculus</i>	Sulfate, sulfite reduction	16S rRNA amplicon sequencing	(62)

## WHY HAVE SULFATE-REDUCING ARCHAEA NOT BEEN DISCOVERED FROM COMMON SOILS?

Today, only four SR archaea genera are cultured: *Archaeoglobus*, *Caldivirga*, *Thermocladium* (40), and *Vulcanisaeta* (156), and none of them are from soil. All four thrive in oil deposits and hot springs. In metagenomic studies, two more sulfate-reducing candidate archaeal phyla, Hydrothermarchaeota and Aigarchaeota, were found to harbor sulfate-reducing machinery (19, 157). Also, some scarce evidence exists that Asgard archaea can be involved in the dissimilatory reduction of thiosulfate (158). Nevertheless, all mentioned archaea thrive or supposedly thrive in diverse extremophilic and/or deep ecosystems. Only one instance of Dsr-containing archaeal Thermoplasmata MAG assembled from soil metagenome was reported by Diao et al. (46). So why evidence for sulfate-reducing archaea in terrestrial sites is so scarce, almost non-existent? The answer most likely lies in evolution. Recent findings suggest that the diversity of sulfur cycle microbiota in the ancient Earth was maintained mainly through intermediate sulfur compounds such as sulfite, rather than sulfate, due to the lack of the latter in the ancient atmosphere and hydrosphere (156, 159–161). Sulfate was initially present in negligible concentrations but began to accumulate only after the occurrence of the global oxidation event (GOE) (162). Given that archaea are intrinsically adapted to chronic energetic stress (163) and that reduction of sulfate, but not sulfite, is ATP dependent (156), it is possible to speculate on the following.

Before GOE, sulfate reduction was not an essential process on early Earth (159, 164). Sulfur cycle microbiota thrived on S intermediates. Among them are sulfite- and thiosulfate-respiring species of *Pyrobaculum*, sulfur-respiring members of Crenarchaeota (165), or the recently discovered sulfite-reducing Diaforarchaea (160) and, possibly, Korarchaeota (166). On the other hand, only a limited bacterial population supposedly metabolized sulfate and could transfer their DSR genes to archaea more than once (156). Furthermore, along with the elevation of oxygen and sulfate content during the GOE, bacteria were the first to take advantage of these electron acceptors. In such an environment, no energetic stress existed for a bacterium that couples the oxidation of a substrate to the reduction of sulfate, and most archaea could not compete with it. More than that, there is evidence that around GOE, another event took place, namely, the decrease in global Ni concentrations, which favored sulfate reducers over Ni-dependent methanogens (167) and probably led to the increase in the worldwide abundance of the former. The pathways of dissimilatory SR were able to evolve extensively and did that to a much more degree in bacteria than archaea. Eventually, the diversity of sulfate-reducing bacteria expanded and moved to the surface, and the first soils began to emerge. It is generally assumed that plants colonized the land surface around 400–500 mya and were the driving force for soil formation. It is then safe to assume that in pre-Devonian times, the diversity of prokaryotes forming (along with plants and fungi) and populating the first soils was mainly composed of bacteria. Bacterial species would inevitably dictate subsequent terrestrial turnover of sulfur. Thus, it is likely that bacteria monopolized dissimilatory sulfate reduction in soils from the very beginning. The exact moment can be set even earlier regarding the abovementioned fungi-SRP associates at rock surfaces and the fact that fungi preceded plants on land. Naturally, such a model could be biased because of insufficient attention of microbiologists to sulfate-reducing archaea in non-extreme environments. It requires rigorous investigation of the sulfur cycle evolution during the start of the Palaeozoic era as well as more intense efforts toward SR-archaea isolation.

## LIFE STRATEGY OF SRPs IN SOILS WITH HIGH OR RAPIDLY CHANGING REDOX POTENTIAL

Based on the studies, it is possible to construct a picture of SRP ecology in soils. First, most SRPs thriving in aerated soils would belong to the bacterial domain. Even with the existence of aerotolerant species, we should expect that SRPs tend to fill anaerobic


microniches in soil, such as soil pores (28), soil pore water, deep horizons, and rhizospheric hotspots where most of the oxygen is readily consumed by aerobes. In these niches, SRPs are highly likely adapted to an oligotrophic, rare biosphere member strategy because the amount of sulfate ions is limited, the competition level among other species is high, and the redox potential is not always low enough. There, a typical sulfate-reducing bacterium (e.g., *Desulfitobacterium* sp., *Desulfosporosinus* sp., or *Desulfovibrio* sp.) probably maintains low-density population and assimilates products of organic matter decomposition, such as different short- and long-chain organic acids and also aromatics and molecular hydrogen. In soil matrix, SRPs will more likely be a part of a multispecies biofilm rather than planktonic form. So, it will be attached to soil particles or to plant roots if it is a part of the rhizosphere (such as species of *Desulfobulbus* and *Desulfovibrio* genera).

If the conditions change, the bacterium can sporulate or go into dormancy, depending on the taxon, facilitating the distribution of its cells across the soil environment. Different mechanisms may promote spreading, such as encapsulation by movable soil particles, flooding events that connect isolated microniches, and traversing the gut of soil invertebrates. For example, Zhou et al. (168) studied iron reduction in earthworms' gut microbial communities during the soil microcosm experiment. They found *Desulfosporosinus*, *Desulfotomaculum*, and *Desulfovibrio* to be important iron reducers in worms' intestines. Once favorable conditions return, inactive cells awaken and start to assimilate the same electron and carbon sources, as it is less likely that their range has changed. What changes more often is the spectrum of available electron acceptors. Apparently, the diversity of acceptors is responsible for the fact that SRPs mediate the remarkable diversity of soil reductive processes. According to the many observations, SRPs are part of the microbial group that reduces a variety of compounds in soil organic matrix, especially humic residues, and a variety of metal cations, especially  $\text{Fe}^{3+}$ . Numerous examples exist of SRP-reducing humic substances (169–171), and sulfate reducers such as *Desulfosporosinus* are found in humic-enriched consortia as a minor but an inevitable part (172). Moreover, several SRP genera are capable of extracellular electron transfer, such as *Desulfuromonas*, *Desulfitobacterium*, *Desulfotomaculum*, *Desulfosporosinus*, and *Desulfobulbus* (39, 144). FeS nanoparticles synthesized by sulfate reducers can facilitate the acquisition and transport of electrons from solid donors (173).

Thus, typical soil sulfate reducers appear to act as an electron relay system between accumulated small molecular weight organics and multiple electron acceptors (Fig. 2). What differs soil SRPs from that of marine and sediment origin is another mode of life, which includes (i) very limited, compared to water, concentrations of sulfate; (ii) more frequent and prolonged exposure to oxygen; and (iii) availability of unique electron donors and acceptors; and (iv) different set of possible syntrophic partners. These constraints determine the small size (down to being the “rare” biosphere members) of SRP populations, high cellular activity without proliferation, the potential role of viruses, and increased metabolic flexibility.

A much more understudied group is microbes capable of dissimilatory sulfate reduction but not constrained by oxygen or other factors due to another taxonomic position. The most prominent are acidobacteria. Almost all sulfate reducers oxidize volatile organics such as short-chain fatty acids, some can process more complex compounds, and it seems that acidobacterial members can couple sulfate reduction to plant-derived polysaccharide catabolism. It is plausible to say that peat soil sulfate reducers are active participants in diverse biomass decomposition and provide nutrients and conditions for other microorganisms. Nothing, however, is known about different lineages in an ecological sense. According to ecophysiological and metagenomic-oriented studies (19, 46), the genetic basis for sulfate reduction has such soil-occurring taxa as Verrucomicrobiota, Armatimonadota, Nitrospirata, Nitrospinota, Planctomycetota, and even Actinomycetota. The actual presence and ecological role of sulfate reduction in the metabolism of these taxa should be the focus of future studies in soil microbial ecology.





**FIG 2** Overview of terrestrial ecosystems where different SRP genera occur according to numerous studies.

## CONCLUSION AND FUTURE PERSPECTIVES

Sulfate-reducing bacteria are a diverse group of prokaryotes that mediate a remarkable number of processes in ecosystems at the global scale. Based on the studies reviewed in this manuscript, it becomes evident that SRPs actively contribute to terrestrial environments as rare soil microorganisms. The four most prevalent soil sulfate-reducing taxa are *Desulfitobacterium*, *Desulfosporosinus*, *Desulfuromonas*, and *Desulfotomaculum*. It seems that these taxa have adapted to various soil types, regardless of contaminants, aeration levels, or vegetation cover. Their ecological role in soils can be summarized as the ability to transfer the electrons from various donors to various acceptors. SRPs contribute to soil organic matter turnover and tightly connect carbon and sulfur cycles. Recent advances in metagenomic and metatranscriptomic approaches let us step forward in the study of sulfate reducers and realize that due to lateral gene transfer, the ability to reduce sulfur compounds spans way beyond previously known species.

We believe that future research in the field of SRPs should focus on the following aspects:

1. The specific mechanisms through which soil SRPs contribute to soil biogeochemical cycling and soil organic matter decomposition.
2. The habitat preferences and ecological behaviors of non-canonical sulfate reducers, such as Acidobacteriota, Verrucomicrobiota, and Planctomycetota.
3. The existence and role of sulfate-reducing archaea in temperate soil ecosystems.

The results may broaden our understanding of microbial processes involved in sulfate transformation in soil and pave the way for more efficient utilization of soil resources and bioremediation practices.

## ACKNOWLEDGMENTS

The study was carried out in the Laboratory «Soil Health» of the Southern Federal University with the financial support of the Ministry of Science and Higher Education of the Russian Federation, agreement no. 075-15-2022-1122, and in the Laboratory of Molecular Genetics of Microbial Consortia of the Southern Federal University, funded by the Strategic Academic Leadership Program of the Southern Federal University ("Priority 2030").

## AUTHOR AFFILIATION

<sup>1</sup>Southern Federal University, Rostov-on-Don, Russia

## AUTHOR ORCID*s*

Konstantin A. Demin  <http://orcid.org/0000-0003-1853-0475>

## FUNDING

Funder	Grant(s)	Author(s)
Ministry of Science and Higher Education of the Russian Federation (Minobrnauki of Russia)	075-15-2022-1122	Konstantin A. Demin Tatiana M. Minkina Andrey V. Gorovtsov
Ministry of Science and Higher Education of the Russian Federation (Minobrnauki of Russia)	Priority 2030	Konstantin A. Demin Evgeniya V. Prazdnova

## AUTHOR CONTRIBUTIONS

Konstantin A. Demin, Conceptualization, Formal analysis, Investigation, Visualization, Writing – original draft, Writing – review and editing | Evgeniya V. Prazdnova, Writing – original draft, Writing – review and editing | Tatiana M. Minkina, Funding acquisition, Project administration, Writing – review and editing | Andrey V. Gorovtsov, Supervision, Writing – original draft, Writing – review and editing

## REFERENCES

- Adam PS, Borrel G, Brochier-Armanet C, Gribaldo S. 2017. The growing tree of Archaea: new perspectives on their diversity, evolution and ecology. *ISME J* 11:2407–2425. <https://doi.org/10.1038/ismej.2017.122>
- Probst AJ, Auerbach AK, Moissl-Eichinger C. 2013. Archaea on human skin. *PLoS One* 8:e65388. <https://doi.org/10.1371/journal.pone.0065388>
- Rasheed PA, Jabbar KA, Mackey HR, Mahmoud KA. 2019. Recent advancements of nanomaterials as coatings and biocides for the inhibition of sulfate reducing bacteria induced corrosion. *Curr Opin Chem Eng* 25:35–42. <https://doi.org/10.1016/j.coche.2019.06.003>
- Tripathi AK, Thakur P, Saxena P, Rauniyar S, Gopalakrishnan V, Singh RN, Gadhamshetty V, Gnimpieba EZ, Jasthi BK, Sani RK. 2021. Gene sets and mechanisms of sulfate-reducing bacteria biofilm formation and quorum sensing with impact on corrosion. *Front Microbiol* 12:754140. <https://doi.org/10.3389/fmicb.2021.754140>
- Zan F, Hao T. 2020. Sulfate in anaerobic co-digester accelerates methane production from food waste and waste activated sludge. *Bioresour Technol* 298:122536. <https://doi.org/10.1016/j.biortech.2019.122536>
- Sato Y, Hamai T, Hori T, Aoyagi T, Inaba T, Kobayashi M, Habe H, Sakata T. 2019. *Desulfosporosinus* spp. were the most predominant sulfate-reducing bacteria in pilot-and laboratory-scale passive bioreactors for acid mine drainage treatment. *Appl Microbiol Biotechnol* 103:7783–7793. <https://doi.org/10.1007/s00253-019-10063-2>
- Muyzer G, Stams AJM. 2008. The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev Microbiol* 6:441–454. <https://doi.org/10.1038/nrmicro1892>
- Canfield DE, Farquhar J. 2009. Animal evolution, bioturbation, and the sulfate concentration of the oceans. *Proc Natl Acad Sci U S A* 106:8123–8127. <https://doi.org/10.1073/pnas.0902037106>
- Shah Walter SR, Jaekel U, Osterholz H, Fisher AT, Huber JA, Pearson A, Dittmar T, Girguis PR. 2018. Microbial decomposition of marine dissolved organic matter in cool oceanic crust. *Nature Geosci* 11:334–339. <https://doi.org/10.1038/s41561-018-0109-5>
- Burton ED, Johnston SG, Kocar BD. 2014. Arsenic mobility during flooding of contaminated soil: the effect of microbial sulfate reduction. *Environ Sci Technol* 48:13660–13667. <https://doi.org/10.1021/es503963k>
- Hansel CM, Fendorf S, Jardine PM, Francis CA. 2008. Changes in bacterial and archaeal community structure and functional diversity along a geochemically variable soil profile. *Appl Environ Microbiol* 74:1620–1633. <https://doi.org/10.1128/AEM.01787-07>

12. Pester M, Knorr K-H, Friedrich MW, Wagner M, Loy A. 2012. Sulfate-reducing microorganisms in wetlands—fameless actors in carbon cycling and climate change. *Front Microbiol* 3:72. <https://doi.org/10.3389/fmicb.2012.00072>
13. Pester M, Bittner N, Deevong P, Wagner M, Loy A. 2010. A 'rare biosphere' microorganism contributes to sulfate reduction in a peatland. *ISME J* 4:1591–1602. <https://doi.org/10.1038/ismej.2010.75>
14. Liu P, Conrad R. 2017. *Syntrophobacteraceae* - affiliated species are major propionate - degrading sulfate reducers in paddy soil. *Environ Microbiol* 19:1669–1686. <https://doi.org/10.1111/1462-2920.13698>
15. Manz W, Eisenbrecher M, Neu TR, Szewzyk U. 1998. Abundance and spatial organization of Gram-negative sulfate-reducing bacteria in activated sludge investigated by *in situ* probing with specific 16S rRNA targeted oligonucleotides. *FEMS Microbiol Ecol* 25:43–61. <https://doi.org/10.1111/j.1574-6941.1998.tb00459.x>
16. Karri S, Sierra-Alvarez R, Field JA. 2006. Toxicity of copper to acetoclastic and hydrogenotrophic activities of methanogens and sulfate reducers in anaerobic sludge. *Chemosphere* 62:121–127. <https://doi.org/10.1016/j.chemosphere.2005.04.016>
17. Zhang H, Song S, Jia Y, Wu D, Lu H. 2019. Stress-responses of activated sludge and anaerobic sulfate-reducing bacteria sludge under long-term ciprofloxacin exposure. *Water Res* 164:114964. <https://doi.org/10.1016/j.watres.2019.114964>
18. Vasconcelos C, McKenzie JA. 2000. Sulfate reducer—s—dominant players in a low-oxygen world? *Science* 290:1711–1712. <https://doi.org/10.1126/science.290.5497.1711>
19. Anantharaman K, Hausmann B, Jungbluth SP, Kantor RS, Lavy A, Warren LA, Rappé MS, Pester M, Loy A, Thomas BC, Banfield JF. 2018. Expanded diversity of microbial groups that shape the dissimilatory sulfur cycle. *ISME J* 12:1715–1728. <https://doi.org/10.1038/s41396-018-0078-0>
20. Marietou A. 2016. Nitrate reduction in sulfate-reducing bacteria. *FEMS Microbiol Lett* 363:fnw155. <https://doi.org/10.1093/femsle/fnw155>
21. Sisler FD, ZoBell CE. 1951. Nitrogen fixation by sulfate-reducing bacteria indicated by nitrogen/argon ratios. *Science* 113:511–512. <https://doi.org/10.1126/science.113.2940.511>
22. Bertics VJ, Löscher CR, Salonen I, Dale AW, Gier J, Schmitz RA, Treude T. 2013. Occurrence of benthic microbial nitrogen fixation coupled to sulfate reduction in the seasonally hypoxic Eckernförde Bay, Baltic Sea. *Biogeosciences* 10:1243–1258. <https://doi.org/10.5194/bg-10-1243-2013>
23. Sayavedra L, Li T, Bueno Batista M, Seah BKB, Booth C, Zhai Q, Chen W, Narbad A. 2021. *Desulfovibrio diazotrophicus* sp. nov., a sulfate - reducing bacterium from the human gut capable of nitrogen fixation. *Environ Microbiol* 23:3164–3181. <https://doi.org/10.1111/1462-2920.15538>
24. Lin L-H, Wang P-L, Rumble D, Lippmann-Pipke J, Boice E, Pratt LM, Sherwood Lollar B, Brodie EL, Hazen TC, Andersen GL, DeSantis TZ, Moser DP, Kershaw D, Onstott TC. 2006. Long-term sustainability of a high-energy, low-diversity crustal biome. *Science* 314:479–482. <https://doi.org/10.1126/science.1127376>
25. Kunapuli U, Jahn MK, Lueders T, Geyer R, Heipieper HJ, Meckenstock RU. 2010. *Desulfotobacterium aromaticivorans* sp. nov. and *Geobacter toluenoxydans* sp. nov., iron-reducing bacteria capable of anaerobic degradation of monoaromatic hydrocarbons. *Int J Syst Evol Microbiol* 60:686–695. <https://doi.org/10.1099/ijs.0.003525-0>
26. Musat F, Galushko A, Jacob J, Widdel F, Kube M, Reinhardt R, Wilkes H, Schink B, Rabus R. 2009. Anaerobic degradation of naphthalene and 2 - methylnaphthalene by strains of marine sulfate - reducing bacteria. *Environ Microbiol* 11:209–219. <https://doi.org/10.1111/j.1462-2920.2008.01756.x>
27. Jochum LM, Schreiber L, Marshall IPG, Jørgensen BB, Schramm A, Kjeldsen KU. 2018. Single-cell genomics reveals a diverse metabolic potential of uncultivated *Desulfatiglans*-related deltaproteobacteria widely distributed in marine sediment. *Front Microbiol* 9:2038. <https://doi.org/10.3389/fmicb.2018.02038>
28. Dhar K, Panneerselvan L, Subashchandrabose SR, Venkateswarlu K, Megharaj M. 2023. Anaerobic degradation of naphthalene and pyrene by sulfate-reducing cultures enriched from former manufactured gas plant soil. *Microb Ecol* 86:271–281. <https://doi.org/10.1007/s00248-022-02042-4>
29. Qian Z, Tianwei H, Mackey HR, van Loosdrecht MCM, Guanghao C. 2019. Recent advances in dissimilatory sulfate reduction: from metabolic study to application. *Water Res* 150:162–181. <https://doi.org/10.1016/j.watres.2018.11.018>
30. Hockin SL, Gadd GM. 2007. Bioremediation of metals and metalloids by precipitation and cellular binding, p 405–434. In *Sulphate-reducing bacteria: environmental and engineered systems*
31. Li J, Zhao W, Du H, Guan Y, Ma M, Rennenberg H. 2022. The symbiotic system of sulfate-reducing bacteria and clay-sized fraction of purplish soil strengthens cadmium fixation through iron-bearing minerals. *Sci Total Environ* 820:153253. <https://doi.org/10.1016/j.scitotenv.2022.153253>
32. Labrenz M, Banfield JF. 2004. Sulfate-reducing bacteria-dominated biofilms that precipitate ZnS in a subsurface circumneutral-pH mine drainage system. *Microb Ecol* 47:205–217. <https://doi.org/10.1007/s00248-003-1025-8>
33. Castillo J, Pérez-López R, Caraballo MA, Nieto JM, Martins M, Costa MC, Olías M, Cerón JC, Tucoulou R. 2012. Biologically-induced precipitation of sphalerite-wurtzite nanoparticles by sulfate-reducing bacteria: implications for acid mine drainage treatment. *Sci Total Environ* 423:176–184. <https://doi.org/10.1016/j.scitotenv.2012.02.013>
34. Peltier E, Ilipilla P, Fowle D. 2011. Structure and reactivity of zinc sulfide precipitates formed in the presence of sulfate-reducing bacteria. *Appl Geochem* 26:1673–1680. <https://doi.org/10.1016/j.apgeochem.2011.04.024>
35. Kang S, Van Nostrand JD, Gough HL, He Z, Hazen TC, Stahl DA, Zhou J. 2013. Functional gene array-based analysis of microbial communities in heavy metals-contaminated lake sediments. *FEMS Microbiol Ecol* 86:200–214. <https://doi.org/10.1111/1574-6941.12152>
36. Lovley DR, Widman PK, Woodward JC, Phillips EJ. 1993. Reduction of uranium by cytochrome c3 of *Desulfovibrio vulgaris*. *Appl Environ Microbiol* 59:3572–3576. <https://doi.org/10.1128/aem.59.11.3572-3576.1993>
37. Bagwell CE, Liu X, Wu L, Zhou J. 2006. Effects of legacy nuclear waste on the compositional diversity and distributions of sulfate-reducing bacteria in a terrestrial subsurface aquifer. *FEMS Microbiol Ecol* 55:424–431. <https://doi.org/10.1111/j.1574-6941.2005.00039.x>
38. Barton LL, Torres FAT, Xu H, Zocco TG. 2015. Metabolism of metals and metalloids by the sulfate-reducing bacteria, p 57–83. In *Bacteria-metal interactions*
39. Lovley DR, Holmes DE. 2022. Electromicrobiology: the ecophysiology of phylogenetically diverse electroactive microorganisms. *Nat Rev Microbiol* 20:5–19. <https://doi.org/10.1038/s41579-021-00597-6>
40. Barton LL, Fauque GD. 2022. Sulfate-reducing bacteria and archaea. Springer, Cham.
41. Li J-T, Jia P, Wang X-J, Ou S-N, Yang T-T, Feng S-W, Lu J-L, Fang Z, Liu J, Liao B, Shu W-S, Liang J-L. 2022. Metagenomic and metatranscriptomic insights into sulfate-reducing bacteria in a revegetated acidic mine wasteland. *NPJ Biofilms Microbiomes* 8:71. <https://doi.org/10.1038/s41522-022-00333-9>
42. George PBL, Coelho KP, Creer S, Lebron I, Robinson DA, Jones DL. 2021. Decoupled richness of generalist anaerobes and sulphate - reducing bacteria is driven by pH across land uses in temperate soils. *Eur J Soil Sci* 72:2445–2456. <https://doi.org/10.1111/ejss.13040>
43. Barton LL, Duarte AG, Staicu LC. 2023. Genomic insight into iron acquisition by sulfate-reducing bacteria in microaerophilic environments. *Biomaterials* 36:339–350. <https://doi.org/10.1007/s10534-022-00410-8>
44. Gao P, Zhang X, Huang X, Chen Z, Marietou A, Holmkvist L, Qu L, Finster K, Gong X. 2023. Genomic insight of sulfate reducing bacterial genus *Desulfotomaculum* reveals their metabolic versatility in biogeochemical cycling. *BMC Genomics* 24:209. <https://doi.org/10.1186/s12864-023-09297-2>
45. Gibson GR. 1990. Physiology and ecology of the sulphate-reducing bacteria. *J Appl Bacteriol* 69:769–797. <https://doi.org/10.1111/j.1365-2672.1990.tb01575.x>
46. Diao M, Dykstra S, Koeksoy E, Ngugi DK, Anantharaman K, Loy A, Pester M. 2023. Global diversity and inferred ecophysiology of microorganisms with the potential for dissimilatory sulfate/sulfite reduction. *FEMS Microbiol Rev* 47:fuad058. <https://doi.org/10.1093/femsre/fuad058>

47. Lanthier M, Villemur R, Lépine F, Bisaillon J-G, Beaudet R. 2001. Geographic distribution of *Desulfotobacterium frappieri* PCP-1 and *Desulfotobacterium* spp. in soils from the province of Quebec, Canada. *FEMS Microbiol Ecol* 36:185–191. <https://doi.org/10.1111/j.1574-6941.2001.tb00839.x>
48. Sitte J, Akob DM, Kaufmann C, Finster K, Banerjee D, Burkhardt E-M, Kostka JE, Scheinost AC, Büchel G, Küsel K. 2010. Microbial links between sulfate reduction and metal retention in uranium-and heavy metal-contaminated soil. *Appl Environ Microbiol* 76:3143–3152. <https://doi.org/10.1128/AEM.00051-10>
49. Li M, Yao J, Sunahara G, Hawari J, Duran R, Liu J, Liu B, Cao Y, Pang W, Li H, Li Y, Ruan Z. 2022. Novel microbial consortia facilitate metalliferous immobilization in non-ferrous metal (loid) s contaminated smelter soil: efficiency and mechanisms. *Environ Pollut* 313:120042. <https://doi.org/10.1016/j.envpol.2022.120042>
50. Yan X, Gao B, Wang J, Zhu X, Zhang M. 2023. Insights into remediation effects and bacterial diversity of different remediation measures in rare earth mine soil with SO<sub>4</sub><sup>2-</sup> and heavy metals. *Front Microbiol* 14:1050635. <https://doi.org/10.3389/fmicb.2023.1050635>
51. Burkhardt E-M, Bischoff S, Akob DM, Büchel G, Küsel K. 2011. Heavy metal tolerance of Fe (III)-reducing microbial communities in contaminated creek bank soils. *Appl Environ Microbiol* 77:3132–3136. <https://doi.org/10.1128/AEM.02085-10>
52. Falk N, Chaganti SR, Weisener CG. 2018. Evaluating the microbial community and gene regulation involved in crystallization kinetics of ZnS formation in reduced environments. *Geochim Cosmochim Acta* 220:201–216. <https://doi.org/10.1016/j.gca.2017.09.039>
53. Mingo FS, Diekert G, Studenik S. 2016. Enrichment of *Desulfotobacterium* spp. from forest and grassland soil using the O-demethylation of phenyl methyl ethers as a growth-selective process. *Microbiology (Reading)* 162:224–235. <https://doi.org/10.1099/mic.0.000218>
54. He W-J, Shi M-M, Yang P, Huang T, Yuan Q-S, Yi S-Y, Wu A-B, Li H-P, Gao C-B, Zhang J-B, Liao Y-C. 2020. Novel soil bacterium strain *Desulfotobacterium* sp. PGC-3-9 detoxifies trichothecene mycotoxins in wheat via de-epoxidation under aerobic and anaerobic conditions. *Toxins (Basel)* 12:363. <https://doi.org/10.3390/toxins12060363>
55. Li Y, Yang G, Yao S, Zhuang L. 2021. *Paradesulfotobacterium ferrireducens* gen. nov., sp. nov., a Fe (III)-reducing bacterium from petroleum-contaminated soil and reclassification of *Desulfotobacterium aromaticivorans* as *Paradesulfotobacterium aromaticivorans* comb. nov. *Int J Syst Evol Microbiol* 71:005025. <https://doi.org/10.1099/ijsem.0.005025>
56. Huang Y, Dai Z, Lin J, Qi Q, Luo Y, Dahlgren RA, Xu J. 2021. Contrasting effects of carbon source recalcitrance on soil phosphorus availability and communities of phosphorus solubilizing microorganisms. *J Environ Manage* 298:113426. <https://doi.org/10.1016/j.jenvman.2021.113426>
57. Chauhan A, Layton AC, Vishnivetskaya TA, Williams D, Pfiffner SM, Rekepalli B, Stackhouse B, Lau MCY, Phelps TJ, Myktyczuk N, Ronholm J, Whyte L, Onstott TC, Saylor GS. 2014. Metagenomes from thawing low-soil-organic-carbon mineral cryosols and permafrost of the Canadian high Arctic. *Genome Announc* 2:10–1128. <https://doi.org/10.1128/genomeA.01217-14>
58. Chen S, Zhang C, Qiu L, Li Q, Zhang K, Luo H. 2022. Biogeochemical transformation of sulfur and its effects on arsenic mobility in paddy fields polluted by acid mine drainage. *Chemosphere* 293:133605. <https://doi.org/10.1016/j.chemosphere.2022.133605>
59. Genovese M, Denaro R, Cappello S, Di Marco G, La Spada G, Giuliano L, Genovese L, Yakimov MM. 2008. Bioremediation of benzene, toluene, ethylbenzene, xylenes - contaminated soil: a biopile pilot experiment. *J Appl Microbiol* 105:1694–1702. <https://doi.org/10.1111/j.1365-2672.2008.03897.x>
60. Mangse G, Werner D, Meynet P, Ogbaga CC. 2020. Microbial community responses to different volatile petroleum hydrocarbon class mixtures in an aerobic sandy soil. *Environ Pollut* 264:114738. <https://doi.org/10.1016/j.envpol.2020.114738>
61. Gagliano AL, Tagliavia M, D'Alessandro W, Franzetti A, Parello F, Quatrini P. 2016. So close, so different: geothermal flux shapes divergent soil microbial communities at neighbouring sites. *Geobiology* 14:150–162. <https://doi.org/10.1111/gbi.12167>
62. Sagova-Mareckova M, Zadorova T, Penizek V, Omelka M, Tejnecky V, Pruchova P, Chuman T, Drabek O, Buresova A, Vanek A, Kopecky J. 2016. The structure of bacterial communities along two vertical profiles of a deep colluvial soil. *Soil Biol Biochem* 101:65–73. <https://doi.org/10.1016/j.soilbio.2016.06.026>
63. Miceli JF, Parameswaran P, Kang D-W, Krajmalnik-Brown R, Torres CI. 2012. Enrichment and analysis of anode-respiring bacteria from diverse anaerobic inocula. *Environ Sci Technol* 46:10349–10355. <https://doi.org/10.1021/es301902h>
64. Ramos LR, Vollú RE, Jurelevicius D, Rosado AS, Seldin L. 2019. Firmicutes in different soils of Admiralty Bay, King George Island, Antarctica. *Polar Biol* 42:2219–2226. <https://doi.org/10.1007/s00300-019-02596-z>
65. Kim O-S, Chae N, Lim HS, Cho A, Kim JH, Hong SG, Oh J. 2012. Bacterial diversity in ornithogenic soils compared to mineral soils on King George Island, Antarctica. *J Microbiol* 50:1081–1085. <https://doi.org/10.1007/s12275-012-2655-7>
66. Denger K, Stackebrandt E, Cook AM. 1999. *Desulfonisporea thiosulfatigenes* gen. nov., sp. nov., a taurine-fermenting, thiosulfate-producing anaerobic bacterium. *Int J Syst Bacteriol* 49:1599–1603. <https://doi.org/10.1099/00207713-49-4-1599>
67. Cheung MK, Wong CK, Chu KH, Kwan HS. 2018. Community structure, dynamics and interactions of bacteria, archaea and fungi in subtropical coastal wetland sediments. *Sci Rep* 8:14397. <https://doi.org/10.1038/s41598-018-32529-5>
68. Pradhan S, Srinivas TNR, Pindi PK, Kishore KH, Begum Z, Singh PK, Singh AK, Pratibha MS, Yasala AK, Reddy GSN, Shivaji S. 2010. Bacterial biodiversity from Roopkund glacier, Himalayan mountain ranges, India. *Extremophiles* 14:377–395. <https://doi.org/10.1007/s00792-010-0318-3>
69. Wang W, Wang H, Feng Y, Wang L, Xiao X, Xi Y, Luo X, Sun R, Ye X, Huang Y, Zhang Z, Cui Z. 2016. Consistent responses of the microbial community structure to organic farming along the middle and lower reaches of the Yangtze River. *Sci Rep* 6:35046. <https://doi.org/10.1038/srep35046>
70. Dukunde A, Schneider D, Schmidt M, Veldkamp E, Daniel R. 2019. Tree species shape soil bacterial community structure and function in temperate deciduous forests. *Front Microbiol* 10:1519. <https://doi.org/10.3389/fmicb.2019.01519>
71. Miletto M, Loeb R, Antheunisse AM, Bodelier PLE, Laanbroek HJ. 2010. Response of the sulfate-reducing community to the re-establishment of estuarine conditions in two contrasting soils: a mesocosm approach. *Microb Ecol* 59:109–120. <https://doi.org/10.1007/s00248-009-9614-9>
72. Nesbø CL, Dlutek M, Zhaxybayeva O, Doolittle WF. 2006. Evidence for existence of mesotogas, members of the order *Thermotogales* adapted to low-temperature environments. *Appl Environ Microbiol* 72:5061–5068. <https://doi.org/10.1128/AEM.00342-06>
73. Nesbø CL, Charchuk R, Pollo SMJ, Budwill K, Kublanov IV, Haverkamp THA, Foght J. 2019. Genomic analysis of the mesophilic *Thermotogae* genus *Mesotoga* reveals phylogeographic structure and genomic determinants of its distinct metabolism. *Environ Microbiol* 21:456–470. <https://doi.org/10.1111/1462-2920.14477>
74. Pedersen K, Ekendahl S. 1992. Incorporation of CO<sub>2</sub> and introduced organic compounds by bacterial populations in groundwater from the deep crystalline bedrock of the Stripa mine. *J Gen Microbiol* 138:369–376. <https://doi.org/10.1099/00221287-138-2-369>
75. Pedersen K. 1996. Investigations of subterranean bacteria in deep crystalline bedrock and their importance for the disposal of nuclear waste. *Can J Microbiol* 42:382–391. <https://doi.org/10.1139/m96-054>
76. Pedersen K. 1997. Microbial life in deep granitic rock. *FEMS Microbiol Rev* 20:399–414. [https://doi.org/10.1016/S0168-6445\(97\)00022-3](https://doi.org/10.1016/S0168-6445(97)00022-3)
77. Drake H, Ivarsson M, Bengtson S, Heim C, Siljeström S, Whitehouse MJ, Broman C, Belivanova V, Åström ME. 2017. Anaerobic consortia of fungi and sulfate reducing bacteria in deep granite fractures. *Nat Commun* 8:55. <https://doi.org/10.1038/s41467-017-00094-6>
78. Drake H, Whitehouse MJ, Heim C, Reiners PW, Tillberg M, Hogmalm KJ, Dopson M, Broman C, Åström ME. 2018. Unprecedented 34S - enrichment of pyrite formed following microbial sulfate reduction in fractured crystalline rocks. *Geobiology* 16:556–574. <https://doi.org/10.1111/gbi.12297>
79. Bomberg M, Miettinen H, Kietäväinen R, Purkamo L, Ahonen L, Vikman M. 2021. Microbial metabolic potential in deep crystalline bedrock, p 41–70. In *The microbiology of nuclear waste disposal*. Elsevier.
80. Purkamo L, Bomberg M, Nyyssönen M, Kukkonen I, Ahonen L, Kietäväinen R, Itävaara M. 2013. Dissecting the deep biosphere: retrieving authentic microbial communities from packer-isolated deep



- crystalline bedrock fracture zones. *FEMS Microbiol Ecol* 85:324–337. <https://doi.org/10.1111/1574-6941.12126>
81. Pedersen K. 1999. Subterranean microorganisms and radioactive waste disposal in Sweden. *Eng Geol* 52:163–176. [https://doi.org/10.1016/S0013-7952\(99\)00004-6](https://doi.org/10.1016/S0013-7952(99)00004-6)
  82. Chi Fru E, Athar R. 2008. *In situ* bacterial colonization of compacted bentonite under deep geological high-level radioactive waste repository conditions. *Appl Microbiol Biotechnol* 79:499–510. <https://doi.org/10.1007/s00253-008-1436-z>
  83. Haynes HM, Pearce CI, Boothman C, Lloyd JR. 2018. Response of bentonite microbial communities to stresses relevant to geodisposal of radioactive waste. *Chem Geol* 501:58–67. <https://doi.org/10.1016/j.chemgeo.2018.10.004>
  84. Safonov AV, Boguslavsky AE, Gaskova OL, Boldyrev KA, Shvartseva OS, Khvashchevskaya AA, Popova NM. 2021. Biogeochemical modelling of uranium immobilization and aquifer remediation strategies near NCCP sludge storage facilities. *Appl Sci* 11:2875. <https://doi.org/10.3390/app11062875>
  85. Krivonogov S, Maltsev A, Zelenina D, Safonov A. 2023. Microbial diversity and authigenic mineral formation of modern bottom sediments in the littoral zone of Lake Issyk-Kul, Kyrgyz Republic (Central Asia). *Biology* 12:642. <https://doi.org/10.3390/biology12050642>
  86. Kudo A, Zheng J, Cayer I, Fujikawa Y, Asano H, Arai K, Yoshikawa H, Ito M. 1996. Behaviour of plutonium interacting with bentonite and sulfate-reducing anaerobic bacteria. *MRS Proc* 465:465. <https://doi.org/10.1557/PROC-465-879>
  87. Maanoja S, Lakaniemi A-M, Lehtinen L, Salminen L, Auvinen H, Kokko M, Palmroth M, Muuri E, Rintala J. 2020. Compacted bentonite as a source of substrates for sulfate-reducing microorganisms in a simulated excavation-damaged zone of a spent nuclear fuel repository. *Appl Clay Sci* 196:105746. <https://doi.org/10.1016/j.clay.2020.105746>
  88. Wang H, Guo CL, Yang CF, Lu GN, Chen MQ, Dang Z. 2016. Distribution and diversity of bacterial communities and sulphate-reducing bacteria in a paddy soil irrigated with acid mine drainage. *J Appl Microbiol* 121:196–206. <https://doi.org/10.1111/jam.13143>
  89. Liu J, Wan Y, Wei X, She J, Ouyang Q, Deng P, Hu H, Zhang X, Fang M, Wei X, Liu W, Gong J, Wang J. 2023. Microbial diversity in paddy rhizospheric soils around a large industrial thallium-containing sulfide utilization zone. *Environ Res* 216:114627. <https://doi.org/10.1016/j.envres.2022.114627>
  90. Xiao W, Zhang Q, Zhao S, Chen D, Gao N, Huang M, Ye X. 2023. Citric acid secretion from rice roots contributes to reduction and immobilization of Cr (VI) by driving microbial sulfur and iron cycle in paddy soil. *Sci Total Environ* 854:158832. <https://doi.org/10.1016/j.scitotenv.2022.158832>
  91. Zhang Q, Chen H-F, Huang D-Y, Guo X-B, Xu C, Zhu H-H, Li B, Liu T-T, Feng R-W, Zhu Q-H. 2022. Sulfur fertilization integrated with soil redox conditions reduces Cd accumulation in rice through microbial induced Cd immobilization. *Sci Total Environ* 824:153868. <https://doi.org/10.1016/j.scitotenv.2022.153868>
  92. Huang L, Wang X, Chi Y, Huang L, Li WC, Ye Z. 2021. Rhizosphere bacterial community composition affects cadmium and arsenic accumulation in rice (*Oryza sativa* L.). *Ecotoxicol Environ Saf* 222:112474. <https://doi.org/10.1016/j.ecoenv.2021.112474>
  93. Shi W, Song W, Zheng J, Luo Y, Qile G, Lü S, Lü X, Zhou B, Lü C, He J. 2021. Factors and pathways regulating the release and transformation of arsenic mediated by reduction processes of dissimilated iron and sulfate. *Sci Total Environ* 768:144697. <https://doi.org/10.1016/j.scitotenv.2020.144697>
  94. Xu X, Wang P, Zhang J, Chen C, Wang Z, Kopittke PM, Kretschmar R, Zhao F-J. 2019. Microbial sulfate reduction decreases arsenic mobilization in flooded paddy soils with high potential for microbial Fe reduction. *Environ Pollu* 251:952–960. <https://doi.org/10.1016/j.envpol.2019.05.086>
  95. An X, Wang Z, Teng X, Zhou R, Wang X, Xu M, Lian B. 2022. Rhizosphere bacterial diversity and environmental function prediction of wild salt-tolerant plants in coastal silt soil. *Ecol Indic* 134:108503. <https://doi.org/10.1016/j.ecolind.2021.108503>
  96. Rolando JL, Kolton M, Song T, Kostka JE. 2022. The core root microbiome of *Spartina alterniflora* is predominated by sulfur-oxidizing and sulfate-reducing bacteria in Georgia salt marshes, USA. *Microbiome* 10:37. <https://doi.org/10.1186/s40168-021-01187-7>
  97. Higashioka Y, Kojima H, Watanabe T, Fukui M. 2015. Draft genome sequence of *Desulfatitalea tepidiphila* S28bFT. *Genome Announc* 3:e01326-15. <https://doi.org/10.1128/genomeA.01326-15>
  98. Thajudeen J, Yousuf J, Veetil VP, Varghese S, Singh A, Abdulla MH. 2017. Nitrogen fixing bacterial diversity in a tropical estuarine sediments. *World J Microbiol Biotechnol* 33:41. <https://doi.org/10.1007/s11274-017-2205-x>
  99. Sidorenko OD. 1989. Role of sulphate-reducing bacteria in rhizosphere of rice and flooded soil in rice fields, p 349–354. In *Developments in soil science*. Vol. 18. Elsevier.
  100. Lin H, Shi J, Chen X, Yang J, Chen Y, Zhao Y, Hu T. 2010. Effects of lead upon the actions of sulfate-reducing bacteria in the rice rhizosphere. *Soil Biol Biochem* 42:1038–1044. <https://doi.org/10.1016/j.soilbio.2010.02.023>
  101. Baumgartner LK, Reid RP, Dupraz C, Decho AW, Buckley DH, Spear JR, Przekop KM, Visscher PT. 2006. Sulfate reducing bacteria in microbial mats: changing paradigms, new discoveries. *Sediment Geol* 185:131–145. <https://doi.org/10.1016/j.sedgeo.2005.12.008>
  102. Dolla A, Fournier M, Dermoun Z. 2006. Oxygen defense in sulfate-reducing bacteria. *J Biotechnol* 126:87–100. <https://doi.org/10.1016/j.jbiotec.2006.03.041>
  103. Berndt WL, Vargas JM. 2006. Dissimilatory reduction of sulfate in black layer. *HortSci* 41:815–817. <https://doi.org/10.21273/HORTSCI.41.3.815>
  104. Uroz S, Buée M, Murat C, Frey-Klett P, Martin F. 2010. Pyrosequencing reveals a contrasted bacterial diversity between oak rhizosphere and surrounding soil. *Environ Microbiol Rep* 2:281–288. <https://doi.org/10.1111/j.1758-2229.2009.00117.x>
  105. Li X, Li Y, Zhu X, Gui X, Ma C, Peng W, Li Y, Zhang Y, Huang W, Hua D, Jia S, Wu M. 2022. Evaluation of the cadmium phytoextraction potential of tobacco (*Nicotiana tabacum*) and rhizosphere micro-characteristics under different cadmium levels. *Chemosphere* 286:131714. <https://doi.org/10.1016/j.chemosphere.2021.131714>
  106. Amora-Lazcano E, Azcón R. 1997. Response of sulphur cycling microorganisms to arbuscular mycorrhizal fungi in the rhizosphere of maize. *Appl Soil Ecol* 6:217–222. [https://doi.org/10.1016/S0929-1393\(97\)00011-5](https://doi.org/10.1016/S0929-1393(97)00011-5)
  107. Hausmann B, Knorr K-H, Schreck K, Tringe SG, Glavina Del Rio T, Loy A, Pester M. 2016. Consortia of low-abundance bacteria drive sulfate reduction-dependent degradation of fermentation products in peat soil microcosms. *ISME J* 10:2365–2375. <https://doi.org/10.1038/ismej.2016.42>
  108. Hausmann B, Pelikan C, Rattei T, Loy A, Pester M. 2019. Long-term transcriptional activity at zero growth of a cosmopolitan rare biosphere member. *mBio* 10:e02189-18. <https://doi.org/10.1128/mBio.02189-18>
  109. Shade A, Hogan CS, Klimowicz AK, Linske M, McManus PS, Handelsman J. 2012. Culturing captures members of the soil rare biosphere. *Environ Microbiol* 14:2247–2252. <https://doi.org/10.1111/j.1462-2920.2012.02817.x>
  110. Karnachuk OV, Rusanov II, Panova IA, Kadnikov VV, Avakyan MR, Ikkert OP, Lukina AP, Beletsky AV, Mardanov AV, Knyazev YV, Volochaev MN, Pimenov NV, Ravin NV. 2023. The low-temperature germinating spores of the thermophilic *Desulfofundulus* contribute to an extremely high sulfate reduction in burning coal seams. *Front Microbiol* 14:1204102. <https://doi.org/10.3389/fmicb.2023.1204102>
  111. Orellana R, Arancibia A, Badilla L, Acosta J, Arancibia G, Escar R, Ferrara G, Seeger M. 2021. Ecophysiological features shape the distribution of prophages and CRISPR in sulfate reducing prokaryotes. *Microorganisms* 9:931. <https://doi.org/10.3390/microorganisms9050931>
  112. Wang L, Wang Y, Huang X, Ma R, Li J, Wang F, Jiao N, Zhang R. 2022. Potential metabolic and genetic interaction among viruses, methanogen and methanotrophic archaea, and their syntrophic partners. *ISME Commun* 2:50. <https://doi.org/10.1038/s43705-022-00135-2>
  113. Dalcin Martins P, Danczak RE, Roux S, Frank J, Borton MA, Wolfe RA, Burris MN, Wilkins MJ. 2018. Viral and metabolic controls on high rates of microbial sulfur and carbon cycling in wetland ecosystems. *Microbiome* 6:138. <https://doi.org/10.1186/s40168-018-0522-4>
  114. Ji M, Fan X, Cornell CR, Zhang Y, Yuan MM, Tian Z, Sun K, Gao R, Liu Y, Zhou J. 2023. Tundra soil viruses mediate responses of microbial communities to climate warming. *mBio* 14:e0300922. <https://doi.org/10.1128/mbio.03009-22>



115. Gadd GM. 2017. Geomicrobiology of the built environment. *Nat Microbiol* 2:16275. <https://doi.org/10.1038/nmicrobiol.2016.275>
116. Nuppenen-Puputti M, Kietäväinen R, Purkamo L, Rajala P, Itävaara M, Kukkonen I, Bomberg M. 2020. Rock surface fungi in deep continental biosphere—exploration of microbial community formation with subsurface *in situ* biofilm trap. *Microorganisms* 9:64. <https://doi.org/10.3390/microorganisms9010064>
117. Edgcomb VP, Teske AP, Mara P. 2022. Microbial hydrocarbon degradation in Guaymas Basin—exploring the roles and potential interactions of fungi and sulfate-reducing bacteria. *Front Microbiol* 13:831828. <https://doi.org/10.3389/fmicb.2022.831828>
118. Biebl H, Pfennig N. 1978. Growth yields of green sulfur bacteria in mixed cultures with sulfur and sulfate reducing bacteria. *Arch Microbiol* 117:9–16. <https://doi.org/10.1007/BF00689344>
119. Panova IA, Rusanov II, Kadnikov VV, Latygolets EA, Avakyan MR, Ivanov MV, Zyusman VC, Kovaleva AA, Ravin NV, Pimenov NV, Karnachuk OV. 2020. Sulfate reduction in underground horizons of a flooded coal mine in Kuzbass. *Microbiology* 89:542–550. <https://doi.org/10.1134/S0026261720050185>
120. Fuseler K, Krekeler D, Sydow U, Cypionka H. 1996. A common pathway of sulfide oxidation by sulfate-reducing bacteria. *FEMS Microbiol Lett* 144:129–134. <https://doi.org/10.1111/j.1574-6968.1996.tb08518.x>
121. Thorup C, Schramm A, Findlay AJ, Finster KW, Schreiber L. 2017. Disguised as a sulfate reducer: growth of the deltaproteobacterium *Desulfurivibrio alkaliphilus* by sulfide oxidation with nitrate. *mBio* 8:10–1128. <https://doi.org/10.1128/mBio.00671-17>
122. Fukui M, Teske A, Assmus B, Muyzer G, Widdel F. 1999. Physiology, phylogenetic relationships, and ecology of filamentous sulfate-reducing bacteria (genus *Desulfonema*). *Arch Microbiol* 172:193–203. <https://doi.org/10.1007/s002030050760>
123. Orsi WD, Barker Jørgensen B, Biddle JF. 2016. Transcriptional analysis of sulfate reducing and chemolithoautotrophic sulfur oxidizing bacteria in the deep seafloor. *Environ Microbiol Rep* 8:452–460. <https://doi.org/10.1111/1758-2229.12387>
124. Tare V, Sabumon PC. 1995. Application of sulfate-reducing and sulfide-oxidizing bacterial symbiosis for wastewater treatment. *Water Qual Res J* 30:305–324. <https://doi.org/10.2166/wqrj.1995.029>
125. Yamamoto-Ikemoto R, Matsui S, Komori T, Bosque-Hamilton E. 1998. Interactions between filamentous sulfur bacteria, sulfate reducing bacteria and poly-P accumulating bacteria in anaerobic-oxic activated sludge from a municipal plant. *Water Sci Technol* 37:599–603. <https://doi.org/10.2166/wst.1998.0725>
126. Meyer DD, de Andrade PAM, Durrer A, Andreote FD, Corção G, Brandelli A. 2016. Bacterial communities involved in sulfur transformations in wastewater treatment plants. *Appl Microbiol Biotechnol* 100:10125–10135. <https://doi.org/10.1007/s00253-016-7839-3>
127. Wang R, Xu S, Jiang C, Zhang Y, Bai N, Zhuang G, Bai Z, Zhuang X. 2019. Impacts of human activities on the composition and abundance of sulfate-reducing and sulfur-oxidizing microorganisms in polluted river sediments. *Front Microbiol* 10:231. <https://doi.org/10.3389/fmicb.2019.00231>
128. Ende FP, Meier J, Gernerden H. 1997. Syntrophic growth of sulfate-reducing bacteria and colorless sulfur bacteria during oxygen limitation. *FEMS Microbiol Ecol* 23:65–80. <https://doi.org/10.1111/j.1574-6941.1997.tb00392.x>
129. Liao P, Kim C, Saxton MA, Malkin SY. 2022. Microbial succession in a marine sediment: inferring interspecific microbial interactions with marine cable bacteria. *Environ Microbiol* 24:6348–6364. <https://doi.org/10.1111/1462-2920.16230>
130. Xia FF, Su Y, Wei XM, He YH, Wu ZC, Ghulam A, He R. 2014. Diversity and activity of sulphur - oxidizing bacteria and sulphate - reducing bacteria in landfill cover soils. *Lett Appl Microbiol* 59:26–34. <https://doi.org/10.1111/lam.12240>
131. Jiang L, Cai C, Zhang Y, Mao S, Sun Y, Li K, Xiang L, Zhang C. 2012. Lipids of sulfate-reducing bacteria and sulfur-oxidizing bacteria found in the Dongsheng uranium deposit. *Chin Sci Bull* 57:1311–1319. <https://doi.org/10.1007/s11434-011-4955-4>
132. Plugge CM, Zhang W, Scholten JCM, Stams AJM. 2011. Metabolic flexibility of sulfate-reducing bacteria. *Front Microbiol* 2:81. <https://doi.org/10.3389/fmicb.2011.00081>
133. Zhao R, Biddle JF. 2021. Helarchaeota and co-occurring sulfate-reducing bacteria in seafloor sediments from the Costa Rica Margin. *ISME Commun* 1:25. <https://doi.org/10.1038/s43705-021-00027-x>
134. Krukenberg V, Harding K, Richter M, Glöckner FO, Gruber-Vodicka HR, Adam B, Berg JS, Knittel K, Tegetmeyer HE, Boetius A, Wegener G. 2016. *Candidatus Desulfoferriplus auxilii*, a hydrogenotrophic sulfate-reducing bacterium involved in the thermophilic anaerobic oxidation of methane. *Environ Microbiol* 18:3073–3091. <https://doi.org/10.1111/1462-2920.13283>
135. Sedano-Núñez VT, Boeren S, Stams AJM, Plugge CM. 2018. Comparative proteome analysis of propionate degradation by *Syntrophobacter fumaroxidans* in pure culture and in coculture with methanogens. *Environ Microbiol* 20:1842–1856. <https://doi.org/10.1111/1462-2920.14119>
136. Woodcroft BJ, Singleton CM, Boyd JA, Evans PN, Emerson JB, Zayed AAF, Hoelzle RD, Lamberton TO, McCalley CK, Hodgkins SB, Wilson RM, Purvine SO, Nicora CD, Li C, Frolking S, Chanton JP, Crill PM, Saleska SR, Rich VI, Tyson GW. 2018. Genome-centric view of carbon processing in thawing permafrost. *Nature* 560:49–54. <https://doi.org/10.1038/s41586-018-0338-1>
137. Fadhlou K, Ben Hania W, Armougom F, Bartoli M, Fardeau M-L, Erauso G, Brasseur G, Aubert C, Hamdi M, Brochier-Armanet C, Dolla A, Ollivier B. 2018. Obligate sugar oxidation in *Mesotoga* spp., phylum *Thermotogae*, in the presence of either elemental sulfur or hydrogenotrophic sulfate - reducers as electron acceptor. *Environ Microbiol* 20:281–292. <https://doi.org/10.1111/1462-2920.13995>
138. Steger D, Wentrup C, Braunecker C, Deevong P, Hofer M, Richter A, Baranyi C, Pester M, Wagner M, Loy A. 2011. Microorganisms with novel dissimilatory (bi) sulfite reductase genes are widespread and part of the core microbiota in low-sulfate peatlands. *Appl Environ Microbiol* 77:1231–1242. <https://doi.org/10.1128/AEM.01352-10>
139. Zhu X, Chen L, Pan H, Wang L, Zhang X, Wang D. 2022. Diversity and biogenesis contribution of sulfate-reducing bacteria in arsenic-contaminated soils from realgar deposits. *Environ Sci Pollut Res Int* 29:31110–31120. <https://doi.org/10.1007/s11356-022-18595-3>
140. Vigneron A, Cruaud P, Alsop E, de Rezende JR, Head IM, Tsensmetz N. 2018. Beyond the tip of the iceberg; a new view of the diversity of sulfite- and sulfate-reducing microorganisms. *ISME J* 12:2096–2099. <https://doi.org/10.1038/s41396-018-0155-4>
141. Zhou J, He Q, Hemme CL, Mukhopadhyay A, Hillesland K, Zhou A, He Z, Van Nostrand JD, Hazen TC, Stahl DA, Wall JD, Arkin AP. 2011. How sulphate-reducing microorganisms cope with stress: lessons from systems biology. *Nat Rev Microbiol* 9:452–466. <https://doi.org/10.1038/nrmicro2575>
142. Joshua MM. 2017. Genomic and transcriptomic evidence for niche partitioning among sulfate-reducing bacteria in redox-stratified cyanobacterial mats of the Middle Island Sinkhole
143. Marshall IPG, Ren G, Jaussi M, Lomstein BA, Jørgensen BB, Røy H, Kjeldsen KU. 2019. Environmental filtering determines family-level structure of sulfate-reducing microbial communities in subsurface marine sediments. *ISME J* 13:1920–1932. <https://doi.org/10.1038/s41396-019-0387-y>
144. Ishii S, Suzuki S, Tenney A, Norden-Krichmar TM, Nealson KH, Bretschger O. 2015. Microbial metabolic networks in a complex electrogenic biofilm recovered from a stimulus-induced metatranscriptomics approach. *Sci Rep* 5:14840. <https://doi.org/10.1038/srep14840>
145. Edwardson CF, Hollibaugh JT. 2017. Metatranscriptomic analysis of prokaryotic communities active in sulfur and arsenic cycling in Mono Lake, California, USA. *ISME J* 11:2195–2208. <https://doi.org/10.1038/ismej.2017.80>
146. Yin Z, Ye L, Jing C. 2022. Genome-resolved metagenomics and metatranscriptomics reveal that aquificae dominates arsenate reduction in tengchong geothermal springs. *Environ Sci Technol* 56:16473–16482. <https://doi.org/10.1021/acs.est.2c05764>
147. Ayala-Muñoz D, Macalady JL, Sánchez-España J, Falagán C, Couradeau E, Burgos WD. 2022. Microbial carbon, sulfur, iron, and nitrogen cycling linked to the potential remediation of a meromictic acidic pit lake. *ISME J* 16:2666–2679. <https://doi.org/10.1038/s41396-022-01320-w>
148. Tan S, Liu J, Fang Y, Hedlund BP, Lian Z-H, Huang L-Y, Li J-T, Huang L-N, Li W-J, Jiang H-C, Dong H-L, Shu W-S. 2019. Insights into ecological role of a new deltaproteobacterial order *Candidatus Acidulodesulfobacterales* by metagenomics and metatranscriptomics. *ISME J* 13:2044–2057. <https://doi.org/10.1038/s41396-019-0415-y>
149. Hao L, Michaelsen TY, Singleton CM, Dottorini G, Kirkegaard RH, Albertsen M, Nielsen PH, Dueholm MS. 2020. Novel syntrophic bacteria in full-scale anaerobic digesters revealed by genome-centric

- metatranscriptomics. ISME J 14:906–918. <https://doi.org/10.1038/s41396-019-0571-0>
150. Coskun ÖK, Özen V, Wankel SD, Orsi WD. 2019. Quantifying population-specific growth in benthic bacterial communities under low oxygen using H<sub>2</sub>18O. ISME J 13:1546–1559. <https://doi.org/10.1038/s41396-019-0373-4>
  151. Flieder M, Buongiorno J, Herbold CW, Hausmann B, Rattei T, Lloyd KG, Loy A, Wasmund K. 2021. Novel taxa of Acidobacteriota implicated in seafloor sulfur cycling. ISME J 15:3159–3180. <https://doi.org/10.1038/s41396-021-00992-0>
  152. Hausmann B, Pelikan C, Herbold CW, Köstlbacher S, Albertsen M, Eichorst SA, Glavina Del Rio T, Huemer M, Nielsen PH, Rattei T, Stingl U, Tringe SG, Trojan D, Wentrup C, Woebken D, Pester M, Loy A. 2018. Peatland *Acidobacteria* with a dissimilatory sulfur metabolism. ISME J 12:1729–1742. <https://doi.org/10.1038/s41396-018-0077-1>
  153. Högfors-Rönholm E, Lundin D, Brambilla D, Christel S, Lopez-Fernandez M, Lillhonga T, Engblom S, Österholm P, Dopson M. 2022. *Gallionella* and *Sulfuricella* populations are dominant during the transition of boreal potential to actual acid sulfate soils. Commun Earth Environ 3:304. <https://doi.org/10.1038/s43247-022-00642-z>
  154. Padhy SR, Bhattacharyya P, Nayak AK, Dash PK, Roy KS, Baig MJ, Mohapatra T. 2020. Key metabolic pathways of sulfur metabolism and bacterial diversity under elevated CO<sub>2</sub> and temperature in lowland rice: a metagenomic approach. Geomicrobiol J 37:13–21. <https://doi.org/10.1080/01490451.2019.1657992>
  155. Rosado-Porto D, Ratering S, Moser G, Deppe M, Müller C, Schnell S. 2022. Soil metatranscriptome demonstrates a shift in C, N, and S metabolisms of a grassland ecosystem in response to elevated atmospheric CO<sub>2</sub>. Front Microbiol 13:937021. <https://doi.org/10.3389/fmicb.2022.937021>
  156. Chernyh NA, Neukirchen S, Frolov EN, Sousa FL, Miroshnichenko ML, Merkel AY, Pimenov NV, Sorokin DY, Ciordia S, Mena MC, Ferrer M, Golyshin PN, Lebedinsky AV, Cardoso Pereira IA, Bonch-Osmolovskaya EA. 2020. Dissimilatory sulfate reduction in the archaeon '*Candidatus Vulcanisaeta moutnovskia*' sheds light on the evolution of sulfur metabolism. Nat Microbiol 5:1428–1438. <https://doi.org/10.1038/s41564-020-0776-z>
  157. Carr SA, Jungbluth SP, Eloe-Fadros EA, Stepanauskas R, Woyke T, Rappé MS, Orcutt BN. 2019. Carboxydrotrophy potential of uncultivated Hydrothermarchaeota from the seafloor crustal biosphere. ISME J 13:1457–1468. <https://doi.org/10.1038/s41396-019-0352-9>
  158. MacLeod F, Kindler GS, Wong HL, Chen R, Burns BP. 2019. Asgard archaea: diversity, function, and evolutionary implications in a range of microbiomes. AIMS Microbiol 5:48–61. <https://doi.org/10.3934/microbiol.2019.1.48>
  159. Farquhar J, Wu N, Canfield DE, Oduro H. 2010. Connections between sulfur cycle evolution, sulfur isotopes, sediments, and base metal sulfide deposits. Econ Geol 105:509–533. <https://doi.org/10.2113/gsecongeo.105.3.509>
  160. Colman DR, Lindsay MR, Amenabar MJ, Fernandes-Martins MC, Roden ER, Boyd ES. 2020. Phylogenomic analysis of novel Diaforarchaea is consistent with sulfite but not sulfate reduction in volcanic environments on early Earth. ISME J 14:1316–1331. <https://doi.org/10.1038/s41396-020-0611-9>
  161. Hahn CR, Farag IF, Murphy CL, Podar M, Elshahed MS, Youssef NH. 2022. Microbial diversity and sulfur cycling in an early earth analogue: from ancient novelty to modern commonality. mBio 13:e0001622. <https://doi.org/10.1128/mbio.00016-22>
  162. Fakhraee M, Hancisse O, Canfield DE, Crowe SA, Katsev S. 2019. Proterozoic seawater sulfate scarcity and the evolution of ocean–atmosphere chemistry. Nat Geosci 12:375–380. <https://doi.org/10.1038/s41561-019-0351-5>
  163. Valentine DL. 2007. Adaptations to energy stress dictate the ecology and evolution of the Archaea. Nat Rev Microbiol 5:316–323. <https://doi.org/10.1038/nrmicro1619>
  164. Halevy I, Johnston DT, Schrag DP. 2010. Explaining the structure of the Archean mass-independent sulfur isotope record. Science 329:204–207. <https://doi.org/10.1126/science.1190298>
  165. Schmidt TM, ed. 2019. Encyclopedia of microbiology. Academic Press.
  166. McKay LJ, Dlakić M, Fields MW, Delmont TO, Eren AM, Jay ZJ, Klingensmith KB, Rusch DB, Inskeep WP. 2019. Co-occurring genomic capacity for anaerobic methane and dissimilatory sulfur metabolisms discovered in the Korarchaeota. Nat Microbiol 4:614–622. <https://doi.org/10.1038/s41564-019-0362-4>
  167. Fontecilla-Camps JC. 2022. Nickel and the origin and early evolution of life. Metallomics 14:4. <https://doi.org/10.1093/mtomcs/mfac016>
  168. Zhou G-W, Yang X-R, Sun A-Q, Li H, Lassen SB, Zheng B-X, Zhu Y-G. 2019. Mobile incubator for iron (III) reduction in the gut of the soil-feeding earthworm *Pheretima guillelmi* and interaction with denitrification. Environ Sci Technol 53:4215–4223. <https://doi.org/10.1021/acs.est.8b06187>
  169. Field JA, Cervantes FJ. 2005. Microbial redox reactions mediated by humus and structurally related quinones, p 343–352. In Use of humic substances to remediate polluted environments: from theory to practice. Vol. 52.
  170. Martinez CM, Alvarez LH, Celis LB, Cervantes FJ. 2013. Humus-reducing microorganisms and their valuable contribution in environmental processes. Appl Microbiol Biotechnol 97:10293–10308. <https://doi.org/10.1007/s00253-013-5350-7>
  171. Piché-Choquette S, Constant P. 2019. Molecular hydrogen, a neglected key driver of soil biogeochemical processes. Appl Environ Microbiol 85:e02418-18. <https://doi.org/10.1128/AEM.02418-18>
  172. Dey S, Awata T, Mitsushita J, Zhang D, Kasai T, Matsuura N, Katayama A. 2021. Promotion of biological nitrogen fixation activity of an anaerobic consortium using humin as an extracellular electron mediator. Sci Rep 11:6567. <https://doi.org/10.1038/s41598-021-85955-3>
  173. Deng X, Dohmae N, Kaksonen AH, Okamoto A. 2020. Biogenic iron sulfide nanoparticles to enable extracellular electron uptake in sulfate-reducing bacteria. Angew Chem Int Ed Engl 59:5995–5999. <https://doi.org/10.1002/anie.201915196>