



Soil microalgae and cyanobacteria: the biotechnological potential in the maintenance of soil fertility and health

Sudharsanam Abinandan, Suresh R. Subashchandrabose, Kadiyala Venkateswarlu & Mallavarapu Megharaj

To cite this article: Sudharsanam Abinandan, Suresh R. Subashchandrabose, Kadiyala Venkateswarlu & Mallavarapu Megharaj (2019): Soil microalgae and cyanobacteria: the biotechnological potential in the maintenance of soil fertility and health, Critical Reviews in Biotechnology, DOI: [10.1080/07388551.2019.1654972](https://doi.org/10.1080/07388551.2019.1654972)

To link to this article: <https://doi.org/10.1080/07388551.2019.1654972>



Published online: 27 Aug 2019.



Submit your article to this journal [↗](#)



View related articles [↗](#)




View Crossmark data [↗](#)

REVIEW ARTICLE



Soil microalgae and cyanobacteria: the biotechnological potential in the maintenance of soil fertility and health

Sudharsanam Abinandan^a, Suresh R. Subashchandrabose^{a,b}, Kadiyala Venkateswarlu^c and Mallavarapu Megharaj^{a,b} 

^aGlobal Centre for Environmental Remediation (GCER), Faculty of Science, University of Newcastle, Callaghan, Australia; ^bCooperative Research Centre for Contamination Assessment and Remediation of Environment (CRC CARE), University of Newcastle, Callaghan, Australia; ^cDepartment of Microbiology, Sri Krishnadevaraya University, Anantapuramu, India

ABSTRACT

The soil microbiota plays a major role in maintaining the nutrient balance, carbon sink, and soil health. Numerous studies reported on the function of microbiota such as plant growth-promoting bacteria and fungi in soil. Although microalgae and cyanobacteria are ubiquitous in soil, very less attention has been paid on the potential of these microorganisms. The indiscriminate use of various chemicals to enhance agricultural productivity led to serious consequences like structure instability, accumulation of toxic contaminants, etc., leading to an ecological imbalance between soil, plant, and microbiota. However, the significant role of microalgae and cyanobacteria in crop productivity and other potential options has been so far undermined. The intent of the present critical review is to highlight the significance of this unique group of microorganisms in terms of maintaining soil fertility and soil health. Beneficial soil ecological applications of these two groups in enhancing plant growth, establishing interrelationships among other microbes, and detoxifying chemical agents such as insecticides, herbicides, etc. through mutualistic cooperation by synthesizing enzymes and phytohormones are presented. Since recombinant technology involving genomic integration favors the development of useful traits in microalgae and cyanobacteria for their potential application in improvement of soil fertility and health, the merits and demerits of various such advanced methodologies associated in harnessing the biotechnological potential of these photosynthetic microorganisms for sustainable agriculture were also discussed.

ARTICLE HISTORY

Received 18 March 2019
Revised 21 June 2019
Accepted 24 July 2019

KEYWORDS

Microalgae; cyanobacteria; biotechnological potential; soil health; soil fertility; phytohormones; biological soil crusts; extracellular polysaccharides

Introduction

The human population is predicted to be over 9 billion by 2050 and this may result in a 70% increase in demand for food production [1]. Development and application of sustainable farming with the perspective of ecological and nutritional balance to feed human population have been the challenges to scientists in the 21st century [2]. The lack of interest to understand the natural phenomenon of sustainable farming paved the way for intensive agriculture for enhanced productivity [3]. Many agricultural soils lack major nutrients such as nitrogen, phosphorus, potassium, and iron, resulting in the consumption of ~200 million metric tons of chemical fertilizers [4,5]. Thus, there will be a huge investment in fossil fuels to produce the required amounts of fertilizers, which is economically tedious besides causing environmental pollution. The overwhelming response towards sustainable agriculture for the maintenance of soil fertility and a healthy

ecosystem with low inputs gained much attention in recent years. The rationale of sustainable agriculture is critically dependent on soil, which is the major component of crop production and a complex life reservoir that needs nutrients for its prolonged stability and productivity [6]. Since soil serves as the largest sink for organic carbon and helps in regulation of fluxes of greenhouse gases, it is important to understand the soil ecology for ecosystem sustainability and restoration of habitats [7]. The role of soil microbiota is to increase organic compounds such as carbon, nitrogen, phosphates, and potassium, thereby improving soil health and quality [8].

Recently, Locey and Lennon [9] combined the scaling law with the lognormal model of biodiversity and predicted that the Earth is the home to more than 1.0 trillion microbial species. Of these, bacteria, fungi, actinomycetes, microalgae, and cyanobacteria are intimately associated and thrive in the uppermost part of the

soil, known as the biological soil crust (BSC), which plays a major role in improving soil fertility and agricultural productivity [10]. These crusts possess a diversity of microorganisms such as plant growth-promoting bacteria (PGPB), fungi, lichens, microalgae, and cyanobacteria. Among the diverse microbiota in the soil, microalgae and cyanobacteria are the primary producers in the food chain that can grow in intimidating environments such as arid, semi-arid and wetland ecosystems. Cyanobacteria have evolved over two billion years and are responsible for photogenic oxygenation. Certain cyanobacteria without the heterocyst can also perform nitrogen fixation, and some belong to chemo-heterotrophs. The soil microalgae and cyanobacteria release nutritive products, organic products and active components that can serve as food for other organisms like bacteria, fungi, and invertebrates.

A perusal of the database from 1908 to 2018, retrieved from the Scopus, indicates that both microalgae and cyanobacteria received less attention when compared with fungi and rhizobia in soils (Figure 1). A total of only 2312 documents have been published on microalgae and cyanobacteria during 1919–2018, mostly originating from countries like USA, China, Germany, India, etc. Again, the quantum of research concerning microalgae as compared to cyanobacteria has been very limited, and never gained momentum! However, the significant contribution in this area of research was in agriculture and environment, which accounted for 43.2 and 45.3%, respectively. It is worthwhile noting that cyanobacteria, especially with their nitrogen-fixing ability, and species of microalgae such

as *Chlamydomonas*, *Scenedesmus* and *Chlorococcum* that produce polysaccharides, have a better ability compared to fungi and other microbes in soil [11]. Nevertheless, the diaspora of research on soil microalgae has decreased substantially when compared with other microbes. These “beneficial microorganisms” have great potential in nutrient recycling, organic waste decomposition, detoxification of harmful chemicals, control of plant pathogens, and the production of metabolites such as hormones, vitamins and enzymes [12–14]. Although the information on the population dynamics of microalgae and cyanobacteria helps to understand their role in soil ecology, the beneficial effects of these microorganisms are of utmost importance in soil ecosystems and have not been fully described so far. In this review, we critically discuss the potential of microalgal and cyanobacterial crusts in colonizing degraded soils, and their implication in the maintenance of soil fertility and health.

Biological soil crusts (BSCs) – role in the survival of microalgae and cyanobacteria

Earth has different habitats and drylands cover 45% of the terrestrial (semi-arid and sub-humid) ecosystem and is anticipated to expand in the near future [15]. Furthermore, elevated concentrations of CO₂ in the atmosphere, volcanic eruptions, UV radiation, and temperature could play a major role in the destruction of biota by nutrient limitation, and an unfavorable habitat for microbial activity thereby leading to soil infertility [16,17]. The primary organisms that colonized after




Figure 1. Analysis of research-based documents powered by SCOPUS accessed on 15 March 2019. Scattered plot shows the overall documents published on fungi, rhizobia, cyanobacteria and microalgae in soil. The bar diagram (inset) indicates the number of articles published on cyanobacteria and microalgae by top 10 countries.

volcanic eruptions were microalgae that formed a gelatinous sheath [18]. Over the soil's surface, microalgae together with cyanobacteria initiate the formation of BSCs. Cyanobacteria also synthesize phytochelatin molecules for their defense against UV-B radiations that are more prevalent in desert regions [19,20]. Treves et al. [21] observed desert soil microalgae modifying photosynthetic functions such as PSII activities that will withstand elevated radiations. Heterocystous cyanobacteria possess thick extracellular sheaths that prevent the damage of nucleic acids from radiation and help to survive in extreme environments [22]. Lynch and Bragg [23] hypothesized that survival of microalgae in dryland silt soils would be delicate as the moisture plays an important role. Besides, the algae cyanobacterial crusts require only small amounts of moisture to initiate their living compared to other microbes [24]. Subsequently, by being confined to the top layer of soil, these crusts form into mounds thereby prevent moisture evaporation and shelter other crusts dominated by lichens and mosses [25]. The major factors that influence algal crust formation in the desert are the wind, water, and dust. Due to the wind factor, microalgae tend to adapt by distributing themselves in deeper layers of the sand and move to the surface when it is wet [26]. Under these conditions, they also tend to secrete copious amounts of hydrophobic amino acids in order to prevent the soil from erosion by wind and water and they will be utilized by other organisms as an organic source [27]. Bhatnagar et al. [28] observed considerable diversity of algal crusts in a desert due to its infrequent distribution. Similarly, Zhang et al. [29] suggested that mineral contents along with soil texture were responsible for the growth at extreme habitats and the heterogeneity of algal crust distribution in the desert. Similarly, Hu and Liu [30] explored the relationship between the five algae from a desert habitat, which consists of high molecular weight proteoglycans, different monosaccharides and methylated monomers in secreted extracellular polysaccharide (EPS). However, the amount of the initial inoculum and soil properties must ensure the trend-setting ability of microalgae. Trivedi et al. [31] reported that cyanobacterial abundance was significant in natural as well as agroecosystems. Based on analysis of the scientific database of NCBI on soil algae, predominant groups present in different zones such as arid, semi-arid, wetland were: *Oscillatoriales*, *Nostocales*, *Chroococcales*, *Synechococcales*, *Chroococcidiopsidales*, *Pleurocapsales*, *Microcoleaceae*, *Chlorellales*, etc.

In eroded and degraded lands, microalgae help to bind soil particles by forming sheaths and filaments [32]. Due to the oxygenic photosynthesis mediated by algae, the soil becomes aerobic by absorbing O₂

through diffusion that is required for initial setting. For instance, the relatively slow formation of the crust by cyanobacterial inoculation in burned soils compared to unburned soils due to the surface disturbance caused by heating [33]. In contrast, Acea et al. [34] observed that inoculation of heated soil samples with cyanobacteria resulted in proliferation and crust formation. Most of the algae are isothermal in nature, and the survival is dependent on protoplasm but not insulation layers [35]. Due to modified light intensity, temperature and moisture levels in arid environments, the diaphanous strata are the favorable habitats for algae [36]. In addition, algae form an association with minerals to produce water-stable organomineral aggregates [37].

Kheirfam et al. [38] demonstrated that the application of native soil cyanobacteria in erosion-prone zone increased the chemical properties such as carbon, nitrogen and organic matter. Furthermore, the synthesis of intracellular metabolites such as superoxide dismutase, proline, and carotenoids tend to help microalgae in drought-mediated stress conditions [39]. The secretion of these metabolites increases the soil compressive strength to remain stable even after cell lysis [40]. Under desiccation stress, the algal filaments are subjected to an increase in surface volume ratio in order to uptake nutrients and prevent loss of water from the cell [41]. Cellular structures such as zygotes, heterocysts, resting spores, and akinetes enable them to survive, but not necessarily withstand high temperatures [42]. Spores and cysts are morphologically distinct that help microalgae in actively dispersing over long distances [43]. By developing cysts, a microalga, *Protosiphon botryoides*, survived 50 years in soil without moisture and was active again [44].

Since topsoil dries out rapidly at high temperatures, suppressed growth of microalgae is resumed after adaptation by changing the physiological conditions [45]. Holzinger and Lütz [20] observed that under desiccation stress, a green alga, *Klebsormidium crenulatum*, was able to tolerate as a result of subsequent changes in the cell wall, cytoplasm, and mitochondria. Furthermore, algae survive photosensitivity during desiccation by enduring inside the microsites within crust formation [46]. Microalgae and cyanobacteria can resurrect once there are no supporting factors for the growth despite the presence of CO₂ and nitrogen. Thus, it is clear that microalgae and cyanobacteria are inclined to adapt to extreme conditions with various physical, metabolic and other forms of alterations mediated by immunity and change the surface quality of soils and provide organic and inorganic bioavailable nutrients to other organisms.

Soil microalgae/cyanobacteria – a very helpful factor for soil fertility

Agricultural lands are susceptible to the loss of soil fertility due to the practice of intensive farming practices, which may lead to 30% of the cultivable land being subject to soil degradation. Among other soil microbiota, microalgae alone account for 27% of the total biomass in agricultural land [47]. As evidenced by their response in different agro-ecosystems, cyanobacteria are the most suitable group of microorganisms to consider as a soil bioindicator of land use [48]. The information available in the literature signifies the potential of microalgae and cyanobacteria to improve various parameters related to soil health and fertility (Table 1). Cyanobacteria and green algae tend to enrich organic plant production particularly during climate change [54]. In addition, an increase in oxidizable C by way of releasing exopolysaccharides during algal cell lysis increases soil organic matter [55]. The soil organic matter is the readily available form of C for the growth of soil microbiota [56]. The cyanobacterial inoculation can increase soil organic carbon (SOC), total availability of nutrients in the topsoil. It is observed that cyanobacteria inoculated in a pot without seeding showed a higher content of carbon, nitrogen, and minerals than seeded pots [57]. However, the plant can also induce priming of soil organic matter (SOM) mineralization, which is due to factors that influence plant growth [58]. Yet, the bioavailability of a complex SOM fraction to support the growth of microbiota is scarce as its solubility and reactivity vary greatly depending upon the environment.

It is also postulated that algae compete with higher plants and prevent leaching of minerals and maintain the reserves for bioavailability [59]. Cyanobacteria such as *Nostoc* sp. are able to colonize plant root systems, and by a tight association facilitate the transfer of nutrients as well as metabolites with the help of hormogonia [60]. The use of native cyanobacterial strains as an alternative supplementation for nitrogen improved paddy yield [61]. Similarly, the use of cyanobacterial biofilms under non-flooded conditions improved N₂ fixation and phosphate solubilization thereby increasing the growth of crop plants [62]. Cyanobacteria not only improved the paddy yield but also showed the highest increase in oxidizable C and soluble C in postharvest soil compared to preharvest soil [63]. In extreme habitats, genera of Cyanophyceae cause retention of silt and clay with an increase in organic C (300%) and N (400%) [64]. Cyanobacterial inoculation contributes to the yield of grains besides leaving C residues that can enhance plant growth without the help of manure [65]. Interestingly, the predominantly growing native algae can reduce the diversity of other diazotrophic organisms and enhance nitrogen fixation [66]. However, the algal inoculation followed by population build-up in the soil will vary depending on the habitats under field conditions. For instance, the density of algae in arable soils ($<1 \times 10^2 \text{ g}^{-1}$) was less than water-logged soils ($>1 \times 10^7 \text{ g}^{-1}$) [67,68]. Mayland et al. [69] observed a two-fold nitrogen content increase in nutrient-limiting semi-arid soil indicating the role of cyanobacteria with active heterocyst.

Table 1. The potential of microalgae and cyanobacteria to improve various properties related to soil fertility.

Soil type	Major soil characteristics	Cyanobacterial/ microalgal inoculants	Soil fertility improvement	Reference
Sandstone, granite, schist and lime	Poor micronutrients	<i>Oscillatoria</i> sp., <i>Nostoc</i> sp. and <i>Scytonema</i> sp.	Formation of microbial crusts with high carbon and nitrogen content.	[34]
Silt loam soil	Poor micronutrient availability	<i>Anabaena</i> sp. and <i>Providencia</i> sp.	Iron concentration in soil was >2–3 folds.	[10]
Desert soil	Low organic carbon and total nitrogen	<i>Microcoleus vaginatus</i> Gom. and <i>Scytonema javanicum</i> Born et Flah	>5-Fold increments of soil organic carbon and nitrogen.	[49]
Sterile soil	Low nutrients	<i>Chlorella</i> sp., <i>Scenedesmus</i> sp., <i>Chlorococcum</i> sp., and <i>Chroococcus</i> sp.	Microbial biomass carbon.	[50]
Clay loam	Low nitrogen	<i>Chlorella vulgaris</i>	Improved nitrogen and soil enzyme activities.	[51]
Semi-arid soil	Organically poor	<i>Anabaena doliolum</i> HH-209, <i>Cylindrospermum sphaerica</i> HH-202, and <i>Nostoc calcicola</i> HH-201	Improved carbon and nitrogen mineralization by promoting soil microbial activities.	[52]
Ferruginous tropical soil	Poorly aggregated soils	<i>Nostoc</i> sp.	Improvement of aggregate stability.	[53]
Glade soil	Poor nutrients and hydro-physical parameters	<i>Tribonema minus</i> , <i>Choricystis minor</i> , and <i>Klebsormidium subtile</i>	Increase in organic carbon content, water drop penetration and decreasing evaporation during dry periods.	[25]

Table 2. Potential traits of soil microorganisms and their merits and demerits to improve soil fertility.

Microorganisms	Potential soil traits	Advantages	Limitations	References
Microalgae	Carbon fixation, extracellular polysaccharides, and initiation of BSCs.	Carbon supplementation to promote soil health. Extra cellular polysaccharides for soil aggregation. Enhancing hydraulic conductivity.	Soil moisture. Carbon limitation. Requirement of pH, mostly slight acidic to basic.	[45] [78] [79]
Cyanobacteria	Nitrogen fixation through heterocyst and non-heterocyst, CO ₂ fixation, and extracellular polysaccharides.	Carbon and nitrogen supply for soil fertility. Extra cellular polysaccharides for soil aggregation. IAA production.	Phosphorus limitation. Decreased N ₂ -fixing ability. Oxygen affects N ₂ fixation.	[80] [81] [82] [83]
PGPR	Development of resistance to pathogens, plant growth promotion, phosphate solubilization, and osmotic stress mitigation.	Fix N ₂ into plants through complex enzymes system. Phosphate mineralization. Organic acid for solubilization of potassium.	Requires phosphates and carbon for efficient activity.	[84] [85–88]
Fungi	Decomposing old recalcitrant soil organic matter, and increased nutrient availability to plants.	Accommodating bacterial enzymes to fix N ₂ without oxygen influence. Entrapment of soil particles. Mucilage, polysaccharides, and soil protein production.	Requires organic carbon for survival. Prefer acid pH.	[5,89]

The essential nutrients next to nitrogen to enrich crops are Phosphorus and potassium. In general, the upper layer in soil consists of organic Phosphorus that ranges from 20–80% of the total phosphorus [70]. In addition, Phosphorus availability to plants decreases in the highly weathered soil and is heavily dependent on microbial activity in such soils [71]. Many studies reported that phosphate-solubilizing bacteria play a major role in converting insoluble phosphate to soluble Phosphorus [72]. Nevertheless, soil microalgae accumulate inorganic phosphate and produce polyphosphates, the easily available form of phosphate to plants [73]. In addition, cyanobacteria can also release enzymes that degrade inorganic phosphate in the soil to increase its bioavailability to plants [74]. During phosphate-starved conditions, cyanobacteria are capable of solubilizing mineral rock containing phosphate by phthalic acid production [75]. However, nitrogen fixation in cyanobacteria ceased in the presence of soluble phosphate whereas the insoluble phosphate helped to maintain the rate of nitrogen fixation [76]. Other possible mechanisms include chelation and phosphate solubilization through the release of organic acids by cyanobacteria [77]. Table 2 presents the merits and demerits of potential traits in microalgae, cyanobacteria, plant growth-promoting rhizobacteria (PGPR) and fungi to improve soil fertility.

New advancements are very helpful to understand the mechanism and response of algae to soil function, and are more attractive for scaling heights and achieve more promising results. The use of cyanobacterial biofilm fertilizer gauge increased attraction due to a reduction in the quantity of chemical fertilizer, cost savings, etc. Cyanobacterial biofilm formulation of

Anabaena-Trichoderma viride resulted in the production of maize hybrids with an associated saving of 60 kg ha⁻¹ nitrogen and an increase of 20–60 kg ha⁻¹ available nitrogen in the soil [90]. The biofilm of *Phormidium* sp. showed higher phosphatase activity under different phosphate regimes [91]. Similarly, the biofilms of *Anabaena-Serratia*, *Anabaena-Pseudomonas* exhibited high acetylene reduction and phosphate activities during wheat cultivation [92]. Prasanna et al. [93] reported a 12–25% increase in yield of soybeans with the use of *Anabaena-T. viride* biofilm formulation along with the compost (paddy straw and vermiculate) as a carrier. Under the SRI system, biofilm formulation of cyanobacteria increased the bioavailability of nitrogen and phosphates to rice plants that resulted in savings of 60 kg N ha⁻¹ season⁻¹ [66,94]. Biofilm formulation increased the levels of soil micronutrients such as zinc (15–41%) and iron (13–46%) over controls in *Oryza sativa* [95]. The biofilms consisting of *Anabaena laxa* and *Rhizobium* sp. showed the higher activity of nitrogenase which resulted in 1724 kg ha⁻¹ yield in chickpea [96]. Microalgae/cyanobacteria as biofilm vectors show a higher degree for nutrient enrichment compared to other suppliers.

Soil health maintenance by microalgae/cyanobacteria – an underexplored option

According to Nkonya et al. [97], an increase in land degradation stretched to 30% compared to the total land area in the world, costing about US\$300 billion to improvise the degraded cover. Other subjugated degraded soils include acid sulfate soils, salinity soils, arid cover, etc. that are prevalent in certain countries




Figure 2. Formation of microalgal/cyanobacterial-based biological soil crusts. (a) Wind drives soil particles from top soil; (b) Infiltration of rain droplets into the soil layer; (c) Colonization of cyanobacteria and microalgae to form a sheath; (d) Soil and water entrapment in sheath to form a cyanobacterial/microalgal crust; (e) Extracellular polysaccharides (EPS) are produced from the crust; and (f) The mound formed acts as a run-off zone to divert rainwater into soil and enhances water infiltration.

such as India and Australia. The use of improved practices along with the application of fertilizers can improve the decreasing rate of land degradation [98]. However, it is evident that the routine application of synthetic fertilizers adversely affects the arable soil characteristics. Organic fertilizer amendment reduced the relative abundance of algal communities even after a year of application [99]. Long-term (82 years) applications of chemical and organic fertilizers in a rice field affected the diversity of eukaryotic microorganisms [100]. This situation, therefore, necessitates the use of alternative approaches like the application of promising microorganisms to restore soil characteristics. Induction of BSCs helps to initiate the restoration of microbial diversity in soil and vegetation cover. Hence, overwhelming interest in developing BSCs to improve the soil quality and its characteristics is vital for restoring regions with soil degradation over the world [101,102]. For instance, the combined application of cyanobacterial formulations with natural chemical additives will help to form BSCs at a faster rate than with the natural processes and enhance soil stability and water-holding capacity (WHC) in degraded soils [102].

Soil quality primarily deals with soil aggregate properties such as gradient, texture, water repellence, WHC and hydraulic conductivity [23]. On the other hand, soil health provides the balance amongst organisms within the soil, and between soil organisms and their

environment [103]. Thus, soil health is of utmost importance for the survival of crop plants. It has been established that algae are responsible for this in the crust formation and for inducing water repellence upon drying. The role of microalgae and cyanobacteria in the formation of BSCs and their implication in soil health is depicted in Figure 2. These crusts also tend to cause retention of clay, which helps in rearranging soil particles for improving soil structure [64]. Furthermore, the hydrophobic compounds synthesized by soil algae play a primary role in establishing water repellence [104]. The hydrophobic derivatives of algal metabolites tend to stabilize water aggregation, bind mineral particles and reduce break down of soil [105]. It should also be noted that filamentous algal crusts can also lead to the clogging of pores, thereby reducing the water infiltration rate [106]. Inevitably, a change in structure and function in soil due to the release of metabolites and hydrophobic derivatives allows preferential flow paths that result in developing wetting instability and improving evaporation loss [107]. These preferential flow paths possess higher substrate and nutrients bioavailability compared to other regions in soil indicating hot spots for elevated microbial activity [108,109]. However, hydrophobic surface chemistry can lead to super hydrophobicity with the help of water droplets that subsequently promotes water repellence [110]. Nevertheless, cyanobacteria and microalgae increase water repellency

during initial crust development and result in slower infiltration due to the crust succession [111].

Microalgae synergistically can also activate quorum sensing among cyanobacteria and other prokaryotes that render them to move or grow towards the substrate and thereby increasing the soil stability [107]. The EPS from cyanobacteria holds water movement in soil by absorption, and thus creates a microenvironment in the soil for water regulation [112]. Similarly, EPS serves as a reservoir of water storage in order to manage the abnormal conditions and can be metabolically active during rewetting [113]. In addition, wetting also increases the size of cyanobacterial colonies by swelling the polysaccharides present in the mucilage and thereby decreasing water runoff in the soil [105]. This makes EPS to increase soil WHC and reduce the trans- evaporation loss to the atmosphere [114]. It was also reported that cyanobacterial inoculation increased WHC, nutrient availability and aggregation of soil particles [115]. Cyanobacterial crusts increase water retention time by decreasing the bulk density and soil porosity along with organic matter and cation- exchange capacity (CEC) [116]. Roger and Reynaud [117] observed that the algal crusts helped prevent erosion from rain and did not slow the water infiltration rate. However, in the presence of cyanobacteria, the soil porosity prevented the damage of water addition due to the breakup effects of periodic wetting [118]. There will be immediate availability of organic matter from cyanobacteria to the food chain, and this contribution is significant [42].

It has also been reported that intensive agricultural practices lead to loss of soil stability, reduction in water infiltration and formation of bio pores, all of which are important for water movement and root development. These associated problems can be abridged by using organic amendments that would help to increase soil quality and properties [119,120]. When applied to the soil, gypsum binds to the polysaccharide sheath of cyanobacteria and protects them from physical break- age [81]. The use of microalgae will also increase the hydraulic conductivity of minimizing other soil amend- ments in order to improve soil fertility [121]. Filamentous algae have the potential in proliferating in soil within 10–50 mm from the surface thereby enhanc- ing the hydrologic condition of the soil [122]. The sheaths from cyanobacterial crusts that are undisturbed for long periods can penetrate up to 10 cm below the soil surface and thus possess ameliorating influences on WHC, CEC and soil stability [81]. Chamizo et al. [123] observed that cyanobacteria in BSCs with higher

electrical conductivity have aggregate water stability and water content at –33 kPa and –1500 kPa, respectively.

Soil enzymes and phytohormones from microalgae/cyanobacteria – role in soil fertility and plant growth

It has been well established that microalgae and cyano- bacteria can produce several important components of varying applications. Enzymes from microalgae and cyanobacteria are exploited to increase soil fertility. The phosphatases are the enzymes that catalyze the hydrolysis of organophosphates in soil and make Phosphorus bioavailable to plants. The presence of microalgae and cyanobacteria increases the metabolic activity of willow plants by inducing the activity of phosphatases (acid and alkaline). For instance, synthesis of alkaline phosphatase, that cleaves the phosphate res- idues of polyphosphate, in cyanobacteria was high under phosphate-deficient conditions [54], and conse- quently helps in increasing the activity of nitrogenase enzyme [124,125]. Even in extreme habitats, there was no reduction in the activity of phosphatase in cyano- bacterial mats whereas other enzymes such as lipase; and glucosidase were significantly affected under these conditions [126,127]. Cyanobacteria also synthesize hydrolytic enzymes that exhibit fungicidal activity such as protease, carboxymethyl cellulase (CMCase), cello- biase and filter paperase (Fpase), all of which are responsible for hastening nutrient recycling of carbon, nitrogen, and phosphate which are necessary for plant growth [84]. Rao and Burns [67] reported a significant increase in the activities of dehydrogenase, urease, and phosphatase with cyanobacterial inoculants that resulted in increased organic matter mineralization.

De Caire et al. [128] reported that *T. tenuis* and *Microchaete tenera* inoculants in soil increased the activ- ities of extracellular enzymes such as glucosidase, phos- phomonoesterase, arylsulfatase, protease, and urease besides the accumulation of intracellular dehydrogenase. The role of enzymes associated with the antioxidant defense mechanisms towards pesticides has been known in microalgae and cyanobacteria as it is important to identify the activities of these resistant enzymes [129]. It is noteworthy that certain enzymes in microalgae are responsible for the degradation of pesticides [12]. For instance, an esterase in the marine cyanobacterium, *Phormidium valderianum* BDU 20041, transformed chlor- pyrifos [130]. Phosphatase enzyme in soil microalgae was not affected by the insecticide, fenamiphos [131]. Such enzymes in microalgae or cyanobacteria, which

Table 3. Enzymatic resistance in cyanobacteria/microalgae towards pesticides as soil contaminants.

Pesticide/group	Cyanobacterium/microalga	Responsible enzyme	Reference
Glyphosate (H)	<i>Anabaena</i> sp. <i>Nostoc</i> sp.	Enol-pyruvyl-shikimate-3-phosphate (EPSP) synthase	[133]
Chlorpyrifos (I)	<i>Spirulina platensis</i>	Alkaline phosphatase	[134]
Organophosphates	<i>Synechococcus</i> sp. PCC7942	Organophosphorus hydrolase	[135]
Fluroxypyr (H)	<i>Chlamydomonas reinhardtii</i>	Glutathione S-transferase	[136]
Chlorpyrifos (I)	<i>Phormidium valderianum</i> BDU 20041	Polyphenol oxidase	[115]
Carbofuran (I)	<i>Nostoc hatei</i>	Glutamine synthetase	[137]
Norflurazon (H)	<i>Synechocystis</i> sp.	Phytoene desaturase	[138]
Metflurazone (H)	<i>Chlorella fusca</i>	Cytochrome P450	[139]
Bentazon(H)	<i>Anabaena cylindrica</i>	Glutathione S-transferase	[140]
Methyl parathion (I)	<i>Anabaena</i> sp. PCC7120	NADP + reductase	[141]
Mancozeb (F)	<i>Nostoc ellipsosporum</i> <i>Tolypothrix tenuis</i>		
Bagalol (F), Thiodan (I)	<i>Scytonema simplex</i>		
Phorate (I)	<i>Tolypothrix tenuis</i>	Isocitrate lyase	[142]

I: Insecticide; F: Fungicide; H: Herbicide.

confer resistance towards pesticides, have a greater role in their biodegradation by fortuitous evolution [132]. Several enzymes of microalgae and cyanobacteria are involved either in biodegradation of environmental pollutants or confer resistance towards such chemicals (Table 3). The presence of these enzymes could help to screen algae-based upon *in vivo* tolerance [133].

Phytohormones are important molecules commonly found in plants and microbes (microalgae, fungi, bacteria, etc.), and are involved in regulating their cellular processes. These phytohormones help to alleviate the abiotic stress in plants and maintain growth [143]. Both cyanobacteria and microalgae, which establish a symbiotic association with plants, synthesize hormones such as auxins (indole acetic acid, IAA), cytokinin, abscisic acid, gibberellins and ethylene [83,144,145]. Moreover, certain phytohormones, produced by cyanobacteria and microalgae, are implicated in improving plant growth (Table 4). It has been established that the synthesis of phytohormones in microalgae is dependent on the bioavailability of precursors, amino acids, from plants [149]. For example, synthesis of auxins in cyanobacteria was only observed with the accumulation of tryptophan, an amino acid, excreted by plants [83]. On the contrary, certain cyanobacteria cannot synthesize IAA even in the presence of tryptophan [150]. However, it has been observed that unicellular cyanobacteria release auxins and filamentous cyanobacteria accumulate them within the cells [151]. In general, phytohormones produced by cyanobacteria during co-cultivation, are transferred to the plants through their roots along with other nutrients [152].

It was also postulated that phytohormones from cyanobacteria or microalgae could influence the endogenous phytohormones of the plant cell [153]. Likewise, cytokinin-binding protein from cyanobacteria was transported into the plant for the synthesis of

endogenous phytohormones [83,154]. Although the transportation of phytohormones into the plant is clearly understood, the molecular mechanism of transportation has yet to be established. The symbiotic association of cyanobacteria and other bacteria is quite common, and the mutualism is for exchanging organic substrates, such as vitamins [62,155–158]. Among other microorganisms, *Nostoc* sp. was high in abundance associating it with the root system of rice plants by developing hormogonia [159]. This association was particularly due to the presence of a gene *ipdc* in the cyanobacterium which is responsible for the IAA synthesis needed for plant growth [151]. A schematic diagram provided in Figure 3 represents the association between a cyanobacterium, microalga, and PGPR in the plant rhizosphere. Meza et al. [160] reported that IAA supplied by a growth-promoting bacterium, *Azospirillum brasilense*, was used by *Chlorella vulgaris* for utilizing inorganic phosphate intracellularly. The intracellular accumulation of ammonium was enhanced by the release of IAA by PGPR [160]. Metabolites such as tryptophan from microalgae will exchange with IAA that is produced from PGPR, indicating an alternative way of obtaining phytohormones by algae [161,162].

Co-cultivation of *A. brasilense* with *Chlorella sorokiniana* resulted in an increase of tryptophan in the microalga and the exudation of IAA from the bacterium during their interaction [163]. A similar observation under stress conditions was where microalgae and bacteria released thiamin during co-cultivation [164]. Thus, the release of phytohormones may be a plausible solution to stress tolerance in microalgae. Similarly, the exogenous role of abscisic acid supported stress tolerance mechanism in *Chlamydomonas reinhardtii* [165]. A green alga, *K. crenulatum*, could significantly withstand desiccation stress by producing phytohormones such as ethylene, cytokinin and abscisic acid [166]. It is also reported that the utilization of phytohormones

Table 4. Phytohormones produced by cyanobacteria/microalgae that enhance plant growth.

Cyanobacterium/microalga	Phytohormone	Plant	Reference
<i>Anabaena vaginicola</i> <i>Cylindrospermum michailovskoense</i> , <i>Nostoc calcicole</i>	Auxins (IAA, indole 3-acetic acid; IBA, indole 3-butyric acid)	<i>Triticum aestivum</i> L.	[146]
<i>Calothrix</i> sp. (403) <i>Calothrix</i> sp. (405) <i>Chlorella</i> sp. (519) <i>Coenochloris</i> sp. (534) <i>Tetracystis</i> sp. (479)	Auxin (IBA) Cytokinins (Kinetin)	<i>Cucumis sativus</i> <i>Glycine max</i>	[147]
<i>Chlamydomonas</i> sp. (460) <i>Chlamydomonas</i> sp. (693) <i>Anabaena oryzae</i> <i>Anabaena doliolum</i> <i>Phormidium fragile</i> <i>Calothrix geitonos</i> <i>Hapalosiphon intricatus</i> <i>Aulosira fertilissima</i> <i>Tolypothrix tenuis</i> <i>Oscillatoria acuta</i> <i>Plectonema boryanum</i> <i>Aphanothece</i> sp. MBDU 515	Auxins (IAA, IBA) Auxin (IAA)	<i>Oryza sativa</i> <i>Arachis hypogaea</i> L. <i>Moringa oleifera</i> Lam.	[13] [148]




Figure 3. Symbiotic relationship between a microalga, cyanobacterium, and PGPR in plant rhizosphere.

improves growth as well as lipids (fatty acids) in microalgae [167–169]. The phytohormones can also detoxify or modify the toxic compounds that inhibit plant and algal growth. For example, phytohormones help to modify the phytotoxicity of heavy metal and prevent algal inhibition [170]. Thus, the coexistence of microalgae and PGPR in plant roots will be not only mutually beneficial but also help to release plant growth-promoting substances.

Genetic engineering potential of microalgae and cyanobacteria for improving soil fertility and health

The use of advanced techniques in algal engineering helps to induce traits for enhancing their role in soil functioning. Figure 4 provides a general schematic

representation of approaches for developing potential traits in microalgae and cyanobacteria to improve soil fertility and health that is essential for sustainable agriculture. Gene transfer is a signature for the manipulation of the rubisco gene from cyanobacterium to plants through endosymbiosis [171]. Even genetic rearrangements have been observed in cyanobacteria from ancient viruses [172]. Nevertheless, a significant paradigm shift is enhanced hydrogenase production by cyanobacteria through genetic manipulation [173]. Such a paradoxical swing can be one of the reasons why the potential of exploring microalgae is still in its infancy. Moreover, earlier research on mutagenic cyanobacteria has suggested that N₂ fixation was higher compared to wild strains especially when ferredoxin hydrogenase was expressed in the mutants [174]. However, iron concentrations, at 1 μM, promoted




Figure 4. Approaches to develop potential traits in microalgae/cyanobacteria for their application to improve soil fertility and health.

ferredoxin biosynthesis but not the growth of mutant strains, suggesting that the FdxH was not replaced by NifJ-flavodoxin. In fact, the *hetR* protein is mainly responsible for regulating N_2 fixation in heterocysts especially in N_2 -fixing dominant strains of *Anabaena* and *Nostoc* [175]. It was also observed that chemical mutagenesis resulted in defects with no heterocyst formation under any conditions. However, Zhou et al. [176] observed that the nitrogen-depleted environment provided higher production of *hetR* protein predominantly located in heterocysts only for a limited time. Furthermore, it was postulated that *hetR* could play other roles in non-heterocystous cyanobacterial species. Nevertheless, additional copies of *hetR* the gene, placed under the promoter, has alleviated the stress and exhibited overexpression of *hetR* protein in *Anabaena* strains [177]. Spiller et al. [178] developed a mutant strain, which can produce the nitrogenase enzyme and excrete ammonium in the environment thus exhibiting excellent fertilizer traits.

The application of recombinant DNA technology involving genomic integration in all the above studies, overproduced heterocysts and reduced further risks of lateral transfer of transgenes. It is also imperative to identify external stress (such as herbicides) and the response of cyanobacteria and microalgae in order to produce resistant biofertilizers. Singh et al. [179] identified hetero-specific DNA-mediated genetic transformation responsible for herbicide resistance in *Gloeocapsa* sp. which could be later expressed in a *Nostoc* species for their application in the field as a biofertilizer. They also suggested that the absence of a hetero-specific barrier in crosses of DNA-mediated genetic

transformation maintains stable expression. Similarly, herbicide-resistant markers could be used in microalgae to resist toxic potential [180]. Chaurasia et al. [181] used the *linA2* gene from *Sphingomonas* and inserted it into an *Anabaena* strain to confer resistance towards the toxicity of lindane through the light-inducible promoter. Both an integrative approach and a light-inducible promoter minimized the chances of lateral gene transfer to non-target organisms and were found suitable for their application in agriculture fields. The induction of gene encoding modified norflurazon-resistant phytoene desaturase (PDS) also enhanced the herbicide resistance efficiently in *Chlorella zofingiensis* by 91%, suggesting that the use of a dominant selectable marker is a pre-requisite for efficient transformation of microorganisms [182]. Similarly, expression of the *hsp* gene to produce a heat shock protein, that serves as the backbone for stress tolerance in certain cyanobacteria with respect to desiccation and temperature, has been well used to improve the strains in biofertilizer application. Nevertheless, expressing the *groESL* operon encoding the heat-shock protein genes enhanced nitrogen fixation levels and photosynthetic activity even in stress-mediating environments [177]. Use of recombinant technology to enhance nitrogenase activity led to catering the demand of nitrogen in rice seedlings. A mutated cyanobacterial strain, resistant to azetidine carboxylate, was used to over-accumulate proline, which helps to withstand salinity stress [183]. The practice of developing spontaneous mutants did not favor the overaccumulation of proline either from stress or resistance. However, toxicity was exerted if the mutant cells are defective also in other assimilatory

Table 5. Potential of genetically modified traits and their merits and demerits in maintaining soil fertility and health.

Microorganism	Genetic modification	Potential trait	Methodology used	Limitations	Reference
<i>Anabaena</i> sp. PCC 7120	<i>fdxH</i> gene	<i>FdxH</i> is necessary for maximum nitrogenase activity and optimal growth under N ₂ -fixing conditions.	Plasmid DNA isolation, restriction enzyme analysis, agarose gel electrophoresis, and cloning.	Iron limitation causes a decrease in growth.	[174]
<i>Synechocystis</i> sp. PCC 6803	<i>nif</i> genes	First example of nitrogenase activity detected in photosynthetic non-diazotroph.	Five-step homologous recombination together with the <i>cnfR</i> gene encoding the transcriptional activator of the <i>nif</i> genes.	Low but significant nitrogenase activity was detected in all transformants.	[188]
<i>Azotobacter</i> sp.	FeSII protein	Forms a protective complex with nitrogenase during periods when nitrogenase is exposed to oxygen.	Subcloning gene and site-directed mutagenesis.	The nitrogenase component bearing the double lysine mutation is degraded much more rapidly under carbon substrate-limited conditions.	[189]
<i>Synechocystis</i> sp. PCC 6803	Hydrogenase genes	Functional way to improve nitrogenase enzyme activity under micro-oxic conditions.	Recombinant plasmids, DNA assembler and Gibson assembly.	Further removal of genes from both directions resulted in a decrease of N ₂ -fixing activity.	[190]
<i>Chlamydomonas reinhardtii</i>	<i>chlL</i> gene	Introduces N ₂ -fixing genes directly into a chloroplast genome.	Plasmid construction and construction of chloroplast transformation vectors.	NifH product can at least partially substitute for the function of the putative "chlorophyll iron protein".	[191]
<i>Azotobacter vinelandii</i>	Fe–Fe nitrogenase system	Metallocluster biosynthesis under molybdenum limiting conditions.	Construction of recombinant plasmids and construction of multiple gene deletion mutants.	Limitation in ratio of ethane/ethylene produced by Fe–Fe nitrogenase is dependent upon electron flux.	[192]

pathways involved in nitrogen [184]. Interestingly, it was also demonstrated that a mutant lacking nitrate accumulation could grow without any damage in an herbicide-treated paddy field [185]. Some mutant strains withstand methylamine (an inhibitor of metabolic activity) that is induced by the action of herbicidal activity [186]. In fact, the spontaneous occurrence of mutants switched off the pathway and conferred resistance against the herbicide. Apart from other activities, the role of EPS in microalgae and cyanobacteria has an extensive function in soil activities. Recently, Liu et al. [187] mutated a microalga, *Cryptocodinium cohnii*, that resulted in higher secretion of EPS compared to the wild type strain indicating that mutagenesis plays a significant role. However, the use of optimal conditions for mutation is required for effective transformation in microbes.

For exploiting the potential traits in maintaining soil fertility and health, several microalgae and cyanobacteria have been genetically modified (Table 5). Although promising results are associated with the use of such advanced techniques, it is essential to overcome certain disadvantages while developing better strains of microalgae and cyanobacteria for their application as potential soil inoculants in a win-win situation.

For instance, soil health improvement through spontaneous gene mutation requires optimization with the reaffirmation of the expressed genes as it results in the shutting of other pathways [183,186,187]. Furthermore, the use of chemicals to mediate mutagenesis is also known to affect the organisms upon genetic modification [175]. However, the approach of recombinant technology for genetic modification has been quite promising, especially in the area of nitrogen fixation by soil microorganisms [177]. Engineering microbes are vulnerable in complex environments due to the presence of other microbes and substrates necessitating proper attention. An integrated approach for genetic modification reduces the risk of gene transfer to other microbes [178,181]. Genetic markers, excess plasmids, stable transformation, and expression are some of the key factors to be addressed in any genetic modification technique. This situation is warranted because the use of microalgal – cyanobacterial consortia in field studies results in more agricultural productivity even in abiotic stress environments and degraded lands. In fact, there is a huge gap since the late 2000s in this area of research because of a shift in interest toward the generation of value-added products such as biofuels. Furthermore, an understanding based on the OMICS

technology to explore the mechanistic and other arenas of mutation is also limited. However, the use of genetically modified organisms remains controversial due to the associated undesirable effects on ecological balance and living beings [193].

Conclusions

Microalgae and cyanobacteria are the primary colonizers that pave the way for other organisms to survive and establish under different climatic and edaphic conditions ranging from desert to wetland. As detailed in the present review, it is very factual that microalgae and cyanobacteria play a major role in maintaining the soil health and fertility necessary for sustained agriculture. The most significant advantage of exploiting microalgae and cyanobacteria is the balance in the dynamics of nutrients such as carbon, nitrogen and phosphate for the productivity of agricultural crops. Metabolic activities, such as the release of phytohormones, enzymes and allelochemicals from microalgae, play a pivotal role in enhancing plant growth, establishing interrelationships among other microbes, and detoxifying chemical agents such as pesticides, herbicides, etc. Moreover, biofertilizer applications using microalgae or cyanobacteria is a promising and beneficial approach for a better nitrogen economy in modern agriculture, reduction in greenhouse gas emissions, and maintenance of soil fertility. Hence, better insights into molecular aspects of microalgae and cyanobacteria are essential in order to realize their full potential in biotechnological applications. The present review underlines the further need to harness soil microalgae and cyanobacteria in the area of environmental biotechnology.

Disclosure statement

Authors declare no conflict of interest.

Funding

SA thankfully acknowledges the University of Newcastle, Australia for providing RTP Scholarship (Commonwealth University Scholarship -APA and IPRS) for Ph.D. program.

ORCID

Mallavarapu Megharaj  <http://orcid.org/0000-0002-6230-518X>

References

- [1] FAO: Food and Agriculture Organization of the United Nations [Internet]. Rome (Italy): FAO. How to feed the World in 2050;2009 [cited 2019 Feb 25]; Available from www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf.
- [2] Prasanna R, Sharma E, Sharma P, et al. Soil fertility and establishment potential of inoculated cyanobacteria in rice crop grown under non-flooded conditions. *Paddy Water Environ*. 2013;11:175–183.
- [3] Chaudhary V, Prasanna R, Nain L, et al. Bioefficacy of novel cyanobacteria-amended formulations in suppressing damping off disease in tomato seedlings. *World J Microbiol Biotechnol*. 2012;28:3301–3310.
- [4] FAO. World fertilizer trends and outlook to 2018. Rome (Italy): FAO; 2015. p. 66.
- [5] Rashid MI, Mujawar LH, Shahzad T, et al. Bacteria and fungi can contribute to nutrients bioavailability and aggregate formation in degraded soils. *Microbiol Res*. 2016;183:26–41.
- [6] Pankhurst CE, Lynch JM, The role of soil microbiology in sustainable intensive agriculture. In: Andrews JH, Tommerup IC, editors. *Advances in Plant Pathology*, Vol. 11. Waltham, MA: Academic Press; 1995. p. 229–247.
- [7] Chen MM, Zhu YG, Su YH, et al. Effects of soil moisture and plant interactions on the soil microbial community structure. *Eur J Soil Biol*. 2007;43:31–38.
- [8] Singh JS, Pandey VC, Singh DP. Efficient soil microorganisms: a new dimension for sustainable agriculture and environmental development. *Agric Ecosyst Environ*. 2011;140:339–353.
- [9] Locey KJ, Lennon JT. Scaling laws predict global microbial diversity. *Proc Natl Acad Sci USA*. 2016;13: 5970–5975.
- [10] Manjunath M, Kanchan A, Ranjan K, et al. Beneficial cyanobacteria and eubacteria synergistically enhance bioavailability of soil nutrients and yield of okra. *Heliyon*. 2016;2:e00066.
- [11] Xiao R, Zheng Y. Overview of microalgal extracellular polymeric substances (EPS) and their applications. *Biotechnol Adv*. 2016;34:1225–1244.
- [12] Megharaj M, Kantachote D, Singleton I, et al. Effects of long-term contamination of DDT on soil microflora with special reference to soil algae and algal transformation of DDT. *Environ Pollut*. 2000;109:35–42.
- [13] Singh DP, Prabha R, Yandigeri MS, et al. Cyanobacteria-mediated phenylpropanoids and phytohormones in rice (*Oryza sativa*) enhance plant growth and stress tolerance. *Anton Van Leeuwen*. 2011;100:557–568.
- [14] Renuka N, Guldhe A, Prasanna R, et al. Microalgae as multi-functional options in modern agriculture: current trends, prospects and challenges. *Biotechnol Adv*. 2018;36:1255–1273.
- [15] Feng S, Fu Q. Expansion of global drylands under a warming climate. *Atmos Chem Phys*. 2013;13: 10081–10094.

- [16] Luo Y, Su B, Currie WS, et al. Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *Bioscience*. 2004;54:731–739.
- [17] Šibanc N, Dumbrell AJ, Mandić-Mulec I, et al. Impacts of naturally elevated soil CO₂ concentrations on communities of soil archaea and bacteria. *Soil Biol Biochem*. 2014;68:348–356.
- [18] Reub M. Notice sur la nouvelle flora de Krakatau. *Ann J Ard Bot Buitenzorg*. 1988;7:221–223.
- [19] Bhargava P, Kumar Srivastava A, Urmil S, et al. Phytochelatin plays a role in UV-B tolerance in N₂-fixing cyanobacterium *Anabaena doliolum*. *J Plant Physiol*. 2005;162:1220–1225.
- [20] Holzinger A, Lütz C. Algae and UV irradiation: Effects on ultrastructure and related metabolic functions. *Micron*. 2006;37:190–207.
- [21] Treves H, Raanan H, Finkel OM, et al. A newly isolated *Chlorella* sp. from desert sand crusts exhibits a unique resistance to excess light intensity. *FEMS Microbiol Ecol*. 2013;86:373–380.
- [22] Garcia-Pichel F, López-Cortés A, Nübel U. Phylogenetic and morphological diversity of cyanobacteria in soil desert crusts from the Colorado Plateau. *Appl Environ Microbiol*. 2001;67:1902–1910.
- [23] Lynch JM, Bragg E. Microorganisms and soil aggregate stability. In: Stewart BA, editor. *Advances in soil science*. New York, NY: Springer; 1985. p. 133–171.
- [24] Lange OL. Photosynthesis of soil-crust biota as dependent on environmental factors. In: Belnap J, Lange OL, editors. *Biological soil crusts: structure, function, and management*. Berlin Heidelberg: Springer; 2003. pp. 217–240.
- [25] Lichner L, Hallett PD, Drongová Z, et al. Algae influence the hydrophysical parameters of a sandy soil. *Catena*. 2013;108:58–68.
- [26] Hu C, Liu Y, Song L, et al. Effect of desert soil algae on the stabilization of fine sands. *J Appl Phycol*. 2002;14:281–292.
- [27] Flaibani A, Olsen Y, Painter TJ. Polysaccharides in desert reclamation: compositions of exocellular proteoglycan complexes produced by filamentous blue-green and unicellular green edaphic algae. *Carb Res*. 1989;190:235–248.
- [28] Bhatnagar A, Makandar MB, Garg MK, et al. Community structure and diversity of cyanobacteria and green algae in the soils of Thar Desert (India). *J Arid Environ*. 2008;72:73–83.
- [29] Zhang B, Zhang Y, Downing A, et al. Distribution and composition of cyanobacteria and microalgae associated with biological soil crusts in the Gurbantunggut desert. China. *Arid Land Res Manage*. 2011;25:275–293.
- [30] Hu CX, Liu YD. Primary succession of algal community structure in desert soil. *Acta Bot Sin*. 2003;45: 917–924.
- [31] Trivedi P, Delgado-Baquerizo M, Anderson IC, et al. Response of soil properties and microbial communities to agriculture: implications for primary productivity and soil health indicators. *Front Plant Sci*. 2016; 7:990.
- [32] Belnap J. The potential roles of biological soil crusts in dryland hydrologic cycles. *Hydrol Process*. 2006;20: 3159–3178.
- [33] Eldridge D, Greene R. Microbiotic soil crusts - a review of their roles in soil and ecological processes in the rangelands of Australia. *Soil Res*. 1994;32: 389–415.
- [34] Acea MJ, Prieto-Fernández A, Diz-Cid N. Cyanobacterial inoculation of heated soils: effect on microorganisms of C and N cycles and on chemical composition in soil surface. *Soil Biol Biochem*. 2003; 35:513–524.
- [35] Brock TD. Primary colonization of surtsey, with special reference to the blue-green algae. *Oikos*. 1973; 24:239–243.
- [36] Cameron RE, Blank GB, United S, et al. *Desert algae: soil crusts and diaphanous substrata as algal habitats*. Pasadena (CA): Jet Propulsion Laboratory, California Institute of Technology; 1966.
- [37] Malam Issa O, Le Bissonnais Y, Défarge C, et al. Role of a cyanobacterial cover on structural stability of sandy soils in the Sahelian part of western Niger. *Geoderma*. 2001;101:15–30.
- [38] Kheirfam H, Sadeghi SH, Homae M, et al. Quality improvement of an erosion-prone soil through microbial enrichment. *Soil Tillage Res*. 2017;165: 230–238.
- [39] Lin CS, Wu JT. Tolerance of soil algae and cyanobacteria to drought stress. *J Phycol*. 2014;50:131–139.
- [40] Xie Z, Liu Y, Hu C, et al. Relationships between the biomass of algal crusts in fields and their compressive strength. *Soil Biol Biochem*. 2007;39:567–572.
- [41] Karsten U, Holzinger A. Light, temperature, and desiccation effects on photosynthetic activity, and drought-induced ultrastructural changes in the green alga *Klebsormidium dissectum* (Streptophyta) from a high alpine soil crust. *Microb Ecol*. 2012;63:51–63.
- [42] Starks TL, Shubert LE, Trainor FR. Ecology of soil algae: a review. *Phycologia*. 1981;20:65–80.
- [43] Geisen S, Mitchell EA, Adl S, et al. Soil protists: a fertile frontier in soil biology research. *FEMS Microbiol Rev*. 2018;42:293–323.
- [44] Lewis LA, Trainor FR. Survival of *Protosiphon botryoides* (Chlorophyceae, Chlorophyta) from a Connecticut soil dried for 43 years. *Phycologia*. 2012; 51:662–665.
- [45] Barclay WR, Lewin RA. Microalgal polysaccharide production for the conditioning of agricultural soils. *Plant Soil*. 1985;88:159–169.
- [46] Gray DW, Lewis LA, Cardon ZG. Photosynthetic recovery following desiccation of desert green algae (Chlorophyta) and their aquatic relatives. *Plant Cell Environ*. 2007;30:1240–1255.
- [47] Megharaj M. Healthy levels of soil algae lift plant growth, December ed. In: *Farming Ahead*, Canberra (Australia): CSIRO; 2001.
- [48] Zancan S, Trevisan R, Paoletti MG. Soil algae composition under different agro-ecosystems in North-Eastern Italy. *Agri Ecosyst Environ*. 2006;112:1–12.
- [49] Wang W, Liu Y, Li D, et al. Feasibility of cyanobacterial inoculation for biological soil crusts formation in desert area. *Soil Biol Biochem*. 2009;41:926–929.

- [50] Renuka N, Prasanna R, Sood A, et al. Exploring the efficacy of wastewater-grown microalgal biomass as a biofertilizer for wheat. *Environ Sci Pollut Res*. 2016; 23:6608–6620.
- [51] Dineshkumar R, Kumaravel R, Gopalsamy J, et al. Microalgae as bio-fertilizers for rice growth and seed yield productivity. *Waste Biomass Valor*. 2018;9: 793–800.
- [52] Nisha R, Kaushik A, Kaushik CP. Effect of indigenous cyanobacterial application on structural stability and productivity of an organically poor semi-arid soil. *Geoderma*. 2007;138:49–56.
- [53] Malam Issa O, Défarge C, Le Bissonnais Y, et al. Effects of the inoculation of cyanobacteria on the microstructure and the structural stability of a tropical soil. *Plant Soil*. 2007;290:209–219.
- [54] Grzesik M, Romanowska-Duda Z. Ability of cyanobacteria and green algae to improve metabolic activity and development of willow plants. *Pol J Environ Stud*. 2015;24:1003–1012.
- [55] De Caire GZ, De Cano MS, Zaccaro De Mulé MC, et al. Exopolysaccharide of *Nostoc muscorum* (Cyanobacteria) in the aggregation of soil particles. *J Appl Phycol*. 1997;9:249–253.
- [56] Li ZP, Han CW, Han FX. Organic C and N mineralization as affected by dissolved organic matter in paddy soils of subtropical China. *Geoderma*. 2010; 157:206–213.
- [57] Maqubela MP, Mnkeni PNS, Issa OM, et al. *Nostoc* cyanobacterial inoculation in South African agricultural soils enhances soil structure, fertility, and maize growth. *Plant Soil*. 2009;315:79–92.
- [58] Paterson E, Thornton B, Midwood AJ, et al. Atmospheric CO₂ enrichment and nutrient additions to planted soil increase mineralisation of soil organic matter, but do not alter microbial utilisation of plant- and soil C-sources. *Soil Biol Biochem*. 2008;40: 2434–2440.
- [59] Shields LM, Durrell LW. Algae in relation to soil fertility. *Bot Rev*. 1964;30:92–128.
- [60] Svircev Z, Tamas I, Nenin P, et al. Co-cultivation of N₂-fixing cyanobacteria and some agriculturally important plants in liquid and sand cultures. *Appl Soil Ecol*. 1997;6:301–308.
- [61] Irisarri P, Gonnet S, Monza J. Cyanobacteria in Uruguayan rice fields: diversity, nitrogen fixing ability and tolerance to herbicides and combined nitrogen. *J Biotechnol*. 2001;91:95–103.
- [62] Prasanna R, Babu S, Bidyarani N, et al. Prospecting cyanobacteria-fortified composts as plant growth promoting and biocontrol agents in cotton. *Ex Agric*. 2015;51:42–65.
- [63] De Mulé MCZ, De Caire GZ, De Cano MS, et al. Effect of cyanobacterial inoculation and fertilizers on rice seedlings and postharvest soil structure. *Commun Soil Sci Plant Anal*. 1999;30:97–107.
- [64] Fletcher JE, Martin WP. Some Effects of algae and molds in the rain-crust of desert soils. *Ecology*. 1948; 29:95–100.
- [65] Subrahmanyam R, Relwani LL, Manna GB. Fertility build-up of rice field soils by blue-green algae. *Proc Nat Sci India B*. 1965;62:252–272.
- [66] Priya H, Prasanna R, Ramakrishnan B, et al. Influence of cyanobacterial inoculation on the culturable microbiome and growth of rice. *Microbiol Res*. 2015; 171:78–89.
- [67] Rao DLN, Burns RG. The effect of surface growth of blue-green algae and bryophytes on some microbiological, biochemical, and physical soil properties. *Biol Fert Soils*. 1990;9:239–244.
- [68] Roger PA, Tirol A, Ardales S, et al. Chemical composition of cultures and natural samples of N₂-fixing blue-green algae from rice fields. *Biol Fert Soils*. 1986;2:131–146.
- [69] Mayland HF, McIntosh TH, Fuller WH. Fixation of isotopic nitrogen on a semiarid soil by algal crust organisms. Madison (WI): Soil Science Society of America Proceedings; 1966.
- [70] Steffens D, Leppin T, Luschin-Ebengreuth N, et al. Organic soil phosphorus considerably contributes to plant nutrition but is neglected by routine soil-testing methods. *Z Pflanzenernähr Bodenk*. 2010;173: 765–771.
- [71] Oberson A, Friesen DK, Rao IM, et al. Phosphorus Transformations in an oxisol under contrasting land-use systems: the role of the soil microbial biomass. *Plant Soil*. 2001;237:197–210.
- [72] Sharma SB, Sayyed RZ, Trivedi MH, et al. Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. SpringerPlus. 2013;2:587.
- [73] Mukherjee C, Chowdhury R, Ray K. Phosphorus recycling from an unexplored source by polyphosphate accumulating microalgae and cyanobacteria – a step to phosphorus security in agriculture. *Front Microbiol*. 2015;6:1421.
- [74] Whitton BA, Grainger SL, Hawley GR, et al. Cell-bound and extracellular phosphatase activities of cyanobacterial isolates. *Microb Ecol*. 1991;21:85–98.
- [75] Yandigeri MS, Meena KK, Srinivasan R, et al. Effect of mineral phosphate solubilization on biological nitrogen fixation by diazotrophic cyanobacteria. *Indian J Microbiol*. 2011;51:48–53.
- [76] Yandigeri MS, Yadav AK, Srinivasan R, et al. Studies on mineral phosphate solubilization by cyanobacteria *Westiellopsis* and *Anabaena*. *Microbiology*. 2011; 80:558–565.
- [77] Singh JS, Kumar A, Rai AN, et al. Cyanobacteria: a precious bio-resource in agriculture, ecosystem, and environmental sustainability. *Front Microbiol*. 2016;7: 529.
- [78] Raven JA. Inorganic carbon acquisition by eukaryotic algae: four current questions. *Photosyn Res*. 2010; 106:123–134.
- [79] Rossi F, Potrafka RM, Pichel FG, et al. The role of the exopolysaccharides in enhancing hydraulic conductivity of biological soil crusts. *Soil Biol Biochem*. 2012; 46:33–40.
- [80] Adams DG. Heterocyst formation in cyanobacteria. *Curr Opin Microbiol*. 2000;3:618–624.
- [81] Belnap J, Gardner JS. Soil microstructure in soils of the Colorado Plateau: the role of the cyanobacterium *Microcoleus vaginatus*. *Great Basin Naturalist*. 1993;53: 40–47.

- [82] Bergman B, Gallon JR, Rai AN, et al. N_2 Fixation by non-heterocystous cyanobacteria1. FEMS Microbiol Rev. 1997;19:139–185.
- [83] Sergeeva E, Liaimer A, Bergman B. Evidence for production of the phytohormone indole-3-acetic acid by cyanobacteria. Planta. 2002;215:229–238.
- [84] Burns RG, DeForest JL, Marxsen J, et al. Soil enzymes in a changing environment: current knowledge and future directions. Soil Biol Biochem. 2013;58:216–234.
- [85] Çakmakçı R, Dönmez F, Aydın A, et al. Growth promotion of plants by plant growth-promoting rhizobacteria under greenhouse and two different field soil conditions. Soil Biol Biochem. 2006;38:1482–1487.
- [86] Cassán F, Maiale S, Masciarelli O, et al. Cadaverine production by *Azospirillum brasilense* and its possible role in plant growth promotion and osmotic stress mitigation. Eur J Soil Biol. 2009;45:12–19.
- [87] Compant S, Clément C, Sessitsch A. Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonization, mechanisms involved and prospects for utilization. Soil Biol Biochem. 2010;42:669–678.
- [88] Rodríguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion. Biotechnol Adv. 1999;17:319–339.
- [89] Fontaine S, Henault C, Aamor A, et al. Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming effect. Soil Biol Biochem. 2011;43:86–96.
- [90] Prasanna R, Hossain F, Babu S, et al. Prospecting cyanobacterial formulations as plant-growth-promoting agents for maize hybrids. South Afr J Plant Soil. 2015;32:199–207.
- [91] Ellwood NTW, Di Pippo F, Albertano P. Phosphatase activities of cultured phototrophic biofilms. Water Res. 2012;46:378–386.
- [92] Swarnalakshmi K, Prasanna R, Kumar A, et al. Evaluating the influence of novel cyanobacterial bio-filmed biofertilizers on soil fertility and plant nutrition in wheat. Eur J Soil Biol. 2013;55:107–116.
- [93] Prasanna R, Triveni S, Bidyarani N, et al. Evaluating the efficacy of cyanobacterial formulations and bio-filmed inoculants for leguminous crops. Arch Agron Soil Sci. 2014;60:349–366.
- [94] Prasanna R, Adak A, Verma S, et al. Cyanobacterial inoculation in rice grown under flooded and SRI modes of cultivation elicits differential effects on plant growth and nutrient dynamics. Ecol Eng. 2015;84:532–541.
- [95] Adak A, Prasanna R, Babu S, et al. Micronutrient enrichment mediated by plant-microbe interactions and rice cultivation practices. J Plant Nutr. 2016;39:1216–1232.
- [96] Bidyarani N, Prasanna R, Babu S, et al. Enhancement of plant growth and yields in Chickpea (*Cicer arietinum* L.) through novel cyanobacterial and biofilmed inoculants. Microbiol Res. 2016;188:97–105.
- [97] Nkonya E, Mirzabaev A, Von Braun J. Economics of land degradation and improvement: a global assessment for sustainable development. Cham (Switzerland): Springer; 2015.
- [98] Bruinsma J. World agriculture: towards 2015/2030: Summary Report. Rome (Italy): Food and Agriculture Organization of the United Nations (FAO); 2002. (9251047618).
- [99] Xiong W, Jousset A, Guo S, et al. Soil protist communities form a dynamic hub in the soil microbiome. ISME J. 2018;12:634.
- [100] Murase J, Hida A, Ogawa K, et al. Impact of long-term fertilizer treatment on the microeukaryotic community structure of a rice field soil. Soil Biol Biochem. 2015;80:237–243.
- [101] Park CH, Li X, Jia RL, et al. Effects of superabsorbent polymer on cyanobacterial biological soil crust formation in laboratory. Arid Land Res Manage. 2015;29:55–71.
- [102] Park CH, Li XR, Jia RL, et al. Combined application of cyanobacteria with soil fixing chemicals for rapid induction of biological soil crust formation. Arid Land Res Manage. 2017;31:81–93.
- [103] Brady NC, Weil RR. The nature and properties of soils, 12th ed. Upper Saddle River (New Jersey): Prentice Hall Inc.; 1999.
- [104] Doerr SH, Shakesby RA, Walsh R. Soil water repellency: its causes, characteristics and hydro-geomorphological significance. Earth Sci Rev. 2000;51:33–65.
- [105] Malam Issa O, Défarge C, Trichet J, et al. Microbiotic soil crusts in the Sahel of western Niger and their influence on soil porosity and water dynamics. Catena. 2009;77:48–55.
- [106] Hallett PD, Young IM. Changes to water repellence of soil aggregates caused by substrate-induced microbial activity. Eur J Soil Sci. 1999;50:35–40.
- [107] Morales VL, Parlange JY, Steenhuis TS. Are preferential flow paths perpetuated by microbial activity in the soil matrix? A review. J Hydrol. 2010;393:29–36.
- [108] Bundt M, Albrecht A, Froidevaux P, et al. Impact of preferential flow on radionuclide distribution in soil. Environ Sci Technol. 2000;34:3895–3899.
- [109] Pivetz B, Steenhuis T. Soil matrix and macropore biodegradation of 2, 4-D. J Environ Qual. 1995;24:564–570.
- [110] McHale G, Newton MI, Shirtcliffe NJ. Water-repellent soil and its relationship to granularity, surface roughness and hydrophobicity: A materials science view. Eur J Soil Science. 2005;56:445–452.
- [111] Fischer T, Veste M, Wiehe W, et al. Water repellency and pore clogging at early successional stages of microbiotic crusts on inland dunes, Brandenburg, NE Germany. Catena. 2010;80:47–52.
- [112] Potts M. Desiccation tolerance: a simple process? Trends Microbiol. 2001;9:553–559.
- [113] De Winder GBM. Ecophysiological strategies of drought-tolerant phototrophic micro-organisms in dune soils [Ph. D. thesis]. Amsterdam (Netherlands): University of Amsterdam; 1990. p. 94.
- [114] Mager DM. Cyanobacterial soil crusts: analysing resilience in Kalahari sand soils. In: Degenovine KM, editor. Semi-arid environments: agriculture, water supply and vegetation. Hauppauge (NY): NOVA Scientific Publishers, Inc; 2010.
- [115] Watanabe I, Roger PA. Nitrogen fixation in wetland rice field. New Delhi (India): Oxford and IBH; 1984.

- [116] Mager DM. Extracellular polysaccharides from cyanobacterial soil crusts and their role in dryland surface processes [Ph.D. Thesis]. Manchester (UK): Manchester Metropolitan University; 2009. p. 273.
- [117] Roger PA, Reynaud PA. Free-living blue-green algae in tropical soils In: Dommergues YR, Diem HG, editors. Microbiology of tropical soils and plant productivity. Dordrecht (Netherlands): Springer; 1982. p. 147–168.
- [118] Falchini L, Sparvoli E, Tomaselli L. Effect of *Nostoc* sp. (Cyanobacteria) inoculation on the structure and stability of clay soils. *Biol Fert Soils*. 1996;23:346–352.
- [119] Stark C, Condrón LM, Stewart A, et al. Influence of organic and mineral amendments on microbial soil properties and processes. *Appl Soil Ecol*. 2007;35:79–93.
- [120] Mengual C, Schoebitz M, Azcón R, et al. Microbial inoculants and organic amendment improves plant establishment and soil rehabilitation under semiarid conditions. *J Environ Manage*. 2014;134:1–7.
- [121] Kaushik BD, Murti G. Effect of blue green algae and gypsum application on physicochemical properties of alkali soils. *Phykos*. 1981;20:91–94.
- [122] Williams JD, Dobrowolski JP, West NE. Microbiotic crust influence on unsaturated hydraulic conductivity. *Arid Soil Res Rehab*. 1999;13:145–154.
- [123] Chamizo S, Cantón Y, Miralles I, et al. Biological soil crust development affects physicochemical characteristics of soil surface in semiarid ecosystems. *Soil Biol Biochem*. 2012;49:96–105.
- [124] Reed SC, Cleveland CC, Townsend AR. Functional ecology of free-living nitrogen fixation: a contemporary perspective. *Annu Rev Ecol Evol Syst*. 2011;42:489–512.
- [125] Pérez CA, Thomas FM, Silva WA, et al. Patterns of biological nitrogen fixation during 60,000 years of forest development on volcanic soils from south-central Chile. *New Zealand J Ecol*. 2014;38:189–200.
- [126] Dagmara SÂ, Jaroslav V, Eliska RÂ. Extracellular enzyme activities in benthic cyanobacterial mats: comparison between nutrient-enriched and control sites in marshes of northern Belize. *Aquat Microb Ecol*. 2006;44:11–20.
- [127] Marxsen J, Zoppini A, Wilczek S. Microbial communities in streambed sediments recovering from desiccation. *FEMS Microbiol Ecol*. 2010;71:374–386.
- [128] De Caire GZ, De Cano MS, Palma RM, et al. Changes in soil enzyme activities following additions of cyanobacterial biomass and exopolysaccharide. *Soil Biol Biochem*. 2000;32:1985–1987.
- [129] Sethunathan N, Megharaj M, Chen ZL, et al. Algal degradation of a known endocrine disrupting insecticide, α -endosulfan, and its metabolite, endosulfan sulfate, in liquid medium and soil. *J Agric Food Chem*. 2004;52:3030–3035.
- [130] Palanisami S, Prabakaran D, Uma L. Fate of few pesticide-metabolizing enzymes in the marine cyanobacterium *Phormidium valderianum* BDU 20041 in perspective with chlorpyrifos exposure. *Pestic Biochem Physiol*. 2009;94:68–72.
- [131] Megharaj M, Singleton I, Kookana R, et al. Persistence and effects of fenamiphos on native algal populations and enzymatic activities in soil. *Soil Biol Biochem*. 1999;31:1549–1553.
- [132] Megharaj M, Ramakrishnan B, Venkateswarlu K, et al. Bioremediation approaches for organic pollutants: a critical perspective. *Environ Int*. 2011;37:1362–1375.
- [133] Forlani G, Pavan M, Gramek M, et al. Biochemical bases for a widespread tolerance of cyanobacteria to the phosphonate herbicide glyphosate. *Plant Cell Physiol*. 2008;49:443–456.
- [134] Thengodkar RRM, Sivakami S. Degradation of chlorpyrifos by an alkaline phosphatase from the cyanobacterium *Spirulina platensis*. *Biodegradation*. 2010;21:637–644.
- [135] Chungjatupornchai W, Fa-Aroonsawat S. Biodegradation of organophosphate pesticide using recombinant cyanobacteria with surface-and intracellular-expressed organophosphorus hydrolase. *J Microbiol Biotechnol*. 2008;18:946–951.
- [136] Zhang S, Qiu CB, Zhou Y, et al. Bioaccumulation and degradation of pesticide fluroxypyr are associated with toxic tolerance in green alga *Chlamydomonas reinhardtii*. *Ecotoxicology*. 2011;20:337–347.
- [137] Jha NM, Mishra KS. Biological responses of cyanobacteria to insecticides and their insecticide degrading potential. *Bull Environ Contam Toxicol*. 2005;75:374–381.
- [138] Martinezferez I, Vioque A, Sandmann G. Mutagenesis of an amino acid responsible in phytoene desaturase from *Synechocystis* for binding of the bleaching herbicide norflurazon. *Pestic Biochem Physiol*. 1994;48:185–190.
- [139] Thies F, Backhaus T, Bossmann B, et al. Xenobiotic biotransformation in unicellular green algae. Involvement of cytochrome P450 in the activation and selectivity of the pyridazinone pro-herbicide metflurazon. *Plant Physiol*. 1996;112:361–370.
- [140] Galhano V, Peixoto F, Gomes-Laranjo J. Bentazon triggers the promotion of oxidative damage in the Portuguese rice field cyanobacterium *Anabaena cylindrica*: response of the antioxidant system. *Environ Toxicol*. 2010;25:517–526.
- [141] Barton JW, Kuritz T, O'Connor LE, et al. Reductive transformation of methyl parathion by the cyanobacterium *Anabaena* sp. strain PCC7120. *Appl Microbiol Biotechnol*. 2004;65:330–335.
- [142] Debnath M, Mandal NC, Ray S. Effect of fungicides and insecticides on growth and enzyme activity of four cyanobacteria. *Indian J Microbiol*. 2012;52:275–280.
- [143] Wani SH, Kumar V, Shriram V, et al. Phytohormones and their metabolic engineering for abiotic stress tolerance in crop plants. *Crop J*. 2016;4:162–176.
- [144] Hartung W, Gimmler H. A stress physiological role for abscisic acid (ABA) in lower plants. In: Behnke HD, Lüttge U, Esser K, Kadereit JW, Runge M, editors. Progress in botany: structural botany physiology genetics taxonomy geobotany, Berlin (Germany): Springer; 1994. p. 157–173.
- [145] Tarakhovskaya ER, Maslov YI, Shishova MF. Phytohormones in algae. *Russ J Plant Physiol*. 2007;54:163–170.

- [146] Shariatmadari Z, Riahi H, Abdi M, et al. Impact of cyanobacterial extracts on the growth and oil content of the medicinal plant *Mentha piperita* L. J Appl Phycol. 2015;27:2279–2287.
- [147] Stirk WA, Ördög V, Van Staden J, et al. Cytokinin- and auxin-like activity in cyanophyta and microalgae. J Appl Phycol. 2002;14:215–221.
- [148] Gayathri M, Kumar PS, Prabha AML, et al. In vitro regeneration of *Arachis hypogaea* L. and *Moringa oleifera* Lam. using extracellular phytohormones from *Aphanothece* sp. MBDU 515. Algal Res. 2015;7:100–105.
- [149] Zhao Y. Auxin Biosynthesis: A simple two-step pathway converts tryptophan to indole-3-acetic acid in plants. Mol Plant. 2012;5:334–338.
- [150] Prasanna R, Joshi M, Rana A, et al. Modulation of IAA production in cyanobacteria by tryptophan and light. Pol J Microbiol. 2010;59:99–105.
- [151] Hussain A, Shah ST, Rahman H, et al. Effect of IAA on in vitro growth and colonization of *Nostoc* in plant roots. Front Plant Sci. 2015;6:46.
- [152] Mazhar S, Cohen JD, Hasnain S. Auxin producing non-heterocystous cyanobacteria and their impact on the growth and endogenous auxin homeostasis of wheat. J Basic Microbiol. 2013;53:996–1003.
- [153] Glick BR, Patten CL, Holguin G, et al. Biochemical and genetic mechanisms used by plant growth promoting bacteria. London (UK): Imperial College Press; 1999.
- [154] Shevchenko GV, Karavaiko NN, Selivankina SY, et al. Possible involvement of cyanobacteria in the formation of plant hormonal system. Russ J Plant Physiol. 2014;61:154–159.
- [155] Ramanan R, Kim BH, Cho DH, et al. Algae–bacteria interactions: evolution, ecology and emerging applications. Biotechnol Adv. 2016;34:14–29.
- [156] Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, et al. Algae acquire vitamin B12 through a symbiotic relationship with bacteria. Nature. 2005;438:90–93.
- [157] Kazamia E, Czesnick H, Nguyen TTV, et al. Mutualistic interactions between vitamin B₁₂-dependent algae and heterotrophic bacteria exhibit regulation. Environ Microbiol. 2012;14:1466–1476.
- [158] Kim HJ, Choi YK, Jeon HJ, et al. Growth promotion of *Chlorella vulgaris* by modification of nitrogen source composition with symbiotic bacteria, *Microbacterium* sp. HJ1. Biomass Bioener. 2015;74:213–219.
- [159] Nilsson M, Rasmussen U, Bergman B. Competition among symbiotic cyanobacterial *Nostoc* strains forming artificial associations with rice (*Oryza sativa*). FEMS Microbiol Lett. 2005;245:139–144.
- [160] Meza B, de-Bashan LE, Bashan Y. Involvement of indole-3-acetic acid produced by *Azospirillum brasilense* in accumulating intracellular ammonium in *Chlorella vulgaris*. Res Microbiol. 2015;166:72–83.
- [161] De-Bashan LE, Antoun H, Bashan Y. Involvement of indole-3-acetic acid produced by the growth-promoting bacterium *Azospirillum* sp. in promoting growth of *Chlorella vulgaris*. J Phycol. 2008;44:938–947.
- [162] De-Bashan LE, Hernandez JP, Bashan Y. The potential contribution of plant growth-promoting bacteria to reduce environmental degradation – a comprehensive evaluation. Appl Soil Ecol. 2012;61:171–189.
- [163] Palacios OA, Bashan Y, Schmid M, et al. Enhancement of thiamine release during synthetic mutualism between *Chlorella sorokiniana* and *Azospirillum brasilense* growing under stress conditions. J Appl Phycol. 2016;28:1521–1531.
- [164] Palacios OA, Gomez-Anduro G, Bashan Y, et al. Tryptophan, thiamine and indole-3-acetic acid exchange between *Chlorella sorokiniana* and the plant growth-promoting bacterium *Azospirillum brasilense*. FEMS Microbiol Ecol. 2016;92:fiw077.
- [165] Yoshida K, Igarashi E, Mukai M, et al. Induction of tolerance to oxidative stress in the green alga, *Chlamydomonas reinhardtii*, by abscisic acid. Plant Cell Environ. 2003;26:451–457.
- [166] Holzinger A, Becker B. Desiccation tolerance in the streptophyte green alga *Klebsormidium*: the role of phytohormones. Comm Int Biol. 2015;8:1059978.
- [167] Bajguz A, Piotrowska-Niczyporuk A. Synergistic effect of auxins and brassinosteroids on the growth and regulation of metabolite content in the green alga *Chlorella vulgaris* (Trebouxiophyceae). Plant Physiol Biochem. 2013;71:290–297.
- [168] Salama ES, Kabra AN, Ji MK, et al. Enhancement of microalgae growth and fatty acid content under the influence of phytohormones. Bioresour Technol. 2014;172:97–103.
- [169] Yu XJ, Sun J, Sun YQ, et al. Metabolomics analysis of phytohormone gibberellin improving lipid and DHA accumulation in *Aurantiochytrium* sp. Biochem Eng J. 2016;112:258–268.
- [170] Piotrowska-Niczyporuk A, Bajguz A, Zambrzycka E, et al. Phytohormones as regulators of heavy metal biosorption and toxicity in green alga *Chlorella vulgaris* (Chlorophyceae). Plant Physiol Biochem. 2012;52:52–65.
- [171] Raven JA, Giordano M, Beardall J, et al. Algal evolution in relation to atmospheric CO₂: carboxylases, carbon-concentrating mechanisms and carbon oxidation cycles. Philos Trans R Soc Lond, B, Biol Sci. 2012;367:493–507.
- [172] Haselkorn R. Developmentally regulated gene rearrangements in prokaryotes. Annu Rev Genet. 1992;26:113–130.
- [173] Tamagnini P, Axelsson R, Lindberg P, et al. Hydrogenases and hydrogen metabolism of cyanobacteria. Microbiol Mol Biol Rev. 2002;66:1–20.
- [174] Masepohl B, Schölisch K, Görlitz K, et al. The heterocyst-specific fdxH gene product of the cyanobacterium *Anabaena* sp. PCC 7120 is important but not essential for nitrogen fixation. Mol Gen Genet. 1997;253:770–776.
- [175] Buikema WJ, Haselkorn R. Isolation and complementation of nitrogen fixation mutants of the cyanobacterium *Anabaena* sp. strain PCC 7120. J Bacteriol. 1991;173:1879–1885.
- [176] Zhou R, Cao Z, Zhao J. Characterization of HetR protein turnover in *Anabaena* sp. PCC 7120. Arch Microbiol. 1998;169:417–426.
- [177] Chaurasia AK, Apte SK. Improved eco-friendly recombinant *Anabaena* sp. strain PCC7120 with enhanced

- nitrogen biofertilizer potential. *Appl Environ Microbiol.* 2011;77:395–399.
- [178] Spiller H, Latorre C, Hassan M, et al. Isolation and characterization of nitrogenase-derepressed mutant strains of cyanobacterium *Anabaena variabilis*. *J Bacteriol.* 1986;165:412–419.
- [179] Singh DT, Nirmala K, Modi DR, et al. Genetic transfer of herbicide resistance gene(s) from *Gloeocapsa* spp. to *Nostoc muscorum*. *Mol Gen Genet.* 1987;208:436–438.
- [180] Malik J, Barry G, Kishore G. The herbicide glyphosate. *Biofactors.* 1989;2:17–25.
- [181] Chaurasia AK, Adhya TK, Apte SK. Engineering bacteria for bioremediation of persistent organochlorine pesticide lindane (γ -hexachlorocyclohexane). *Bioresour Technol.* 2013;149:439–445.
- [182] Liu J, Sun Z, Gerken H, et al. Genetic engineering of the green alga *Chlorella zofingiensis*: a modified norflurazon-resistant phytoene desaturase gene as a dominant selectable marker. *Appl Microbiol Biotechnol.* 2014;98:5069–5079.
- [183] Singh AK, Chakravarthy D, Singh TPK, et al. Evidence for a role for L-proline as a salinity protectant in the cyanobacterium *Nostoc muscorum*. *Plant Cell Environ.* 1996;19:490–494.
- [184] Rai AK, Tiwari SP. NO_3^- nutrition and salt tolerance in the cyanobacterium *Anabaena* sp. PCC 7120 and mutant strains. *J Appl Microbiol.* 1999;86:991–998.
- [185] Vaishampayan A, Sinha RP, Gupta AK, et al. N_2 -fixing rice-field cyanobacterial mutant resistant against uracil herbicide. *Ecohydrol Hydrobiol.* 2003;3:417–423.
- [186] Vaishampayan A, Sinha RP, Gupta AK, et al. A cyanobacterial mutant resistant against a bleaching herbicide. *J Basic Microbiol.* 2000;40:279–288.
- [187] Liu B, Sun Z, Ma X, et al. Mutation breeding of extracellular polysaccharide-producing microalga *Cryptocodinium cohnii* by a novel mutagenesis with atmospheric and room temperature plasma. *Int J Mol Sci.* 2015;16:8201–8212.
- [188] Tsujimoto R, Kotani H, Yokomizo K, et al. Functional expression of an oxygen-labile nitrogenase in an oxygenic photosynthetic organism. *Sci Rep.* 2018;8:7380.
- [189] Lou J, Moshiri F, Johnson MK, et al. Mutagenesis studies of the FeSII protein of *Azotobacter vinelandii*: roles of histidine and lysine residues in the protection of nitrogenase from oxygen damage. *Biochemistry.* 1999;38:5563–5571.
- [190] Liu D, Liberton M, Yu J, et al. Engineering nitrogen fixation activity in an oxygenic phototroph. *mBio.* 2018;9:e01029–18.
- [191] Cheng Q, Day A, Dowson-Day M, et al. The *Klebsiella pneumoniae* nitrogenase Fe protein gene (*nifH*) functionally substitutes for the *chlL* gene in *Chlamydomonas reinhardtii*. *Biochem Biophys Res Commun.* 2005;329:966–975.
- [192] Yang J, Xie X, Wang X, et al. Reconstruction and minimal gene requirements for the alternative iron-only nitrogenase in *Escherichia coli*. *Proc Natl Acad Sci USA.* 2014;111:E3718–E3725.
- [193] Liu L, Bilal M, Duan X, et al. Mitigation of environmental pollution by genetically engineered bacteria—current challenges and future perspectives. *Sci Tot Environ.* 2019;667:444–454.