Max 2-SAT: Biased Formulas, the Race for Every Branching Rule, and the Adaptive Direction

8 October 2026

Working notes

Abstract

We take stock of the notes written since 22 September, keep one framework, and prove one new theorem inside it. The framework is the search with pruning clauses and the sum check, counted by Knuth's identity and analysed through the race identity $Z = S + Q - U$. We show that this analysis does not need a fixed order: along the uniform walk of any branching rule, adaptive or not, the gap of every variable performs a fair ± 1 walk, the shortfall is a fair-coin thinning and Q is a martingale. The theorem is for *biased* formulas: if a linear fraction of the variables have an effective gap at least ε times their own degree, and the degrees have bounded spread, then Max 2-SAT is solved in time $O^*(2^{(1-\gamma)n})$ and polynomial space with $\gamma = \varepsilon^2\eta^2/(3000R)$, at every clause density, by every branching rule. No cancellation and no spectral condition is used: a gap potential $\Phi = \sum_v |\Delta_v|$ is a submartingale, and bias beats the uncertainty of the last variables. This settles, for a different exponent and without the measure, the accumulation question left open in the note of 6–7 October. We then organize the remaining work into three regimes, biased, balanced and aligned, each with its own tool, and describe the adaptive rule that the race suggests: branch on the clause with the largest shortfall whose bad values disagree most with the current aggregate of bad values. The slack of the summed inequality and its exact completion count are proposed as the measure; they generalize the clause weights $r - \log_\lambda V(r,k)$ to overlapping clauses with an exact Pascal recursion.

1 Where we stand

Goal. A deterministic polynomial-space algorithm for Max 2-SAT in time $O^*(c^n)$, $c < 2$, for formulas of every clause density; failing that, natural classes of formulas of unbounded density on which such a bound is proved for every member.

What is proved. Table 1 lists the pieces by the note that contains them. The rules are all sound for every branching order: domination, pruning clauses with cuts and propagation [2], sums and weighted sums of pruning clauses [3], chained pair decisions [3], the three-state compression of near-twin pairs [12], and counting over exact interaction classes [4]. The running-time results are of four kinds.

What is open. The measure-and-conquer analysis with clause weights [3, Sections 6–8] is tight step by step but needs plain steps to be rare along the tilted walk; the accumulation over many clauses ([3, Problem 17]) is open. The balanced regime needs cancellation of bad values at the free variables, which random signs provide and which the pair counts of [10] certify; no clause-level condition without a quadratic form is known. The aligned regime

Result	Class	Tool
$2^{H((1-\varepsilon)/2)n}$, tight [1, 2]	ηn variables with gap $\geq \varepsilon n$	level budget; order-free with pruning clauses
Key Lemma, $\gamma < 0.0236$ [6, 7]	random signs on any graph, any density	race identity, late signs, Knuth
spectral class [8]	$\|J\|, \|L\| \leq K\sqrt{d}$, $\sum h^2 \leq Knd$	same race, Talagrand
early enumeration [9]	random signs, degree spread D/d	count completions of the summed inequality
counting criterion [10, 11]	(C1) pair counts, (C2) gaps, (C3′)	same, deterministic
near twins [12]	βn disjoint r-near-twin pairs	compression, $2^n(3/4)^{\beta n/(r+1)}$
exact classes [4]	t interaction classes, exceptions E	counting ones, $2^{\|E\|}(n+1)^t$
this note, Thm. 5	ηn variables with gap $\geq \varepsilon D_v$, degree spread R	race for every rule, gap potential

Table 1: Running-time results. The first row needs gaps linear in n; the new row needs gaps linear in the variable's own degree, so it covers every density including bounded degree.

(Max-Cut on K_m, the circulant $C_{n,c}$) defeats every rule implied by single flips [3, Section 9]; near-twin compression and exact classes solve the extreme cases, and the inverse question, whether a failure of cancellation forces such structure, is open [12].

What to park. The learning dichotomy [5]: packing and core certificates, exposure, the entropy dichotomy. Its two lasting points are the K_m barrier (also in [3]) and the joint refutation test, which is the linear-relaxation cut of [3, Proposition 7]. From the frontier draft [4] we keep the full-phase weights, the weighted-count certificate and exact interaction classes; the coverage dichotomy is subsumed by the three regimes below. The spectral class [8] is superseded by [10, 11] and, for the biased regime, by Theorem 5, which needs no norm at all.

2 The race for every branching rule

Setting. The formula is canonical with $B = 1$: unit clauses with multiplicity, at most one binary clause per pair [2, Section 2]. For a variable v let D_v be its binary degree, u_v its unit multiplicity after cancelling opposite units, and $h_v = \Delta_v(F)$ its effective gap at the root: positive minus negative binary occurrences plus twice the unit imbalance. Then $|h_v| \leq D_v + 2u_v$. Write $\bar{d} = \frac{1}{n}\sum_v D_v$ and, for a set S, $\bar{d}_S = \frac{1}{|S|}\sum_{v\in S} D_v$. For $a \in \{\pm 1\}^n$ the score is $c_0 + \frac{1}{4}\sum_v h_v a_v - \frac{1}{4}\sum_{u<v} J_{uv} a_u a_v$ with $J_{uv} \in \{0, \pm 1\}$ [8, Definition 1].

The search. At a node some free variable x is chosen by a *branching rule*, a deterministic function of the node; both values are tried. Let Δ be the effective gap of x at the node, $s = \operatorname{sgn}\Delta$ with $\operatorname{sgn} 0 = +1$ its preferred value in ± 1 notation, and $p = (D^f - |\Delta|)/2$ the smaller deficit, where D^f is the number of free neighbours. The child with $a_x = -s$ receives the pruning clause $\langle L_x \leq p - 1\rangle$ over the free neighbours of x [2, Section 8]; the other child receives nothing. A node is cut when a clause is violated (this includes $p \leq 0$) or when the sum check fires: $\sum_{w \text{ free}} \min\{m_w^0, m_w^1\} > \sum_c k_c$ over the live non-vacuous clauses, with k_c the remaining allowance of c and m_w^b the number of such clauses in which b is the bad value of w [3, Section 5]. No domination and no propagation. Soundness for every rule: each branching adds a clause to one child only [2, Proposition 20], and the sum is implied by the clauses.

Proposition 1 (Knuth's identity, any rule)**.** *Let the walk choose a uniform child at every live node, and let π_t be the probability that it is alive after t steps. The tree has $2^t\pi_t$ live nodes at depth t and at most $1 + 2n \max_t 2^t\pi_t$ nodes in all. Hence the search takes time $O^*(\max_t 2^t\pi_t)$ and polynomial space.*

Proof. Every live node has two children and is reached by the walk with probability 2^{-t}, whatever the rule. □

Write x_t for the variable chosen at step t of the walk, σ_t for its value, $a_t = 2\sigma_t - 1$, Δ_t for its gap at that node and $\xi_t = [a_t = -\operatorname{sgn}\Delta_t]$ for the event that a clause is created. Let H_t be the history $(x_1, \sigma_1, \dots, x_t, \sigma_t)$. The *uncut process* runs the walk through all n steps, creates every clause and keeps it. On the event that the walk is alive at depth T, the uncut process and the search agree up to depth T.

Lemma 2 (The race)**.** *At any node, over all clauses created so far,*

$$\sum_{w \text{ free}} \min\{m_w^0, m_w^1\} - \sum_c k_c = S + Q - U,$$

$$S = \sum_c \Big(\frac{|\Delta_c|}{2} + 1\Big), \qquad Q = \sum_c \Big(j_c - \frac{a_c}{2}\Big), \qquad U = \frac{1}{2}\sum_{w \text{ free}} |m_w^1 - m_w^0|,$$

where Δ_c is the gap at the creation of c, a_c the number of its support variables assigned since, and j_c the number of those that took their bad value. If the walk is alive at depth T, then $Z_T = S_T + Q_T - U_T \le 0$.

Proof. This is [6, Lemma 2, Remark 3]: the identity is algebra at one node and does not mention the order; removing vacuous clauses only strengthens the check, so the check of the search fires whenever the check over all created clauses fires. □

Lemma 3 (The values are the coins)**.** *For every branching rule:*

(a) Given H_{t-1}, the variable x_t, its gap Δ_t, the counts $m_{x_t}^b$ and the free set are determined, and a_t is uniform on ± 1. Hence ξ_t is a fair coin given H_{t-1}.

(b) The gap of a free variable v is $\Delta_v(t) = h_v - \sum_{s \le t,\, x_s \sim v} J_{vx_s} a_s$. When a neighbour x_s of v is assigned, Δ_v changes by ± 1, each with probability $\frac{1}{2}$ given H_{s-1}. So, sampled at the times its neighbours are assigned, Δ_v is h_v plus a simple symmetric random walk.

(c) $Q_t - Q_{t-1} = \frac{1}{2} a_t b_{x_t}$ with $b_{x_t} = m_{x_t}^1 - m_{x_t}^0$ taken at depth $t-1$; so Q is a martingale with $|Q_t - Q_{t-1}| \le \frac{1}{2} D_{x_t}$.

Proof. (a) The rule is a function of the node, and σ_t is a fresh uniform bit. (b) The coefficient of a_v in the score after substituting the assigned variables is $\frac{1}{4}(h_v - \sum J_{vx_s} a_s)$, and in the residual canonical formula it is $\frac{1}{4}\Delta_v$ [8, Lemma 4(a)]; the increment $-J_{vx_s} a_s$ is ± 1 with probability $\frac{1}{2}$ because a_s is uniform given H_{s-1} and $J_{vx_s} = \pm 1$ is fixed. A ± 1 sequence in which every term is uniform given the past is a sequence of independent fair coins. (c) Assigning x_t raises a_c by one for every created clause containing it and j_c by $[\sigma_t$ bad in $c]$, which sums to $\frac{1}{2} a_t b_{x_t}$; b_{x_t} is determined by H_{t-1} and $|b_{x_t}|$ is at most the number of created clauses containing x_t, at most D_{x_t}. □

This is the deterministic form of the late-signs lemma of [7]: there the unseen coins were the signs, here they are the values. The difference is that the bad values of the free variables are now fixed by the formula, so U_T is not a sum of coins. For biased formulas we do not need it to be.

3 Biased formulas

Definition 4 (Biased formulas)**.** Let $\varepsilon, \eta \in (0,1]$ and $R \geq 1$. A canonical formula with $B = 1$ is (ε, η, R)-*biased* if there is a set S of at least ηn variables such that

(B1) every $v \in S$ has $|h_v| \geq \varepsilon D_v$: its majority literal exceeds its minority literal by an ε-fraction of its degree, counting a unit clause as two occurrences;

(B2) $\bar{d}_S \geq 1$ and $\sum_{v \in V} (D_v + u_v)^2 \leq R\, n\, \bar{d}_S^2$: no part of the formula is much denser than the biased variables.

Condition (B1) is the gap condition of [3] in its relative form, where the gap is compared with the variable's own degree and not with n; it allows every density, from bounded degree to complete graphs. Condition (B2) holds with $R = 1$ for regular graphs without unit clauses and with $R = O(1)$ whenever all degrees are within a constant factor of $\bar{d}_S$. The set S need not be known to the algorithm.

Theorem 5 (Biased formulas, every branching rule)**.** *Let F be (ε, η, R)-biased. For every branching rule, the search of Section 2 solves Max 2-SAT on F in polynomial space and time*

$$O^*\big(2^{(1-\gamma)n}\big), \qquad \gamma = \frac{\varepsilon^2 \eta^2}{3000\, R}.$$

The constant is not optimized; the order ε^2 is right (Remark 9). The proof uses three estimates along the walk of Lemma 3, one for each term of the race, and a potential that turns the initial gaps into a lower bound on the gaps at branching time.

Lemma 6 (Gap potential)**.** *Fix a set S and a branching rule, and let the uncut process apply the rule to every node, cut or not. Put $\Phi_t = \sum_{v \in S \text{ free at } t} |\Delta_v(t)|$ and $G_t = \sum_{s \leq t,\, x_s \in S} |\Delta_s|$. Then*

$$G_T = \Phi_0 - \Phi_T + \sum_{t \leq T} z_t + M_T,$$

where $z_t \geq 0$ is the number of free neighbours $v \in S$ of x_t with $\Delta_v(t-1) = 0$, and M is a martingale with respect to (H_t) with $|M_t - M_{t-1}| \leq 2D_{x_t}$. Consequently

$$\Pr\big[G_T < \Phi_0 - \Phi_T - x\big] \leq \exp\Big(-\frac{x^2}{8 \sum_{t \leq T} D_{x_t}^2}\Big).$$

Proof. At step t the variable x_t leaves the free set, taking $|\Delta_t|$ with it if $x_t \in S$, and each free neighbour $v \in S$ of x_t changes its gap by ± 1. So $\Phi_t - \Phi_{t-1} = -[x_t \in S]\,|\Delta_t| + Y_t$ with $Y_t = \sum_v (|\Delta_v(t)| - |\Delta_v(t-1)|)$ over these neighbours. By Lemma 3(b) each term is $|\Delta \pm 1| - |\Delta|$ with a uniform sign given H_{t-1}, whose conditional mean is $[\Delta = 0]$; hence $\mathbb{E}[Y_t \mid H_{t-1}] = z_t$. Summing over $t \leq T$ gives $\Phi_T - \Phi_0 = -G_T + \sum_t Y_t$, which is the identity with $M_T = \sum_{t \leq T} (Y_t - z_t)$. The number of terms in Y_t is at most D_{x_t}, so $Y_t \in [-D_{x_t}, D_{x_t}]$ and $z_t \in [0, D_{x_t}]$, whence $|Y_t - z_t| \leq 2D_{x_t}$. The tail is the Azuma–Hoeffding inequality [13, 14]. □

The potential says that bias cannot be destroyed: assigning a neighbour moves a gap by one step of a fair walk, which does not lower $|\Delta_v|$ in expectation, whatever the rule does. The term $\sum_t z_t$ is the source of the $\sqrt{d}$ gaps of balanced formulas (a walk at 0 moves away); for biased formulas we drop it.

Lemma 7 (Shortfall and martingale)**.** *For every branching rule, with $V = \sum_v ((D_v + u_v)^2 + 1)$ and $E = \sum_v D_v^2$:*

(a) $\Pr\big[S_T \leq \frac{1}{4} G_T^{\text{all}} + \frac{T}{2} - \alpha\big] \leq e^{-\alpha^2/(2V)}$*, where $G_T^{\text{all}} = \sum_{t \leq T} |\Delta_t| \geq G_T$;*

(b) $\Pr[Q_T \le -\kappa] \le e^{-2\kappa^2/E}$;

(c) $U_T \le \frac{1}{2}\sum_{w \text{ free at } T} D_w$ *for every walk.*

Proof. (a) $S_T = \sum_{t \le T} X_t$ with $X_t = \xi_t(|\Delta_t|/2 + 1) \ge 0$. By Lemma 3(a), $\mathbb{E}[X_t \mid H_{t-1}] = |\Delta_t|/4 + \frac{1}{2}$ and $\mathbb{E}[X_t^2 \mid H_{t-1}] = \frac{1}{2}(|\Delta_t|/2+1)^2 \le (D_{x_t}+u_{x_t})^2+1 =: v_t$, using $|\Delta_t| \le |h_{x_t}|+D_{x_t} \le 2(D_{x_t}+u_{x_t})$ and $(a+1)^2 \le 2a^2+2$. For $\lambda \ge 0$, $e^{-\lambda X} \le 1 - \lambda X + \lambda^2 X^2/2$ gives $\mathbb{E}[e^{-\lambda X_t} \mid H_{t-1}] \le \exp(-\lambda \mathbb{E}[X_t \mid H_{t-1}] + \lambda^2 v_t/2)$, so $\exp(-\lambda \sum_{s \le t}(X_s - \mathbb{E}[X_s \mid H_{s-1}]) - \frac{\lambda^2}{2}\sum_{s \le t} v_s)$ is a supermartingale, and $\lambda = \alpha/V$ with $\sum_t v_t \le V$ gives the bound [15]. (b) Lemma 3(c) and Azuma–Hoeffding with $\sum_t (D_{x_t}/2)^2 \le E/4$. (c) $|m_w^1 - m_w^0|$ is at most the number of created clauses containing w, at most D_w. □

Proof of Theorem 5. Write $d = \bar{d}_S$ and $\theta = \varepsilon\eta$. Put $\delta = \theta^2/(256R)$ and $T = \lceil (1-\delta)n \rceil$, so that the free set W at depth T has at most δn variables. By Proposition 1 and $\max_t 2^t \pi_t \le \max\{2^T, 2^n \pi_T\}$ (as π_t is non-increasing), it suffices to show $\pi_T \le 3e^{-cn}$ with $c = \theta^2/(4096R)$: then $\gamma = \min\{\delta, c/\ln 2\} \ge \theta^2/(3000R)$.

Sizes. By (B2), $E = \sum_v D_v^2 \le Rnd^2$ and $V = \sum_v ((D_v+u_v)^2+1) \le Rnd^2 + n \le 2Rnd^2$. By Cauchy–Schwarz, any set W of at most δn variables has $\sum_{w \in W}(D_w + u_w) \le \sqrt{|W| \sum_v (D_v+u_v)^2} \le \sqrt{\delta R}\, nd = \frac{\theta}{16}\, nd$.

The potential. $\Phi_0 = \sum_{v \in S} |h_v| \ge \varepsilon \sum_{v \in S} D_v = \varepsilon |S| d \ge \theta n d$ by (B1). At depth T, $|\Delta_w(T)| \le |h_w| + D_w \le 2(D_w + u_w)$, so $\Phi_T \le 2\sum_{w \in W}(D_w + u_w) \le \frac{\theta}{8} nd$. Take $x = \frac{\theta}{4} nd$ in Lemma 6: except with probability $\exp(-x^2/(8E)) \le \exp(-\theta^2 n/(128R))$,

$$G_T \;\ge\; \theta nd - \tfrac{\theta}{8} nd - \tfrac{\theta}{4} nd = \tfrac{5}{8}\,\theta nd.$$

The three terms. Take $\alpha = \kappa = \frac{\theta}{32} nd$ in Lemma 7. Except with probability $e^{-\alpha^2/(2V)} \le e^{-\theta^2 n/(4096R)}$ we have $S_T \ge \frac{1}{4} G_T - \alpha \ge \frac{5}{32}\theta nd - \frac{1}{32}\theta nd = \frac{4}{32}\theta nd$, and except with probability $e^{-2\kappa^2/E} \le e^{-\theta^2 n/(512R)}$ we have $Q_T \ge -\frac{1}{32}\theta nd$. For every walk, $U_T \le \frac{1}{2}\sum_{w \in W} D_w \le \frac{1}{32}\theta nd$.

The race. Outside the three exceptional events, $Z_T = S_T + Q_T - U_T \ge \frac{4-1-1}{32}\theta nd > 0$, so by Lemma 2 the walk is not alive at depth T. Hence $\pi_T \le 3e^{-\theta^2 n/(4096R)}$. □

Remark 8 (Every density, every rule). Only (B1) and (B2) enter; D_v may be 3 or $n-1$. The branching rule is arbitrary, so the theorem covers in particular the adaptive rule of [3, Section 6] run without domination and propagation: for formulas in which every variable has ratio at most $\beta_0 < \frac{1}{2}$, which are $(1-2\beta_0, 1, R)$-biased, the tree has $2^{(1-\gamma)n}$ nodes whatever the plain steps do. This answers the accumulation question of [3, Problem 17, open question 1] in the weak form "some constant below 2"; the exponent $H(\beta')$ hoped for there remains open and would need the measure.

Remark 9 (Order of the exponent). The circulant formula $C_{n,c}$ with gap εn is $(\varepsilon, 1, 1)$-biased, and every rule of [2] has exactly $\binom{n-1}{a} = 2^{H((1-\varepsilon)/2)n - O(\log n)}$ leaves on it; the sum check does not change this [3, Section 5]. So no analysis of this search can give $\gamma > 1 - H((1-\varepsilon)/2) = \varepsilon^2/(2\ln 2) + O(\varepsilon^4)$, and the order ε^2 in Theorem 5 is right. The constant 3000 comes from three union-bounded tails and the crude bound $|\Delta_t| \le 2(D+u)$; a single exponential moment of Z_T would improve it.

Remark 10 (Why the sum check is needed). For a bounded-degree variable, the probability that it is undominated after its predecessors are assigned is a constant, so the level budget of [2] cannot certify an exponential saving at the last levels; and a single pruning clause compares a shortfall $\varepsilon D_x/2$ with its whole free support, which is decided only at the end. In the race the shortfalls of all clauses add up, while only the variables still free at depth T carry uncertainty; bias of order $\varepsilon\bar{d}$ per clause beats uncertainty of order $\bar{d}$ per free variable as soon as fewer than $\varepsilon n/2$ variables are free. This is the "accumulation over many clauses" that the measure with nearly disjoint counted clauses cannot do [3, Remark 12].

Remark 11 (Better constants from the order). With the fixed order by decreasing degree and $S = V$, the last δn variables have $\sum_{w \in W} D_w \le \delta n \bar{d}$ without the square root, so δ can be taken of order ε/R and the margin is linear in ε; only the tails keep the ε^2. With partial bias, put S last. The theorem is stated for every rule because the adaptive direction of Section 5 needs that freedom.

Remark 12 (What is not covered). Domination and propagation are sound but change which values are coins; the count is for the plain search. Complementary pairs ($B = 2$) give steps ± 2 and change constants once the race identity with multiplicities is written out [8, Section 4]. Weighted clauses are not covered.

		$d = 12$	32	128	150
random signs, $\varepsilon = \frac{1}{4}$	single clauses	0.86	0.77	0.57	0.60
	with the sum check	0.90	0.91	0.89	0.93
random signs, $\varepsilon = \frac{1}{2}$	single clauses	0.90	0.86	0.81	0.78
	with the sum check	0.92	0.93	0.94	0.95
aligned, $\varepsilon = \frac{1}{4}$	single clauses	0.74	0.65	0.51	0.55
	with the sum check	0.72	0.72	0.51	0.47
aligned, $\varepsilon = \frac{1}{2}$	single clauses	0.84	0.73	0.68	0.70
	with the sum check	0.85	0.76	0.76	0.69

Table 2: Saving $n - \log_2(\text{tree size})$ as a fraction of n on biased formulas with $n = 300$ (fixed order, population splitting with $M = 40$). Each pair carries a clause with probability $d/(n-1)$; every variable has $|h_v| \ge \varepsilon D_v$. "Aligned" means that every clause is $(u \vee v)$ or $(\neg u \vee \neg v)$ after renaming each variable to its majority literal, the type of $C_{n,c}$.

Table 2 shows the two faces of the theorem. With random signs the sum check keeps the saving flat in d while single clauses decay, exactly as in the balanced case [6, Table 2], but at a much higher level. On aligned formulas the sum has no opposite literals to cancel and gives nothing; the saving is then the bias alone, and it is still flat in d.

4 Three regimes, one framework

The race separates the formulas by what wins it. Write N_x^f for the free neighbours of x at a node, and for a free variable w let $b_w = m_w^1 - m_w^0$ be the *aggregate*: the excess of live clauses in which $w = 1$ is bad over those in which $w = 0$ is bad.

1. **Biased.** Gaps of order εD give a shortfall of order $\varepsilon \bar{d}$ per clause, which beats the trivial uncertainty D_w of each free variable once few variables are free. Theorem 5: every rule, every density, no cancellation.

2. **Balanced, unaligned.** Gaps are of order $\sqrt{D}$ (the term $\sum_t z_t$ of Lemma 6: a gap at 0 moves away), so the shortfall is of order $\sqrt{d}$ per clause and the race is won only if $|b_w|$ is of order $\sqrt{m_w}$ rather than m_w: the bad values of the clauses containing w must cancel. Random signs give this [7]; for a fixed formula the aggregate pair-count condition (C1) of [10] certifies it, and Lemma 6 replaces the gap-profile condition (C3) without Talagrand's inequality and without (C1), under the degree-spread condition (B2) alone (Section 4.1).

3. **Aligned.** Many clauses agree about the bad values of many variables, so b_w is of order m_w and the sum is useless; by the barrier of [3, Section 9] no rule implied by single

flips helps. The exits are structural: near-twin compression [12] and exact interaction classes [4].

The research direction is then a single question.

Problem 13 (Dichotomy at a node)**.** *Find a polynomial-time test at a node, in terms of gaps, aggregates b_w and agreement counts between signed neighbourhoods, with the following property. Either the test names a free variable whose pruning clause raises Z by a constant fraction of its support size, so that the race is won along the walk; or it names a set of variables whose signed neighbourhoods agree on all but a constant fraction of their supports, so that near-twin compression or class counting solves the residual formula within the budget. Then prove that one of the two happens at every node of every formula of density $\omega(1)$.*

The near-twin note proves the second exit for pairs and leaves open whether a failure of cancellation forces linearly many near-twin pairs [12, Section 5]. The zero-gap Hadamard example of [10, Section 6] shows that pairwise cancellation and alignment are not the only obstacles: there the bad values cancel exactly, but no clause has a shortfall. Lemma 6 explains it: with $h = 0$ and no earlier neighbours, $\sum_t z_t$ is the only source of gaps, and the left side, branched first, never has an assigned neighbour.

4.1 The balanced regime without Talagrand

The deterministic criterion of [10] needs (C1) for two things: the martingale Q, the uncertainty U, and, through the addendum [11], the concentration of the total gap. The potential gives the last one directly.

Proposition 14. *For a fixed order, $\mathbb{E}\, G_T^{\mathrm{all}} \geq \sum_{t \leq T} \max\{|h_t|, \sqrt{D_t^- /3}\}$, where D_t^- counts the neighbours of x_t before it, and for every $x > 0$*

$$\Pr[G_T^{\mathrm{all}} < \mathbb{E}\, G_T^{\mathrm{all}} - x] \leq \exp\Big(-\frac{x^2}{8 \sum_v D_v^2}\Big).$$

Proof. The mean is the computation of [11, Proposition 1]: Δ_t is h_t minus a symmetric sum of D_t^- fair signs, so $\mathbb{E}|\Delta_t| \geq |h_t|$ by Jensen and $\mathbb{E}|\Delta_t| \geq \mathbb{E}|X_t| \geq \sqrt{D_t^- /3}$ by Hölder. For a fixed order G_T^{all} is a function of the independent values $a_1, \ldots, a_T$, and changing a_u changes the gap of each later neighbour of x_u by 2, so G_T^{all} changes by at most $2D_{x_u}$. McDiarmid's inequality with these differences gives the tail. For an adaptive rule the same bound follows from Lemma 6, whose martingale has the same increments. □

So under (B2) the condition (C3) of [10] follows from (C3′) with $\eta = g^2/(8R)$, in place of $g^2/(128\tau K)$ in [11], and without the norm bound. What (C1) is still needed for is Q, whose steps are $\frac{1}{2} b_{x_t}$, and U_T: both are sums of bad values, which for a fixed formula are not coins. In clause terms, (C1) says that for every weighting of the branched variables the weighted agreement counts between their signed neighbourhoods stay bounded by Kd times the weight. This is the one place where a quadratic form has not been replaced by a count of clauses; the row-sum version fails on random formulas [11, Remark 5]. We regard it as the definition of "unaligned" until Problem 13 replaces it by an adaptive test.

5 The adaptive rule that the race suggests

Everything the race needs is visible at a node and countable in clause terms.

5.1 The node

Let W be the free variables, $\mathcal{C}$ the live clauses with allowances k_c, and for $w \in W$ let m_w^0, m_w^1 count the clauses of $\mathcal{C}$ in which 0, respectively 1, is the bad value of w. The *aggregate* is $b_w = m_w^1 - m_w^0$ and the *slack* is

$$B = \sum_{c \in \mathcal{C}} k_c - \sum_{w \in W} \min\{m_w^0, m_w^1\} = U - S - Q \ \geq 0,$$

the node being cut exactly when $B < 0$. For a free variable x with gap Δ_x, preferred value $s = \operatorname{sgn} \Delta_x$ and free neighbours N_x^f, the pruning clause created in the child $a_x = -s$ has bad values $\beta_x(w) = -sJ_{xw} \in \{\pm 1\}$ for $w \in N_x^f$ [8, Lemma 4(b)]. Its *agreement* with the aggregate is

$$\begin{aligned} A_x &= \sum_{w \in N_x^f} \left(|b_w + \beta_x(w)| - |b_w| \right) \\ &= \#\{w : \beta_x(w) = \operatorname{sgn} b_w \text{ or } b_w = 0\} - \#\{w : \beta_x(w) = -\operatorname{sgn} b_w \neq 0\}, \end{aligned}$$

and its *value* is $\nu_x = |\Delta_x|/2 + 1 - A_x/2$.

Proposition 15 (Child slacks)**.** *Branching on x gives the children*

$$B_{\text{pref}} = B - \tfrac{1}{2}(|b_x| + s\, b_x), \qquad B_{\text{pruned}} = B - \tfrac{1}{2}(|b_x| - s\, b_x) - \nu_x.$$

Proof. In the child $a_x = a$, U loses $\frac{1}{2}|b_x|$ and Q gains $\frac{1}{2}ab_x$ (Lemma 3(c)), so B loses $\frac{1}{2}(|b_x| + ab_x)$. The pruned child also adds $|\Delta_x|/2 + 1$ to S and $\frac{1}{2}A_x$ to U. □

So exactly one child, the one in which x takes its majority bad value, pays $|b_x|$; this is a Pascal step, tight for every weight. The pruned child pays in addition the value of the new clause, which is positive when the gap is large and the clause disagrees with the aggregate on its free support, and negative when the clause agrees with it.

5.2 The three steps of [3] in these terms

- *Case 2, inside the clauses.* The variable with the largest $|b_x|$: the most contested free variable, where the live clauses disagree most about the bad value. This generalizes "branch on the clause most likely to be violated" [3, Section 4] from one clause to all of them.
- *Case 1, a new clause.* The variable with the largest ν_x: a large gap, and bad values that disagree with the current aggregate. "Counting only sufficiently new clauses" [3, Section 6] becomes a number: a clause is new to the extent that A_x is below its support size, and it is worth ν_x units of slack.
- *A plain step.* $b_x = 0$ and $\nu_x \leq 0$ for every free x: no contested variable and no clause that beats the aggregate.

One-step greed, maximizing $|b_x| + \nu_x$, is the wrong currency early in the tree: a fresh clause has a large free support and $b \approx 0$ there, so $A_x \approx |N_x^f|$ and $\nu_x < 0$ for every balanced x, although the race is won in the end. The reason is timing [2, Remark 29]: the support of a clause is paid into U at creation and repaid through Q, as fair coins, when the support is assigned. The right currency is what is left at the end.

5.3 A reserve

Fix in advance a set W_0 of δn variables, the *reserve*, to be assigned last. This is the set L of [1, Section 7], with a weaker demand: not that every reserved variable be dominated, but that the summed inequality over W_0 have few completions [9]. Branch on $V \setminus W_0$ by any rule; at depth $n - \delta n$ enumerate the completions of the summed inequality over W_0, or continue the search inside W_0.

Proposition 16 (What the reserve costs)**.** *Let the rule on $V \setminus W_0$ be arbitrary. For the uniform walk let $s = S + Q$ at depth $n - \delta n$, where Q is over the assigned support variables, and let $E_W = \sum_{w \in W_0} b_w^2$ be the energy of the aggregate on the reserve. Then the number of completions of the summed inequality over W_0 at that node is at most $2^{\delta n}\rho$, where $\rho = \exp(-2s^2/E_W)$ if $s > 0$ and $\rho = 1$ otherwise, and the total work of the search is $O^*\big(2^{(1-\delta)n} + 2^n\,\mathbb{E}\rho\big)$.*

Proof. A completion $a \in \{\pm 1\}^{W_0}$ satisfies the summed inequality iff $s + \frac{1}{2}\sum_{w \in W_0} a_w b_w \le 0$; for uniform a this is a Hoeffding tail [10, Lemma 1]. Knuth's identity gives the total. □

So the rule on $V \setminus W_0$ should make s^2/E_W large: large shortfalls, and an aggregate on the reserve that stays balanced. Creating the clause of x changes s by $|\Delta_x|/2 + 1$ and E_W by $|N_x^f \cap W_0| + 2\sum_{w \in N_x^f \cap W_0} b_w \beta_x(w)$, where the last sum is the correlation of the new clause with the aggregate on the reserve, the sum of the pair counts C_{xc} of [10, (1)] over the live clauses c.

Rule R (designed cancellation). Write $\rho_x = \sum_{w \in N_x^f \cap W_0} b_w \beta_x(w)$ for the correlation of the new clause with the aggregate on the reserve. To first order, creating the clause of x changes s^2/E_W by $(s/E_W)\big(|\Delta_x| + 2 - (s/E_W)(|N_x^f \cap W_0| + 2\rho_x)\big)$. Among the free variables outside the reserve, branch on the x that maximizes

$$|\Delta_x| + 2 - \frac{s}{E_W}\big(|N_x^f \cap W_0| + 2\rho_x\big)\,:$$

a large gap, and bad values that disagree with the aggregate on the reserve. When every free x has a large positive ρ_x, prefer the largest $|b_x|$ instead. On biased formulas any rule wins (Theorem 5); on balanced formulas the correlation term is the only control over E_W we have, and it is a count of agreeing minus disagreeing bad values, exactly the quantity whose aggregate bound is (C1).

The exit. If at some node every free x outside the reserve has correlation at least $(1 - \kappa)|N_x^f \cap W_0|$ with the aggregate, then every remaining row J_{x,W_0} agrees with $-s_x \operatorname{sgn}(b)$ on all but a κ-fraction of its support: the clauses between the remaining variables and the reserve are, after renaming the reserve by $\operatorname{sgn} b$, all of the same type up to exceptions. This is the near-twin structure of [12] with respect to W_0; what it does not control are the clauses inside the remaining set and inside the reserve. Making this exit an algorithm with a certified cost is Problem 13 in its concrete form.

6 Measures

The analyses above use five quantities as measures. None of them is additive over variables, and that is the point: additive measures certify a saving branch by branch, which fails for linear gaps [2, Section 7.2].

1. **Knuth's measure** $\max_t (t + \log_2 \pi_t)$. It is exact for every search in which every live node has two children, it absorbs plain steps, and it reduces the analysis to one survival probability. Theorem 5 is an analysis of this measure; so is the Key Lemma [7].

2. **The volume of the summed inequality.** At a node orient every free w so that its cheaper value is 0, let $c_w = |b_w|$ and $V(c, B) = \#\{z \in \{0,1\}^W : \sum_w c_w z_w \le B\}$, computable by the knapsack dynamic program in polynomial time [4, Section 3]. Put

$$\mu_\lambda = \#\{w \in W : b_w = 0\} + \log_\lambda V(c, B),$$

   so that $\mu_\lambda = n$ at the root. For $c_w = 1$ on r variables and $B = k$ this is $r - e_\lambda(r, k)$ with the weights $e_\lambda(r, k) = r - \log_\lambda V(r, k)$ of [3, Theorem 10]; the measure extends them to overlapping clauses by summing them first. A step on a contested variable w satisfies $V(c, B) = V(c \setminus w, B) + V(c \setminus w, B - c_w)$, so it is tight for every λ (Pascal's rule); a step on an uncontested variable has drops 1 and 1; a creation is admissible at base λ iff it raises $e_\lambda = |W| - \log_\lambda V$ by at least $\log_\lambda \frac{1}{\lambda - 1}$, which is the condition of [3, Lemma 7] with the sum in place of one clause. So "a clause counts if it is sufficiently new" has an exact meaning: it counts if it lowers the volume of the summed inequality by a factor $\lambda^{-c(\lambda)}$. With $\lambda = 2$ this is the certificate $N_{\mathbf{1}}$ of [4] and the quantity that the reserve enumeration pays [9]; with $\lambda < 2$ it is the measure of [3], and the plain steps are again the obstacle.

3. **The gap potential** $\Phi = \sum_{v \text{ free}} |\Delta_v|$, a submartingale along the walk of every rule (Lemma 6). It certifies future shortfall: the bias in the formula is conserved in expectation and is spent, $|\Delta_t|$ at a time, when variables are branched.

4. **Shortfall and energy on the reserve**, the pair (s, E_W) of Proposition 16 with the bound $2^{|W_0|} e^{-2s^2/E_W}$ and the exact transitions $s \mapsto s \pm \frac{1}{2} b_w$, $E_W \mapsto E_W - b_w^2$ of [10, Section 7]. This is the measure for designed cancellation.

5. **Alignment.** The number of distinct signed neighbourhood types on the reserve, or the number of near-twin pairs; the measure of the structural exit, with cost $(n + 1)^t$ for t exact classes [4] and $(3/4)^{m/(r+1)}$ for m near-twin pairs [12].

The first and the third are what Theorem 5 needs. The second and the fourth are the two faces of the sum check, exact and Gaussian. The fifth is what is left when the sum fails.

7 Cleaning up: what to keep, and in which order

Core, to be read in this order.

1. [2]: the rules (domination, pruning clauses), the level budget, the tight bound for absolute linear gaps and the lower bound $C_{n,c}$.

2. [3]: pair decisions, the sum rule and its characterization as the linear-relaxation cut, the clause weights, averaging over paths, the barrier for single-flip rules.

3. [6, 7]: the race identity, Knuth's identity and the Key Lemma for random signs.

4. [9, 10, 11]: early enumeration, the deterministic counting criterion and its checkable form.

5. [12] and [4, Section 5]: the structural exits.

6. This note: the race for every rule, Theorem 5, the three regimes, the reserve rule and the measures.

Parked. [5] entirely, except its K_m example, which is in [3], and its joint refutation test, which is [3, Proposition 7]. [4] except Sections 3 and 5. [8]: its theorem is a special case of the counting criterion of [10, 11] with (C1) in spectral form, and the biased part of its class is covered by Theorem 5 without any norm.

One framework. All kept results are statements about one search, pruning clauses with the sum check (domination, propagation and pair decisions as sound additions), counted by Knuth's identity and analysed by the race $Z = S + Q - U$ along the uniform walk. The three regimes of Section 4 say which term wins. The open problem is Problem 13.

Next steps, in order.

1. Write Theorem 5 for $B = 2$ and unit clauses in the input, which needs the race identity with multiplicities in written form ([8, Section 4], item 1), and optimize the constant with a single exponential moment of Z_T.

2. Test Rule R against the fixed order on balanced random formulas and on formulas with planted aligned groups, measuring s^2/E_W at the reserve; the prediction is a constant-factor gain in the exponent on random formulas and a visible exit condition on planted ones.

3. Prove Problem 13 for the reserve: either Rule R keeps $E_W \le C\, s\, \sqrt{|W_0|}$, or the rows J_{x,W_0} of the remaining variables fall into few signed types. The first alternative is a vector balancing statement with chosen order and forced signs; the second is the inverse question of [12].

4. Partial bias with the adaptive measure: recover the exponent $\max\{1-\eta, H((1-\varepsilon)/2)\}$ of [2, Theorem 11] in the relative setting, using Theorem 5 for the biased variables and the clause weights for the rest.

8 A plan for all formulas of growing density

The target is every formula with $m \le \rho n$, $\rho = \omega(1)$. Two facts fix the shape of any such result. Formulas with $2^{\,n - O(n \log d/d)}$ strict local maxima exist at every degree d (disjoint copies of Max-Cut on K_{d+1}, and Hastings' connected versions), and every rule implied by single flips keeps all of them [3, Section 9]. So the algorithm needs two engines and the proof needs a dichotomy: a *race engine* (pruning clauses and the sum check, which wins whenever the bad values cancel) and an *aggregation engine* (counting over groups of variables that act alike), and a lemma that every node of every formula is handled by one of them. Theorem 5 and the Key Lemma are the race engine on the two kinds of formulas where cancellation is free or random. What follows is the plan for the rest.

8.1 Cancellation in clause terms: the gap signs

Along the uniform walk, with $g_s = \operatorname{sgn} \Delta_s$ the sign of the gap of x_s at its branching time and ξ_s the fair coin of Lemma 3(a),

$$b_w = \sum_{s \sim w,\, s \le T} \xi_s\, J_{sw}\, (-g_s) = -\tfrac{1}{2} \sum_{s \sim w,\, s \le T} J_{sw} g_s \;+\; M_w,$$

where M_w is a martingale with at most m_w steps of size $\frac{1}{2}$. So $|b_w|$ is of order $\sqrt{m_w}$ exactly when the gap signs of the neighbours of w, weighted by J_{sw}, cancel. The gap sign g_s is the

sign of $h_s - \sum_{u<s} J_{su}a_u$, a function of the values of the earlier neighbours of x_s; two gap signs $g_s, g_{s'}$ are correlated through the earlier neighbours they share, and by a Littlewood–Offord argument (condition on the shared coins; $|\mathbb{E}\,\mathrm{sgn}(Z+c)| \le \Pr[|Z| \le |c|]$ for the private sum Z)

$$|\mathbb{E}[g_s g_{s'}]| \le C\,\frac{|\langle J_{s,\cdot}, J_{s',\cdot}\rangle_<| + 1}{\sqrt{D_s^- D_{s'}^-}},$$

where $\langle J_{s,\cdot}, J_{s',\cdot}\rangle_<$ is the *signed codegree* of s and s' over the earlier variables: common neighbours with agreeing parities minus those with opposite parities. Summing over the pairs of neighbours of w,

$$\mathbb{E}\Big(\sum_{s\sim w} J_{sw} g_s\Big)^2 \;\lesssim\; m_w + \frac{C}{d} \sum_{s\neq s'\in N(w)} |\langle J_{s,\cdot}, J_{s',\cdot}\rangle|,$$

and to leading order, with signs, the second term is the signed count of 4-cycles through w: pairs of clauses at w and pairs of clauses at a common neighbour u, counted with the parity of the cycle, which is the diagonal entry $(J^4)_{ww}$. This is the combinatorial content of the spectral condition: $\sum_w (J^4)_{ww} = \mathrm{tr}\, J^4 \le \|J\|^2\, nd$, so $\|J\| \le K\sqrt{d}$ bounds the average signed 4-cycle count through a variable by $K^2 d^2$, against d^3 for K_{d+1} and for $C_{n,c}$.

Definition 17 (Alignment of a variable)**.** $\mathrm{al}(w) = \sum_{s\neq s'\in N(w)} J_{sw}J_{s'w}\langle J_{s,\cdot}, J_{s',\cdot}\rangle$, the number of even 4-cycles through w minus the number of odd ones; and $\mathrm{al}^+(w)$ the same sum with absolute values.

Target T1 (the race engine for fixed formulas). Every formula with $\mathrm{al}^+(w) \le \kappa d_w d$ for every w, or with $\mathrm{al}(w) \le \kappa d_w d$ under a control of the lower-order terms, is solved by the search of Section 2 in time $2^{(1-\gamma(\kappa))n}$, at every density. The proof is that of the Key Lemma [7] with the sign-correlation bound in place of the late signs, together with the shortfall of Proposition 14. The class contains random formulas, Max-Cut with arbitrary signs on every graph of codegree $O(1)$ (for example girth at least 5), and every formula with $\|J\| \le K\sqrt{d}$ on average; its definition is a local count. The absolute version is enough for $d \le \sqrt{n}$; for denser formulas the signed version is needed, and its error terms are the first thing to work out.

8.2 Aggregation when the gap signs are correlated

If $\mathrm{al}^+(w)$ is large, then many pairs s, s' of neighbours of w have signed codegree of order d: similar signed neighbourhoods, hence correlated gap signs, which is what defeats the sum. The two exits we have need almost identical neighbourhoods: near-twin compression needs discrepancy $r = O(1)$ [12], exact classes need discrepancy 0 [4]. The missing tool is an aggregation for groups of variables with pairwise signed codegree $(1-\kappa)d$: write $J = A + N$ with A constant on the blocks of the groups and N the discrepancies, enumerate the group counts ($2^{O(n \log L/L)}$ choices for groups of size L), which fixes the contribution of A, and search the residual formula N, which is unaligned. The obstacle is soundness: pruning clauses come from single flips, and a flip changes a group count. Inside a count-constrained subproblem the pruning rule must therefore come from *swaps*, a variable and a group-mate flipped in opposite directions; this is the class-flip rule of [3, Proposition 2], whose window ([3, Proposition 1]) is the pruning clause of a swap, with the shared neighbourhood cancelled and a cardinality condition on the discrepancies.

Target T2. Unions of aligned blocks (cliques with constant J, or copies of one signed neighbourhood) joined by a residual of degree r: time $2^{o(n)}$ times the cost of the residual search. For $r = O(1)$ this is [12]; for $r = \omega(1)$ it needs swap pruning on the residual with the sum check, analysed by the race with the count constraints as live clauses (they add to m_w^0 and m_w^1 equally, so they do not disturb b_w).

8.3 The dichotomy and its first test

Target T3. If the average of $\mathrm{al}^+(w)$ over the free variables exceeds $\kappa d_w d$ at a node, then a linear number of free variables lie in groups of pairwise signed codegree at least $(1-\kappa')d$. This is a statement about the graph of pairs with large signed codegree, of Kővári–Sós–Turán or dependent-random-choice type, and it is the risky step: a blow-up in which each variable has L copies with their neighbourhoods re-signed at random on a $(1-\kappa)$-fraction of the positions has pairwise codegree κd for κ well below 1, defeats the sum for $L \to \infty$ (the aggregate on a free variable grows like $\sqrt{Lm_w}$), and is not compressible by near twins. The first experiment is therefore the blow-up family with $L \in \{2, 4, 8, 16\}$ and $\kappa \in \{0.5, 0.8, 0.95\}$: measure $E_W = \sum_w b_w^2$ and the margin of the race, then the same with count enumeration and swap pruning. Its outcome decides the form of T3.

Milestones. (M1) T1 in the absolute form, $d \le \sqrt{n}$: the first worst-case class of growing density defined by a local count and strictly beyond random signs. (M2) T2 with swap pruning, first for bounded r where it must reproduce [12], then for $r = o(\sqrt{d})$. (M3) The blow-up experiment and T3 in the form it supports. (M4) The algorithm: at every node compute $\mathrm{al}^+(w)$ for the free variables, run the race engine when it is small and otherwise extract correlated groups and aggregate; the proof is T1+T2+T3. If T3 fails, the fallback is a theorem for the union of the T1 class and the near-twin class, both of unbounded density and both defined by counts of clauses. The measures of Section 6 are the quantities this plan monitors: $\mathrm{al}^+(w)$ decides the engine, E_W tracks the race, and the group size L prices the aggregation.

9 Checks

`explore_biased_class.py` reuses the search and the population splitting estimator of `explore_key_lemma.py`.

- Without arguments (minutes): Table 2, with the verification that every generated formula satisfies (B1) for the stated ε (the minimum ratio $|h_v|/D_v$ is printed).
- `terms` (a minute): the race terms at depth $0.8n$ and $0.9n$ on biased formulas with $n = 1000$, $d \in \{6, 30, 200, 500\}$ and $\varepsilon \in \{\frac{1}{4}, \frac{1}{2}\}$, in units of $n\bar{d}$: S_T between 0.08 and 0.30, U_T between 0.004 and 0.04, Q_T within ± 0.002. The race is won by a factor 5 to 20 at every density, which is the mechanism of the proof; the proof's margin $\frac{2}{32}\varepsilon\eta$ is far below it.

References

[1] Exact Max 2-SAT with an absolute linear polarity gap. Working note, 23 September 2026 (`max2sat_absolute_linear_gap_proof.tex`).

[2] Max 2-SAT and the domination rule: results and ideas so far. Working note, 5 October 2026 (`max2sat_results_and_ideas.tex`).

[3] Max 2-SAT with pruning clauses: pair decisions, sums, and a measure. Working note, 6–7 October 2026 (`max2sat_pairs_sums_measure.tex`).

[4] Exploring the branching frontier for Max 2-SAT. Draft, 7 October 2026 (`max2sat_branching_frontier.tex`).

[5] A learning and coverage dichotomy for Max 2-SAT. Draft, 7 October 2026 (`max2sat_learning_dichotomy.tex`).

[6] A simplified plan for Max 2-SAT: one search, one count, one Key Lemma. Working note, 8 October 2026 (`max2sat_plan_key_lemma.tex`).

[7] The Key Lemma for random signs. Working note, 8 October 2026 (`max2sat_key_lemma_proof.tex`).

[8] Max 2-SAT below 2^n on unaligned formulas. Working note, 8 October 2026 (`max2sat_explicit_class.tex`).

[9] Max 2-SAT by early constraint-guided enumeration. Working note, 8 October 2026 (`max2sat_early_enumeration.tex`).

[10] Replacing random signs by neighborhood counts. Working note, 8 October 2026 (`max2sat_counting_conditions.tex`).

[11] A checkable replacement for the gap-profile condition. Addendum, 8 October 2026 (`max2sat_counting_addendum.tex`).

[12] Near-twin neighborhoods as a structural exit. Working note, 8 October 2026 (`max2sat_near_twin_pairs.tex`).

[13] K. Azuma. Weighted sums of certain dependent random variables. *Tôhoku Mathematical Journal* 19(3):357–367, 1967.

[14] W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association* 58(301):13–30, 1963.

[15] A. Maurer. A bound on the deviation probability for sums of non-negative random variables. *Journal of Inequalities in Pure and Applied Mathematics* 4(1), Article 15, 2003.