Max 2-SAT with Pruning Clauses: Pair Decisions, Sums, and a Measure

6 October 2026, revised 7 October 2026

Working notes

Abstract

This note continues [**?**]. It records the approach we now pursue, the partial results, and the questions they raise; the revision of 7 October incorporates results of the draft [**?**]. (i) Pair decisions $u = v$ versus $u \neq v$ can be chained if every condition in a $\neq$ child flips the whole class of variables tied together by earlier decisions; flipping single variables loses optima, already for two clauses. (ii) Branching on variables of the active pruning clauses makes clauses cut earlier but does not change how the saving decays with the degree. (iii) Adding up the active pruning clauses gives a sound cut rule in which opposite literals of a shared variable cancel. On random formulas it saves about $0.54n$ at every density we tried, up to $n = 800$, while single-clause checks fall to $0.19n$. Weighted sums cut exactly when the linear relaxation of the active clauses is infeasible. (iv) For a measure-and-conquer analysis with weights on pruning clauses we compute the optimal weights. Capped at one, the branching number depends on the absolute allowance of the clause and tends to 2. Without the cap the weights have a closed form, the clause length minus the logarithm of the number of permitted completions, and the branching number depends only on the relative allowance β and tends to $2^{H(\beta)}$, which is tight. (v) The bound needs plain steps to be rare along one random walk, not on every path. (vi) For formulas in which every variable has a linear gap relative to its degree, we prove the parts that concern branching on near-dominated variables and, for the rule without release, the resolution of one clause. What remains is the accumulation over many clauses; in experiments plain steps cost at most a constant factor, even on the circulant family. (vii) All these rules except pair decisions keep every strict local maximum, so the number of strict local maxima bounds the size of their search trees from below. Reaching $(2 - \delta)^n$ at growing density therefore needs a rule that moves several variables at once, such as class decisions.

1 The approach

Goal. As in [**?**]: a deterministic algorithm for Max 2-SAT with running time $O^*(c^n)$, $c < 2$, in polynomial space, for formulas of every clause density. The tools are DPLL with the domination rule [**?**, **?**] and pruning clauses [**?**].

Plan. We analyse the search with a measure in the spirit of measure and conquer: the number of free variables minus a weight for each counted pruning clause. A clause is counted only if it brings enough variables that no other counted clause contains. The branching steps fall into three kinds.

- *Case 1*: branch on a variable outside the counted clauses whose pruning clause will be counted. Variables close to domination give strong clauses.

- *Case 2*: branch on a variable of a counted clause. In one child the clause gets shorter; in the other it gets tighter, and when its allowance is used up it propagates, fixing its remaining variables.
- *Plain steps*: neither is possible.

The analysis is amortized over general scenarios rather than an exact case analysis. It has a cold start: in a balanced formula the first pruning clauses are weak, and strong ones appear only after many branchings. The plan is therefore to analyse nodes with suitable structural properties and to show that many nodes have them.

Three kinds of nodes. The results below suggest a split.

1. Many free variables are close to domination: the measure, with cases 1 and 2 (Sections **??**–**??**).
2. The formula is balanced, and the active clauses carry different literals on shared variables: sums of pruning clauses (Section **??**).
3. Aligned structure, as in near-twins and the circulant family $C_{n,c}$ of [**?**]: pair decisions (Section **??**).

Section **??** sharpens this split: every rule except pair decisions keeps all strict local maxima, so on formulas with many of them only rules that move several variables at once can help.

2 Setting

We use the notation of [**?**, Section 2] with $B = 1$: a canonical formula has at most one binary clause per pair. For a literal ℓ the deficit is $p_\ell = d_2(\neg\ell) - d_1(\ell) + d_1(\neg\ell)$, the free degree of a variable v is $D_v = p_v + p_{\neg v}$, and v is dominated when $\min\{p_v, p_{\neg v}\} \le 0$. We call $\min\{p_v, p_{\neg v}\}/D_v$ the *ratio* of v; it is at most $\frac{1}{2}$.

When the search branches on x, let ℓ be the literal with the smaller deficit $p = p_\ell \ge 1$. The child with ℓ true is the *preferred* child. The other child, the *pruned* child, receives the pruning clause $P_x = \langle L_x \le p - 1\rangle$, where L_x consists of the literals $\neg m$ for the clauses $(\ell \vee m)$ and m for the clauses $(\neg\ell \vee m)$. By the flip identity [**?**, Lemma 18] an assignment that violates P_x is not worse after flipping x, and the search with domination, cuts and propagation is exact for every branching rule [**?**, Proposition 20].

Clause states. At a node, a pruning clause is in *state* (r, k) if r of its literals are unassigned and at most k of them may still become true. Branching on x creates the state $(D_x, p - 1)$. When one of its literals is assigned, the state moves to $(r - 1, k)$ if the literal is false and to $(r - 1, k - 1)$ if it is true. The clause is *vacuous* when $k \ge r$, it *propagates* when $k = 0$, setting all its remaining literals false, and it is violated when $k < 0$, in which case the node is cut.

3 Pair decisions

3.1 One decision

For a complete assignment α and a set C of variables, let α^C be α with all variables of C flipped and $G_C(\alpha) = \text{score}(\alpha^C) - \text{score}(\alpha)$; write G_u for $G_{\{u\}}$. A *pair decision* on unassigned variables u, v has the children $u = v$ and $u \ne v$. The first is explored in full. In the second

we keep only the assignments with $G_u(\alpha) < 0$ and $G_v(\alpha) < 0$; this is sound, because if $G_u(\alpha) \geq 0$, then $\alpha^{\{u\}}$ has $u = v$ and is at least as good.

Let α have $u = 1$ and $v = 0$; the other case is symmetric. By the flip identity, $G_u(\alpha) < 0$ is the pruning clause of u with respect to $\neg u$, and $G_v(\alpha) < 0$ is the pruning clause of v with respect to v. Call a common neighbour w of u and v *shared* if its clause with u and its clause with v have the same signs, and let ρ_u, ρ_v be the numbers of the remaining clauses of u and of v; a clause on $\{u, v\}$ counts for both. On a shared w the two pruning clauses contain opposite literals of w.

Proposition 1 (Window)**.** *Write $G_u(\alpha) = \sigma + a$ and $G_v(\alpha) = -\sigma + b$, where σ is the contribution of the clauses with shared neighbours and a, b are the contributions of the remaining clauses, so that $|a| \leq \rho_u$ and $|b| \leq \rho_v$. The child $u \neq v$ keeps α only if $b + 1 \leq \sigma \leq -a - 1$. In particular:*

1. *$a + b \leq -2$, a condition on the non-shared clauses alone; it is the sum of the two pruning clauses, in which the shared part cancels;*
2. *σ takes at most $\rho_u + \rho_v - 1$ values;*
3. *if $\rho_u + \rho_v \leq 1$, the child is empty and u, v can be merged.*

Proof. For a shared w, the clause of w with v at $v = 0$ is the same function of w as the clause of w with u at $u = 0$, and likewise at the value 1. So flipping v from 0 to 1 changes it by minus the amount by which flipping u from 1 to 0 changes the clause of w with u. This gives the form of G_u and G_v; the rest is arithmetic. □

3.2 Chaining decisions

Along a path, decided pairs tie variables into classes. Write $[u]$ for the class of u; the decisions fix every relation inside it, and flipping a whole class preserves these relations.

Proposition 2 (Chained pair decisions)**.** *Run the search of [?] on classes: after a pair decision on u, v the two classes are merged in both children, with $u = v$ or $u = \neg v$; value branchings, domination and pruning clauses apply to whole classes; and in the child $u \neq v$ only the assignments with $G_{[u]}(\alpha) < 0$ and $G_{[v]}(\alpha) < 0$ are kept, for the classes at the time of the decision. Then the search is exact.*

Proof. As in [**?**, Proposition 20], order the complete assignments by the first decision at which their paths part, preferring the child $u = v$ of a pair decision and the preferred child of a value branching, and let α^* be the largest optimal assignment. Suppose α^* is cut in the child $u \neq v$ because $G_{[u]}(\alpha^*) \geq 0$. Flipping all of $[u]$ changes no relation inside $[u]$, and no earlier decision relates $[u]$ to another class, since such a decision would have merged them; also no variable of $[u]$ is assigned above the decision. So $\alpha^{*[u]}$ follows the path of α^* to the decision, takes the child $u = v$ there, and is optimal, a contradiction. The case $[v]$ and the other kinds of cuts are handled as in [**?**]. □

Equivalently, a pair decision removes one free variable in each child, as a value branching does, and the formula over merged variables turns a clause inside a class into a unit clause or a constant.

Single flips are not enough. Take $(x_0 \vee x_1)$, $(\neg x_1 \vee \neg x_2)$ and decide $x_0?x_1$, then $x_1?x_2$. The optimum 2 is reached by 010, 100, 101 and 110 (values of $x_0x_1x_2$). The assignment 110 lies in the child $x_1 \neq x_2$ below $x_0 = x_1$. Flipping x_1 alone gives 100, also optimal, so 110 would be cut on the grounds that 100 lies in the child $x_1 = x_2$. But 100 has $x_0 \neq x_1$ and lies below the other child of the first decision, where flipping x_1 gives back 110 and cuts 100. The two cuts point at each other. Also 010 and 101 are cut in this way, and only 000 and 111, of score 1, survive. Flipping the class $\{x_0, x_1\}$ instead turns 110 into 000, a strict loss, and 110 survives.

On chains $x_0?x_1$, $x_1?x_2, \ldots$ in the circulant formulas $C_{9,2}$, $C_{11,3}$, $C_{13,3}$, the single-flip version loses the optimum (it finds 18, 33, 39 instead of 28, 43, 60), while the class version keeps it, and only 10, 10 and 14 complete assignments pass all its conditions.

Remark 3. In $C_{n,c}$ consecutive variables are near-twins: their environments differ in two neighbours. That so few complete assignments pass suggests that pair decisions attack the family that defeats every single-variable rule [**?**, Theorem 13]. These are complete assignments, not tree nodes, however: the class conditions involve the whole boundary of a class, and when they can be checked during the search is open.

4 Branching strategies and timing

Measuring trees. For $n \geq 50$ we estimate tree sizes by population splitting [**?**], a variant of Knuth's estimator [**?**]: keep M random live nodes of each branching depth, expand all their children, estimate the number of live nodes of the next depth, and resample. The estimate is unbiased. On formulas with $n = 30$ and 36 it agrees with the exact count within 0.3 bits, already for $M = 8$. We report the *saving* $n - \log_2(\text{tree size})$ as a fraction of n, averaged over two random formulas, for random balanced formulas: each pair carries a clause with probability $d/(n-1)$, with uniformly random signs.

Branching inside clauses. The rule *inside clauses* branches on an unassigned variable of the active pruning clause that is most likely to be violated, i.e. whose state (r, k) maximizes $\Pr[\mathrm{Bin}(r, \frac{1}{2}) \geq k + 1]$. Table **??** compares it with a random static order.

average degree d	16	32	64	128	200
random order	0.64	0.51	0.37	0.31	0.26
inside clauses	0.80	0.64	0.48	0.39	0.30
random order and the sum rule (Section **??**)	0.69	0.64	0.62	0.54	0.54

Table 1: Saving as a fraction of n, for $n = 400$.

The strategy helps at every density, by about a quarter, but the saving still decreases with d. For the random order it is close to $1.55\, n\sqrt{(\ln d)/d}$, the timing estimate of [**?**, Remark 29].

Remark 4 (The timing in clause terms). A balanced variable with D free neighbours and imbalance M, typically about $\sqrt{d}$, gets a clause allowing about $(D - M)/2$ of its D literals to be true. Under random values about half of the assigned literals are true, so the clause can cut only once about $D - M$ of its literals are assigned. Branching inside a clause brings it there faster, but each such branching opens a new clause whose support is almost entirely new: in a random formula two neighbourhoods share about d^2/n variables. So few clauses are near the end of their supports at the same time. Random formulas have no near-twins, so assuming that every two variables have different neighbourhoods does not remove this effect.

The *coverage strategy*, which branches where the new clause brings the most new variables and otherwise on clause variables adjacent to uncovered ones, has not been tested yet.

5 Sums of pruning clauses

Let I be the set of active pruning clauses at a node that are not vacuous; vacuous clauses can never be violated and would only weaken a sum. For an unassigned variable w, let m_w^1 and m_w^0 be the numbers of clauses of I in which $w = 1$, respectively $w = 0$, makes a literal true, and let k_i be the allowance of clause i.

Proposition 5 (Sum rule)**.** *If*

$$\sum_{w \text{ unassigned}} \min\{m_w^0, m_w^1\} > \sum_{i \in I} k_i,$$

then no completion of the node satisfies all clauses of I, and the node can be cut.

Proof. A completion that satisfies the clauses makes at most $\sum_i k_i$ further literals of them true. Whatever value w gets, it makes at least $\min\{m_w^0, m_w^1\}$ of their literals true. □

The sum is implied by the clauses, so the search stays exact; we checked the optimum against brute force on 150 formulas with $n \le 12$, in 42 of which the rule changed the tree. A variable shared with opposite literals contributes in advance, whatever its value; one shared with the same literal contributes nothing. This is the algorithmic side of [**?**, Lemma 27]: shared opposite literals give a smaller joint survival probability than independent clauses. Proposition **??**(1) is the case of two clauses.

Example. $\langle\{w_1, \dots, w_7\} \le 3\rangle$ and $\langle\{\neg w_1, \dots, \neg w_7\} \le 3\rangle$ say that at most three of the w_i are 1 and at most three are 0, which is impossible. Neither clause is violated or propagates at the start. A search with single-clause checks and propagation branches until three equal values appear and has 20 leaves. In the sum every w_i contributes $\min\{1, 1\} = 1$, so $7 > 3 + 3$ and the node is cut at once.

Remark 6 (Why the sum decides early)*.* This is a heuristic count. Suppose a path carries k active clauses, created at variables of imbalance about $\sqrt{d}$, so that every bound lies about $\sqrt{d}/2$ below the expected count of its clause. These shortfalls point the same way and add up to order $k\sqrt{d}$. In a random formula an unassigned w lies in about $m_w = kd/n$ of the clauses, with independent literal signs, so the uncertain part $\sum_w |m_w^1 - m_w^0|$ over the u unassigned variables is of order $u\sqrt{kd/n}$. The sum decides once this falls below the shortfall, i.e. once $u \lesssim \sqrt{kn}$. Since k grows in proportion to the number of assigned variables, this happens while a constant fraction of the variables is unassigned, and d cancels.

n	50	100	200	400	600	800
single-clause checks	0.48	0.39	0.29	0.26	0.22	0.19
with the sum rule	0.53	0.44	0.51	0.54	0.54	0.58

Table 2: Saving as a fraction of n on dense random formulas ($d \approx n/2$), random order. For $n = 600$ and 800: one formula and $M = 40$.

With single-clause checks the saving grows roughly like $2\sqrt{n \ln n}$; with the sum rule it stays near $n/2$ (Table **??**), and across the degrees of Table **??** it flattens at about $0.54n$. On

the aligned formulas $C_{n,c}$, $15 \le n \le 21$, which have no opposite literals, the sum removes at most 5% of the nodes.

Any nonnegative combination of the active clauses is also sound, and the family of all of them has an exact description [**?**].

Proposition 7 (Sums are linear-relaxation cuts)**.** *Some nonnegative combination of the active clauses $L_i \le k_i$ cuts a node, in the sense of Proposition* **??***, if and only if the linear relaxation $\{x \in [0,1]^r : L_i(x) \le k_i \text{ for all } i\}$ is empty.*

Proof. With weights $a_i \ge 0$ the rule cuts iff the minimum of $\sum_i a_i L_i$ over $\{0,1\}^r$ exceeds $\sum_i a_i k_i$. The function is affine, so this is also its minimum over $[0,1]^r$, and a point of the relaxation rules the cut out. Conversely, if the relaxation is empty, the image of the box under $x \mapsto (L_i(x))_i$, a compact convex set, misses the closed convex set $\{y : y_i \le k_i\}$. A separating hyperplane must have nonnegative coefficients, since the second set is unbounded below in every coordinate, and these coefficients give a cut. □

So sums see exactly what the linear relaxation sees, although the separating weights may be large. Contradictions of parity type are invisible to them: $x + y = y + z = z + x = 1$, written as six cardinality clauses, has the fractional solution $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ but no Boolean one [**?**]. Counting the assignments that satisfy a weighted sum, instead of only testing it, gives a graded certificate that satisfies a branching inequality, together with a sufficient condition for a running-time bound in terms of the progress of this count along a random walk [**?**]. In these terms Remark **??** says, heuristically, that on random formulas the relaxation of the active clauses becomes infeasible while about half of the variables are free.

6 A measure for pruning clauses

6.1 Counting, measure and rule

A clause is counted only on its new variables: when it is created, delete the literals of variables that lie in the support of a counted clause. Deleting literals keeps a pruning clause valid, since it can only lower the count, so the counted clauses are valid and have pairwise disjoint supports. The search may keep the full clauses for cuts; the measure uses only the counted ones. Let

$$\mu = n_{\text{free}} - \sum_{\text{counted } C} \varepsilon(r_C, k_C),$$

where (r_C, k_C) is the state of C and $\varepsilon \ge 0$ is a weight to be chosen. The rule:

1. if a counted clause is active, branch on one of its variables (case 2);
2. otherwise, if a free variable has a *feasible* new clause (defined below), branch on it (case 1);
3. otherwise branch on any variable (a plain step).

A counted clause in a state of weight 0 is *released*: it no longer counts.

6.2 Steps and weights

A step is *admissible* for $\lambda \in (1,2)$ if its children have measure drops d_0, d_1 with $\lambda^{-d_0} + \lambda^{-d_1} \le 1$.

- Case 1 on x with new degree D and deficit p: drops 1 and $1 + \varepsilon(D, p-1)$, or 1 and $1 + D$ if $p = 1$. This is admissible iff $\varepsilon(D, p-1) \ge c(\lambda) := \log_\lambda \frac{1}{\lambda - 1}$; we call the new clause feasible in that case.

- Case 2 in state (r,k): the false child has drop $1+V_f-\varepsilon(r,k)$ with $V_f=\varepsilon(r-1,k)$, or $V_f=0$ if $(r-1,k)$ is vacuous or released; the true child has drop $1+V_t-\varepsilon(r,k)$ with $V_t=\varepsilon(r-1,k-1)$, and $V_t=r-1$ if $k=1$, when the other $r-1$ literals are set false.
- A plain step has drops of at least 1.

Further variables fixed in a child only add to its drop, as long as no forced transition lowers a weight by more than one.

Lemma 8 (Optimal weights)**.** *For $\lambda\in(1,2)$ and a cap $\kappa\in(0,\infty]$ define, for $r=2,3,\dots$ and $1\le k<r$,*

$$\varepsilon(r,k)=\max\Big\{0,\ \min\{\kappa,\ 1-\log_\lambda(\lambda^{-V_f}+\lambda^{-V_t})\}\Big\},$$

with V_f,V_t as above. These are the largest weights, bounded by κ, for which every case-2 step in a state of positive weight is admissible. Moreover $\varepsilon(r,k)\le 1+\min\{V_f,V_t\}$, so no transition lowers a weight by more than one, forced assignments never increase μ, and $\varepsilon(r,k)\le r-1$, so $\mu\ge 0$.

Proof. The step in state (r,k) is admissible iff $\lambda^{\varepsilon(r,k)}(\lambda^{-1-V_f}+\lambda^{-1-V_t})\le 1$, which is the bound on $\varepsilon(r,k)$; it depends only on shorter states, and larger children allow a larger parent. Since $\lambda^{-V_f}+\lambda^{-V_t}\ge\lambda^{-\min\{V_f,V_t\}}$, we get $\varepsilon(r,k)\le 1+\min\{V_f,V_t\}$; a forced literal then changes μ by -1 plus a weight decrease of at most one. Induction on r with $V_f\le r-2$ gives $\varepsilon(r,k)\le r-1$. □

6.3 With and without the cap

Proposition 9 (Cap one)**.** *With $\kappa=1$, for large r the weights approach $1,1-c,1-2c,\dots$ for $k=1,2,3,\dots$, where $c=c(\lambda)$. So case 1 with allowance $k_0=p-1$ is admissible for large D iff $\lambda^{1/k_0}(\lambda-1)\ge 1$.*

Proof. For large r, $V_t=r-1\to\infty$ in state $(r,1)$, so its weight is the cap 1. For $k\ge 2$ a stationary weight ε_k must satisfy $\lambda^{1-\varepsilon_k}\ge\lambda^{-\varepsilon_k}+\lambda^{-\varepsilon_{k-1}}$, i.e. $\varepsilon_k\le\varepsilon_{k-1}-c$. Case 1 needs $1-(k_0-1)c\ge c$, i.e. $k_0c\le 1$. □

allowance $k_0=p-1$	1	2	3	4	8	16
λ with cap one	1.618	1.756	1.820	1.857	1.922	1.959

Table 3: The cap $\varepsilon<1$: the branching number depends only on k_0.

So with the cap λ tends to 2, roughly as $2-(\ln 2)/k_0$: the k_0 units of allowance share one unit of weight. Only variables a bounded number of units away from domination help, and closeness to domination in the relative sense, $p\le\beta D$, gives nothing for large degrees.

Without the cap the weights have a closed form [**?**]. Let $V(r,k)=\sum_{j=0}^{\min\{r,k\}}\binom{r}{j}$ be the number of completions that a clause in state (r,k) permits.

Theorem 10 (No cap)**.** *The weights $e_\lambda(r,k)=r-\log_\lambda V(r,k)$ satisfy the recursion of Lemma* **??** *with equality and without the clipping at zero, if a vacuous state keeps its value $e_\lambda(r,k)=r(1-\log_\lambda 2)<0$ instead of 0. With these weights every case-2 step is tight, no transition lowers a weight by more than one, and creating a clause in state (D,k_0) is admissible if and only if*

$$V(D,k_0)\le(\lambda-1)\lambda^D.$$

Hence, for $k_0 \le \beta D$ with $\beta < \frac{1}{2}$ and any $\lambda > 2^{H(\beta)}$, every creation with $D \ge \ln \frac{1}{\lambda-1} / (\ln \lambda - H(\beta) \ln 2)$ is admissible, and the smallest admissible λ for $k_0 = \lfloor \beta D \rfloor$ tends to $2^{H(\beta)}$ as $D \to \infty$.

Proof. By Pascal's rule $V(r,k) = V(r-1,k) + V(r-1,k-1)$, so $\lambda^{-e_\lambda(r-1,k)} + \lambda^{-e_\lambda(r-1,k-1)} = \lambda^{1-e_\lambda(r,k)}$, which is the recursion with equality; for $k = 1$ the true child has $V(r-1,0) = 1$, i.e. weight $r-1$, as for propagation. A transition changes the weight by $-1 + \log_\lambda \frac{V(r,k)}{V(r-1,k')} \ge -1$, since a restriction cannot increase V. Case 1 is admissible iff $\lambda^{-1} + \lambda^{-1-D} V(D,k_0) \le 1$, which is the stated condition. The threshold follows from $V(D,k_0) \le 2^{H(\beta)D}$, and the limit from $V(D,k_0) \ge \binom{D}{k_0} \ge 2^{H(k_0/D)D}/(D+1)$. □

The weights of Lemma **??** with $\kappa = \infty$ are these weights with *release*: where e_λ would be negative the clause is released and its weight is 0. Clipping only raises weights, so it can only lower the branching numbers, slightly at finite D (Table **??**). With release, every creation with $k_0 \le \beta D$ and $D \le 400$ is already admissible at $\lambda = 2^{H(\beta)}$, for $\beta \in \{\frac{1}{8}, \frac{1}{4}, 0.3, \frac{3}{8}, 0.45\}$.

β		$D = 32$	64	128	256	512	$2^{H(\beta)}$
1/8	with release	1.411	1.419	1.429	1.439	1.446	1.458
	without	1.431	1.437	1.443	1.448	1.452	
1/4	with release	1.668	1.693	1.713	1.728	1.739	1.755
	without	1.695	1.715	1.730	1.740	1.746	
3/8	with release	1.849	1.877	1.898	1.913	1.923	1.938
	without	1.873	1.897	1.912	1.923	1.929	

Table 4: Without the cap: the branching number for creating a clause with $k_0 = \beta D$, with release (Lemma **??**) and without (Theorem **??**).

For example, with release at $\lambda = 1.643$ the weights of the states $(16,k)$ are 10.34, 6.22, 3.10, 0.89 for $k = 1, \dots, 4$, those of $(12,k)$ are 6.89, 3.38, 1.00, and those of $(8,k)$ are 3.66, 1.03. At fixed allowance a longer clause weighs more; at fixed margin $r - 2k$ a shorter one does.

Remark 11 (Tightness). If every variable has ratio β, the bound $2^{H(\beta)n}$ is that of [**?**, Theorem 11] with $\beta = (1-\varepsilon)/2$, and it is attained on $C_{n,c}$ by every branching rule with domination and pruning clauses [**?**, Theorem 13, Observation 22]. So the step analysis is optimal for this class.

Remark 12 (Why one might cap, and why not to). In base 2 a disjoint block saves exactly $\log_2 \frac{2}{1+\theta} < 1$ bit [**?**, Proposition 25], since the preferred child carries no clause: a clause is worth less than one bit to the whole tree. This suggests $\varepsilon < 1$, so that the variable assigned when a clause disappears pays for it. In an analysis by branching numbers, however, the weight must also pay for the steps inside the clause before it propagates, about one per unit of allowance. Without the cap the weight of a tight clause is close to the number of variables it will force: $(16,1)$ weighs 10.34 and forces 15 when it propagates. What the payoff argument needs is Lemma **??**: no transition lowers a weight by more than one.

Remark 13 (What nearly disjoint clauses can certify). Heuristically, for a random formula of degree d: a clause created when its variable has D free neighbours is worth $O(1 + d/D)$, since the imbalance is $O(\sqrt{d})$; at most n_{free}/D disjoint counted clauses coexist; and $D \approx d\, n_{\text{free}}/n$. While n_{free} falls from $2N$ to N, the counted clauses therefore certify $O((n/d)(n/N))$, which pays for all N variables only once $N \lesssim n/\sqrt{d}$. Summing over the stages gives $O(n/\sqrt{d})$ in total, the timing barrier again. Counting only nearly disjoint clauses cannot give c^n for balanced formulas of growing degree.

7 Averaging over paths

Lemma 14 (Averaging)**.** *Let $\lambda \in (1,2]$, let the measure be n at the root and nonnegative everywhere, and let every internal node be either admissible or plain, as in Section* **??***. Let $\mathcal{W}$ be the random walk from the root that at an admissible node moves to child i with probability $q_i \geq \lambda^{-d_i}$, where $q_0 + q_1 \leq 1$ and the walk stops with the remaining probability, and at a plain node moves to each child with probability $\frac{1}{2}$. Let S be the number of plain nodes on $\mathcal{W}$. Then*

$$\#\textit{leaves} \leq \lambda^n \, \mathbb{E}[(2/\lambda)^S].$$

Proof. For a leaf ℓ let $s(\ell)$ be the number of plain nodes on its path and $\Pi(\ell)$ the probability that $\mathcal{W}$ ends at ℓ. The drops along the path add up to at most n, those at plain nodes to at least $s(\ell)$, so the drops at admissible nodes add up to at most $n - s(\ell)$ and $\Pi(\ell) \geq \lambda^{-(n-s(\ell))} 2^{-s(\ell)}$. Hence $\#\text{leaves} = \sum_\ell \Pi(\ell)/\Pi(\ell) \leq \lambda^n \sum_\ell \Pi(\ell)(2/\lambda)^{s(\ell)} \leq \lambda^n \mathbb{E}[(2/\lambda)^S]$. □

Corollary 15. *1. If every path has at most s plain nodes, there are at most $\lambda^{n-s}2^s$ leaves. With the weights of Lemma* **??** *and $\lambda = 2^{H(\beta)}$, this is the bound for the rule of Section* **??** *when every node offers case 1 with $k_0 \leq \beta D$ or case 2, for $D \leq 400$; for larger D take λ slightly above $2^{H(\beta)}$ (Theorem* **??***).*

2. If $\Pr[S > s] \leq \rho$, there are at most $\lambda^{n-s}2^s + \rho\, 2^n$ leaves, since a walk has at most n plain nodes.

3. If $S - \bar{s}$ is sub-Gaussian with variance proxy σ^2, there are at most $\lambda^n (2/\lambda)^{\bar{s}} e^{\theta^2\sigma^2/2}$ leaves, where $\theta = \ln(2/\lambda)$.

Item 2 has the shape of the bound of [**?**]: most branches are good, and the bad ones are exponentially rare. Two points matter.

- The probability is over the tilted walk $\mathcal{W}$, not over uniform assignments. At a case-1 step $\mathcal{W}$ takes the preferred child with probability $1/\lambda$, and it may give the pruned child $1 - 1/\lambda$, which is at least $\lambda^{-1-\varepsilon}$ exactly when the step is admissible.
- The failure probability must be exponentially small, because bad paths together can cost 2^n. In item 3 the price of fluctuations is second order: for $\lambda = 1.755$ and $\sigma^2 = n/4$ the factor is about $2^{0.003n}$.

Variables fixed during plain steps can be credited at rate λ: if $\pi(\ell) \geq s(\ell)$ is the total drop at the plain nodes of the path, the proof gives $\#\text{leaves} \leq \mathbb{E}[\lambda^{n-\pi} 2^S]$. This is how domination after plain steps, the mechanism of [**?**, Theorem 11], enters.

8 Availability in formulas with linear gaps

Assume every variable starts with ratio at most $\beta_0 < \frac{1}{2}$, fix a threshold $\beta' \in (\beta_0, \frac{1}{2})$, let $\lambda = 2^{H(\beta')}$, and let case 1 take the free variable of smallest ratio among those with $p - 1 \leq \beta' D$ and a feasible clause. Since active clauses are finished first, a plain step happens only when no free variable passes the threshold. We want $\mathbb{E}[(2/\lambda)^S] = 2^{o(n)}$.

Each assignment of a neighbour y of v lowers exactly one of the two deficits of v [**?**, Lemma 2]. Call it *helpful* for v if it lowers the smaller one.

Lemma 16 (Window)**.** *Let v start with degree D_0 and smaller deficit $p_0 \leq \beta_0 D_0$. If at least a β_0-fraction of its assigned neighbours were helpful, then the ratio of v is at most β_0.*

Proof. With j assigned neighbours, $h \geq \beta_0 j$ of them helpful, the smaller deficit is at most $p_0 - h \leq \beta_0(D_0 - j)$ and the degree is $D_0 - j$. □

Conversely, assignments that are not helpful lower the other deficit, and once enough of them have occurred, the other literal of v passes the threshold. For $p_0 = \beta_0 D_0$ and no helpful assignments this happens after a fraction $(1 - \beta_0 - \beta')/(1 - \beta')$ of the neighbours, 64% for $\beta_0 = \frac{1}{4}$ and $\beta' = 0.3$. So v fails the threshold only inside a bounded window, and a plain step needs every free variable inside its window at the same time.

Lemma 17 (Case-1 steps)**.** *Let $\mathcal{W}$ take the preferred child of a case-1 step with probability $1/\lambda$ and the pruned child with probability $1 - 1/\lambda$. Then every neighbour of the branched variable is helped with probability at least $1 - 1/\lambda$, and $1 - 2^{-H(\beta)} \geq \beta$ for $0 < \beta \leq \frac{1}{2}$, with equality only at $\beta = \frac{1}{2}$.*

Proof. Exactly one value of the branched variable helps a given neighbour, and both values have probability at least $1 - 1/\lambda$. The inequality is $H(\beta) \geq \log_2 \frac{1}{1-\beta}$, which after expanding H reads $\beta \log_2 \frac{1-\beta}{\beta} \geq 0$. □

With $\lambda = 2^{H(\beta')}$ each case-1 step helps every neighbour with probability at least $\beta' > \beta_0$. If all assignments were case-1 steps, the Azuma–Hoeffding inequality would bound the probability that a variable with m assigned neighbours has a helpful fraction below β_0 by $e^{-2(\beta' - \beta_0)^2 m}$, and Lemma **??** would keep it at the threshold.

The obstacle. In a pruned child, $\mathcal{W}$ then resolves the clause of the branched variable x, which allows at most a β'-fraction of the neighbours of x to take their bad values. These assignments follow the clause, not a coin. In aligned formulas, where after renaming every clause is $(x \vee y)$ or $(\neg x \vee \neg y)$, the bad value of y for the clause of x is its minority value, which is the helpful value for every other neighbour of y. So the other variables see a helpful fraction of about the ratio of x, which is below their own if x has the smallest ratio. The per-variable argument does not cover this, and on $C_{n,c}$ plain steps do occur.

Measurements. On random formulas with linear gaps, with biased signs or aligned, $n \in \{200, 400\}$, d from 20 to 200, $\beta_0 = \frac{1}{4}$ and $\beta' = 0.26$, no walk had a plain step. Even with $\beta' = \beta_0$ on dense aligned formulas, a walk had at most 16 plain steps and $\log_2 \mathbb{E}[(2/\lambda)^S] = 0.10$. Table **??** shows the extreme case.

	$\beta' - \beta_0 = 0.02$		$\beta' - \beta_0 = 0.05$	
n	walks with a plain step	$\log_2 \mathbb{E}[(2/\lambda)^S]$	walks with a plain step	$\log_2 \mathbb{E}[(2/\lambda)^S]$
101	20%	0.49	9%	0.29
201	19%	1.32	12%	0.17
401	23%	2.01	17%	0.55
801	26%	2.73	11%	0.63

Table 5: Plain steps along $\mathcal{W}$ on $C_{n,c}$ with $\varepsilon = 0.5$ ($\beta_0 \approx \frac{1}{4}$), 150 walks (80 for $n = 801$).

So the bound is λ^n up to a factor below 8, that is $2^{H(\beta')n}$. For gaps that are linear in n this exponent is already known for every order [**?**, Theorems 11 and 24]; what is new here is that the gaps are relative to each variable's own degree, so the formulas may have any density, and that the rule is adaptive.

Why, heuristically. In random formulas the variables reach their windows at different times, and the rule takes the best free variable, so one outside its window is always available. In $C_{n,c}$ all variables move together, but $\mathcal{W}$ follows typical surviving assignments. For a uniformly random assignment satisfying a clause, the remaining allowance divided by the remaining length is a martingale, so the bad fraction stays near the ratio of the clause, the minority fraction stays near β_0, and only fluctuations reach the window.

Problem 18 (The missing lemma)**.** *Show that while $\mathcal{W}$ resolves a counted clause created in state (D, k_0), the fraction of its literals that become true is at least $k_0/D - \eta$, except with probability $e^{-\Omega(\eta^2 D)}$.*

For the rule with release this is open. Without release it holds, in a stronger form.

Proposition 19 (One clause without release)**.** *Use the weights e_λ of Theorem **??**, keep every counted clause until its support is assigned, and postpone domination and new clauses while it is active. Then inside the clause $\mathcal{W}$ makes all $V(D, k_0)$ permitted completions equally likely, and for $0 \le k_0 < D/2$, $1 \le t \le k_0$ and $q = k_0/(D - k_0 + 1)$ the number J of its literals that become true satisfies*

$$\Pr[J \le k_0 - t] \le q^t.$$

Proof. With tight weights, $\mathcal{W}$ moves to a child with probability $\lambda^{-\mathrm{drop}} = V(\text{child})/V(\text{parent})$, and these factors telescope to $1/V(D, k_0)$ along every permitted completion. For $j \le k_0$, $\binom{D}{j-1} = \frac{j}{D-j+1}\binom{D}{j} \le q\binom{D}{j}$, hence $\binom{D}{i} \le q^t\binom{D}{i+t}$ for $i \le k_0 - t$; summing over i and dividing by $V(D, k_0)$ gives the bound [**?**]. □

So Problem **??** holds for this rule, with $e^{-\Omega(\eta D)}$ in place of $e^{-\Omega(\eta^2 D)}$: J ends within $O(1)$ of the allowance. Given J, the true literals form a uniformly random subset of the support. Hence a variable with s neighbours in the support, for each of which one value is designated as helpful when the clause is created, sees a helpful fraction of at least $k_0/D - \eta$ among them, whatever the designations, except with probability $q^{\lceil \eta D/2 \rceil} + 2e^{-\eta^2 s/8}$ [**?**]. The price of giving up release is a slightly larger branching number (Table **??**).

This suggests how to finish the argument. By Lemma **??** a variable stays at the threshold while at least a β_0-fraction of its assigned neighbours helped it; case-1 steps help with probability at least β' (Lemma **??**), and a clause passes on a helpful fraction of at least its ratio minus η. So in case 1 we should take the largest ratio below β', not the smallest: when this ratio is close to β', every helpful fraction exceeds β_0, and the creation is still feasible. What remains open is the accumulation of these bounds over many adaptively chosen clauses, in particular for small overlaps s, and the assignments that domination forces between clauses. We have not tested the largest-ratio choice; on random formulas there are no plain steps to remove, and in $C_{n,c}$ all variables have the same ratio.

9 A barrier for single-flip rules

Call a complete assignment a *strict local maximum* if flipping any single variable strictly lowers the score.

Proposition 20 (Strict local maxima are never cut)**.** *Consider a search whose fixings and cuts are domination, pruning clauses with propagation, and sums or weighted sums of active pruning clauses; the full phases and counting certificates of [**?**] are included. For every branching rule, every strict local maximum is a leaf of its search tree. Hence the tree has at least as many leaves as the formula has strict local maxima.*

Proof. Let α be a strict local maximum and follow its path. If domination fixes a literal ℓ, flipping α towards ℓ does not lower the score by the flip identity, so α already has ℓ true. If α enters a pruned child, flipping the branched variable strictly lowers the score, so by the flip identity α satisfies the new pruning clause. Hence α satisfies every clause on its path, agrees with every propagation they cause, and satisfies every sum of them and every count that includes their common solutions. □

Many strict local maxima. Max-Cut on K_m, written with the clauses $(x_i \vee x_j)$ and $(\neg x_i \vee \neg x_j)$ for every pair, has $\binom{m}{m/2}$ strict local maxima for even m, the balanced cuts. Disjoint copies with $d = m - 1$ have $2^{n-O(n(\log d)/d)}$ of them, which is the order of the saving of [**?**]. Disjoint copies are separated by splitting into components, but Hastings shows that his upper bounds on the number of local maxima are tight, up to polylogarithmic factors for dense formulas and up to a factor of order d for sparse ones [**?**]. So single-flip reasoning, however refined, cannot by itself give $(2 - \delta)^n$ at growing density. In particular the progress condition of [**?**] fails on such formulas, since the count at every depth is at least the number of strict local maxima.

What can cut them. Pair decisions can: in a $\neq$ child they compare an assignment with the flip of a whole class, a move of several variables. On K_8, K_{10} and K_{12} the chained class conditions of Section **??** leave 36, 94 and 276 complete assignments, against 70, 252 and 924 strict local maxima. For exact twins this is visible directly: merged with $u = \neg v$, the two clauses of a twin pair with a common neighbour w add up to one plus the indicator of the literal of w, independent of u. So the merged variable keeps only the clauses between u and v, domination fixes it, and the two orientations of the pair are never both explored. Hastings also constructs sparse formulas with many maxima under moves of up to k variables [**?**], so moves of bounded size may meet a similar barrier; class flips are not of bounded size.

A second obstacle. The circulant formulas $C_{9,2}$, $C_{11,3}$ and $C_{13,3}$ have no strict local maxima at all, yet on $C_{n,c}$ every rule of [**?**] costs $\binom{n-1}{a}$ leaves [**?**, Theorem 13, Observation 22]. Their leaves are only one-sided local maxima: the preferred child is never pruned. The class conditions reduce these three formulas to 10, 10 and 14 complete assignments.

Consequence for the plan. On formulas with few strict local maxima, single-flip rules and their sums can in principle suffice, and the question is when they cut (Sections **??** and **??**). On formulas with many, or where one-sided pruning is the bottleneck, a rule that moves several variables at once is necessary, and among our tools only class decisions qualify.

10 Numerical checks

The scripts are in this folder.

- `explore_pair_chain.py` (seconds): the two-clause example, the chains in $C_{n,c}$ and in Max-Cut on K_m, and the numbers of strict local maxima (Sections **??** and **??**).
- `explore_sum_rule.py`: `check` (seconds) compares the optimum with brute force on 150 formulas, with and without the sum rule and for both strategies, and the estimator with exact counts for $n = 30, 36$; `density` and `dense` (tens of minutes) give Tables **??** and **??** up to $n = 400$.

- `explore_clause_weights.py` (two seconds): Proposition **??**, the rows with release of Table **??**, the weight table, and the check that no transition lowers a weight by more than one. The rows without release solve $V(D, k_0) = (\lambda - 1)\lambda^D$ (Theorem **??**).
- `explore_availability.py` (ten seconds): the measurements of Section **??** for $n \leq 401$.

The entries for $n = 600, 800$ in Table **??** and for $n = 801$ in Table **??** are single runs of the same code with larger n.

11 Open questions and new directions

1. **Accumulation over clauses.** Combine Proposition **??** with Lemmas **??** and **??** over many adaptively chosen clauses, including the assignments forced by domination between them, or settle Problem **??** for the rule with release. This would give $2^{H(\beta')n}$, for every $\beta' > \beta_0$ and every density, for formulas in which every variable has a gap linear in its degree, with an adaptive rule.

2. **Partial gaps.** If only ηn variables have gaps, plain steps on the others are unavoidable. Does the refined form of Lemma **??**, which credits domination during plain steps, recover the exponent $\max\{1 - \eta, H((1 - \varepsilon)/2)\}$ of [**?**, Theorem 11]?

3. **Ending the cold start.** Find structural properties that produce near-dominated variables in balanced formulas. A candidate is coverage: a set S whose variables have most of their neighbours outside S. After a random assignment of the rest, a variable of S is dominated once its imbalance, a ± 1 walk over its outside neighbours, exceeds its degree inside S; by the Erdős–Littlewood–Offord bound [**?**] this fails with probability $O(b/\sqrt{D})$ for inside degree b and outside degree D. So inside degree up to $\varepsilon\sqrt{D}$ suffices, not only the bounded degree needed for independence [**?**], provided the fields of S are jointly spread out; near-twins have identical fields, which is where that assumption is needed. This is the idea behind Hastings' bound [**?**], and in pseudo-random formulas such sets have size $O(n/\sqrt{d})$, so coverage alone gives degree-dependent savings.

4. **Counting overlapping clauses.** By Remark **??** the measure must count groups of overlapping clauses. The weighted-sum counts of [**?**] provide such a joint weight, with a branching inequality and a running-time bound under a condition on their progress. The open part is to show linear progress before the last levels for an interesting class, starting with pseudo-random formulas.

5. **The sum rule.** Prove the behaviour of Table **??** on random formulas (Remark **??**); try the best weighted sums, which by Proposition **??** are the linear-relaxation cuts, and integer reasoning beyond them for contradictions of parity type; choose orders that maximize cancellation; test formulas with planted aligned groups, where cancellation is partial.

6. **A dichotomy at each node.** Make the three kinds of nodes of Section **??** precise, so that every node offers either a strong clause, a decisive sum, or an aligned structure for pair decisions. By Section **??**, a natural dividing line is whether the residual formula has few or many strict local maxima, with one-sided pruning as a second obstacle.

7. **Pair decisions.** By Proposition **??** they are the only rule here that can cut strict local maxima. Find orders in which the class conditions of Proposition **??** are checked early; decide whether class decisions can reach the bound $O^*((n + 1)^t)$ of exact interaction classes [**?**] and degrade gracefully when the classes are only approximate; and check

whether Hastings' sparse formulas with many maxima under moves of several variables [**?**] also defeat class flips.

8. **Strategies.** Test the coverage strategy of Section **??**, and the largest-ratio choice of Section **??** on formulas with partial gaps.

References

[1] A literal-deficit measure for Max 2-SAT. Working note, 22 September 2026 (`max2sat_literal_deficit_proof.tex`).

[2] Exact Max 2-SAT with an absolute linear polarity gap. Working note, 23 September 2026 (`max2sat_absolute_linear_gap_proof.tex`).

[3] Max 2-SAT and the domination rule: results and ideas so far. Working note, 5 October 2026 (`max2sat_results_and_ideas.tex`).

[4] Exploring the branching frontier for Max 2-SAT: full clause phases, joint completion bounds, and coverage. Draft, 7 October 2026 (`max2sat_branching_frontier.tex`).

[5] P. Erdős. On a lemma of Littlewood and Offord. *Bulletin of the American Mathematical Society* 51(12):898–902, 1945.

[6] A. Golovnev and K. Kutzkov. New exact algorithms for the 2-constraint satisfaction problem. *Theoretical Computer Science* 526:18–27, 2014.

[7] M. B. Hastings. Local maxima and improved exact algorithm for MAX-2-SAT. arXiv:1610.07100, 2016.

[8] D. Aldous and U. Vazirani. "Go with the winners" algorithms. In *Proc. 35th FOCS*, pp. 492–501, 1994.

[9] D. E. Knuth. Estimating the efficiency of backtrack programs. *Mathematics of Computation* 29(129):121–136, 1975.