A Simplified Plan for Max 2-SAT: One Search, One Count, One Key Lemma

8 October 2026

Working notes

Abstract

We select one direction from the notes [1, 2, 3, 4]: DPLL with the pruning clauses of [1] and the sum check of [2], along a fixed variable order. Knuth's identity turns its running time into one probability, that a uniformly random partial assignment survives t steps. An exact identity shows that the sum check is a race between the accumulated shortfall of the clauses, a martingale, and the uncertainty left on the free variables. Their expected values all scale like $\sqrt{d}$, which explains why the sum check, unlike single clauses, cuts a typical partial assignment after about $0.4n$ steps at every density, and why the search saves between $0.50n$ and $0.62n$ in our runs up to $n = 1600$. We state the one result we need, a Key Lemma for random formulas of every density; the companion note [5] proves it with $\gamma \approx 0.02$, for every graph with random signs. The worst case reduces to the same argument under three cancellation conditions, together with aggregation for formulas that violate them. Everything else is parked.

1 The choice

We keep one search: the pruning clauses of [1] with the sum check of [2]. Three reasons.

1. It is the only mechanism we have found that beats the timing barrier [1, Remark 29]: single clauses cut only near the end of their supports, while the sum decides early at every density (Tables 1 and 2).

2. Its proofs are short. Soundness is already proved, and the running time is a single probability (Section 3).

3. It isolates what remains. The sum fails exactly on aligned structure, where every rule implied by single flips fails [2, Section 9], so the worst case needs one more ingredient, aggregation, and nothing else (Section 7).

The redundancy question of [4]. In this view a new clause need not add information by itself. The sum adds up the shortfall of every active clause, whether or not the clause is implied by the others. What makes the sum decisive is cancellation: two clauses that contain opposite literals of a shared variable make one of the two literals true whatever its value. Only overlaps with the same literals defeat the sum, and if they dominate, the formula is aligned.

2 The search

The formula is canonical with $B = 1$, notation as in [1, Section 2]. Fix the order $x_1, \dots, x_n$. A node at depth $t-1$ has $x_1, \dots, x_{t-1}$ assigned and branches on x_t, trying both values. Let p be the smaller deficit of x_t at the node and ℓ the literal attaining it (ties: the positive literal). The child in which ℓ is false receives the pruning clause $P_t = \langle L_t \le p-1 \rangle$ over the free neighbours of x_t; the other child receives nothing. A node is cut when

- some pruning clause has more true literals than it allows, or
- the *sum check* fires: $\sum_{w \text{ free}} \min\{m_w^0, m_w^1\} > \sum_{c \in I} k_c$, where I is the set of live pruning clauses that are not vacuous, k_c is the allowance of c, and m_w^b is the number of clauses of I in which the value b of w makes a literal true [2, Section 5].

There is no domination and no propagation; both are sound and help in practice, but the analysis is for this plain version. If $p = 0$, the pruned child is cut at once. Soundness: each branching adds a clause to one child only, so [1, Proposition 20] applies, and the sum is implied by the clauses.

3 Counting nodes

Proposition 1 (Knuth's identity [6])**.** *Let σ be uniform in $\{0,1\}^n$ and let S_t be the probability that the partial assignment $\sigma_1, \dots, \sigma_t$ survives all checks at depths $1, \dots, t$, with $S_0 = 1$. The search tree has exactly $2^t S_t$ live nodes at depth t, and at most $1 + 2n \max_t 2^t S_t$ nodes in total. Hence the search runs in time $O^*(\max_t 2^t S_t)$ and polynomial space.*

Proof. The checks at depth s depend only on $\sigma_1, \dots, \sigma_s$, so the live nodes at depth t correspond to the surviving prefixes, each of probability 2^{-t}. Every node other than the root is a child of a live node, and each live node has two children. □

So the whole running-time analysis is one probability estimate for a uniformly random partial assignment. No measure, no clause weights, no tilted walk and no availability argument are needed.

4 The sum check as a race

Let c be the clause created at step i. Write D_c for its support size, the number of free neighbours of x_i at that time, and Δ_c for the effective gap of x_i then, so that its threshold is $p_c = (D_c - |\Delta_c|)/2$ by [1, (1)]. At a later node, a_c of its support variables are assigned and j_c of them took the value that makes their literal true. For a set I of clauses and a free variable w put $b_w = m_w^1 - m_w^0$.

Lemma 2 (The race)**.** *For every set I of pruning clauses that are not violated,*

$$\sum_{w \text{ free}} \min\{m_w^0, m_w^1\} - \sum_{c \in I} k_c = S(I) + Q(I) - U(I),$$

where

$$S(I) = \sum_{c \in I} \Big(\frac{|\Delta_c|}{2} + 1 \Big), \qquad Q(I) = \sum_{c \in I} \Big(j_c - \frac{a_c}{2} \Big), \qquad U(I) = \frac{1}{2} \sum_{w \text{ free}} |b_w|.$$

So the sum check over I fires if and only if $S(I) + Q(I) > U(I)$.

Proof. $\min\{m_w^0, m_w^1\} = \frac{1}{2}(m_w^0 + m_w^1) - \frac{1}{2}|b_w|$, and $\sum_w (m_w^0 + m_w^1) = \sum_c (D_c - a_c)$. With $k_c = p_c - 1 - j_c = \frac{1}{2}(D_c - |\Delta_c|) - 1 - j_c$, each clause contributes $\frac{1}{2}(D_c - a_c) - k_c = \frac{1}{2}|\Delta_c| + 1 + j_c - \frac{1}{2}a_c$. □

S is the *shortfall*: every clause belongs to a variable set against its majority, and its bound lies $|\Delta_c|/2 + 1$ below the expected count of its support. U is the *uncertainty*: whatever values the free variables take, the clauses that disagree about a variable cancel, and only $|b_w|$ is left open. Q records how far the assigned support variables deviated from fair coins. The identity was checked in exact arithmetic on 7,259 random states.

Remark 3 (Which clauses to sum). Removing a vacuous clause v never weakens the check: its term $\frac{1}{2}(D_v - a_v) - k_v$ is at most $-\frac{1}{2}(D_v - a_v)$, while removing it changes U by at most $\frac{1}{2}(D_v - a_v)$. So the check of the search fires whenever the check over all non-violated clauses fires. For the analysis we take I_t to be all clauses created up to depth t and write S_t, Q_t, U_t and $Z_t = S_t + Q_t - U_t$. A partial assignment that survives depth t has $Z_t \le 0$.

Proposition 4 (One step)**.** *Along a uniformly random σ, at step t:*

1. *$Q_t - Q_{t-1} = \frac{1}{2}(2\sigma_t - 1)\, b_{x_t}$, with b_{x_t} taken at depth $t-1$; so Q is a martingale.*
2. *U loses $\frac{1}{2}|b_{x_t}|$, because x_t is no longer free.*
3. *With probability $\frac{1}{2}$, when σ_t is the non-preferred value, a clause is created: S grows by $\frac{1}{2}|\Delta_t| + 1$, and b_w changes by ± 1 for each of its support variables w.*

In the random-sign model of Section 5, the signs added to the b_w in item 3 are independent fair coins given the history, since the sign of w in its clause with x_t plays no role before step t. Hence

$$\mathbb{E}\big[Z_t - Z_{t-1} \mid \sigma_1, \ldots, \sigma_{t-1}\big] = \frac{|\Delta_t|}{4} + \frac{1}{2} + \frac{|b_{x_t}|}{2} - \frac{z_t}{4},$$

where z_t is the number of free neighbours w of x_t with $b_w = 0$.

Proof. Assigning x_t raises a_c for the clauses containing it and raises j_c for those in which σ_t is the bad value, which gives item 1. Items 2 and 3 follow from the definitions. Adding an independent fair sign to b_w raises $|b_w|$ by one in expectation if $b_w = 0$ and leaves it unchanged otherwise. □

5 The Key Lemma

Model. In $R(n, d)$ each of the $\binom{n}{2}$ pairs carries one clause with probability $d/(n-1)$, independently, with independent uniform signs. By symmetry the fixed order does not matter.

Key Lemma 5. There are constants $\gamma > 0$ and d_0 such that for all n and all $d_0 \le d \le n - 1$, with probability $1 - o(1)$ over $F \in R(n, d)$,

$$\max_{0 \le t \le n} 2^t S_t(F) \le 2^{(1-\gamma)n}.$$

The Key Lemma is proved in [5, Corollary 3] for every $\gamma < 0.0236$. By Proposition 1, the search of Section 2 solves Max 2-SAT on $R(n, d)$ in time $O^*(2^{(1-\gamma)n})$ and polynomial space, with probability $1 - o(1)$, for every density. This is an average-case statement, and its novelty still has to be checked against the literature.

		$d = 16$	64	256	$(n-1)/2$
$n = 400$	single clauses	0.19	0.52	0.76	0.72
	with the sum check	0.19	0.41	0.43	0.43
$n = 1600$	single clauses	0.11	0.49	0.72	0.84
	with the sum check	0.11	0.39	0.42	0.44

Table 1: Median depth, as a fraction of n, at which a uniformly random partial assignment is cut; 30 assignments per entry.

		$d = 64$	256	$(n-1)/2$
$n = 400$	single clauses	0.36	0.20	0.26
	with the sum check	0.52	0.51	0.54
$n = 800$	single clauses	0.40	0.24	0.21
	with the sum check	0.56	0.53	0.50
$n = 1600$	single clauses	0.43	0.24	0.16
	with the sum check	0.62	0.56	0.56

Table 2: Saving $n - \log_2(\text{tree size})$ as a fraction of n, estimated by population splitting (unbiased; one formula per entry).

Evidence. Table 1 gives the depth at which a uniformly random partial assignment is cut, Table 2 the size of the whole tree, both in exactly this setting.

With single clauses the cut moves towards the last level as d grows; with the sum check it stays near $0.4n$, and the saving stays between $0.50n$ and $0.62n$.

Why the density cancels. For $d \to \infty$ use the approximations that a free variable at time τn lies in about $\tau d/2$ clauses, that Δ_t has variance about $d(1+\tau)$, and that b_w is a sum of independent signs. Then Proposition 4 gives drift $\sqrt{d}\,\mu(\tau)$ per step with

$$\mu(\tau) = \frac{1}{4}\sqrt{\frac{2(1+\tau)}{\pi}} + \frac{1}{2}\sqrt{\frac{\tau}{\pi}} - \frac{1-\tau}{4\sqrt{\pi\tau}},$$

and $Z_{\tau n} \approx n\sqrt{d}\,F(\tau)$ with

$$F(\tau) = \int_0^\tau \mu = \frac{1}{\sqrt{\pi}}\Big(\frac{\sqrt{2}}{6}\big((1+\tau)^{3/2} - 1\big) + \frac{\tau^{3/2} - \tau^{1/2}}{2}\Big).$$

Early on, new clauses meet fresh variables and U grows faster than S; the drift turns positive at $\tau_0 \approx 0.14$, and F changes sign at $\tau^* \approx 0.46$, independently of d. The measured cuts at 0.39–0.44 come slightly earlier, because single clauses and the removal of vacuous clauses also act. A single clause, by contrast, compares an imbalance of order $\sqrt{d}$ with a remaining support of order d, and so decides only at depth $1 - O(1/\sqrt{d})$.

6 How it is proved

The proof in [5] is simpler than the outline we had here (drift, noise, combine).

1. **One depth.** $\max_t 2^t S_t \le \max\{2^T, 2^n S_T\}$, so it suffices to show that a random partial assignment survives one depth $T = \tau n$ only with probability e^{-cn}; then $\gamma = \min\{1 - \tau,\, c/\ln 2\}$.

2. **Late signs.** The search never uses the signs of x_t before it reaches x_t. So, given the past, Δ_t is a sum of unseen signs (± 1 over the 2-clauses of x_t, ± 2 over its unit clauses), the bad values of x_t in the clauses that contain it are unseen fair coins, and at depth T so are the bad values of the free variables.

3. **Three tails.** Szarek's inequality and a one-sided bound for nonnegative increments give the shortfall; Q is a walk with fair steps $\pm\frac{1}{2}$ (Azuma); U is controlled by Jensen's and McDiarmid's inequalities. No drift analysis of U is needed, because U is evaluated only at depth T, where the bad values of the free variables are still unseen.

The graph enters only through a few degree sums, so the result holds for every graph with random signs whose margin is of order $n\sqrt{d}$: every $\gamma < 0.0236$ for $R(n, d)$, and every $\gamma < 0.0132$ for every d-regular graph in any order [5, Corollaries 3 and 4]. The constant is far below the measured $0.50n$–$0.62n$ and the Gaussian estimate of about $0.2n$ from the drift above. The loss comes from four separately added tails, from Szarek's constant, and from using a single depth. A single exponential moment of Z_t near depth $0.5n$ is the natural way to improve it.

7 The worst case

The proof of [5] uses random signs in three places: the size and concentration of the shortfall, the step size of the martingale Q, and the size of the uncertainty U. For a fixed formula, σ alone keeps Q a martingale and gives the shortfall its mean, but nothing more. In particular our earlier claim that the shortfall part holds for every formula is true only for its mean.

Cancellation conditions [5, Section 6]. The proof goes through verbatim for a fixed formula if, except with probability $e^{-\Omega(n)}$ over σ: (W1) U_T is a small constant times $n\sqrt{d}$; (W2) $\sum_{t \le T} b_{x_t}^2 = O(nd)$; (W3) the effective gaps keep their typical size, $\sum_t |\Delta_t| = \Omega(n\sqrt{d})$ and $\sum_t \Delta_t^2 = O(nd)$. (W1) and (W2) say that the clauses containing a variable disagree about its value often enough that $|b_w|$ is of order $\sqrt{m_w}$ rather than m_w, at the free and at the assigned variables.

When it fails. If cancellation fails, many clauses agree about the values of many variables: the formula is aligned, as in the circulant formulas $C_{n,c}$, where every clause has the same bad value at every variable. By the barrier of [2, Section 9], no rule implied by single flips can help there: Max-Cut on K_m has $\binom{m}{m/2}$ strict local maxima, and $C_{n,c}$ has exponentially many optimal assignments; even comparisons in both children with fixed tie-breaking keep 126 assignments of $C_{9,2}$, more than the 70 leaves of plain domination. Both are solved by aggregation: counting the ones within exact interaction classes [3, Section 5].

The worst-case target. (i) The Key Lemma under (W1)–(W3); this is the remark above. (ii) Formulas that violate them have enough aligned structure for aggregation to pay. (iii) A rule that decides between the two at a node. Part (ii) is the hard part; it is a statement about formulas whose active clauses agree about their variables, not about arbitrary graphs. By Corollary 4 of [5] the graph alone is never the obstacle.

8 Parked

Clause-weight measures and the availability analysis [2, Sections 6–8]; packing and core certificates, coverage, exposure, the entropy dichotomy and the desired frontier theorem [3, 4];

comparisons in both children [4]; pair decisions [2, Section 3] except as a possible form of aggregation. None of them was needed for the Key Lemma. The barrier and the linear-relaxation characterization of sums [2] are kept, because they define the worst case and the strongest version of the check.

9 Checks

- `explore_death_depth.py` (about 15 seconds): Table 1.
- `explore_key_lemma.py`: `check` verifies Lemma 2 in exact arithmetic; `400 50` and similar give the rows of Table 2 (minutes for $n \leq 800$, longer for 1600).
- `explore_key_lemma_proof.py`: the checks of [5].

References

[1] Max 2-SAT and the domination rule: results and ideas so far. Working note, 5 October 2026 (`max2sat_results_and_ideas.tex`).

[2] Max 2-SAT with pruning clauses: pair decisions, sums, and a measure. Working note, 6–7 October 2026 (`max2sat_pairs_sums_measure.tex`).

[3] Exploring the branching frontier for Max 2-SAT. Draft, 7 October 2026 (`max2sat_branching_frontier.tex`).

[4] A learning and coverage dichotomy for Max 2-SAT. Draft, 7 October 2026 (`max2sat_learning_dichotomy.tex`).

[5] The Key Lemma for random signs. Working note, 8 October 2026 (`max2sat_key_lemma_proof.tex`).

[6] D. E. Knuth. Estimating the efficiency of backtrack programs. *Mathematics of Computation* 29(129):121–136, 1975.