Weighted and Round-Based Early Enumeration

Random signs on every graph, Max-Cut on spectral expanders, and count guessing without a given decomposition

12 October 2026

Abstract

Two modifications of early enumeration remove the main restrictions of the earlier class theorems. Weighting each learned constraint by the inverse square root of the degree gives a fixed exponential saving for Max 2-SAT with random literal signs on *every* interaction graph, with arbitrary unit clauses and several clauses per edge allowed; the earlier bound required a bounded ratio of maximum to minimum degree. Fixing the preferred values of the branching variables only between a constant number of rounds makes the analysis depend on the ordinary spectral norm of the remainder instead of the norm of its triangular part. As a consequence, Max-Cut with arbitrary unit clauses is solved in time $O^*(c^n)$, $c < 2$, and polynomial space on every d-regular graph whose nontrivial eigenvalues are $O(\sqrt{d})$, for all $d_0 \le d \le (1-\varepsilon)n$. This includes Paley graphs, random regular graphs and Ramanujan graphs. Finally, the counts of count guessing can be replaced by projections onto eigenvectors, guessed on a polynomial grid; no decomposition has to be given. All algorithms are deterministic, exact on every input, and use polynomial space.

1 Results

Write the score of a formula as $F(x) = c + \sum_u h_u x_u + \sum_{u<v} J_{uv} x_u x_v$ on $x \in \{0,1\}^n$, and $\mathrm{gain}_t(x) = F(x^{t\to 1}) - F(x^{t\to 0}) = h_t + \sum_{w\neq t} J_{tw} x_w$. An equal-polarity clause, $(u \vee v)$ or $(\neg u \vee \neg v)$, contributes -1 to J_{uv}, a mixed one $+1$.

- **Random signs on every graph (Theorem 4).** Fix any graph G and any multiset of unit clauses. Put on each edge a random multiset of at most m_0 clauses whose law is invariant under complementing either endpoint, for example one clause with uniform literal signs, or each of the four clauses independently with probability q. Weighted early enumeration enumerates at most $3 \cdot 2^n e^{-c_0 n}$ assignments in expectation, with $c_0 > 0$ depending only on m_0 and a non-degeneracy constant κ, not on G.

- **Round-based enumeration (Theorem 9).** The class of the count-guessing note, with its triangular-norm condition replaced by $\|N\|^2 \le K\, D_N$ for the ordinary operator norm of the remainder, and with the aligned part an arbitrary low-rank matrix $\sum_k \lambda_k v_k v_k^{\mathsf{T}}$.

- **Max-Cut on spectral expanders (Corollary 12).** Every d-regular graph with all nontrivial eigenvalues at most $\lambda_0\sqrt{d}$ in absolute value, $d_0(\lambda_0) \le d \le (1-\varepsilon)n$, with both clauses $(u \vee v), (\neg u \vee \neg v)$ on each edge and arbitrary unit clauses: time $O^*(c^n)$ with $c = c(\lambda_0, \varepsilon) < 2$.

- **Spectral count guessing (Corollary 13, Proposition 14).** Guessing the projections of x onto the top eigenvectors of J on a grid of polynomial size handles planted low-rank parity structure without knowing it.

2 Preliminaries

Enumeration. All algorithms branch on a prefix of T variables, learn affine constraints on the way, and at each prefix enumerate the completions of one summed affine inequality. Such solutions can be listed with polynomial delay, and the cost is $O^*(2^T + W)$, where W is the number of complete assignments satisfying the summed inequality of their own prefix [4, 7]. For uniform x, $W = 2^n \mathbb{P}[x \text{ passes}]$.

Probabilistic tools.

(T1) *Azuma–Hoeffding.* If $X_1, X_2, \dots$ are martingale differences and X_i lies, given the past, in an interval of length ℓ_i, then $\mathbb{P}[\sum_i X_i \le -a] \le e^{-2a^2/\sum_i \ell_i^2}$.

(T2) *Exponential supermartingale.* If, given the past, X_i is a Rademacher sum with variance proxy V_i (so $\mathbb{E}e^{\lambda X_i} \le e^{\lambda^2 V_i/2}$) and $\sum_i V_i \le V$ always, then $\mathbb{P}[\sum_i X_i \le -a] \le e^{-a^2/(2V)}$.

(T3) *Talagrand.* If f is convex and ℓ-Lipschitz on $[-1,1]^T$ and u is uniform on $\{\pm 1\}^T$, then $\mathbb{P}[f \le \mathbb{E}f - r - 8\sqrt{\pi}\,\ell] \le 4e^{-r^2/(16\ell^2)}$ [9].

(T4) *Hanson–Wright.* There is an absolute constant $c_{\mathrm{HW}} > 0$ such that for a symmetric matrix M with zero diagonal and a uniform $\beta \in \{\pm 1\}^m$, $\mathbb{P}[\beta^{\mathsf{T}} M \beta \ge s] \le \exp(-c_{\mathrm{HW}} \min\{s^2/\|M\|_F^2,\, s/\|M\|\})$ [8].

(T5) *Fourth moment.* A Rademacher sum Z satisfies $\mathbb{E}|Z| \ge \sqrt{\operatorname{Var} Z/3}$.

Lemma 1 (Symmetric shift)**.** *If Z is symmetric, $a \in \mathbb{R}$ and $K > 0$, then $\mathbb{E}\min\{|a+Z|, K\} \ge \frac{1}{2}\mathbb{E}\min\{|Z|, K\}$.*

Proof. The function $\phi(z) = \min\{z, K\}$ is concave and nondecreasing on $[0,\infty)$ with $\phi(0) = 0$, hence subadditive. So $\phi(|a+Z|) + \phi(|a-Z|) \ge \phi(|a+Z| + |a-Z|) \ge \phi(2|Z|) \ge \phi(|Z|)$. Take expectations and use that $a+Z$ and $a-Z$ have the same law. □

3 Random signs on every graph

Definition 2 (Complementation-invariant random signs)**.** Let $G = (V, E)$ be a graph without isolated vertices, $n = |V|$, with any fixed order $1, \dots, n$ of V, and let $T = \lceil n/2 \rceil$. For each edge $e = \{u, v\}$ let M_e be a random multiset of binary clauses on u, v, independently over the edges, such that

(i) $|M_e| \le m_0$;

(ii) the law of M_e is invariant under replacing the literal of u by its complement in every clause of M_e, and likewise for v;

(iii) $\mathbb{E}\big[(\sum_{c \in M_e} s_u(c))^2\big] \ge \kappa > 0$ for every edge e and endpoint u, where $s_u(c) \in \{\pm 1\}$ is the sign of the literal of u in c.

The unit clauses form an arbitrary multiset, fixed independently of the M_e.

One clause per edge with independent uniform literal signs has $m_0 = \kappa = 1$. Including each of the four clauses on every edge independently with probability q has $m_0 = 4$ and $\kappa = 4q(1-q)$; with $q = \frac{1}{4}$, this is the model in which the four clauses of each pair of a fixed set are sampled with probability $\frac{1}{4}$.

Algorithm A (weighted early enumeration). For a clause $c \in M_{tv}$, the *partner literal* is its literal on v. Let b_t be the number of unit clauses (t) minus the number of unit clauses $(\neg t)$, and $G_{tv}(x_v) = \sum\{s_t(c) : c \in M_{tv},$ the partner literal of c is false under $x_v\}$. Then $\mathrm{gain}_t(x) = b_t + \sum_{v\sim t} G_{tv}(x_v)$. Branch on $x_1, \dots, x_T$. At step t let

$$\Delta_t = b_t + \sum_{v<t,\, v\sim t} G_{tv}(x_v) + \frac{1}{2} \sum_{v>t,\, v\sim t} \sum_{c\in M_{tv}} s_t(c),$$

the expected gain of x_t when the later variables are uniform, and $p_t = [\Delta_t \geq 0]$. If $x_t \neq p_t$, then t is a *creator*. At each prefix of length T enumerate the completions satisfying

$$\sum_{t \text{ creator}} \omega_t((2p_t - 1)\,\mathrm{gain}_t(x) + 1) \leq 0, \qquad \omega_t = \deg_G(t)^{-1/2}, \tag{1}$$

and evaluate F on each.

Lemma 3 (Exactness)**.** *Among the optima, let x° maximize, lexicographically, the vector of indicators $[x_t = p_t]$, $t \leq T$, along its own path. Then $(2p_t - 1)\mathrm{gain}_t(x^\circ) \leq -1$ for every creator t on its path, so x° satisfies* (1) *and is enumerated.*

Proof. Scores are integers. If $(2p_t - 1)\mathrm{gain}_t(x^\circ) \geq 0$ for a creator t, flipping x°_t to p_t gives an optimum with the same path before t and indicator one at t, which is lexicographically larger. □

Theorem 4 (Random signs on every graph)**.** *In the model of Definition 2, Algorithm A is exact on every input, and for every graph, every order and every multiset of unit clauses the expected number of enumerated assignments is at most $3 \cdot 2^n e^{-c_0 n}$, where*

$$c_0 = \frac{c_1^2}{2704\, m_0^2}, \qquad c_1 = \frac{\sqrt{\kappa}}{2\sqrt{1 + 4m_0^4/\kappa^2}}.$$

Hence the expected running time is $O^*(2^{n/2} + 2^n e^{-c_0 n})$, and the running time exceeds $O^*(2^n e^{-c_0 n/2})$ with probability at most $3e^{-c_0 n/2}$.

For one uniformly signed clause per edge, $c_0 = 1/54080$. The earlier bound for this model, Theorem 4.1 of [7], has base $2e^{-d/(1250D)}$ with d, D the minimum and maximum degree, requires $d \geq 6$, and excludes unit clauses.

Proof. Step 0: race identity. Let x be uniform and independent of the formula. Write $\mathrm{gain}_t(x) = \Delta_t + \sum_{v>t, v\sim t} \xi_{tv}$ with $\xi_{tv} = G_{tv}(x_v) - \frac{1}{2}\sum_{c\in M_{tv}} s_t(c)$. For a creator, $(2p_t - 1)\Delta_t = |\Delta_t|$. Hence x passes (1) if and only if $S + Q \leq 0$, where, with I the set of creators,

$$S = \sum_{t\in I} \omega_t(|\Delta_t| + 1), \qquad Q = \sum_{t\in I} \omega_t(2p_t - 1) \sum_{v>t,\, v\sim t} \xi_{tv}.$$

Step 1: late signs. By (ii), M_e has the law of a multiset obtained from an independent copy σ_e of M_e, the *skeleton*, by multiplying the signs of u in all its clauses by an independent fair sign $\rho_{u,e}$, and those of v by $\rho_{v,e}$. Put $\theta_{u,e} = \rho_{u,e}(2x_u - 1)$. The literal of u in c is false under x_u exactly when $\sigma_u(c)\theta_{u,e} = -1$. All $\theta_{u,e}$ are independent fair signs, independent of the skeletons and of x, because the ρ's are. Reveal the skeletons first; then, for $v = 1, \dots, n$, reveal the signs $\rho_{v,e}$ ($e \ni v$) and x_v. Let $\mathcal{F}_v$ be the information after step v.

For $e = \{t, v\}$ we have $G_{tv}(x_v) = \rho_{t,e}\, a_e$ with $a_e = \sum\{\sigma_t(c) : \sigma_v(c)\theta_{v,e} = -1\}$. Hence $\Delta_t = b_t + Z_t$, where

$$Z_t = \sum_{v<t} \rho_{t,e}\, a_e + \sum_{v>t} \rho_{t,e}\, a'_e, \qquad a'_e = \tfrac{1}{2}\textstyle\sum_{c\in M_e} \sigma_t(c),$$

and the coefficients a_e, a'_e are $\mathcal{F}_{t-1}$-measurable. Given $\mathcal{F}_{t-1}$, Z_t is a Rademacher sum with variance $Y_t = \sum_{v<t} a_e^2 + \sum_{v>t} a_e'^2 \le m_0^2 \deg(t)$, and x_t is uniform and independent of Δ_t, so t is a creator with probability $\frac{1}{2}$. For $v > t$, $\xi_{tv} = \rho_{t,e}\,\theta_{v,e}\,\eta_e$ with $\eta_e = \sum\{\sigma_t(c) : \sigma_v(c) = -1\} - a'_e$ and $|\eta_e| \le m_0/2$.

Step 2: the shortfall. Let $\tilde{Y}_t = [t \in I]\,\omega_t \min\{|\Delta_t|, m_0\sqrt{\deg t}\} \in [0, m_0]$; then $S \ge \sum_{t\le T} \tilde{Y}_t$. By Lemma 1, (T5) and $(z-K)_+ \le z^2/(4K)$,

$$\mathbb{E}[\tilde{Y}_t \mid \mathcal{F}_{t-1}] \ge \tfrac{1}{2}\omega_t \cdot \tfrac{1}{2}\Big(\sqrt{Y_t/3} - \frac{Y_t}{4m_0\sqrt{\deg t}}\Big) \ge \tfrac{1}{4}\omega_t\sqrt{Y_t}\Big(\tfrac{1}{\sqrt{3}} - \tfrac{1}{4}\Big) \ge \mu_t := \frac{\omega_t\sqrt{Y_t}}{13},$$

using $Y_t \le m_0^2 \deg t$. By (T1), $\mathbb{P}[\sum_{t\le T}\tilde{Y}_t < \sum_{t\le T}\mu_t - a] \le e^{-2a^2/(Tm_0^2)}$.

Step 3: the mean shortfall. For $v < t$, a_e takes the values $\alpha_\pm = \sum\{\sigma_t(c) : \sigma_v(c) = \mp 1\}$ with probability $\frac{1}{2}$ each, so $\mathbb{E}a_e^2 \ge \frac{1}{4}\mathbb{E}(\alpha_+ + \alpha_-)^2 = \frac{1}{4}\mathbb{E}(\sum_c \sigma_t(c))^2 \ge \kappa/4$; also $\mathbb{E}a_e'^2 \ge \kappa/4$. The summands of Y_t are independent over the edges and lie in $[0, m_0^2]$, so $\mathbb{E}Y_t \ge \kappa\deg(t)/4$ and $\operatorname{Var} Y_t \le m_0^4 \deg(t)/4$. Hence

$$\mathbb{E}\sqrt{Y_t} \ge \frac{(\mathbb{E}Y_t)^{3/2}}{(\mathbb{E}Y_t^2)^{1/2}} = \frac{\sqrt{\mathbb{E}Y_t}}{\sqrt{1 + \operatorname{Var} Y_t/(\mathbb{E}Y_t)^2}} \ge c_1\sqrt{\deg t},$$

and $\mathbb{E}\sum_{t\le T}\mu_t \ge c_1 T/13$. The sum $\sum_{t\le T}\mu_t$ is a function of the independent skeletons and signs θ; those of one edge change at most two terms, each by at most $\omega m_0/13$. Since $\sum_e(\omega_u+\omega_v)^2 \le 2\sum_v \omega_v^2 \deg v = 2n$, McDiarmid's inequality gives $\mathbb{P}[\sum_{t\le T}\mu_t < c_1T/13 - b] \le e^{-169b^2/(m_0^2 n)}$.

Step 4: the interaction term. Order the summands of Q by their later endpoint v. The summand for the edge $\{t, v\}$ is $[t \in I]\,\omega_t(2p_t - 1)\rho_{t,e}\eta_e \cdot \theta_{v,e}$; its coefficient is known before step v and $\theta_{v,e}$ is a fresh fair sign, independent of the other signs revealed at step v. By (T1), with interval lengths $\omega_t m_0$ and $\sum_{t\le T}\omega_t^2 \deg(t) \le T$, $\mathbb{P}[Q \le -c] \le e^{-2c^2/(m_0^2 T)}$.

Step 5. Take $a = b = c = c_1T/52$. Outside the three exceptional events, $S + Q \ge c_1T/13 - 3c_1T/52 = c_1T/52 > 0$. With $T \ge n/2$ the three exponents are at least $c_1^2T/(1352m_0^2) \ge c_0 n$, so the three probabilities are at most $e^{-c_0 n}$ each, and $\mathbb{E}W \le 3 \cdot 2^n e^{-c_0 n}$. Markov's inequality gives the tail statement. □

Corollary 5 (Arbitrary supports)**.** *Fix any set of variable pairs and include each of the four clauses on each of these pairs independently with probability $q \in (0, 1)$; add arbitrary unit clauses (variables in no pair are set by their unit clauses). Then Max 2-SAT is solved in expected time $O^*(2^n e^{-c_0(q)n})$ with $c_0(q) > 0$ depending only on q. No condition on the density, the degree sequence or the order is needed.*

Remark 6 (Complementary pairs)*.* Condition (iii) fails when every edge carries a complementary pair, as in Max-Cut with random renamings (a $\pm J$ spin glass). Then the later-neighbour part of Y_t vanishes and the shortfall comes from earlier neighbours only. With a uniformly random order, $\mathbb{E}\sqrt{Y_t}$ is of order $\sqrt{(t/n)\deg t}$ and the same proof should go through; we do not write it out.

4 Round-based enumeration with guessed projections

Setting. Let $P = \sum_{k=1}^r \lambda_k v_k v_k^{\mathsf{T}}$ with $\lambda_k \in \mathbb{R}$ and $v_k \in \mathbb{R}^n$, and let N be the off-diagonal part of $J - P$, so $N_{uu} = 0$ and $J_{uv} = P_{uv} + N_{uv}$ for $u \ne v$. Put $y_k(x) = \langle v_k, x\rangle$. For each k fix a finite grid $\mathcal{Y}_k \subset \mathbb{R}$ such that every $y_k(x)$, $x \in \{0,1\}^n$, is within $\delta_k \ge 0$ of a grid point, and let

$$s_t = \sum_k |\lambda_k|\,|v_k(t)|\,(\delta_k + |v_k(t)|).$$

Let $T = \lfloor \tau n \rfloor$, and split the prefix into $\ell \ge 2$ consecutive *rounds* $B_1, \dots, B_\ell$ of the branching order; the remaining variables form round $\ell + 1$. Write $\rho(t)$ for the round of t.

Algorithm C. For each $\hat{y} \in \mathcal{Y} = \prod_k \mathcal{Y}_k$ run the following. Put $h^{\hat{y}} = h + \sum_k \lambda_k \hat{y}_k v_k$ and $F'_{\hat{y}}(x) = c + \langle h^{\hat{y}}, x\rangle + \sum_{u<v} N_{uv} x_u x_v$. Branch on the prefix. For $t \in B_j$ let

$$\Delta_t = h_t^{\hat{y}} + \sum_{s:\rho(s)<j} N_{ts} x_s + \frac{1}{2} \sum_{w \neq t:\rho(w)\geq j} N_{tw}$$

and $p_t = [\Delta_t \geq 0]$; the preferred value of t depends only on the earlier rounds. If $x_t \neq p_t$, then t is a creator and $\gamma_t(x) = (2p_t - 1)(F'_{\hat{y}}(x^{t\to 1}) - F'_{\hat{y}}(x^{t\to 0}))$. At each prefix enumerate the completions of

$$\sum_{t \text{ creator}} (\gamma_t(x) - s_t) \leq 0, \tag{2}$$

evaluate F on them, and return the best assignment over all runs.

Lemma 7 (Flip identity)**.** *If $|y_k(x) - \hat{y}_k| \leq \delta_k$ for all k, then for every t the gains of x_t under F and under $F'_{\hat{y}}$ differ by at most s_t.*

Proof. $\mathrm{gain}_t(x) = h_t + \sum_w N_{tw} x_w + \sum_{w\neq t} P_{tw} x_w$ and $\sum_{w\neq t} P_{tw} x_w = \sum_k \lambda_k v_k(t)\, y_k(x^{t\to 0})$. The gain under $F'_{\hat{y}}$ replaces the last sum by $\sum_k \lambda_k v_k(t) \hat{y}_k$. Finally $|y_k(x^{t\to 0}) - \hat{y}_k| \leq |y_k(x) - \hat{y}_k| + |v_k(t)|$. □

Lemma 8 (Exactness)**.** *Every optimum x^* is enumerated in a run whose grid point is within δ_k of $y_k(x^*)$ for all k.*

Proof. Every flip at x^* changes F by at most zero, so by Lemma 7 $\gamma_t(x^*) \leq s_t$ for every creator on its path. No tie convention is needed. □

Theorem 9 (Round-based enumeration)**.** *Let $D_N = \frac{1}{n}\sum_{u\neq v} N_{uv}^2 > 0$ and $W_t = \sum_{s:\rho(s)<\rho(t)} N_{ts}^2$, and suppose*

$$\|N\|^2 \leq K\, D_N, \tag{C1}$$

$$\sum_{t:\, 2\leq\rho(t)\leq\ell} \sqrt{W_t} \geq 4\sqrt{3}\, g\, n \sqrt{D_N}, \tag{C2}$$

$$\sum_{t\leq T} s_t \leq \tfrac{1}{8} g\, n \sqrt{D_N}, \tag{C3}$$

with $\|N\|$ the operator norm, and $n \geq 256\pi\tau(\ell-1)K/g^2$. Put $K' = (\ell-1)K$ and $\alpha = \max\{2, 9\tau\sqrt{K'}/(4g)\}$. Then every run of Algorithm C enumerates at most $(6 + 2\ell)\, 2^n e^{-c_* n}$ assignments, where

$$c_* = \min\Big\{ \frac{g^2}{32\tau\alpha^2 K'},\ \frac{g^2}{512\ell^2\tau K},\ c_{\mathrm{HW}} \frac{g^2}{256\ell^2\tau K},\ c_{\mathrm{HW}} \frac{g}{16\ell\sqrt{K}} \Big\}.$$

Algorithm C is exact and runs in polynomial space and time $O^*(|\mathcal{Y}|\,(2^{\tau n} + 2^n e^{-c_* n}))$.

Condition (C1) involves only the ordinary operator norm of the remainder, not the norm of its triangular part as in the count-guessing note [2]. The price is that preferred values adapt to earlier rounds only.

Lemma 10 (Race identity with rounds)**.** *Let x be uniform and $u = 2x - 1$. For a creator $t \in B_j$, $\gamma_t(x) = |\Delta_t| - \frac{1}{2} u_t \sum_{w\neq t:\rho(w)\geq j} N_{tw} u_w$. Hence x passes (2) if and only if $S' + Q_{\mathrm{same}} + Q_{\mathrm{later}} \leq 0$, where*

$$S' = \sum_{t\in I} (|\Delta_t| - s_t), \qquad Q_{\mathrm{same}} = -\frac{1}{2} \sum_{j\leq\ell} \sum_{t\in I\cap B_j} \sum_{w\in B_j\setminus\{t\}} u_t N_{tw} u_w,$$

$$Q_{\mathrm{later}} = -\frac{1}{2} \sum_{w:\rho(w)\geq 2} u_w \kappa_w, \qquad \kappa_w = \sum_{t\in I:\, \rho(t)<\rho(w)} u_t N_{tw}.$$

Proof. For $\rho(w) \geq j$ write $x_w = \frac{1}{2} + \frac{1}{2}u_w$; then the gain of x_t under $F'_{\hat{y}}$ is $\Delta_t + \frac{1}{2}\sum_{w\neq t:\rho(w)\geq j} N_{tw}u_w$. For a creator, $2p_t - 1 = -u_t$ and $(2p_t-1)\Delta_t = |\Delta_t|$. □

Lemma 11 (Gap profile with rounds)**.** *Let L be the $T\times T$ matrix with $L_{ts} = N_{ts}$ if $\rho(s) < \rho(t)$ and 0 otherwise, and $H_t = h_t^{\hat{y}} + \frac{1}{2}\sum_{w\neq t} N_{tw}$. Then $\Delta_t = H_t + \frac{1}{2}(Lu)_t$, $\|L\|^2 \leq (\ell-1)\|N\|^2 \leq K'D_N$, and, with $M = \alpha\sqrt{K'D_N}$,*

$$\mathbb{P}\Big[\sum_{t\leq T} \min\{|\Delta_t|, M\} < \tfrac{3}{4}g\,n\sqrt{D_N}\Big] \leq 4e^{-g^2n/(16\tau K')}.$$

Proof. The rows of L in round j form the block $N_{B_j\times B_{<j}}$ of norm at most $\|N\|$, so $\|Lu\|^2 = \sum_j \|N_{B_j\times B_{<j}}u\|^2 \leq (\ell-1)\|N\|^2\|u\|^2$. The rest is the proof of the gap-profile lemma in [2], with K' in place of K. Split $t\leq T$ into Small $= \{|H_t| \leq M/2\}$ and Large. For $t\in$ Small, $(a-b)_+ \leq a^2/(4b)$ gives $\min\{|\Delta_t|, M\} \geq |\Delta_t| - (Lu)_t^2/(8M)$, a total loss of at most $K'D_NT/(8M)$. For $t\in$ Large with $|(Lu)_t| \leq M/2$ the clipped gap is at least $M/4$, and at most $4T/\alpha^2$ indices have $|(Lu)_t| > M/2$. Each $(Lu)_t$ is a symmetric Rademacher sum with variance W_t, so by (T5) $\mathbb{E}|\Delta_t| \geq \frac{1}{2}\sqrt{W_t/3}$; also $M/4 \geq \frac{1}{2}\sqrt{W_t/3}$. By (C2), $\mathbb{E}G + \frac{M}{4}|\text{Large}| \geq 2gn\sqrt{D_N}$ for $G = \sum_{\text{Small}}|\Delta_t|$, while the two losses total at most $\frac{9}{8}T\sqrt{K'D_N}/\alpha \leq \frac{1}{2}gn\sqrt{D_N}$. The function G is convex with Lipschitz constant $\ell_G \leq \frac{1}{2}\sqrt{TK'D_N}$; (T3) with $r = \frac{1}{2}gn\sqrt{D_N}$, and $8\sqrt{\pi}\,\ell_G \leq \frac{1}{4}gn\sqrt{D_N}$ by the bound on n, gives the claim. □

Proof of Theorem 9. Exactness is Lemma 8. Fix a run and a uniform x.

Shortfall. Let $Y_t = [t\in I]\min\{|\Delta_t|, M\}$. Given the variables of the rounds before j, the values Δ_t, $t\in B_j$, are fixed and the u_t, $t\in B_j$, are independent and uniform, so the Y_t are independent with values in $[0,M]$ and means $\frac{1}{2}\min\{|\Delta_t|, M\}$. By (T1), $\sum_t Y_t \geq \frac{1}{2}\sum_t \min\{|\Delta_t|, M\} - a$ except with probability $e^{-2a^2/(TM^2)}$. Take $a = \frac{1}{8}gn\sqrt{D_N}$. With Lemma 11 and (C3), $S' \geq \sum_t Y_t - \sum_{t\leq T} s_t \geq \frac{1}{8}gn\sqrt{D_N}$ outside two events.

Later interactions. In round $j\geq 2$, the increment of Q_{later} is $-\frac{1}{2}\sum_{w\in B_j} u_w\kappa_w$, where the κ_w depend only on earlier rounds and the u_w are fresh. With $v_t = u_t[t\in I]$, $\sum_{w\in B_j}\kappa_w^2 = \|N^{\mathsf{T}}_{B_{<j}\times B_j}v\|^2 \leq \|N\|^2T$, so $\sum_w \kappa_w^2 \leq \ell K D_N T$ on every path. By (T2) with $V = \frac{1}{4}\ell K D_N T$, $\mathbb{P}[Q_{\text{later}} \leq -b] \leq e^{-2b^2/(\ell K D_N T)}$. Take $b = \frac{1}{32}gn\sqrt{D_N}$.

Same-round interactions. Fix j and condition on the earlier rounds. Put $c_t = 2p_t - 1$ and $\beta_t = -c_tu_t$ for $t\in B_j$; the β_t are independent uniform signs, $t\in I$ if and only if $\beta_t = 1$, and $u_t = -c_t\beta_t$. With the matrix $M^{(j)}_{tw} = c_tc_wN_{tw}$ on B_j, which has zero diagonal and $\|M^{(j)}\| \leq \|N\|$,

$$Q_{\text{same},j} = -\frac{1}{2}\sum_{t\neq w}\frac{1+\beta_t}{2}\beta_t\beta_wM^{(j)}_{tw} = -\frac{1}{4}\beta^{\mathsf{T}}M^{(j)}\beta - \frac{1}{4}\sum_w r_w\beta_w, \qquad r = M^{(j)}\mathbf{1}.$$

Here $\|r\|^2 \leq \|N\|^2|B_j|$ and $\|M^{(j)}\|_F^2 \leq |B_j|\|N\|^2$. Let $e = gn\sqrt{D_N}/(64\ell)$. By (T1) the linear part is below $-e$ with probability at most $e^{-8e^2/(|B_j|KD_N)}$, and by (T4) the quadratic part is below $-e$ with probability at most $\exp(-c_{\text{HW}}\min\{16e^2/(|B_j|KD_N), 4e/\sqrt{KD_N}\})$. Summing over the rounds, $Q_{\text{same}} \geq -\frac{1}{32}gn\sqrt{D_N}$ outside 2ℓ events.

Conclusion. Outside all exceptional events, $S' + Q_{\text{same}} + Q_{\text{later}} \geq (\frac{1}{8} - \frac{1}{32} - \frac{1}{32})gn\sqrt{D_N} > 0$. Using $|B_j| \leq T \leq \tau n$, the exponents of the exceptional probabilities are $g^2n/(16\tau K')$, $g^2n/(32\tau\alpha^2K')$, $g^2n/(512\ell\tau K)$, $g^2n/(512\ell^2\tau K)$, $c_{\text{HW}}g^2n/(256\ell^2\tau K)$ and $c_{\text{HW}}gn/(16\ell\sqrt{K})$, each at least c_*n (the first because $\alpha\geq 2$). With the factor 4 of Lemma 11, their total probability is at most $(6+2\ell)e^{-c_*n}$. □

5 Max-Cut with arbitrary unit clauses on spectral expanders

For a d-regular graph G let $\lambda(G) = \max_{i \geq 2} |\lambda_i(A_G)|$.

Corollary 12 (Max-Cut on spectral expanders)**.** *Fix $\lambda_0 \geq 1$ and $0 < \varepsilon \leq \frac{1}{2}$. There are n_0 and $c = c(\lambda_0, \varepsilon) < 2$ such that the following holds for every d-regular graph G on $n \geq n_0$ vertices with*

$$64\lambda_0^2 \leq d \leq (1-\varepsilon)n, \qquad \lambda(G) \leq \lambda_0\sqrt{d},$$

and every multiset of unit clauses. The formula consisting of $(u \vee v)$ and $(\neg u \vee \neg v)$ for every edge uv of G, together with the unit clauses, is solved by a deterministic algorithm in polynomial space and time $O^*(c^n)$.

Without unit clauses this is Max-Cut on G, since the edge contributes $1 + [x_u \neq x_v]$. With unit clauses it is Max-Cut with arbitrary external fields.

Proof. Apply Theorem 9 with $r = 1$, $v_1 = \mathbf{1}$, $\lambda_1 = \beta := -2d/(n-1)$, exact counts $\mathcal{Y}_1 = \{0, \ldots, n\}$ and $\delta_1 = 0$, so $s_t = |\beta|$; with $\tau = \frac{1}{2}$, $\ell = 2$, and B_1, B_2 the first and second $n/4$ variables of any order. Here $J = -2A_G$ and $N = -2A_G - \beta(\mathbf{1}\mathbf{1}^{\mathsf{T}} - I)$.

(C1): on $\mathbf{1}$, N acts as $-2d - \beta(n-1) = 0$; on $\mathbf{1}^{\perp}$ its eigenvalues are $-2\lambda_i + \beta$. So $\|N\| \leq 2\lambda_0\sqrt{d} + 2 \leq 3\lambda_0\sqrt{d}$. An edge has $N_{uv} = -2(1 - d/(n-1))$ and a non-edge $N_{uv} = -\beta$, so $D_N = 4d(1 - d/(n-1)) \geq 2\varepsilon d$ for $n \geq 2/\varepsilon$. Thus (C1) holds with $K = 9\lambda_0^2/(2\varepsilon)$.

(C2): since $1 - d/(n-1) \geq \varepsilon/2$, an edge from $t \in B_2$ to B_1 contributes at least ε^2 to W_t. With $\deg_{B_1}(t) \leq d$,

$$\sum_{t \in B_2} \sqrt{W_t} \geq \varepsilon \sum_{t \in B_2} \sqrt{\deg_{B_1}(t)} \geq \frac{\varepsilon\, e(B_1, B_2)}{\sqrt{d}} \geq \frac{\varepsilon}{\sqrt{d}}\Big(\frac{dn}{16} - \frac{\lambda(G)\, n}{4}\Big) \geq \frac{\varepsilon n\sqrt{d}}{32},$$

by the expander mixing lemma $|e(S, S') - d|S||S'|/n| \leq \lambda(G)\sqrt{|S||S'|}$ for disjoint S, S', and $\sqrt{d} \geq 8\lambda_0$. Since $\sqrt{D_N} \leq 2\sqrt{d}$, (C2) holds with $g = \varepsilon/(256\sqrt{3})$.

(C3): $\sum_{t \leq T} s_t = T|\beta| \leq 2d \leq \frac{1}{8} gn\sqrt{2\varepsilon d}$ once $\sqrt{n} \geq 16\sqrt{2}/(g\sqrt{\varepsilon})$, because $d \leq n$.

All parameters depend only on λ_0 and ε, and $|\mathcal{Y}| = n + 1$. □

Examples.

- *Paley graphs* P_q, $q \equiv 1 \pmod 4$ prime: $d = (q-1)/2$ and the nontrivial eigenvalues are $(-1 \pm \sqrt{q})/2$, so $\lambda_0 = 1$ and $\varepsilon = \frac{1}{2}$ work for $q \geq 13$.
- *Random d-regular graphs*: $\lambda(G) = O(\sqrt{d})$ with high probability, for fixed d by Friedman's theorem and for every $d \leq n/2$ by the result of Tikhomirov and Youssef [10].
- *Ramanujan graphs* that are not bipartite, such as the Lubotzky–Phillips–Sarnak graphs, with $\lambda(G) \leq 2\sqrt{d-1}$.

The numerical check in `explore_paley_maxcut.py` found the constant of the triangular condition of [2] to be about 1.3 on Paley graphs in a random order; Theorem 9 makes that condition unnecessary.

Why no simpler algorithm is known for these classes.

- Fixing all variables outside an independent set S costs $O^*(2^{n-|S|})$. By the Hoffman bound, $\alpha(G) \leq n\lambda(G)/(d + \lambda(G)) \leq \lambda_0 n/\sqrt{d}$, so the saving vanishes as d grows.
- Degree-based polynomial-space algorithms give $O^*(2^{(1-2/\Delta)n})$ for Max-Cut [3] and $O^*(2^{n(1-\alpha \ln d/d)})$ for Max 2-CSP [5]; both savings vanish as d grows.

- Expanders have treewidth linear in n, so dynamic programming over decompositions gives no saving, and also needs exponential space.
- The automorphism groups of Paley and Cayley graphs have polynomial size and give only polynomial savings.
- Williams' algorithm [11] beats 2^n on every instance, but uses exponential space. Whether general Max-Cut admits a polynomial-space algorithm faster than 2^n is open [6].

We found no polynomial-space algorithm with a fixed saving for Max-Cut on these families when d grows; this search was not exhaustive. For $d < 64\lambda_0^2$ the interaction graph has bounded degree, and polynomial-space algorithms for Max 2-CSP of bounded degree [5] give a fixed saving there, so the saving can be made uniform over all $d \le (1-\varepsilon)n$.

6 Count guessing without a given decomposition

Corollary 13 (Spectral count guessing)**.** *Let $v_1, \dots, v_r$ be orthonormal and $\lambda_1, \dots, \lambda_r$ real, for example approximate eigenpairs of J of largest absolute value, and use the grids $\mathcal{Y}_k = \delta\mathbb{Z} \cap [-\|v_k\|_1 - \delta, \|v_k\|_1 + \delta]$. Then $|\mathcal{Y}| \le (2\sqrt{n}/\delta + 3)^r$ and $\sum_{t \le T} s_t \le \sum_k |\lambda_k|(\delta\sqrt{n} + 1)$. Hence, if* (C1) *and* (C2) *hold for N and $\sum_k |\lambda_k|(\delta\sqrt{n}+1) \le \frac{1}{8} gn\sqrt{D_N}$, Max 2-SAT is solved in polynomial space and time $O^*((2\sqrt{n}/\delta + 3)^r (2^{\tau n} + 2^n e^{-c_* n}))$.*

Proof. $\sum_t s_t = \sum_k |\lambda_k|(\delta\|v_k\|_1 + \|v_k\|_2^2)$ and $\|v_k\|_1 \le \sqrt{n}$. Exactness holds for any vectors (Lemma 8). □

The algorithm needs no planted structure as input: it computes approximate eigenpairs of J and checks the conditions. The next proposition shows that random formulas whose parity means have low rank satisfy them.

Proposition 14 (Planted low-rank parity structure)**.** *Let the coefficients J_{uv}, $u < v$, be independent with $|J_{uv} - \mathbb{E}J_{uv}| \le B$ and $\nu \le \operatorname{Var} J_{uv} \le \chi\nu$, and suppose $\mathbb{E}J_{uv} = \bar{A}_{uv}$ for $u \ne v$, where $\bar{A}$ is a symmetric matrix of rank at most r with $|\bar{A}_{uu}| \le B_0$ and $\|\bar{A}\| \le \mu n$. Let P collect the eigenpairs of J with $|\lambda| > \theta := 4\sqrt{\chi\nu n}$, and split the prefix into two rounds of size $T/2$. There is $C = C(\chi, r, \tau, B_0)$ such that, if $n\nu \ge C(B^2 \log n + \mu^2)$, then with probability $1 - o(1)$ the conditions of Corollary 13 hold with $K = O(\chi)$, $g = \Omega(\tau^{3/2}/\sqrt{\chi})$ and the grid spacing $\delta = gn\sqrt{D_N}/(16\sqrt{n}\,\Lambda)$, $\Lambda = \sum_k |\lambda_k|$, which the algorithm computes from the data. The running time is $O^*(n^{O(r)}\, 2^{(1-\gamma)n})$ with $\gamma > 0$ depending only on χ and τ.*

Proof. Let X be the off-diagonal matrix of the $J_{uv} - \mathbb{E}J_{uv}$. By the norm bound of Bandeira and van Handel [1], applied as in Proposition 12 of [2], $\|X\| \le 3\sqrt{\chi\nu n}$ with probability $1 - o(1)$. Then $J = \bar{A} - \operatorname{diag}(\bar{A}) + X$ has at most r eigenvalues of absolute value above θ (Weyl), so P has rank $r' \le r$, and $J - P$ has all eigenvalues in $[-\theta, \theta]$. Hence $\bar{A} - P = (J - P) - X + \operatorname{diag}(\bar{A})$ has rank at most $2r$ and norm at most $\theta + \|X\| + B_0$, and

$$N = X + \operatorname{offdiag}(\bar{A} - P), \qquad \|N\| \le \|X\| + 2(\theta + \|X\| + B_0) = O(\sqrt{\chi\nu n}).$$

The low-rank part has Frobenius norm at most $\sqrt{2r}(\theta + \|X\| + B_0) = O(\sqrt{r\chi\nu n})$, against $\|X\|_F \ge \sqrt{(1-\eta)n(n-1)\nu}$ by Bernstein's inequality, so $D_N = (1 - o(1))\frac{1}{n}\|X\|_F^2 \ge (1 - 2\eta)(n - 1)\nu$ and (C1) holds with $K = O(\chi)$. For (C2), Bernstein's inequality gives $\sum_{s \in B_1} X_{ts}^2 \ge (1 - \eta)|B_1|\nu$ for all $t \in B_2$, and the low-rank part changes $\sum_{t \in B_2} \sqrt{W_t}$ by at most $\sqrt{T} \cdot O(\sqrt{r\chi\nu n}) = O(n\sqrt{r\chi\nu})$, which is negligible against $|B_2|\sqrt{(1-\eta)|B_1|\nu} = \Theta(\tau^{3/2} n^{3/2}\sqrt{\nu})$. For (C3), the choice of δ gives $\Lambda\delta\sqrt{n} = \frac{1}{16} gn\sqrt{D_N}$, so it remains to check $\Lambda \le \frac{1}{16} gn\sqrt{D_N}$. Each of the at most r selected eigenvalues has $|\lambda_k| \le \|J\| \le \mu n + B_0 + \|X\|$, and $\sqrt{D_N} \ge (1 - o(1))\sqrt{n\nu}$; so this holds once $n\nu \ge (32r/g)^2\mu^2$ and n is large. The number of runs is $(2\sqrt{n}/\delta + 3)^r = (32\Lambda/(g\sqrt{D_N}) + 3)^r = n^{O(r)}$, since $\Lambda/\sqrt{D_N} = O(r\mu\sqrt{n/\nu} + r\sqrt{\chi})$ and $\nu \ge B^2 \log n/n$. □

This removes the assumption "given the decomposition" from the count-guessing theorem for planted bisections, planted group structure, and any other parity pattern whose mean has bounded rank. Approximate eigenpairs suffice because Lemma 8 holds for any vectors and the conditions are checked for the vectors actually used.

7 What remains

- *Complementary pairs* ($\pm J$ spin glasses on a fixed graph) need the random-order version of Theorem 4.
- *Irregular expanders.* Theorem 9 measures $\|N\|$ against the average D_N; for very irregular degree sequences a weighted version, as in Theorem 4, is the natural next step.
- *Biased random formulas at density* $\omega(1) \le d < C \log n$, by branching first on the few vertices of large degree.
- *Aligned structure of high rank*, such as noisy blow-ups, where the counts of many small groups would have to be guessed.
- The literature search behind Section 5 should be extended before any claim of novelty.

The script `explore_weighted_rounds.py` checks exactness of Algorithms A and C, including the spectral grid, against exhaustive search: no optimum was lost on 200 formulas for Algorithm A (which enumerated 8.6% of all assignments on average), nor on 120 formulas each for exact counts and spectral grids in Algorithm C (24% and 21% enumerated per run). These small instances test correctness only, not the asymptotic bounds.

References

[1] A. S. Bandeira and R. van Handel. Sharp nonasymptotic bounds on the norm of random matrices with independent entries. *Ann. Probab.* 44(4):2479–2506, 2016.

[2] *Count guessing makes constraints useful*, working note, 11 October 2026 (`max2sat_count_guessing.tex`).

[3] F. Della Croce, M. J. Kaminski and V. Th. Paschos. An exact algorithm for MAX-CUT in sparse graphs. *Oper. Res. Lett.* 35(3):403–408, 2007.

[4] *Max 2-SAT by early constraint-guided enumeration*, working note, 8 October 2026 (`max2sat_early_enumeration.tex`).

[5] A. Golovnev and K. Kutzkov. New exact algorithms for the 2-constraint satisfaction problem. *Theoret. Comput. Sci.* 526:18–27, 2014.

[6] K. Lalovic. Exact algorithms for MaxCut on split graphs. arXiv:2405.20599, 2024.

[7] *Dense Max 2-SAT in polynomial space*, research draft, 12 October 2026 (`dense_max2sat_paper.tex`).

[8] M. Rudelson and R. Vershynin. Hanson–Wright inequality and sub-gaussian concentration. *Electron. Commun. Probab.* 18(82):1–9, 2013.

[9] M. Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. *Publ. Math. IHÉS* 81:73–205, 1995.

[10] K. Tikhomirov and P. Youssef. The spectral gap of dense random regular graphs. *Ann. Probab.* 47(1):362–419, 2019.

[11] R. Williams. A new algorithm for optimal 2-constraint satisfaction and its implications. *Theoret. Comput. Sci.* 348(2–3):357–365, 2005.