Exploring the Branching Frontier for Max 2-SAT

Full clause phases, joint completion bounds, and coverage

7 October 2026

Abstract

We develop two concrete variations on the branching program of the October 5–6 notes. First, resolving a pruning clause completely gives an explicit measure based on its exact completion count. Its analysis walk is uniform on the permitted assignments, giving a simple concentration lemma. This changes the algorithm and does not prove the earlier missing lemma for clauses that are released. Second, counting the assignments satisfying a weighted sum of active pruning clauses gives an overlap-aware, polynomial-time certificate when the integer weights are polynomially bounded. Inherited certificates give a branching inequality and an explicit variable-selection score. We prove a conditional theorem identifying the cumulative progress still needed for a fixed base below two. We also formalize when neighborhood coverage helps, why it need not yield small separators, and how exact interaction classes offer a different structural exit. No density-uniform running-time theorem is claimed.

1 Target and status

The target is a deterministic polynomial-space algorithm with time $O^*((2-\delta)^n)$ for a constant $\delta > 0$ independent of a growing density bound $\rho(n)$. Here $\rho = m_2/n$ counts binary clauses per variable; the average occurrence degree is $d = 2\rho$. An exponent saving that tends to zero with d does not meet this target, even if it is a useful improvement. The first milestone could be all instances with $\rho(n) = \log\log n$; $\rho = O(n)$ covers the full simple unweighted setting.

The notes already identify three difficulties: obtaining useful gaps, overlap between pruning supports, and detecting contradictions before the search reaches almost all variables. Our contributions below are mathematical derivations within that program, with small exact checks. They have not been independently refereed, and no novelty claim relative to all literature is made.

The polynomial-space baselines we checked have degree-dependent savings [3, 4]; these support the distinction above. Hastings also gives instances with very many strict local optima. Consequently, a bound based only on enumerating all local optima cannot provide a uniform constant saving. Our web check did not locate a newer theorem settling the target; it is not an exhaustive literature survey.

We use the deficit notation of the earlier notes. In a residual formula, $p = \min(p_x, p_{\neg x})$, and D is the free binary degree. For $B = 1$, branching against the literal with smaller deficit creates the valid constraint $\langle L_x \leq p-1 \rangle$ on D distinct variables. The preferred child receives no new constraint. The soundness arguments in the earlier notes apply to the modified strategies considered here.

2 Resolve a pruning clause as one phase

2.1 An explicit branching rule and measure

Define the exact cardinality volume

$$V(r,k) = \sum_{j=0}^{\min\{r,k\}} \binom{r}{j}, \qquad V(r,k) = 0 \quad \text{if } k < 0.$$

Consider this variant of the rule. At a phase boundary choose an undominated x. Explore its preferred child recursively. In the other child enumerate all assignments of its D neighbors satisfying the new clause, then recurse on the remaining variables. During this enumeration postpone other domination fixings and additional pruning clauses. Earlier constraints may be retained, but for the clean phase-distribution argument do not condition the enumeration on them. They can be checked at the phase boundary. This may do more work, but it preserves exactness. Combinations enumerate the allowed assignments in polynomial space with polynomial overhead per output; there is no need to scan all 2^D assignments.

Proposition 1 (Full-phase branching)**.** *For $k = p - 1$, the phase has one child with a decrease of one free variable, and $V(D,k)$ children with a decrease of $D+1$. It is admissible at base $1 < \lambda < 2$ whenever*

$$\lambda^{-1} + V(D,k)\lambda^{-(D+1)} \le 1, \quad \textit{equivalently} \quad V(D,k) \le (\lambda - 1)\lambda^D. \tag{1}$$

If every phase satisfies this condition, the search runs in $O^*(\lambda^n)$ time and polynomial space.

Proof. These are exactly the enumerated alternatives. Induct on the number of free variables using (1). The work enumerating phase alternatives is polynomial per child, and there are at most n levels of recursive phases on a path. Retained constraints can only remove alternatives. □

For an equivalent binary-step analysis put

$$e_\lambda(r,k) = r - \log_\lambda V(r,k), \qquad \mu = u + \sum_C \log_\lambda V(r_C, k_C) = n_{\text{free}} - \sum_C e_\lambda(r_C, k_C), \tag{2}$$

where u counts uncovered free variables and the counted supports are disjoint. A single active clause is sufficient for the phase algorithm. Weights e_λ are permitted to be negative. This is intentional: a weak residual constraint carries the cost of completing its support. The measure is still nonnegative because each live clause has $V \ge 1$.

Lemma 2 (Binary steps and forced assignments)**.** *Resolving a clause variable satisfies the branching inequality with equality for the measure* (2)*. A forced assignment never increases μ. These statements do not require $e_\lambda \ge 0$.*

Proof. Use $V(r,k) = V(r-1,k) + V(r-1,k-1)$. If the literal is set to b, the measure drop is $\log_\lambda(V(r,k)/V(r-1,k-b))$. Summing $\lambda^{-\text{drop}}$ over the nonempty children gives one. Restriction cannot increase V. For an uncovered variable the drop is one. At $k = 0$ there is one completion, so propagating the remaining literals also respects the measure. □

The same inequality shows that a transition cannot decrease e_λ by more than one. At a vacuous state $e_\lambda = r(1 - \log_\lambda 2) < 0$. Keeping the clause until its support is completed is permitted. Releasing it would reduce the measure, but would change the distribution used next.

2.2 A concentration lemma without an unproved walk assumption

Proposition 3 (Distribution inside an isolated full phase)**.** *Choose each child during clause resolution with probability equal to its remaining completion count divided by that of its parent. The resulting assignment is uniform among the $V(D,k)$ permitted assignments. If J is its number of true pruning literals and $0 \le k < D/2$, then for $1 \le t \le k$,*

$$\Pr[J \le k - t] \le \left(\frac{k}{D-k+1}\right)^t. \tag{3}$$

Proof. The child probabilities telescope to $1/V(D,k)$ at every completion. For $q = k/(D-k+1)$ and $j \le k$, $\binom{D}{j-1}/\binom{D}{j} = j/(D-j+1) \le q$. Thus $\binom{D}{i} \le q^t\binom{D}{i+t}$ for $i \le k-t$. Sum and divide by $V(D,k)$. □

For fixed $k/D \le \beta < 1/2$, deviations of ηD below the allowance have probability at most $\exp(-\Omega_\beta(\eta D))$. This proves a phase-level statement at least as strong as the sought concentration scale, *for this modified full-resolution rule.* It does not establish that statement for the October 6 walk, whose zero-weight states are released. It also does not establish the availability of subsequent good phases.

The mean satisfies

$$\mathbb{E}[J] \ge k - \frac{q}{1-q}.$$

By symmetry each support literal is true with probability $\mathbb{E}[J]/D = k/D - O_\beta(1/D)$ when $k/D \le \beta < 1/2$. For another variable, its helpful values need not agree with all of these literals, and it may share only part of the support. The following bound handles both issues for a single complete phase.

Lemma 4 (Helpful updates on any fixed overlap)**.** *Fix $s \ge 1$ variables of the clause support and designate one helpful value for each, with arbitrary signs. Let Y be the number taking that value in a uniform permitted completion. If $0 < \eta < k/D < 1/2$ and $q = k/(D-k+1)$, then*

$$\Pr\left[Y < \left(\frac{k}{D} - \eta\right)s\right] \le q^{\lceil \eta D/2 \rceil} + 2e^{-\eta^2 s/8}. \tag{4}$$

Proof. By (3), except with probability at most the first term, $J/D \ge k/D - \eta/2$. Condition on such a value $J = j$ and write $p = j/D$. The true positions form a uniform j-subset. Split the designated positions into a where true is helpful and $b = s - a$ where false is helpful. Their helpful counts have means pa and $(1-p)b$, whose sum is at least ps, since $p < 1/2$. Each count is a sample without replacement from a zero-one population. Hoeffding's bound [5] applies to each. For their sum to be below $(k/D - \eta)s$, one of the two counts must be at least $\eta s/4$ below its mean. The two lower-tail bounds are at most $e^{-\eta^2 s^2/(8a)}$ and $e^{-\eta^2 s^2/(8b)}$, omitting an empty group. Each is at most $e^{-\eta^2 s/8}$. A union bound proves the claim; no independence between the two counts is assumed. □

The designation is fixed at the beginning of the phase, for example toward a chosen deficit of a variable that remains outside its support. This is conditional on the phase-start history and imposes no alignment assumption. It is not a theorem about many adaptive phases: their ratios k/D may be too low, intersections may be small, and the deficit orientation may change.

2.3 Entropy scale and a remaining availability problem

For $k \le \beta D$ with $0 < \beta < 1/2$, $V(D,k) \le 2^{H(\beta)D}$. Therefore, for any $2^{H(\beta)} < \lambda < 2$, every such phase with

$$D \ge \frac{\ln(1/(\lambda-1))}{\ln\lambda - H(\beta)\ln 2} \tag{5}$$

is feasible. Small degrees can be checked directly or handled separately. Do not infer uniform finite-size feasibility at the exact entropy base from an asymptotic estimate alone.

For $k = D/4$, the numerical roots of (1) are:

D	16	32	64	128
full-phase base	1.67236	1.69519	1.71536	1.73019

They approach $2^{H(1/4)} \simeq 1.75477$. These are local branching numbers, not global bounds without an availability argument. The earlier optimized release rule can have smaller finite-size branching numbers.

Allow plain steps and use the previous averaging construction. At a feasible phase give the preferred child probability $1/\lambda$ and spread the remaining probability uniformly across its $V(D,k)$ alternatives. Each alternative then has probability at least $\lambda^{-(D+1)}$ by (1). The proof of the earlier averaging lemma gives

$$L \le \lambda^n \mathbb{E}[(2/\lambda)^S]$$

for the number S of plain steps on this phase walk. The exponentially weighted cost of plain steps remains the actual obstacle. A small empirical mean for S is insufficient.

A useful next experiment is to choose the largest feasible ratio below a fixed threshold, rather than the smallest ratio. A very strong clause can force values unfavorable to its neighbors; a ratio near the threshold passes on a less biased distribution. This is a motivated strategy, not a proved universal improvement.

3 A joint measure from exact weighted-sum counts

3.1 The certificate

At a search state let the active pruning inequalities be $L_i(x) \le k_i$, $i = 1, \dots, m$, over r free variables. For nonnegative integer weights a_i define

$$N_a = \#\{x \in \{0,1\}^r : \ \textstyle\sum_i a_i L_i(x) \le \sum_i a_i k_i\}. \tag{6}$$

Every completion satisfying all active clauses is counted by N_a. Unlike an additive clause weight, N_a explicitly incorporates overlapping supports, cancellation and redundant clauses.

For free variable w, let A_w^b be its contribution to $\sum_i a_i L_i$ when it takes value b. Set

$$K = \sum_i a_i k_i - \sum_w \min(A_w^0, A_w^1), \qquad c_w = |A_w^1 - A_w^0|.$$

After orienting each variable so that its cheaper value is zero,

$$N_a = \sum_{j \le K} [z^j] \prod_w (1 + z^{c_w}). \tag{7}$$

This is a one-dimensional integer dynamic program. It takes $O(r(1 + \sum_w c_w))$ arithmetic operations, with counts having at most $r + 1$ bits. If clause multiplicities and the weights a_i are bounded by polynomials in the input size, both time and space are polynomial. Merely bounding the *bit length* of arbitrary huge weights is not enough.

The unweighted sum cut is exactly the test $N_{\mathbf{1}} = 0$. The certificate gives useful information before that cut becomes possible. For a single cardinality clause it gives its exact binomial volume, including the roughly one-half factor for a nearly balanced clause that a Chernoff bound can miss.

3.2 Inheritance makes the branching inequality rigorous

Theorem 5 (Joint completion certificate)**.** *Store a weight vector a and $U = N_a$ at every live node. Cut if $U = 0$. For each child, evaluate the inherited vector after substitution and any polynomial-size list of additional candidate vectors, and store the smallest count. Extend the inherited vector by zero on newly created pruning clauses. Then, after branching and any further sound fixings,*

$$U_0 + U_1 \le U. \tag{8}$$

Also $U \ge 1$ at every kept node and $U = 2^n$ at an unconstrained root.

Proof. For a fixed vector, assignments counted at the parent partition according to the value of the branching variable: $N_a = N_{a,0} + N_{a,1}$. Further fixings restrict these sets. A new constraint has inherited weight zero and therefore does not invalidate the old certificate. Minimization over a list containing the inherited vector cannot increase either child count. These facts prove (8). The counts are nonnegative integers and upper-bound all jointly feasible completions, so a zero count justifies a cut. □

The value U here is part of the augmented search state, together with its witness. We do not claim that globally minimizing (6) over all bounded integer vectors is polynomial-time. Polynomial candidate lists and bounded rounds of coordinate improvement suffice for the theorem. If every state were optimized independently by a heuristic without retaining the inherited witness, (8) would not be guaranteed.

Useful candidates are the zero vector, each single clause, all clauses with weight one, promising clause pairs, and bounded coordinate modifications. A linear program or a smooth Chernoff optimization can propose weights; the final integer count still certifies the result exactly.

3.3 A concrete strategy and the missing global theorem

The measure $\log_2 U$ has a valid base-two branching inequality. It does not yet give a base below two. For candidate variable x, form its two children, including its new pruning clause, and calculate

$$I_x = \log_2 U - \log_2(U_{x=0} + U_{x=1}) \ge 0. \tag{9}$$

Choose the variable maximizing this certified decrease. If the denominator is zero, both children can be discarded. Polynomially many candidates and certificate evaluations preserve polynomial time per visited node and polynomial space. This is an explicit branching strategy optimizing a proved certificate; its worst-case advantage is still unproved.

It is essential to count work spent forming all candidate children. A smaller selected search tree need not give a faster implementation at small sizes. Fixed polynomial lookahead overhead is harmless for an eventual exponential-base theorem, but matters in experiments.

Theorem 6 (A sufficient cumulative-progress condition)**.** *Use the certificate search above, with polynomial work per visited node. Normalize child probabilities to $U_i/(U_0 + U_1)$ whenever the sum is nonzero. Absorb successful terminal nodes with $U = 1$, and regard a node with no kept children as death. Let G_t be the sum of* (9) *along this walk through branching depth t, with $G_t = +\infty$ after death.*

Suppose, for constants $\alpha, \gamma, \xi > 0$ and $t_0 = \lfloor (1-\alpha)n \rfloor$,

$$\Pr[G_{t_0} < \gamma n] \le 2^{-\xi n}.$$

Then the search runs in

$$O^*\left(2^{(1-\min\{\alpha,\gamma,\xi\})n}\right).$$

Proof. Write $R_v = (U_{v0} + U_{v1})/U_v = 2^{-I_v}$. For a frontier node w, the probability of reaching it under normalized transitions times $\prod_{v \prec w} R_v$ is $U_w/2^n$. Hence the total certificate mass at depth t is $2^n \mathbb{E}[2^{-G_t}]$. At t_0 this is at most $2^n(2^{-\gamma n} + 2^{-\xi n})$. Subsequent total mass cannot increase, and each kept node has integer mass at least one. There are at most n branching levels. The earlier levels have at most 2^{t_0+1} nodes. Discarded children and polynomial lookahead add only polynomial overhead to these counts. □

This is a precise replacement for "encounter many variables with gaps". The required quantity is linear cumulative *joint* progress, early enough in the tree, with exponentially controlled bad paths under the stated walk. Repeatedly creating almost redundant clauses need not contribute to it. So far no condition depending only on $\rho(n) = \omega(1)$ proves this premise.

3.4 A real limitation of weighted sums

Suppose the relaxation $0 \le x_w \le 1$, $L_i(x) \le k_i$ is feasible. For any nonnegative weights a, its feasible fractional point gives

$$\min_{x \in \{0,1\}^r} \sum_i a_i L_i(x) = \min_{x \in [0,1]^r} \sum_i a_i L_i(x) \le \sum_i a_i k_i.$$

The equality holds because the objective is linear on a box. Therefore $N_a \ge 1$ for every weighted sum, even if the Boolean conjunction is infeasible. For example, $x + y = y + z = z + x = 1$ has the fractional solution $(1/2, 1/2, 1/2)$ but no Boolean solution. Encode each equality by its two cardinality inequalities. No nonnegative weighted-sum cut detects the contradiction at the root.

Conversely, separating the image of the box from the lower orthant shows that an infeasible linear relaxation admits a nonnegative combination whose minimum exceeds its bound. Such multipliers need not lie in our small candidate list or have small numerical values. Thus weighted sums need either branching or stronger integer reasoning in the remaining cases.

4 What coverage actually certifies

Let C be the union of selected active supports and let $S \subseteq V \setminus C$ be a set of candidate variables.

Proposition 7 (Exact and approximate coverage)**.** *If $N(v) \subseteq C$ for every $v \in S$, then S is independent and the residual Max 2-SAT instance can be solved in time $O^*(2^{n-|S|})$ and polynomial space. More generally, if every $v \in S$ has at most b neighbors outside C, then a polynomial-time construction gives an independent set of size at least $|S|/(b+1)$, and time $O^*(2^{n-|S|/(b+1)})$ follows.*

Proof. In the first case no edge has both endpoints in S. Assign its complement and optimize every variable of S independently. In the second case $G[S]$ has maximum degree at most b; a greedy independent-set construction removing a chosen vertex and its neighbors proves the claim. □

Here n is the number of variables in the residual subproblem. The statement does not require a small separator. For instance, $K_{t,t}$ has exact coverage of one side by the other, a linear independent set, and no sublinear balanced separator. Conversely, if C contains nearly every variable, coverage by C says little. Also, $b = \varepsilon d$ is not enough for a constant saving when d grows: the independent-set guarantee can still be sublinear.

A useful coverage dichotomy therefore must quantify both the number of *uncovered centers* and the absolute amount of their remaining internal interaction. Merely counting covered neighbors, or measuring a covered fraction, misses these quantities.

If a candidate cut C really does leave components K_j with $\max_j |K_j| = q$, independent optimization after enumerating C gives time $O^*(2^{|C|+q})$. Inherited pruning clauses can couple these components. One safe terminal action is to discard those auxiliary constraints and solve the residual *soft* Max 2-SAT objective exactly. Every returned assignment is still an assignment of the original formula, and the enlarged subtree contains every originally retained completion. This is valid for this optimization search, though it forfeits the discarded pruning information.

5 An alternative structural exit: interaction classes

Small separators are not the only useful outcome of aligned neighborhoods. Write a residual objective in Boolean quadratic form

$$f(x) = c + \sum_i h_i x_i + \sum_{i<j} J_{ij} x_i x_j.$$

Suppose the remaining variables can be partitioned into t classes such that J_{ij} depends only on the classes of i, j, including a uniform coefficient for distinct variables in the same class. The h_i can differ.

Proposition 8 (Exact interaction classes)**.** *Such an instance is solvable in polynomial space and time $O^*(\prod_{a=1}^{t}(|C_a|+1)) \le O^*((n+1)^t)$. If the property holds after deleting an exception set E, time $O^*(2^{|E|}(n+1)^t)$ suffices.*

Proof. Enumerate the number k_a of ones in each class. All quadratic contributions are determined: they are $J_{aa}\binom{k_a}{2}$ within a class and $J_{ab}k_ak_b$ across classes. The best linear contribution selects the k_a largest h_i in each class. Enumerating E changes only fields on the remaining vertices and a constant. □

Thus $|E| \le (1-\delta)n$ and $t\log_2(n+1) = o(n)$ give a fixed-base improvement. This extends the complete aligned example in the earlier notes and can apply with very large separators. It requires *exact signed interaction classes.* Shared supports, large Gram correlations, approximate low rank, and a small number of active rows do not alone prove this property. Turning approximate alignment into a small exception set or a manageable residual interaction graph is a separate open direction.

6 Checks and what the experiments do not establish

The reproducible script is `explore_branching_frontier.py`. Its output is `branching_frontier_checks.json`. No existing notes or experiments were modified. The script uses exact integer counts, except for reported branching roots and numerical checks of logarithmic identities.

- The full-phase search matches brute-force optimum on 80 random $B = 1$ formulas with $3 \le n \le 10$, exercising 103 phase nodes.
- There are 41,650 checks of (3) and 3,599 checks of forced measure transitions, including negative-weight states.
- On 150 random cardinality systems with $3 \le n \le 9$, every tested certificate upper-bounds the brute-force joint completion count. All 918 variable-splitting identities and inequalities pass, as do 150 tests adding an extra constraint in one child.

- The small candidate optimizer improves over the best single-clause bound in 55 of these 150 systems, and over the all-ones weighted sum in 103. These are certificate comparisons on synthetic systems, not Max 2-SAT running-time estimates.
- A contradictory pair on seven variables plus a weak clause on 20 new variables has all-ones count 134,215,040. Weights $(1, 1, 0)$ give zero. The odd-cycle equality example remains positive for every tested vector with weights in $\{0, 1, 2\}$; its actual joint count is zero.

We also implemented one-step candidate branching using (9) and compared it with static branching on six random $B = 1$ formulas, edge probability 0.7. Both versions use the same completion certificates and domination. For speed, their weight candidates are just zero, singletons, all ones, and the inherited vector. All optimum values match brute force.

	kept nodes visited		candidate children evaluated	
n	static	volume strategy	static	volume strategy
12	14	6	22	56
12	16	7	28	86
12	4	2	4	22
16	91	32	162	694
16	40	19	72	412
16	107	51	190	866

The selected trees are smaller, but the lookahead performs more work. These tiny instances establish implementability and provide correctness checks; they do not show an asymptotic improvement, nor even a wall-clock speedup. They are also too small to test a growing-density claim.

7 The next mathematical questions

Availability under complete phases. Use the single-phase overlap bound (4) to study drift across adaptively chosen phases. Investigate largest-feasible-ratio branching and bound the exponential moment of plain steps. The unresolved issue is to control the chosen clause ratios and accumulation of local errors; full phases do not provide this global drift theorem for free.

Joint progress before the last levels. Prove the cumulative-progress premise in the theorem for an interesting class, initially pseudorandom signed instances. The critical quantitative question is whether useful inequalities add linear total information before branching depth $(1 - \alpha)n$. Uniform assignments and the certificate walk are different distributions; substituting one for the other is not valid.

From coverage failure to a certified structural exit. Track the size of the uncovered candidate set, its internal degree, the components after a prospective cut, and signed interaction classes. A valid dichotomy must either create genuinely new joint progress or provide a quantified residual solver. High overlap alone cannot certify either side.

Beyond one aggregate inequality. The LP-feasible but integer-infeasible example motivates small groups of aggregate inequalities, parity reasoning, or class decisions. A constant number of weighted inequalities can be counted by a multidimensional dynamic program when all coefficient sums are polynomial. Letting this dimension grow raises a polynomial-space issue; the resulting table must not be silently charged as polynomial storage.

The density milestone. For a theorem covering all formulas with a growing density bound, explicitly include $B = 2$ instances such as Max-Cut encodings. The weighted-sum certificate extends to integer multiplicities directly; the exchangeable binomial phase lemma is specifically $B = 1$. A weighted phase has a computable generating function, but no automatic exchangeability. The missing result is a structural/probabilistic argument covering every instance at the selected growing density. None of the checks above supplies it.

References

[1] Max 2-SAT and the domination rule: results and ideas so far. Working note, 5 October 2026.

[2] Max 2-SAT with pruning clauses: pair decisions, sums, and a measure. Working note, 6 October 2026.

[3] A. Golovnev. New upper bounds for MAX-2-SAT and MAX-2-CSP w.r.t. the average variable degree. IPEC 2011. `https://golovnev.org/papers/max2sat.pdf`.

[4] M. B. Hastings. Local maxima and improved exact algorithm for MAX-2-SAT. 2016. `https://arxiv.org/abs/1610.07100`.

[5] W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association* 58(301):13–30, 1963. `https://doi.org/10.1080/01621459.1963.10500830`.